Esketamine-induced post-traumatic stress disorder flashbacks during treatment-resistant depression indication: is it just a side effect?

Maud Rothärmel[1], Lila Mekaoui[2], François Kazour[3], Morgane Herrero[4], Eva-Maria Beetz-Lobono[5], Aiste Lengvenyte[6], Jérôme Holtzmann[7], Philippe Raynaud[8], Macarena Cuenca[9], Samuel Bulteau[10], Pierre de Maricourt[9], Thomas Husson[1], Emilie Olié[6], Bénédicte Gohier[3], Anne Sauvaget[10], Raphaël Gaillard[9], Raphaëlle Richieri[11], David Szekely[5], Ludovic Samalin[12], Olivier Guillin[1,13], Virginie Moulier[1,14], Wissam El-Hage[15, 16], Andrew Laurin[10], Lucie Berkovitch[9,17,18,19]

1. University Department of Psychiatry, Therapeutic Centre of Excellence, Institute of Psychiatry – Rouvray Hospital Centre, Sotteville-lès-Rouen, France;

2. Mental and Brain Illness Clinic, Sainte-Anne Hospital, GHU Paris – Psychiatry and Neurosciences, Paris, France;

3. Department of Psychiatry and Addictology, CHU Angers, Angers, France;

4. Department of Psychiatry, CHU Saint Etienne, Saint Etienne, France;

5. Centre Hospitalier Princesse Grace, Service de psychiatrie, Principauté de Monaco

6. Department of Emergency Psychiatry and Post-Acute Care, CHU Montpellier, France; IGF, Univ. Montpellier, CNRS, INSERM, Montpellier, France

7. Service de Psychiatrie de l'Adulte, Centre Expert Dépression Résistante FondaMental, CHU de Grenoble, hôpital Nord, Grenoble, France;

8. Department of Psychiatry, Thuir Hospital Centre, Thuir, France

9. University Department of Psychiatry, Pôle Hospitalo-Universitaire Psychiatrie Paris 15, Groupe Hospitalier Universitaire Paris, Paris, France;

10. Nantes Université, CHU Nantes, INSERM, MethodS in Patients-centered outcomes and HEalth Research, SPHERE, F-44000 Nantes, France;

11. Service Universitaire de santé mentale et physique, Centre Expert Dépression Résistante, CHU Sainte-Marguerite, APHM, Université Aix-Marseille, Marseille,

France; Aix Marseille Univ, CNRS, Centrale Marseille, Institut Fresnel, Marseille, France.

12. Department of Psychiatry, CHU Clermont-Ferrand, University of Clermont Auvergne, CNRS, Clermont Auvergne INP, Institut Pascal (UMR 6602), Clermont-Ferrand, France;

13. CHU Rouen, Normandy University, Rouen, France;

14. Unité de Recherche Clinique (URC), EPS Ville Evrard, Neuilly-sur-Marne, France;

15. Centre Régional de Psychotraumatologie, Centre Hospitalier Régional

16. Universitaire (CHRU) de Tours, Tours, France, 3 INSERM U1253 Imagerie et Cerveau (iBrain), Tours, France

17. Paris Cité University, Paris, France.

18. Department of Psychiatry, Yale University School of Medicine, 40 Temple Street, New Haven, CT, 06511, United States

19. Saclay CEA Centre, Neurospin, Gif-Sur-Yvette Cedex, France

Correspondence: Maud Rothärmel
maud.rotharmel@ch-lerouvray.fr

University Department of Psychiatry, Therapeutic Centre of Excellence, Institute of Psychiatry – Rouvray Hospital Centre, 4, rue Paul Eluard, 76300 Sotteville-lès-Rouen, France.

Tel: 00 33 2 32 95 68 25, Fax: 00 33 2 32 95 68 24

ABSTRACT

Background: Posttraumatic stress disorder (PTSD) is a severe and frequent affection that is highly comorbid to major depressive disorder. Comorbid PTSD and depression are usually treatment-resistant, with a high risk of functional impairment and suicide. Esketamine nasal spray is a recent validated treatment for treatment-resistant depression (TRD), but its efficacy on comorbid TRD-PTSD remains insufficiently documented. In particular, flashbacks can occur during esketamine administration and their influence on clinical outcomes is unknown.

Objectives: Our main objective was to describe esketamine-induced traumatic flashbacks and their impact on clinical trajectories within a sample of patients with comorbid TRD-PTSD.

Methods: We retrospectively collected clinical data of patients receiving esketamine nasal spray for TRD with comorbid PTSD who experienced at least one flashback of their trauma during esketamine sessions across 11 psychiatric departments.

Results: Between February 2020 and March 2023, 22 adult patients with TRD met inclusion criteria. In sixteen patients (72.7%) flashbacks disappeared as the sessions progressed. In six patients (27.3%), esketamine treatment was stopped because of persistent flashbacks. When esketamine was continued, clinical response was observed both for depression and PTSD (depression response rate: 45.5% and remission rate: 22.7%; PTSD response rate: 45.5% and remission: 18.2%).

Limitations: The retrospective design of the study and the absence of a comparator group are the main limitations of our study.

Conclusions: Our results suggest that the occurrence of esketamine-induced traumatic flashbacks does not hinder clinical response. On the contrary, when managed appropriately and combined with targeted psychotherapy, it could even contribute to positive outcomes.

Key words: treatment-resistant depression, post-traumatic stress disorder, esketamine nasal spray, traumatic flashbacks

Highlights

- Esketamine nasal spray is recently validated for treatment-resistant depression.
- Its efficacy on comorbid post-traumatic stress disorder is poorly documented.
- Traumatic flashbacks can occur during esketamine administration.
- Esketamine-induced traumatic flashbacks does not hinder its clinical response.

Clinical impact statement

Esketamine nasal spray is recently validated for treatment-resistant depression (TRD). Its efficacy on comorbid post-traumatic stress disorder (PTSD) is poorly documented. In this study, we reported the data of 22 adult patients who received esketamine nasal spray for TRD with comorbid PTSD and experienced flashbacks during esketamine sessions. These flashbacks did not appear to be a contreindication to the administration of esketamine and clinical response was observed both for depression and PTSD. Our results suggest that esketamine could be safely administered to patients with comorbid PTSD and TRD and that esketamine could lead to a substantial improvement in this population.

1. Introduction

Posttraumatic stress disorder (PTSD) includes persistent and recurrent symptoms of unwanted intrusive memories, avoidance behavior, changes in physical and emotional reactions, and negative changes in thinking and mood. Among intrusive symptoms, patients may experience flashbacks of the traumatic event where they feel as if it were happening again (American Psychiatric Association, 2013). PTSD lifetime prevalence is high ~8.3% (Kilpatrick et al., 2013). Its risk increases with trauma exposure severity, cumulative traumas, particularly for interpersonal traumas, i.e., physical and sexual assault in the context of relationship (Kessler et al., 2017). PTSD is often comorbid with other mental disorders. In around half of the cases, it is associated with major depressive disorder, which is a high-risk factor for treatment-resistance (Hegel et al., 2005; Kaplan & Klinetob, 2000; Kessler et al., 1995). Additionally, comorbid depression and PTSD induce greater functional impairment (Post et al., 2011) and a three times higher likelihood of suicidality than for depression alone (Karatzias et al., 2019).

Treatment strategies for PTSD include early interventions for prevention (Kearns et al., 2012), trauma-focused therapies (Bisson et al., 2013; Lenz et al., 2017; Watkins et al., 2018), and pharmacological treatments (e.g., antidepressants and α1 blocker) (Burback et al., 2023). Trauma-centered exposure psychotherapy is the first-line recommended therapy and has shown a significant efficacy in treating PTSD (Cusack et al., 2016; Watts et al., 2013). It consists in reactivating the traumatic memory in a safe environment and to learn how to prevent anxious response in order to reconsolidate the memory without its fear component, a phenomenon called fear extinction (Foa et al., 2008). However, up to 50% of patients may not be fully responsive to a correctly conducted psychotherapy when they have access to it (Resick et al., 2017; Steenkamp et al., 2015). By contrast, pharmacological therapy has an efficacy that appears to be modest at most (Watts et al., 2013) and in any case lower than that of psychotherapy (Merz et al., 2019). Innovative treatments for PTSD are therefore strongly needed, especially as current pharmacological and psychological therapies in combination do not appear to sufficiently improve clinical outcomes (Hetrick et al., 2010)

Ketamine is a noncompetitive *N*-methyl-D-aspartate glutamate receptor antagonist whose antidepressant properties at the dose of 0.5 mg/kg have been evidenced about 20 years ago (Berman et al., 2000) and have since been robustly reproduced (Hashimoto, 2019). Ketamine has also shown promising results in comorbid PTSD with treatment-resistant depression (TRD) (Albott et al., 2018) and in chronic PTSD alone (Duek et al., 2021; Feder et al., 2014, 2021; Pradhan et al., 2017), although some replications failed (Abdallah et al., 2022). More recently,

S-enantiomer of ketamine (esketamine), which was validated for the treatment of TRD – but not PTSD –(European Medicines Agency, 2021; U.S. Food and Drug Administration, 2020), was found to improve both PTSD and depressive symptoms in a cohort of patients suffering from comorbid PTSD and TRD (Rothärmel et al., 2022). In this sample, it empirically appeared that patients could experience intense flashbacks during esketamine sessions, usually followed by a strong decrease or even a resolution of their PTSD symptoms.

In this present study, we extend those findings by compiling clinical data of patients who received esketamine nasal spray for TRD with comorbid PTSD and experienced flashbacks during esketamine sessions. Our first objective was to describe esketamine-induced traumatic flashbacks and their potential influence on clinical trajectories. A secondary aim was to provide guidelines to help patients and caregivers to manage these esketamine-induced flashbacks.

2. Material and Methods

2.1. General design and participants

This study was an observational, retrospective and multicentric study comprising patients (21–62 years) with TRD and comorbid PTSD who have received esketamine nasal spray between February 2020 and March 2023. All the patients were treated in compliance with the guidelines provided by the French regulatory authority (ANSM, *Agence Nationale de Sécurité du Médicament et des produits de santé*) and the good clinical practice for TRD patients. Several mental health services from different university hospitals were involved in the recruitment.

The inclusion criteria comprised: *i)* Age≥18 years old; *ii)* Diagnosis of moderate-to-severe TRD, in accordance with the DSM-5 criteria (Tolentino & Schmidt, 2018) for a single or recurrent episode of major depression within unipolar or bipolar disorder; *iii)* Absence of remission after an adequate dose and duration of two or more different antidepressants given during the current episode in case of unipolar disorder; lack of remission on lithium at an adequate plasmatic level (0.8 mEq/L) combined with lamotrigine (50-200 mg per day) or full-dose quetiapine monotherapy (300-600 mg/day) in case of bipolar disorder (Pacchiarotti et al., 2009); *iv)* Comorbid PTSD according to the DSM-5; *v)* Treatment by esketamine with at least one traumatic flashback during esketamine sessions.

The study was conducted in accordance with the Declaration of Helsinki. This protocol has been approved by our local ethics committee (E2022-39, CHU Rouen, July 29th, 2022).

2.2. Study procedures

All participants received esketamine nasal spray in a full or partial hospitalization setting. All patients had baseline blood tests and an electrocardiogram to determine medical stability before the initiation of esketamine. Post-administration monitoring was undertaken by a healthcare professional for at least 120 minutes. Esketamine was initiated at 56 mg and adjusted on an individual basis during the treatment period in accordance with the following treatment guidelines: weeks 1–4: 56 or 84 mg twice weekly; weeks 5–8: 56 or 84 mg once weekly; and week 9 and onwards: 56 or 84 mg once every 1 or 2 weeks (European Medicines Agency, 2021). However, psychiatrists were free to adjust treatment dosage and frequency on an individual basis, depending on the patient's response and tolerance to the treatment.

Anamnestic data were retrospectively collected. They included sociodemographic information, history of depressive disease, treatment history for the current major depressive episode (MDE), comorbidities, level of resistance (Maudsley staging model) and current psychotherapy received by the patient during esketamine treatment (CBT for cognitive behavioral therapy, or EMDR for eye movement desensitization and reprocessing).

Details on life-traumatic events, esketamine-induced trauma flashbacks and adverse events during esketamine sessions were also collected. Psychometric assessments included Montgomery-Åsberg Depression Rating Scale (MADRS) (Montgomery & Asberg, 1979), Clinical Global Impression-Suicide Scale (CGI-SS) (Lindenmayer et al., 2003), and PTSD Checklist for DSM-5 (PCL-5) (Ashbaugh et al., 2016). They were obtained from patients' medical records at the time of esketamine-induced flashbacks. For depression, response to treatment was defined as an improvement of more than 50% compared to baseline MADRS scores (Frank et al., 1991) and remission was defined as the resolution of depressive symptoms, no longer allowing a diagnosis of depression according to the criteria established by the DSM-5, with a MADRS score of less than 10 (McIntyre et al., 2005). For PTSD symptoms, no psychometric assessments were routinely used. We report PCL-5 whenever available and response and remission as assessed by the patient's clinician.

2.3. Statistics

Statistical analyses were conducted using SPSS, version 28 (IBM, Armonk, NY). All tests were two-tailed, with a statistical significance level set at $p<0.05$. Continuous variables are expressed as mean ± standard deviation (SD), while categorical variables are reported as

percentages. Fisher's exact test was used to assess the statistical relationship between two categorical outcomes. For continuous outcomes (number of adverse effects and depression score), Mann-Whitney non-parametric test was used.

3. Results

3.1. Sample characteristics

Eleven clinical centers took part in the study. We included a total of 22 patients (72.7% with unipolar and 27.3% with bipolar disorder) who had experienced at least one flashback of their trauma during an esketamine session. The sociodemographic and clinical characteristics are reported in **Table 1**.

Trauma causing PTSD was reported by patients as being related to sexual abuse in adulthood (n=10), sexual abuse in childhood (n=9), physical violence (n=7), discovery of a person who died (n=4), emotional negligence during childhood (n=1), forced abortion (n=1), children death (n=1) and workplace bullying (n=1).

Table 1. Characteristics of the sample.

	N=22
Sociodemographic characteristics	
Age, mean ± SD (range)	41.8 ± 12.0 (21-62)
Sex (F/M)	19/3
Marital status	
Single	7 (31.8%)
Married	14 (63.6%)
Divorced	1 (4.5%)
Professional occupation	
Sick leave	16 (72.7%)
Unemployed	4 (18.2%)
Student	2 (9.1%)
Current depressive episode	
Duration (months), mean ± SD (range)	35.7 ± 33.2 (3-120)
Maudsley staging level, mean ± SD (range)	8.4 ± 2.3 (3-13)
Psychiatric comorbities	
Unipolar depression	16 (72.7%)
Bipolar I disorder	2 (9.1%)
Bipolar II disorder	4 (18.2%)
Simplex PTSD	3 (13.6%)
Complex PTSD	19 (86.4%)
Anxiety disorder	14 (63.6%)
Eating disorder	9 (40.9%)
Alcohol use disorder	5 (22.7%)
Substance use disorder (other than alcohol)	4 (18.2%)
Previous suicidal attempts	17 (77.3%)

PTSD: Posttraumatic stress disorder; SD: Standard deviation.

3.2. Esketamine-induced traumatic flashbacks

First-induced flashbacks occurred on average during the 4.4th session (SD 5.1, median 2.5, range 1-22): 40.9% of patients had flashbacks during the first session and 77.3% had flashbacks during the first five sessions.

During the week of flashbacks' occurrence, the clinical symptoms were still high for depression (mean MADRS score 29.6 ± 9.8; n=22; range 6-50) and for PTSD (mean PCL-5 score 52.9 ± 15.0; n=9; range 31-73). The mean CGI-SS score was 2.2 ±1.6 (n=20; range 0-5).

The dosage of esketamine during the first session in which the flashbacks occurred was 56 mg in 40.9% of the patients (n=9) and 84 mg in 59.1% of the patients (n=13). Esketamine was administered during an individual session for 81.8% of the patients (n=18) and during a group session (large room with beds separated by screens) for 18.2% of the patients (n=4).

Concomitant medication prescriptions included benzodiazepines (77.3%, n=17) and antipsychotics (54.5%, n=12). All patients except one were taking antidepressants: venlafaxine (n=5), fluoxetine (n=3), duloxetine (n=3), sertraline (n=3), mirtazapine (n=3), clomipramine (n=2), paroxetine (n=2), amitriptyline (n=1) or vortioxetine (n=1). Seven patients (31.8%) were treated by lithium. Regarding psychotherapy, twelve patients (54.5%) were undergoing cognitive-behavioral therapy (CBT) or eye movement desensitization and reprocessing (EMDR).

The other adverse effects induced by esketamine were: depersonalization/derealization (without flashbacks) (59.1%, n=13), nausea and vomiting (27.3%, n=6), sensory hypersensitivity (27.3%, n=6), headache (22.7%, n=5), sedation (13.6%, n=3) and high blood pressure (4.6%, n=1).

3.3. Flashbacks description

The patients experienced flashbacks about the following traumatic events: rape (n=13), physical violence (n=7), discovery of the dead person (n=4), delivery with pregnancy denial (n=1) and forced abortion (n=1). In 36.4% of the patients (n=8), the flashbacks revealed the trauma to the medical team. Less than one third of patients (27.3%, n=6) had flashbacks about events they had forgotten. The flashbacks induced strong anxiety (81.8%, n=18), agitation (31.8%, n=7), or catharsis with crying and feeling of release (54.5%, n=12). According to the patients, the emotional impact was mild (4.5%, n=1), moderate (50.0%, n=11) or severe (45.5%, n=10). The impact was not associated with co-prescribed antipsychotics (Fisher's

exact test; $p=1.00$), depression score (U=58; $p=0.895$), or the number of adverse effects (U=40; $p=0.156$).

In some patients with multiple traumas, we observed that esketamine-induced flashbacks would preferentially start with the most recent trauma and progressively involve older traumas. Interestingly, one patient experienced a flashback where the traumatic scene had a different ending that was much more soothing for her than her usual flashbacks.

Verbatim of patients and caregivers

Patients described very vivid and distressing esketamine-induced flashbacks: "*I didn't expect to relive the events so realistically. I could not get away from the images, I was frightened, I felt like I was present in my memories, as if my eyes were back in the body of the little girl that I was at the time… I could see the images with great precision in terms of colors, light...*"

The reaction of caregivers was considered by patients as essential: "*The support and kindness of the nurse were precious to me. The climate of trust she created allowed me to verbalize what I was experiencing.*"

Caregivers also reported an intense experience: "*She relived traumatic scenes from the past as if they were happening in the present. I tried to put her back in the present, in the office, by pointing out the place, the time and my presence with a reassuring voice. I asked her to verbalize what she was seeing and to try to change the course of the scenario by finding a way out. Once the peak passed, the patient's face relaxed and she reported a decrease in the flow of thoughts and images.*"

3.4. Clinical course

Eighteen patients had finished their esketamine cure when data was collected. For those patients, the average total number of esketamine sessions was 23.2 (SD 15.6, range 4-54) and flashbacks occurred in 50.6% (SD 36.3) of the esketamine sessions.

In 72.7% of the full sample of patients (n=16), flashbacks disappeared as the sessions progressed. But, in six patients (27.3%), esketamine treatment was stopped because of persistent flashbacks on average after 6 ± 3 sessions: 3 patients had a partial response to depression but not to PTSD; 3 patients had no response to depression and PTSD. Three patients wanted to continue working on their traumas in CBT or EMDR after stopping the esketamine sessions.

As regards the clinical outcome, response and remission rates for depression were 45.5% (n=10) and 22.7% (n=5) respectively and response and remission rates for PTSD were 45.5% (n=10) and 18.2% (n=4) respectively, about 6 months after the start of treatment. A significant association was found between responses for depression and for PTSD (Fisher's exact test; $p=0.002$) and between remission for depression and for PTSD (Fisher's exact test; $p<0.001$).

Co-treatment by CBT or EMDR was neither associated with depression response (Fisher's exact test; $p=1.00$) nor with PTSD response (Fisher's exact test; $p=0.675$). Flashbacks impact was significantly associated with PTSD response (Fisher's exact test; $p=0.006$): the responder rate was 91.7% in patients with a light to mild flashbacks impact (n=12) and only 30% in patients with severe impact (n=10).

Patients verbally reported a feeling of catharsis following esketamine-induced flashbacks in 12 cases but this feeling was not associated with PTSD response (Fisher's exact test; $p=0.378$).

4. Discussion

In this retrospective observational study, we report 22 cases of patients with comorbid TRD and PTSD who received esketamine as an antidepressant treatment and experienced trauma flashbacks during at least one esketamine session. Trauma flashbacks are part of PTSD dissociative symptomatology. They can occur spontaneously and significantly contribute to patients' suffering. At first sight, bringing those symptoms back during esketamine session could thus be considered as an adverse effect that should be avoided or impose esketamine cessation. In this cohort, esketamine was indeed interrupted in 27.3% of the patients because of a flashback occurrence during a session. In participants for whom esketamine was continued, the flashbacks gradually disappeared over the course of sessions and the patients exhibited an improvement on both depression and PTSD scores (45.5% of response and ~20% of remission on average) without any severe adverse event. Moreover, the outcomes for depression and PTSD were significantly correlated, suggesting that esketamine could be efficient either on one of these two dimensions with beneficial consequences on the other, or simultaneously on the two dimensions.

Esketamine could contribute to PTSD healing at different levels. First, it seems to help patients to verbalize their trauma. Indeed, 36.4% of them had not previously talked about their trauma to the medical team. More importantly, esketamine may facilitate access to buried memories

since 27.3% of patients reported that they had forgotten the events that reappeared during the session. Thanks to its antidepressant properties, esketamine could decrease the negative effects associated with flashbacks and act as an analgesic molecule favoring desensitization. Finally, derealization occurring during dissociation sometimes places the patient in the position of being a spectator to traumatic reviviscences, which can help to gain emotional distance.

Interestingly, in this case series, the impact of flashbacks seemed to play a role in the subsequent outcome, with a higher PTSD response rate if the impact was light or mild and a lower rate of response when the impact was severe. The profile of these two categories of patients may differ and confounding factors could have contributed both to the intensity of flashbacks impact and to the response rates, independently of any esketamine administration. Nevertheless, this result is in line with the general principles of exposure-based behavioral therapies in PTSD (Rothbaum & Schwartz, 2002): sufficiently long exposure in a safe environment allows for habituation and extinction of fear, while the persistence of anxiety at the end of the session may further sensitize patients. In this regard, we explored whether the co-prescription of antipsychotics could decrease flashbacks impact but did not observe any significant effects. Further studies are needed to assess whether co-prescriptions could help to decrease anxiety, thereby promoting extinction.

Both the access to forgotten trauma during esketamine sessions and the anti-chronologically order in which flashbacks often occur suggest that esketamine may favor memory reactivation, since relatively more recent memories are more prone to destabilization than older ones (Frankland et al., 2006). Finally, flashbacks were sometimes very intense and had a cathartic effect even if this aspect did not seem to be associated with PTSD resolution.

Because of the progressive decrease in flashbacks observed over the course of the sessions in this sample, we suggest that trauma-focused psychotherapy, with exposure-extinction procedure, could be delivered after esketamine administration to potentiate esketamine effect. This proposal is also supported by previously documented neurocognitive properties of ketamine, notably its effects on emotional processing and on synaptic plasticity.

Indeed, first, ketamine has been shown to decrease amygdala reactivity to negative emotional stimuli in healthy controls (Abel et al., 2003; Scheidegger et al., 2016) and in patients with major depressive disorder (C.-T. Li et al., 2016). In patients with PTSD, ketamine's beneficial effects are correlated with an increased top-down inhibition of the amygdala by the ventromedial prefrontal cortex during emotional face-viewing, and ketamine efficacy is higher

in patients with low prefrontal inhibition of amygdala responses to faces at baseline (Norbury et al., 2021). In this view, esketamine would normalize the increased sensitivity to negative emotional stimuli in depression and PTSD (Groenewold et al., 2013; Hayes et al., 2012) by restoring the prefrontal control over the amygdala. Psychotherapeutic techniques could even be proposed during esketamine sessions to better manage the advent of flashbacks and decrease fear memory reconsolidation. For instance, it has been suggested that the link between dissociation and mindfulness may be mediated by attention and emotional acceptance (Vancappel et al., 2023), and that specific exercises targeting both processes would be efficient to treat dissociative symptoms (Bishop et al., 2004). In order to foster acceptance, caregivers can use the "train metaphor" inspired from EMDR techniques (Shapiro, 1989). The patient is invited to consider that he is like the passenger on the train watching the landscape passing by. The landscape can be composed of emotions, sensations, thoughts, and images. The patient is encouraged letting this landscape pass. Then, the patient is invited to pay attention to the present moment by noting what she/he currently perceives with his five senses and from her/his internal experience (emotions, sensations, thoughts). This "dropping anchor" technique (Harris, 2017) derived from Acceptance and Commitment Therapy (ACT) (Hayes et al., 2012), allows the patient to be here and now, intentionally, without filter or judgment.

Second, under esketamine, flashbacks occur when patients are in a different state of mind and their subjective feelings and memories may be more easily reshaped. Importantly, its effect on memory seems to be highly dependent on the delay with encoding. Indeed, several studies found that ketamine could inflate memory retention or impede fear extinction when administered before or right after fear conditioning in experimental psychology paradigms (Corlett et al., 2013; Honsberger et al., 2015; Morena et al., 2017), with potential detrimental effect on PTSD outcomes (Juven-Wetzler et al., 2014; Schönenberg et al., 2005, 2008). On the contrary, we posit that ketamine elicits ancient memories destabilization through subjective experience of dissociation. Indeed, at a neurobiological level, ketamine acts as an NMDA antagonist, resulting in a secondary glutamate release and *in fine* in brain excitation and an increased synaptic plasticity (Duman et al., 2016), notably in the prefrontal cortex (N. Li et al., 2010; Pham & Gardier, 2019). Synaptic plasticity and glutamate receptor modulators have been suggested to play a role in memory destabilization and reconsolidation (Finnie & Nader, 2012; Monfils et al., 2009). More specifically, ketamine seems to facilitate fear memory extinction (Duek et al., 2021; Philippens et al., 2021; Wei et al., 2020) possibly through a decrease of fear

memory reconsolidation (Duclot et al., 2016) or a decreased access to contextual details during memory retrieval (Honey et al., 2005).

In line with previous proposals (Fattore et al., 2018), we therefore suggest that esketamine constitutes a pharmacological tool that could foster trauma-resolution through different mechanisms: 1) destabilizing memory and increasing the advent of flashbacks, 2) enhancing patients' control over their negative emotions during flashbacks, a mechanism that could be reinforced through dedicated psychotherapeutic interventions and 3) reducing fear memory reconsolidation during reactivation and thereby favoring extinction. In this perspective, we strongly recommend that PTSD should not be considered as an exclusion criterion for esketamine treatment and that patients with PTSD receiving esketamine should be informed in advance of the possible advent of flashbacks during the sessions. Then, they could be proposed a psychotherapeutic support during administration to maximize benefits. Caregivers must also be prepared for this type of reaction.

Interestingly, another molecule with a different pharmacological mechanism, 3,4-methylenedioxy-methamphetamine (MDMA), significantly reduces PTSD and depressive symptoms when associated with psychotherapy (Mitchell et al., 2021; Smith et al., 2022). MDMA has specific cognitive effects that are particularly relevant to alleviating PTSD symptoms. For instance, it enhances emotional empathy (Hysek et al., 2014; Kuypers et al., 2017), increases self-compassion (Kamboj et al., 2015) and has several pro-social effects (Bedi et al., 2009; Hysek et al., 2014) that could facilitate the therapeutic alliance and psychotherapeutic work. Like ketamine, it decreases amygdala reactivity to negative emotional stimuli (Bedi et al., 2009; Gamma et al., 2000) and promotes fear memory extinction (Hake et al., 2019; Young et al., 2015), suggesting that those changes play a crucial role in treatment efficacy. The interest of reactivating memories and to modulate them with other therapeutic interventions has been investigated with mixed results, e.g., β-adrenergic receptor antagonist propranolol (Brunet et al., 2018; Raut et al., 2022; Vaiva et al., 2003), transmagnetic stimulation (Isserles et al, 2013, but: Isserles et al., 2021) and electroconvulsivotherapy (Gahr et al., 2014; Kroes et al., 2014, but: Tang et al., 2021).

On another note, ketamine has already been proposed as a "maximizer" of cognitive behavioral therapy in other psychiatric affections, such as obsessive-compulsive disorder (Rodriguez et al., 2016), depression (Bottemanne et al., 2021; Dore et al., 2019; Hasler, 2020) and addictions

(Grabski et al., 2022; Krupitsky et al., 2002), notably through its ability to rewrite maladaptive memories (Das et al., 2019). Our results suggest that it could also be the case for PTSD.

Despite the perspectives that our study provides, several limitations should be mentioned. First, it was not designed to compare the outcomes for patients with comorbid TRD and PTSD based on whether or not they would experience traumatic flashbacks under esketamine; or to assess the prevalence of esketamine-induced trauma flashbacks among this population. Indeed, data were retrospectively collected after the empirical observation that some patients had such flashbacks during an esketamine administration and felt subsequently relieved. Instead, our study intends to highlight the existence of this phenomenon and emphasizes that these patients may have positive outcomes if esketamine is not interrupted. Further studies with a systematic and prospective data collection are required to evaluate the prevalence of esketamine-induced trauma flashbacks and its potential role as a predicting factor of response. Second, our study does not allow to compare esketamine efficacy in comorbid TRD and PTSD and TRD alone. The response rate looks quite similar to that of a real-world study that assessed the efficacy of esketamine in TRD in a French cohort (45.5% in our study versus 47.8% in this study on TRD) (Samalin et al., 2022). By contrast, the rate of remission appears to be lower (22.7% in our study versus 37% in the TRD study), possibly because comorbid PTSD leads to the persistence of residual symptoms that are more intractable. Larger cohorts including both patients with TRD alone and comorbid PTSD and TRD could in the future perform those comparisons.

Conclusion

In summary, this case series shows that esketamine could be safely administered to patients with comorbid PTSD and TRD even if flashbacks occur during the sessions. Comorbid TRD and PTSD usually have a poor prognosis (Karatzias et al., 2019; Panagioti et al., 2012; Post et al., 2011) but our study suggests that esketamine could lead to a substantial improvement in this population (see also: Rothärmel et al., 2022). Indeed, thanks to its specific neurocognitive effects, esketamine may favor the advent of flashbacks but also facilitate trauma-resolution by decreasing negative emotions associated to them and promoting fear extinction, in particular if psychotherapy is simultaneously delivered.

Conflict of interest

Most of the authors of this article received honoraria or consulting fees from Janssen-Cilag (MR, LM, MH, JH, SB, MD, EO, BG, AS, RG, RR, DZ, LS, OG, AL, LB) with no financial or other relationship relevant to the subject of this article.

Acknowledgments

We thank all the participants. We also thank Gérald Adler, Cherifa Benosman, Julien Corniquet, Wendy Chevallier, Graziella Guesdon, Carine Kerninon, Corinne Lagniez and and Julie Mougeolle for assistance in esketamine administration.

LB thanks the *Fondation Bettencourt-Schueller*, the *Philippe Foundation,* the *Foundation L'Oréal-Unesco* and the National Institute of Mental Health (R01MH116038 and U01MH121766) for their support.

Author contributions statements

Conceptualization : MR, LM, LB ; *Formal analysis* : VM ; *Investigation* : MR, LM, ALa, FK, MH, E-VB-L, Ale, JH, PR, MC, TH, RR, ; *Methodology* : MR, LM, LB, AS ; *Project administration* : MR, LM ; *Supervision* : OG, AS, RG, WE, SB, PM, EO, BG, DZ, LS ; *writing original draft* : MR, LB ; *Writing—review & editing* : ALa, SB, AS, RG, WE, TH, LS.

References

Abdallah, C. G., Roache, J. D., Gueorguieva, R., Averill, L. A., Young-McCaughan, S., Shiroma, P. R., Purohit, P., Brundige, A., Murff, W., Ahn, K.-H., Sherif, M. A., Baltutis, E. J., Ranganathan, M., D'Souza, D., Martini, B., Southwick, S. M., Petrakis, I. L., Burson, R. R., Guthmiller, K. B., … Krystal, J. H. (2022). Dose-related effects of ketamine for antidepressant-resistant symptoms of posttraumatic stress disorder in veterans and active duty military : A double-blind, randomized, placebo-controlled multi-center clinical trial. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*. https://doi.org/10.1038/s41386-022-01266-9

Abel, K. M., Allin, M. P. G., Kucharska-Pietura, K., David, A., Andrew, C., Williams, S., Brammer, M. J., & Phillips, M. L. (2003). Ketamine alters neural processing of facial emotion recognition in healthy men : An fMRI study. *NeuroReport*, *14*(3), 387-391.

Albott, C. S., Lim, K. O., Forbes, M. K., Erbes, C., Tye, S. J., Grabowski, J. G., Thuras, P., Batres-Y-Carr, T. M., Wels, J., & Shiroma, P. R. (2018). Efficacy, Safety, and Durability of Repeated Ketamine Infusions for Comorbid Posttraumatic Stress Disorder and Treatment-Resistant Depression. *The Journal of Clinical Psychiatry*, *79*(3), 17m11634. https://doi.org/10.4088/JCP.17m11634

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders : DSM-5* (5th edition). American Psychiatric Association. https://dsm.psychiatryonline.org/doi/book/10.1176/appi.books.9780890425787

Ashbaugh, A. R., Houle-Johnson, S., Herbert, C., El-Hage, W., & Brunet, A. (2016). Psychometric Validation of the English and French Versions of the Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5). *PloS One*, *11*(10), e0161645. https://doi.org/10.1371/journal.pone.0161645

Bedi, G., Phan, K. L., Angstadt, M., & de Wit, H. (2009). Effects of MDMA on sociability and neural response to social threat and social reward. *Psychopharmacology*, *207*(1), 73-83. https://doi.org/10.1007/s00213-009-1635-z

Berman, R. M., Cappiello, A., Anand, A., Oren, D. A., Heninger, G. R., Charney, D. S., & Krystal, J. H. (2000). Antidepressant effects of ketamine in depressed patients. *Biological psychiatry*, *47*(4), 351-354.

Bishop, S. R., Lau, M., Shapiro, S., Carlson, L., Anderson, N. D., Carmody, J., Segal, Z. V., Abbey, S., Speca, M., Velting, D., & Devins, G. (2004). Mindfulness : A proposed operational definition. *Clinical Psychology: Science and Practice*, *11*(3), 230-241. https://doi.org/10.1093/clipsy.bph077

Bisson, J. I., Roberts, N. P., Andrew, M., Cooper, R., & Lewis, C. (2013). Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *The Cochrane Database of Systematic Reviews*, *2013*(12), CD003388. https://doi.org/10.1002/14651858.CD003388.pub4

Bottemanne, H., Baldacci, A., Muller, C., Boyreau, A., & Claret, A. (2021). Psychothérapie augmentée par la kétamine (KAP) dans les troubles de l'humeur : Mode d'emploi. *L'Encéphale*, S0013700621002244. https://doi.org/10.1016/j.encep.2021.08.011

Brunet, A., Saumier, D., Liu, A., Streiner, D. L., Tremblay, J., & Pitman, R. K. (2018). Reduction of PTSD Symptoms With Pre-Reactivation Propranolol Therapy : A Randomized Controlled Trial. *American Journal of Psychiatry*, *175*(5), 427-433. https://doi.org/10.1176/appi.ajp.2017.17050481

Burback, L., Brémault-Phillips, S., Nijdam, M. J., McFarlane, A., & Vermetten, E. (2023). Treatment of Posttraumatic Stress Disorder : A State-of-the-art Review. *Current Neuropharmacology*. https://doi.org/10.2174/1570159X21666230428091433

Corlett, P. R., Cambridge, V., Gardner, J. M., Piggot, J. S., Turner, D. C., Everitt, J. C., Arana, F. S., Morgan, H. L., Milton, A. L., Lee, J. L., Aitken, M. R. F., Dickinson, A., Everitt, B. J., Absalom, A. R., Adapa, R., Subramanian, N., Taylor, J. R., Krystal, J. H., & Fletcher, P. C. (2013). Ketamine Effects on Memory Reconsolidation Favor a Learning Model of Delusions. *PLOS ONE*, *8*(6), e65088. https://doi.org/10.1371/journal.pone.0065088

Cusack, K., Jonas, D. E., Forneris, C. A., Wines, C., Sonis, J., Middleton, J. C., Feltner, C., Brownley, K. A., Olmsted, K. R., Greenblatt, A., Weil, A., & Gaynes, B. N. (2016). Psychological treatments for adults with posttraumatic stress disorder : A systematic review and meta-analysis. *Clinical Psychology Review*, *43*, 128-141. https://doi.org/10.1016/j.cpr.2015.10.003

Das, R. K., Gale, G., Walsh, K., Hennessy, V. E., Iskandar, G., Mordecai, L. A., Brandner, B., Kindt, M., Curran, H. V., & Kamboj, S. K. (2019). Ketamine can reduce harmful drinking by pharmacologically rewriting drinking memories. *Nature Communications*, *10*(1), Art. 1. https://doi.org/10.1038/s41467-019-13162-w

Dore, J., Turnipseed, B., Dwyer, S., Turnipseed, A., Andries, J., Ascani, G., Monnette, C., Huidekoper, A., Strauss, N., & Wolfson, P. (2019). Ketamine Assisted Psychotherapy (KAP) : Patient Demographics, Clinical Data and Outcomes in Three Large Practices Administering Ketamine with Psychotherapy. *Journal of Psychoactive Drugs*, *51*(2), 189-198. https://doi.org/10.1080/02791072.2019.1587556

Duclot, F., Perez-Taboada, I., Wright, K. N., & Kabbaj, M. (2016). Prediction of individual differences in fear response by novelty seeking, and disruption of contextual fear memory reconsolidation by ketamine. *Neuropharmacology*, *109*, 293-305. https://doi.org/10.1016/j.neuropharm.2016.06.022

Duek, O., Li, Y., Kelmendi, B., Amen, S., Gordon, C., Milne, M., Krystal, J. H., Levy, I., & Harpaz-Rotem, I. (2021). Modulating amygdala activation to traumatic memories with a single ketamine infusion. *medRxiv*, 2021.07.07.21260166. https://doi.org/10.1101/2021.07.07.21260166

Duman, R. S., Aghajanian, G. K., Sanacora, G., & Krystal, J. H. (2016). Synaptic plasticity and depression : New insights from stress and rapid-acting antidepressants. *Nature Medicine*, *22*(3), 238-249. https://doi.org/10.1038/nm.4050

European Medicines Agency. (2021). *Spravato 28 mg nasal spray (esketamine), solution. Summary of Product Characteristics.* https://www.ema.europa.eu/en/documents/product-information/spravato-epar-product-information_en.pdf

European Medicines Agency, 2021. Spravato 28 mg nasal spray (esketamine), solution. Summary of Product Characteristics. [WWW Document]. URL

https://www.ema.europa.eu/en/documents/product-information/spravato-epar-product-information_en.pdf. (s. d.).

Fattore, L., Piva, A., Zanda, M. T., Fumagalli, G., & Chiamulera, C. (2018). Psychedelics and reconsolidation of traumatic and appetitive maladaptive memories : Focus on cannabinoids and ketamine. *Psychopharmacology*, *235*(2), 433-445. https://doi.org/10.1007/s00213-017-4793-4

Feder, A., Costi, S., Rutter, S. B., Collins, A. B., Govindarajulu, U., Jha, M. K., Horn, S. R., Kautz, M., Corniquel, M., Collins, K. A., Bevilacqua, L., Glasgow, A. M., Brallier, J., Pietrzak, R. H., Murrough, J. W., & Charney, D. S. (2021). A Randomized Controlled Trial of Repeated Ketamine Administration for Chronic Posttraumatic Stress Disorder. *The American Journal of Psychiatry*, *178*(2), 193-202. https://doi.org/10.1176/appi.ajp.2020.20050596

Feder, A., Parides, M. K., Murrough, J. W., Perez, A. M., Morgan, J. E., Saxena, S., Kirkwood, K., Aan Het Rot, M., Lapidus, K. A. B., Wan, L.-B., Iosifescu, D., & Charney, D. S. (2014). Efficacy of intravenous ketamine for treatment of chronic posttraumatic stress disorder : A randomized clinical trial. *JAMA Psychiatry*, *71*(6), 681-688. https://doi.org/10.1001/jamapsychiatry.2014.62

Finnie, P. S. B., & Nader, K. (2012). The role of metaplasticity mechanisms in regulating memory destabilization and reconsolidation. *Neuroscience & Biobehavioral Reviews*, *36*(7), 1667-1707. https://doi.org/10.1016/j.neubiorev.2012.03.008

Foa, E. B., Chrestman, K. R., & Gilboa-Schechtman, E. (2008). *Prolonged exposure therapy for adolescents with PTSD emotional processing of traumatic experiences, therapist guide*. Oxford University Press.

Frank E, Prien RF, Jarrett RB, Keller MB, Kupfer DJ, Lavori PW, Rush AJ, Weissman MM. (s. d.). *Conceptualization and rationale for consensus definitions of terms in major depressive disorder. Remission, recovery, relapse, and recurrence. Arch Gen Psychiatry. 1991 Sep;48(9):851-5. Doi : 10.1001/archpsyc.1991.01810330075011. PMID: 1929776.*

Frankland, P. W., Ding, H.-K., Takahashi, E., Suzuki, A., Kida, S., & Silva, A. J. (2006). Stability of recent and remote contextual fear memory. *Learning & Memory*, *13*(4), 451-457. https://doi.org/10.1101/lm.183406

Gahr, M., Schönfeldt-Lecuona, C., Spitzer, M., & Graf, H. (2014). Electroconvulsive Therapy and Posttraumatic Stress Disorder : First Experience With Conversation-Based Reactivation of Traumatic Memory Contents and Subsequent ECT-Mediated Impairment of Reconsolidation. *The Journal of Neuropsychiatry and Clinical Neurosciences*, *26*(3), E38-E39. https://doi.org/10.1176/appi.neuropsych.13070159

Gamma, A., Buck, A., Berthold, T., Liechti, M. E., & Vollenweider, F. X. (2000). 3,4-Methylenedioxymethamphetamine (MDMA) modulates cortical and limbic brain activity as measured by [H(2)(15)O]-PET in healthy humans. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *23*(4), 388-395. https://doi.org/10.1016/S0893-133X(00)00130-5

Grabski, M., McAndrew, A., Lawn, W., Marsh, B., Raymen, L., Stevens, T., Hardy, L., Warren, F., Bloomfield, M., Borissova, A., Maschauer, E., Broomby, R., Price, R.,

Coathup, R., Gilhooly, D., Palmer, E., Gordon-Williams, R., Hill, R., Harris, J., … Morgan, C. J. A. (2022). Adjunctive Ketamine With Relapse Prevention–Based Psychological Therapy in the Treatment of Alcohol Use Disorder. *American Journal of Psychiatry*, appi.ajp.2021.21030277. https://doi.org/10.1176/appi.ajp.2021.21030277

Groenewold, N. A., Opmeer, E. M., de Jonge, P., Aleman, A., & Costafreda, S. G. (2013). Emotional valence modulates brain functional abnormalities in depression : Evidence from a meta-analysis of fMRI studies. *Neuroscience and Biobehavioral Reviews*, *37*(2), 152-163. https://doi.org/10.1016/j.neubiorev.2012.11.015

Hake, H. S., Davis, J. K. P., Wood, R. R., Tanner, M. K., Loetz, E. C., Sanchez, A., Ostrovskyy, M., Oleson, E. B., Grigsby, J., Doblin, R., & Greenwood, B. N. (2019). 3,4-methylenedioxymethamphetamine (MDMA) impairs the extinction and reconsolidation of fear memory in rats. *Physiology & Behavior*, *199*, 343-350. https://doi.org/10.1016/j.physbeh.2018.12.007

Harris, Russ. Passez à l'ACT. Pratique de la thérapie d'acceptation et d'engagement, traduction de Penet Claude, traduction de Milleville Laurence. De Boeck Supérieur, 2017. (s. d.).

Hashimoto, K. (2019). Rapid-acting antidepressant ketamine, its metabolites and other candidates : A historical overview and future perspective. *Psychiatry and Clinical Neurosciences*, *73*(10), 613-627. https://doi.org/10.1111/pcn.12902

Hasler, G. (2020). Toward specific ways to combine ketamine and psychotherapy in treating depression. *CNS Spectrums*, *25*(3), 445-447. https://doi.org/10.1017/S1092852919001007

Hayes, J. P., Vanelzakker, M. B., & Shin, L. M. (2012). Emotion and cognition interactions in PTSD : A review of neurocognitive and neuroimaging studies. *Frontiers in Integrative Neuroscience*, *6*, 89. https://doi.org/10.3389/fnint.2012.00089

Hegel, M. T., Unützer, J., Tang, L., Areán, P. A., Katon, W., Noël, P. H., Williams, J. W., & Lin, E. H. B. (2005). Impact of comorbid panic and posttraumatic stress disorder on outcomes of collaborative care for late-life depression in primary care. *The American Journal of Geriatric Psychiatry: Official Journal of the American Association for Geriatric Psychiatry*, *13*(1), 48-58. https://doi.org/10.1176/appi.ajgp.13.1.48

Hetrick SE, Purcell R, Garner B, Parslow R. (s. d.). *Combined pharmacotherapy and psychological therapies for post traumatic stress disorder (PTSD). Cochrane Database Syst Rev. 2010 Jul 7;(7):CD007316. Doi : 10.1002/14651858.CD007316.pub2. PMID: 20614457.*

Honey, G. D., Honey, R. A. E., O'Loughlin, C., Sharar, S. R., Kumaran, D., Suckling, J., Menon, D. K., Sleator, C., Bullmore, E. T., & Fletcher, P. C. (2005). Ketamine Disrupts Frontal and Hippocampal Contribution to Encoding and Retrieval of Episodic Memory : An fMRI Study. *Cerebral Cortex*, *15*(6), 749-759. https://doi.org/10.1093/cercor/bhh176

Honsberger, M. J., Taylor, J. R., & Corlett, P. R. (2015). Memories reactivated under ketamine are subsequently stronger : A potential pre-clinical behavioral model of psychosis. *Schizophrenia research*, *164*(0), 227-233. https://doi.org/10.1016/j.schres.2015.02.009

Hysek, C. M., Schmid, Y., Simmler, L. D., Domes, G., Heinrichs, M., Eisenegger, C., Preller, K. H., Quednow, B. B., & Liechti, M. E. (2014). MDMA enhances emotional empathy

and prosocial behavior. *Social Cognitive and Affective Neuroscience*, *9*(11), 1645-1652. https://doi.org/10.1093/scan/nst161

Isserles, M., Shalev, A. Y., Roth, Y., Peri, T., Kutz, I., Zlotnick, E., & Zangen, A. (2013). Effectiveness of Deep Transcranial Magnetic Stimulation Combined with a Brief Exposure Procedure in Post-Traumatic Stress Disorder – A Pilot Study. *Brain Stimulation*, *6*(3), 377-383. https://doi.org/10.1016/j.brs.2012.07.008

Isserles, M., Tendler, A., Roth, Y., Bystritsky, A., Blumberger, D. M., Ward, H., Feifel, D., Viner, L., Duffy, W., Zohar, J., Keller, C. J., Bhati, M. T., Etkin, A., George, M. S., Filipcic, I., Lapidus, K., Casuto, L., Vaishnavi, S., Stein, A., … Ressler, K. J. (2021). Deep Transcranial Magnetic Stimulation Combined With Brief Exposure for Posttraumatic Stress Disorder : A Prospective Multisite Randomized Trial. *Biological Psychiatry*, *90*(10), 721-728. https://doi.org/10.1016/j.biopsych.2021.04.019

Juven-Wetzler, A., Cohen, H., Kaplan, Z., Kohen, A., Porat, O., & Zohar, J. (2014). Immediate ketamine treatment does not prevent posttraumatic stress responses in an animal model for PTSD. *European Neuropsychopharmacology: The Journal of the European College of Neuropsychopharmacology*, *24*(3), 469-479. https://doi.org/10.1016/j.euroneuro.2013.08.007

Kamboj, S. K., Kilford, E. J., Minchin, S., Moss, A., Lawn, W., Das, R. K., Falconer, C. J., Gilbert, P., Curran, H. V., & Freeman, T. P. (2015). Recreational 3,4-methylenedioxy-N-methylamphetamine (MDMA) or « ecstasy » and self-focused compassion : Preliminary steps in the development of a therapeutic psychopharmacology of contemplative practices. *Journal of Psychopharmacology (Oxford, England)*, *29*(9), 961-970. https://doi.org/10.1177/0269881115587143

Kaplan, M. J., & Klinetob, N. A. (2000). Childhood emotional trauma and chronic posttraumatic stress disorder in adult outpatients with treatment-resistant depression. *The Journal of Nervous and Mental Disease*, *188*(9), 596-601. https://doi.org/10.1097/00005053-200009000-00006

Karatzias, T., Hyland, P., Bradley, A., Cloitre, M., Roberts, N. P., Bisson, J. I., & Shevlin, M. (2019). Risk factors and comorbidity of ICD-11 PTSD and complex PTSD : Findings from a trauma-exposed population based sample of adults in the United Kingdom. *Depression and Anxiety*, *36*(9), 887-894. https://doi.org/10.1002/da.22934

Kearns, M. C., Ressler, K. J., Zatzick, D., & Rothbaum, B. O. (2012). Early interventions for PTSD : A review. *Depression and Anxiety*, *29*(10), 833-842. https://doi.org/10.1002/da.21997

Kessler, R. C., Aguilar-Gaxiola, S., Alonso, J., Benjet, C., Bromet, E. J., Cardoso, G., Degenhardt, L., de Girolamo, G., Dinolova, R. V., Ferry, F., Florescu, S., Gureje, O., Haro, J. M., Huang, Y., Karam, E. G., Kawakami, N., Lee, S., Lepine, J.-P., Levinson, D., … Koenen, K. C. (2017). Trauma and PTSD in the WHO World Mental Health Surveys. *European Journal of Psychotraumatology*, *8*(sup5), 1353383. https://doi.org/10.1080/20008198.2017.1353383

Kessler, R. C., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*, *52*(12), 1048-1060. https://doi.org/10.1001/archpsyc.1995.03950240066012

Kilpatrick, D. G., Resnick, H. S., Milanak, M. E., Miller, M. W., Keyes, K. M., & Friedman, M. J. (2013). National Estimates of Exposure to Traumatic Events and PTSD Prevalence Using DSM-IV and DSM-5 Criteria. *Journal of traumatic stress*, *26*(5), 537-547. https://doi.org/10.1002/jts.21848

Kroes, M. C. W., Tendolkar, I., van Wingen, G. A., van Waarde, J. A., Strange, B. A., & Fernández, G. (2014). An electroconvulsive therapy procedure impairs reconsolidation of episodic memories in humans. *Nature Neuroscience*, *17*(2), 204-206. https://doi.org/10.1038/nn.3609

Krupitsky, E., Burakov, A., Romanova, T., Dunaevsky, I., Strassman, R., & Grinenko, A. (2002). Ketamine psychotherapy for heroin addiction : Immediate effects and two-year follow-up. *Journal of Substance Abuse Treatment*, *23*(4), 273-283. https://doi.org/10.1016/S0740-5472(02)00275-1

Kuypers, K. P., Dolder, P. C., Ramaekers, J. G., & Liechti, M. E. (2017). Multifaceted empathy of healthy volunteers after single doses of MDMA : A pooled sample of placebo-controlled studies. *Journal of Psychopharmacology*, *31*(5), 589-598. https://doi.org/10.1177/0269881117699617

Lenz, A. S., Haktanir, A., & Callender, K. (2017). Meta-Analysis of Trauma-Focused Therapies for Treating the Symptoms of Posttraumatic Stress Disorder. *Journal of Counseling & Development*, *95*(3), 339-353. https://doi.org/10.1002/jcad.12148

Li, C.-T., Chen, M.-H., Lin, W.-C., Hong, C.-J., Yang, B.-H., Liu, R.-S., Tu, P.-C., & Su, T.-P. (2016). The effects of low-dose ketamine on the prefrontal cortex and amygdala in treatment-resistant depression : A randomized controlled study. *Human Brain Mapping*, *37*(3), 1080-1090. https://doi.org/10.1002/hbm.23085

Li, N., Lee, B., Liu, R.-J., Banasr, M., Dwyer, J. M., Iwata, M., Li, X.-Y., Aghajanian, G., & Duman, R. S. (2010). MTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science (New York, N.Y.)*, *329*(5994), 959-964. https://doi.org/10.1126/science.1190287

Merz, J., Schwarzer, G., & Gerger, H. (2019). Comparative Efficacy and Acceptability of Pharmacological, Psychotherapeutic, and Combination Treatments in Adults With Posttraumatic Stress Disorder : A Network Meta-analysis. *JAMA Psychiatry*, *76*(9), 904-913. https://doi.org/10.1001/jamapsychiatry.2019.0951

Mitchell, J. M., Bogenschutz, M., Lilienstein, A., Harrison, C., Kleiman, S., Parker-Guilbert, K., Ot'alora G., M., Garas, W., Paleos, C., Gorman, I., Nicholas, C., Mithoefer, M., Carlin, S., Poulter, B., Mithoefer, A., Quevedo, S., Wells, G., Klaire, S. S., van der Kolk, B., … Doblin, R. (2021). MDMA-assisted therapy for severe PTSD : A randomized, double-blind, placebo-controlled phase 3 study. *Nature Medicine*, *27*(6), Art. 6. https://doi.org/10.1038/s41591-021-01336-3

Monfils, M.-H., Cowansage, K. K., Klann, E., & LeDoux, J. E. (2009). Extinction-Reconsolidation Boundaries : Key to Persistent Attenuation of Fear Memories. *Science*, *324*(5929), 951-955. https://doi.org/10.1126/science.1167975

Montgomery SA, Asberg M. (s. d.). *A new depression scale designed to be sensitive to change. Br J Psychiatry. 1979 Apr;134:382-9. Doi : 10.1192/bjp.134.4.382. PMID: 444788.*

Morena, M., Berardi, A., Peloso, A., Valeri, D., Palmery, M., Trezza, V., Schelling, G., & Campolongo, P. (2017). Effects of ketamine, dexmedetomidine and propofol anesthesia

on emotional memory consolidation in rats : Consequences for the development of post-traumatic stress disorder. *Behavioural Brain Research*, *329*, 215-220. https://doi.org/10.1016/j.bbr.2017.04.048

Norbury, A., Rutter, S. B., Collins, A. B., Costi, S., Jha, M. K., Horn, S. R., Kautz, M., Corniquel, M., Collins, K. A., Glasgow, A. M., Brallier, J., Shin, L. M., Charney, D. S., Murrough, J. W., & Feder, A. (2021). Neuroimaging correlates and predictors of response to repeated-dose intravenous ketamine in PTSD : Preliminary evidence. *Neuropsychopharmacology*, *46*(13), Art. 13. https://doi.org/10.1038/s41386-021-01104-4

Pacchiarotti, I. et al. (2009) 'Treatment-resistant bipolar depression : Towards a new definition', Acta Psychiatrica Scandinavica, 120(6), pp. 429–440. Doi : 10.1111/j.1600-0447.2009.01471.x. (s. d.).

Panagioti, M., Gooding, P. A., & Tarrier, N. (2012). A meta-analysis of the association between posttraumatic stress disorder and suicidality : The role of comorbid depression. *Comprehensive Psychiatry*, *53*(7), 915-930. https://doi.org/10.1016/j.comppsych.2012.02.009

Pham, T. H., & Gardier, A. M. (2019). Fast-acting antidepressant activity of ketamine : Highlights on brain serotonin, glutamate, and GABA neurotransmission in preclinical studies. *Pharmacology & Therapeutics*, *199*, 58-90. https://doi.org/10.1016/j.pharmthera.2019.02.017

Philippens, I. H. C. H. M., Draaisma, L., Baarends, G., Krugers, H. J., & Vermetten, E. (2021). Ketamine treatment upon memory retrieval reduces fear memory in marmoset monkeys. *European Neuropsychopharmacology: The Journal of the European College of Neuropsychopharmacology*, *50*, 1-11. https://doi.org/10.1016/j.euroneuro.2021.04.004

Post, L. M., Zoellner, L. A., Youngstrom, E., & Feeny, N. C. (2011). Understanding the Relationship Between Co-occurring PTSD and MDD : Symptom Severity and Affect. *Journal of anxiety disorders*, *25*(8), 1123-1130. https://doi.org/10.1016/j.janxdis.2011.08.003

Pradhan, B., Wainer, I., Moaddel, R., Torjman, M., Goldberg, M., Sabia, M., Parikh, T., & Pumariega, A. (2017). Trauma Interventions using Mindfulness Based Extinction and Reconsolidation (TIMBER) psychotherapy prolong the therapeutic effects of single ketamine infusion on post-traumatic stress disorder and comorbid depression : A pilot randomized, placebo-controlled, crossover clinical trial. *Asia Pacific Journal of Clinical Trials: Nervous System Diseases*, *2*(3), 80. https://doi.org/10.4103/2542-3932.211589

Raut, S. B., Canales, J. J., Ravindran, M., Eri, R., Benedek, D. M., Ursano, R. J., & Johnson, L. R. (2022). Effects of propranolol on the modification of trauma memory reconsolidation in PTSD patients : A systematic review and meta-analysis. *Journal of Psychiatric Research*, *150*, 246-256. https://doi.org/10.1016/j.jpsychires.2022.03.045

Resick, P. A., Wachen, J. S., Dondanville, K. A., Pruiksma, K. E., Yarvis, J. S., Peterson, A. L., Mintz, J., & and the STRONG STAR Consortium. (2017). Effect of Group vs Individual Cognitive Processing Therapy in Active-Duty Military Seeking Treatment

for Posttraumatic Stress Disorder : A Randomized Clinical Trial. *JAMA Psychiatry*, *74*(1), 28-36. https://doi.org/10.1001/jamapsychiatry.2016.2729

Rodriguez, C. I., Wheaton, M., Zwerling, J., Steinman, S. A., Sonnenfeld, D., Galfalvy, H., & Simpson, H. B. (2016). Can Exposure-Based CBT Extend the Effects of Intravenous Ketamine in Obsessive-Compulsive Disorder? An Open-Label Trial. *The Journal of Clinical Psychiatry*, *77*(3), 7005. https://doi.org/10.4088/JCP.15l10138

Rothärmel, M., Benosman, C., El-Hage, W., Berjamin, C., Ribayrol, D., Guillin, O., Gaillard, R., Berkovitch, L., & Moulier, V. (2022). Efficacy and Safety of Intranasal Esketamine in Patients With Treatment-Resistant Depression and Comorbid Chronic Post-traumatic Stress Disorder : Open-Label Single-Arm Pilot Study. *Frontiers in Psychiatry*, *13*. https://www.frontiersin.org/articles/10.3389/fpsyt.2022.865466

Rothbaum, B. O., & Schwartz, A. C. (2002). Exposure Therapy for Posttraumatic Stress Disorder. *American Journal of Psychotherapy*, *56*(1), 59-75. https://doi.org/10.1176/appi.psychotherapy.2002.56.1.59

Samalin, L., Rothärmel, M., Mekaoui, L., Gaudré-Wattinne, E., Codet, M.-A., Bouju, S., & Sauvaget, A. (2022). Esketamine nasal spray in patients with treatment-resistant depression : The real-world experience in the French cohort early-access programme. *International Journal of Psychiatry in Clinical Practice*, *26*(4), 352-362. https://doi.org/10.1080/13651501.2022.2030757

Scheidegger, M., Henning, A., Walter, M., Lehmann, M., Kraehenmann, R., Boeker, H., Seifritz, E., & Grimm, S. (2016). Ketamine administration reduces amygdalo-hippocampal reactivity to emotional stimulation. *Human Brain Mapping*, *37*(5), 1941-1952. https://doi.org/10.1002/hbm.23148

Schönenberg, M., Reichwald, U., Domes, G., Badke, A., & Hautzinger, M. (2005). Effects of peritraumatic ketamine medication on early and sustained posttraumatic stress symptoms in moderately injured accident victims. *Psychopharmacology*, *182*(3), 420-425. https://doi.org/10.1007/s00213-005-0094-4

Schönenberg, M., Reichwald, U., Domes, G., Badke, A., & Hautzinger, M. (2008). Ketamine aggravates symptoms of acute stress disorder in a naturalistic sample of accident victims. *Journal of Psychopharmacology (Oxford, England)*, *22*(5), 493-497. https://doi.org/10.1177/0269881107082481

Shapiro, F. (1989). Efficacy of the Eye Movement Desensitization procedure in the treatment of traumatic memories. Journal of Traumatic Stress, 2(2), 199–223. Https://doi.org/10.1002/jts.2490020207. (s. d.).

Smith, K. W., Sicignano, D. J., Hernandez, A. V., & White, C. M. (2022). MDMA-Assisted Psychotherapy for Treatment of Posttraumatic Stress Disorder : A Systematic Review With Meta-Analysis. *The Journal of Clinical Pharmacology*, *62*(4), 463-471. https://doi.org/10.1002/jcph.1995

Steenkamp, M. M., Litz, B. T., Hoge, C. W., & Marmar, C. R. (2015). Psychotherapy for Military-Related PTSD : A Review of Randomized Clinical Trials. *JAMA*, *314*(5), 489-500. https://doi.org/10.1001/jama.2015.8370

Tang, V. M., Trought, K., Gicas, K. M., Kozak, M., Josselyn, S. A., Daskalakis, Z. J., Blumberger, D. M., Voineskos, D., Knyahnytska, Y., Pasricha, S., Chung, Y., Zhou, Y., Isserles, M., & Wong, A. H. C. (2021). Electroconvulsive therapy with a memory

reactivation intervention for post-traumatic stress disorder : A randomized controlled trial. *Brain Stimulation*, *14*(3), 635-642. https://doi.org/10.1016/j.brs.2021.03.015

Tolentino JC, Schmidt SL. DSM-5 Criteria and Depression Severity : Implications for Clinical Practice. Front Psychiatry. 2018 Oct 2;9:450. Doi : 10.3389/fpsyt.2018.00450. PMID: 30333763; PMCID: PMC6176119. (s. d.).

U.S. Food and Drug Administration. (2020). *SPRAVATO® (esketamine) nasal spray, CIII. Prescribing Information.* https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/211243lbl.pdf

Vaiva, G., Ducrocq, F., Jezequel, K., Averland, B., Lestavel, P., Brunet, A., & Marmar, C. R. (2003). Immediate treatment with propranolol decreases posttraumatic stress disorder two months after trauma. *Biological Psychiatry*, *54*(9), 947-949. https://doi.org/10.1016/S0006-3223(03)00412-8

Vancappel, A., Hingray, C., Reveillere, C., & El-Hage, W. (2023). DISENTANGLING THE LINK BETWEEN MINDFULNESS AND DISSOCIATION IN PTSD : THE MEDIATING ROLE OF ATTENTION AND EMOTIONAL ACCEPTANCE. *Journal of Trauma & Dissociation*, 1-15. https://doi.org/10.1080/15299732.2023.2231907

Watkins, L. E., Sprang, K. R., & Rothbaum, B. O. (2018). Treating PTSD : A Review of Evidence-Based Psychotherapy Interventions. *Frontiers in Behavioral Neuroscience*, *12*, 258. https://doi.org/10.3389/fnbeh.2018.00258

Watts, B. V., Schnurr, P. P., Mayo, L., Young-Xu, Y., Weeks, W. B., & Friedman, M. J. (2013). Meta-Analysis of the Efficacy of Treatments for Posttraumatic Stress Disorder. *The Journal of Clinical Psychiatry*, *74*(06), e541-e550. https://doi.org/10.4088/JCP.12r08225

Wei, M.-D., Wang, Y.-H., Lu, K., Lv, B.-J., Wang, Y., & Chen, W.-Y. (2020). Ketamine reverses the impaired fear memory extinction and accompanied depressive-like behaviors in adolescent mice. *Behavioural Brain Research*, *379*, 112342. https://doi.org/10.1016/j.bbr.2019.112342

Young, M. B., Andero, R., Ressler, K. J., & Howell, L. L. (2015). 3,4-Methylenedioxymethamphetamine facilitates fear extinction learning. *Translational Psychiatry*, *5*(9), e634. https://doi.org/10.1038/tp.2015.138