Prevalence of Asymptomatic non-Falciparum and Falciparum Malaria in the 2014-15 Rwanda Demographic Health Survey

Claudia Gaither[1], Camille Morgan[2], Rebecca Kirby[3], Corine Karema[4], Pierre Gashema[5], Samuel J. White[1], Hillary M. Topazian[6], David Geibrecht[3], Kyaw Thwai[1], Neeva Wernsman Young[3], Koby Boyter[1], Tharcisse Munyaneza[7], Claude Mambo Muvunyi [8], Jean De Dieu Butera[7], Jeffrey A. Bailey[3*#], Jean-Baptiste Mazarati[4,5*#], Jonathan J. Juliano[1,2,9,10*#]

1: Institute for Global Health and Infectious Diseases, University of North Carolina, Chapel Hill, NC, USA
2: Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA
3: Department of Pathology, Brown University, Providence, RI, USA
4: Quality Equity Health Care, Kigali, Rwanda
5: INES-Ruhengeri, Ruhengeri, Rwanda
6. MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, UK
7: National Reference Laboratory, Rwanda Biomedical Center, Kigali, Rwanda
8: Rwanda Biomedical Centre, Kigali, Rwanda
9: Division of Infectious Diseases, School of Medicine, University of North Carolina, Chapel Hill, NC, USA
10: Curriculum in Genetics and Molecular Biology, School of Medicine, University of North Carolina, Chapel Hill, NC, USA

Running title: Asymptomatic malaria in Rwanda

*: Co-senior authors
#: Corresponding authors:
 Jonathan Juliano
 110 Mason Farm Rd.
 CB#7036
 Chapel Hill, NC 27599
 Email: jjuliano@med.unc.edu

 Jean Baptiste Mazarati
 NM 155 Musanze
 INES
 Ruhengeri, Rwanda
 Email: jmazarati@gmail.com

 Jeffrey A Bailey
 55 Claverick Street, Room 312 & 314
 Brown University
 Providence, RI 02903
 Email: jeffrey_bailey@brown.edu

Abstract

Background: Recent molecular surveillance suggests an unexpectedly high prevalence of non-falciparum malaria in Africa. Malaria control is also challenged by undetected asymptomatic *P. falciparum* malaria resulting in an undetectable reservoir for potential transmission. Context-specific surveillance of asymptomatic *P. falciparum* and non-falciparum species is needed to properly inform malaria control programs.

Methods: We performed quantitative real time PCR for four malaria species in 5,050 primarily adult individuals in Rwanda using the 2014-2015 Demographic Health Survey. We assessed correlates of infection by species to explore attributes associated with each species. Asymptomatic *P. ovale spp.*, *P. malariae, and P. falciparum* malaria infection had broad spatial distribution across Rwanda. *P. vivax* infection was rare.

Results: Overall infection prevalence was 22.3% (95%CI [20.3, 24.3]), with *P. falciparum* and non-falciparum at 16.3% [14.5, 18.1] and 8.0% [6.6, 9.3], respectively. Parasitemias tended to be low and mixed species infections were common, especially where malaria transmission and overall prevalence was the highest. *P. falciparum* infection was associated with lower wealth, rural residence and low elevation. Fewer factors were significantly associated with non-falciparum malaria.

Conclusions: Asymptomatic non-falciparum malaria and *P. falciparum* malaria are common and widely distributed across Rwanda in adults. Continued molecular monitoring, preferably done by the national malaria control program, of *Plasmodium* diversity using routine survey samples is needed to strengthen malaria control.

Key words: Plasmodium; falciparum; asymptomatic; ovale; vivax; malariae

Introduction

In 2023 there were an estimated 263 million malaria cases worldwide, with 94% occurring in the WHO African Region. [1] Human malaria infection is caused primarily by five species of the *Plasmodium* parasite (falciparum, malariae, ovale curtisi, ovale walkeri, and vivax), with most cases and morbidity in Africa attributable to *Plasmodium falciparum.* The clinical burden of non-falciparum species in Africa is not well characterized, but all five species have been reported in multiple countries where they have previously not been considered a threat. Falciparum clearly remains the primary cause of morbidity and mortality, but severe disease has been reported with non-falciparum species as well, including anemia, renal failure and other severe outcomes.[2–5] Increase in non-falciparum prevalence has been observed in areas where falciparum prevalence has steadily declined, presenting opportunities for other Plasmodium species to flourish. [6,7]

Diagnosis of malaria in Africa relies principally on rapid diagnostic tests (RDTs) detecting the antigen *P. falciparum* histidine rich protein 2 (HRP2), often with a second less sensitive band for pan-species lactate dehydrogenase (LDH).[8] Designed to detect clinical malaria, RDTs for malaria surveillance will often miss low density *P. falciparum* infections and infections with non-falciparum species. Alternate approaches are needed to accurately define the full asymptomatic reservoir. Molecular detection, such as through real-time PCR, can identify infections at lower parasitemia levels than RDTs can, and better characterize non-falciparum malaria infections. Molecular approaches are particularly important for use in asymptomatic community surveillance.[9] Asymptomatic infections may have significant but underestimated morbidity,[10] and are often followed by symptomatic infection.[11] In addition, they provide a reservoir for ongoing transmission.[12–14] Thus, characterizing the epidemiology of asymptomatic malaria is important for malaria control and prevention, as malaria elimination requires control of all species of malaria, whether symptomatic or asymptomatic.

Molecular identification of non-falciparum malaria infections (*P. malariae, P. ovale curtisi, P. ovale*

wallikeri, and *P. vivax*) is critical for addressing morbidity from these parasites and for reaching malaria

elimination in settings with falciparum-based diagnostics and treatments. Accurate diagnosis is

imperative for malaria control, as the clinical features and treatments differ. In contrast with *P.*

falciparum infection, *P. vivax* and *P. ovale* cause relapse through the persistence of hypnozoites

(dormant liver stage parasites).[15,16] As hypnozoites do not respond to blood-stage treatment, namely

artemisinin-combination therapies (ACTs), the primary treatment for severe malaria in most countries,

relapsing infections require radical cure.[17] Thus, their presence may require national malaria control

programs to alter therapeutic options in the country. To-date, *P. ovale spp.* and *P. malariae* malaria have

been more frequently described, due in part to wider implementation of highly sensitive nucleic acid

detection, and also likely due to increasing prevalence as *P. falciparum* is controlled as was seen for *P.*

vivax in Southeast Asia.[6,7,18] *P. vivax* is also being reported more frequently due to wider

implementation of molecular diagnostics.[4,19,20]

Rwanda has historically had strong malaria control, leveraging effective antimalarials, insecticide-treated

bed nets, and indoor residual spraying.[21] Malaria cases in Rwanda increased from 48 to 403 per 1,000

between 2012 and 2018, with mortality increasing 41% over the same time.[21] Emerging insecticide

resistance, an increase in irrigated agriculture, and insufficient insecticide-treated bed net coverage,

among other factors, have been attributed to these increases.[21] From 2018 to 2022, malaria

prevalence decreased from an estimated 321 to 76 cases per 1,000, still higher than 2012 estimates,[22]

and with detection of non-falciparum species among clinical malaria patients.[23] There has been one

recent report of *P. vivax* commonly occurring in Hue.[23] These inconsistent declines in prevalence

suggest the need to characterize the epidemiology of asymptomatic and non-falciparum infections in

this period to inform future control efforts.

We used survey, GPS, and biospecimen data collected in the 2014-15 Rwanda Demographic and Health

Survey (RDHS), a large nationally population-representative study, to characterize asymptomatic and

non-falciparum infections across the country's adult population.

Methods

Ethics Statement

Dried blood spots (DBS) were provided by the Ministry of Health- Rwanda. This research was reviewed

and approved by the Rwanda National Ethics Committee (reference 102/RNEC/2023). The University of

North Carolina and Brown University IRBs deemed this non-human subjects research.

Study design and population

The 2014-2015 Rwanda DHS studied 12,699 households from 492 GPS-located clusters from all 30

recognized districts. In 50% of households, a total of 16,930 dried blood spot (DBS) specimens were

collected for HIV testing from men aged 15-59 years and women aged 15-49 years, and a subsample of

children 0-14 years; in the other 50% of households, rapid malaria diagnostic testing was completed on

children aged 6-59 months.[24] Previously, we used these data to estimate clusters that would

represent high and low malaria transmission areas.[25] High prevalence clusters had a RDT or

microscopy positivity rate of >15%. We included 1,887 samples from these 56 high prevalence clusters in

3 regions. In addition to samples from high prevalence areas, a random subset of 3,163 samples from

401 low prevalence clusters were selected. A total of 5,050 DBS samples from 457 out of 492 DHS

clusters were analyzed for four species of malaria infection by real time PCR. (**S1 Figure**).

Species-specific real time PCR

DNA from each sample was extracted from three 6 mm DBS punches using Chelex and screened for four

species of malaria infection using real-time PCR assays.[26] These assays (**S1 Table**) targeted the 18s

genes for *P. malariae*, *P. ovale*, and *P. vivax*, and the varATS repeat in *P. falciparum*.[20,27–29] To allow

for quantification of *P. falciparum*, mock DBS were created using whole blood and cultured 3d7

parasites (MRA-102, BEI Resources, Manasas, VA) and extracted with the same assay used for samples.

Controls for non-falciparum species used serial dilutions of plasmid DNA (MRA-180, MRA-179, MRA-178;

BEI Resources, Manassas VA), with estimates for parasitemia based on an estimated six 18s rRNA gene

copies per parasite.[30] All assays were run for 45 cycles to enable detection of lower density infections.

The high cycle number approach has been previously evaluated for *P. ovale* and *P. vivax*, where assays

were tested against 390 negative controls (human DNA) with no false positives.[31] We had no false

positive results in our non-template controls for this study. In addition, we ran each assay to 45 cycles

against plasmids of the other species to evaluate for cross-reactivity (**S2 Table**). All positive samples

were confirmed by manually reviewing the amplification curves in the machine software. Standard

curves had a minimum r-squared value of 0.95 across all runs. A positive result for each species was

determined using a 45-cycle cutoff, unless otherwise stated. In addition, the assays retain a high

specificity at high cycle number.[32]

Spatial & ecological variables

Deidentified survey and geospatial data from the 2014-2015 Rwanda DHS were matched to PCR data

using DBS sample barcodes. Clusters with individuals positive for any species of malaria infection were

mapped using DHS geospatial coordinates. Individual level covariates assessed for association included

sex, age group, wealth quintile, education level, livestock ownership, source of drinking water, bed net

ownership, whether the household bed net has been treated with long-lasting insecticide (LLIN = long-

lasting insecticide-treated net) and sleeping under a LLIN the night prior to the survey. Cluster level

covariates included region, urban/rural status of place of residence, elevation, month of data collection,

proportion of a given cluster living in a household with a bed net, proportion of the cluster that slept

under an LLIN, land cover, average daily maximum temperature for the current month and precipitation

for the prior month. Land cover estimates were taken from the Regional Center for Mapping Resources

for Development and SERVIR-Eastern and Southern Africa, cluster-level temperature values were

obtained from the 2014-2015 Rwanda DHS geospatial covariates, and precipitation values from the

Climate Hazards Group InfraRed Precipitation with Station (CHIRPS) dataset. [33] Survey clusters were

assigned GPS coordinate values within buffers as described previously in accordance with DHS

specifications.[31,34]

Statistical analysis

We estimated species-specific prevalence, non-falciparum prevalence, and overall *Plasmodium sp.*

prevalence, applying HIV sampling weights, inverse propensity for selection weights, and weights to

account for selection by cluster transmission intensity and the skewed selection of samples from low

and high transmission clusters. [35,36] HIV sampling weights are provided in DHS data and account for

potential selection bias resulting from differential DBS collection among survey participants. The DHS

statistics guide recommends inclusion of individual level sampling weights if participants are sampled in

a structured way (over-sampling from high transmission clusters). These three weights were multiplied

for the final weights used for each participant included in the analysis. We estimated bivariate

associations between each *Plasmodium* species and a variety of covariates available in the DHS and

investigated in other contexts,[31,37] including household wealth, livestock ownership, or bed net use,

using the same combination of weights. We report prevalence differences and 95% confidence intervals

to assess precision. Multivariable analysis was not performed as the combination of weights applied

should produce estimates representative of the full survey and country. We analyzed data using the

survey (4.2.1), *srvyr* (v1.2.0), and *sf* (v1.0-8) packages using R 4.2.1 (R Foundation for Statistical

Computing). Shapefiles of Rwanda district boundaries and water bodies were downloaded from the

OCHA Regional Office for Southern and Eastern Africa database, and borders for neighboring countries

were downloaded from ArcGIS Hub.[38] [39] [40]

Results

Study population characteristics

The study population (n = 5,050, weighted n = 5,099) was 51% (2,609/5,099) female, 32% (1,651/5,099)

aged 15-24 years, 76% (3,538/5,099) lived in rural areas, and 75% (3,831/5,099) had a primary school

education or no education (63% (3,191/5,099) reported primary education, 13% (640/5,099) reported

preschool/none). Only 1.5% (77/5,099) of the study population was aged 0-14 years due to limited

sampling of children in the original DHS, so the results of this survey are only representative of Rwandan

adults. Overall, most (82% (4,196/5,099)) individuals reported a household bed net and 62%

(3,173/5,099) reported sleeping under a long-lasting insecticide treated net the night before the survey.

However, 76% (3,880/5,099) of the study population lived in a household that did not meet the World

Health organization's criteria of at least 1 net per 1.8 household members. For each of our covariates of

interest, the study population was mostly comparable to and representative of the overall DHS

population (**S3 Table**).

Prevalence of P. falciparum *and non-falciparum infection by real time PCR*

A total of 1,478 *P. falciparum*, 283 *P. ovale*, 186 *P. malariae* and 7 *P. vivax* infections were identified. The

overall weighted prevalence of any non-falciparum malaria infection was 8.0% (407/5,099, 95% CI: [6.6,

9.3]) compared to 16.3% (831/5,099 [14.5, 18.1]) for *P. falciparum* and 22.3% (1,137/5,099 [20.3, 24.3])

overall malaria prevalence with HIV sampling, inverse propensity of selection, and transmission intensity

correction weights applied. Species specific weighted prevalences were 3.4% (172/5,099) [2.6, 4.1] and

4.7% (241/5,099) [3.5, 5.9] for *P. malariae* and *P. ovale spp.* Unweighted prevalence for *P. vivax* was

0.14%, (7/5,050) with a manually calculated 95% CI [0.03, 0.25]. Study population characteristics and the

distribution of PCR positives by species are shown in **Table 1**. Unweighted prevalence estimates were

calculated for each species based on false-positive rates of high cycle number PCR (**S4 Table**). Using a

more restrictive cut off of 40 cycles (requiring a higher parasitemia to be positive at approximately 1

parasite per microliter of template DNA), resulted in weighted overall prevalences of 4.3% (218/5,099)

[3.5, 5.1] for any non-falciparum infection, compared to 13.2% (674/5,099) [11.6, 14.8] for *P. falciparum*

and 16.4% (836/5,099) [14.7, 18.1] for overall malaria. Species specific weighted prevalences at this cut

off were 2.7% (139/5,099) [2.0, 3.4] and 1.6% (80/5,099) [1.1, 2.0], and for *P. malariae*, and *P. ovale spp.*

Unweighted prevalence for *P. vivax* was 0.06% (3/5,050 [0.00, 0.13]. The largest difference in estimated

prevalence was for *P. ovale spp.* This is not surprising given the distribution of estimated parasitemia

values (**Figure 1**), showing a lower median parasitemia in the non-falciparum species compared to *P.*

falciparum. District level weighted prevalences and their differences by PCR cut off are shown in **S5**

Table and **S6 Table**. *P. falciparum*, *P. ovale* and *P. malariae* infections were distributed across the

country, while *P. vivax* infections were more localized (**S2 Figure**). District level overall malaria

prevalence is shown in **Figure 2**, while district level prevalences for each species are illustrated in **Figure**

3. Among *P. ovale spp. P. malariae,* and *P. vivax* infections, 48.4%, (137/283) [42.5, 54.4]), 46.8%

(87/186) [39.5,54.2]), and 57% (4/7) [20.2, 88.2], (unweighted counts) were infected with at least one

other species of malaria (**S7 Table**).

Bivariate associations for infection

Bivariate regression models using weighted (as previously described) survey data found multiple

associations for infection with *P. falciparum* malaria, but few for non-falciparum malaria (**Figure 4**).

Covariates with more than two categories were dichotomized to provide interpretable comparisons

across the study population. For example, Anopheles mosquito vectors generally do not thrive at

elevations over 1,500 meters, although this is not a fixed limit. [41] Land cover classifications were

dichotomized to compare less vegetation/more human activity to more vegetation/less human activity.

No adjusted analyses were conducted as we are only exploring the strength of individual associations

based on prevalence measures, and generalizability of participants analyzed to the country or survey as

a whole is provided through weighting. Similar to previous work, a higher prevalence of *P. falciparum*

malaria in our dataset was significantly (at a 0.05 confidence level for all associations) associated with

multiple study covariates related to socioeconomic status (lower wealth quintile, lower education

status, and unpiped drinking water). Secondary or higher education, residence in a household with at

least 1 bed net per 1.8 household members, piped drinking water, continuous increase (and clusters at

or above the study sample average) in prior month's average rainfall, and higher altitude (>1,500

meters) were all associated with significantly lower prevalence of *P. falciparum* malaria. Rural clusters

compared to urban, clusters designated as perennial cropland or grassland (compared to woodland or

sparse forest), lower wealth index (first and second quintiles compared to the upper three), younger age

(15-24 years compared to over 24 years) and residence in clusters with monthly average temperatures

at or above the average for all clusters in analysis were associated with higher *P. falciparum* prevalence.

Fewer associations were found for non-falciparum malaria. No significant associations were found for P.

malariae infection at an alpha level of 0.05. *P. ovale* infection was significantly associated with residence

in clusters at or above the average monthly temperature, while lower prevalence of infection was

associated with younger age in both the binary categorization described above and continuous decrease

in age. Associations for *P. vivax* infection were not attempted due to the limited number of infections in

the survey (n=7).

Discussion

In Rwanda, a setting with robust malaria control efforts but inconsistently declining prevalence, we conducted the largest assessment of asymptomatic and non-falciparum malaria to-date, primarily among adults. We found an asymptomatic malaria weighted prevalence of 22.3%, using PCR, substantially higher than estimates produced using RDT and microscopy. Additionally, we observed 8.0% of the population was infected with a non-falciparum species. While we observed *P. falciparum* prevalence to be 16.3% in adults, malaria detection by microscopy in the DHS reported a 2% malaria prevalence among children age 6-59 months and 0.6% among women age 15-49.[42] This high prevalence in adults found using molecular detection is consistent with a study of school-aged individuals in the Huye District in the same year, finding a 22% prevalence of any malaria (19% *P. falciparum* prevalence) using a combined microscopy and PCR approach.[43]

We used an ultra-sensitive assay for *P. falciparum* to detect primarily asymptomatic, lower density infections, improving the characterization of *P. falciparum* prevalence. While a higher PCR cycle cut-off may cause concern about false-positive detection, our assays have been run extensively at 45 cycles with little to no evidence of false positives[31], and are consistent with estimates using a lower cut off (40 cycles). The largest change in prevalence occurred with *P. ovale spp.*, where estimated prevalence dropped from 4.7% to 1.6% at different cycle cut offs, reflecting the high number of low density infections detected. The relative decrease in prevalence for each species based on cycle cutoff was not always consistent, with some regions having no decline in *P. falciparum* prevalence with large declines in *P. ovale* malaria (e.g. Karongi near Lake Kivu) or the opposite with no decline in non-falciparum but lower *P. falciparum* prevalence estimates (e.g. Rutsiro and Nyamasheke). With asymptomatic *P. falciparum* malaria increasing risk of symptomatic disease at one month and contributing to significant

morbidity, the high observed prevalence of asymptomatic, low-parasitemia infections offers an

important direction for malaria control.[10,11] Estimating prevalence of sub-patent malaria should be

prioritized as a complement to RDT or microscopy detected prevalence to accurately characterize

transmissible reservoirs in endemic regions.

Non-falciparum malaria was detected in 8.0% of individuals nationally, a prevalence not previously

appreciated in the country. *P. ovale spp.* and *P. malariae* were both common in Rwanda (4.7% and 3.3%

prevalence, respectively) and distributed in regions of both high and low transmission. A recent

household survey estimated that *P. falciparum* is responsible for 97% of malaria infections in Rwanda,

with *P. malariae* and *P. ovale spp.* each responsible for 1%–2% of total infections (Uwimana A.

{Unpublished]).[44] *P. vivax* is present but remains relatively uncommon (0.14% unweighted prevalence)

and sporadic, but consistent with reporting of occasional clinical cases. [44] More recently, a cluster of *P.*

vivax was reported in the Huye District.[23] Given distinct treatment approaches by species, the higher

than expected prevalence of *P. ovale spp.* and occasional *P. vivax* infection underscores the need for

molecular monitoring to guide control efforts towards elimination. Regular surveillance for non-

falciparum malaria is needed to inform regional treatment guidelines, particularly if relapsing malaria

species continue to expand their prevalence.

Not surprisingly, mixed species infections were common and widely distributed, but occurred more

commonly in clusters in the south and east where malaria transmission was the highest (**S2 Figure**).

Among *P. ovale spp. P. malariae,* and *P. vivax* infections, 44%, 45% and 57% (unweighted) were infected

with at least one other species of malaria (**S7 Table**). Bivariate associations for mixed infections largely

reflected those observed for *P. falciparum,* albeit with lower magnitude and wider confidence intervals

attributable to the smaller total counts. Mixed infections are often underappreciated by clinicians and

laboratorians and may lead to severe disease complications. A recent meta-analysis suggested that

patients with mixed infections have a higher proportion of pulmonary complications and multiple organ

failure than patients with *P. falciparum* infection alone.[45] The impact of mixed species infections on

clinical malaria outcomes in Rwanda is unknown and requires additional evaluation in symptomatic

infections, which were not included in this study.

The associations we observed between *P. falciparum* infection and lower household wealth, no bed net

use and lower elevation are consistent with previous studies using these data, with malaria positivity

defined by RDT or microscopy.[46,47] We observed malaria prevalence was highest in the South and

East as expected, but asymptomatic infection remained common in other areas. Like similar studies,

associations of covariates for non-falciparum malaria were few and traditional risk factors for *P.*

falciparum were less strongly associated with non-falciparum malaria.[20,28,31,37] This raises concern

for how the control program can target non-falciparum infections without better diagnostics in the

community. The reasons for the relative lack of risk factors, especially for *P. ovale spp.*, remains unclear.

Relapsing malaria, caused by *P. ovale* and *P. vivax,* may not be associated with typical covariates due to

the inability to discern between incident or relapse infections in the study. Additionally, non-falciparum

infections are under-detected in this sample compared to falciparum due to the ultra-sensitive assay

used for falciparum detection, which may bias estimates and increase uncertainty for associations with

non-falciparum infections.

While this data reflects the epidemiology from nearly a decade ago, it remains important for malaria

control. The baseline set here is useful for understanding the interventions that have been used in

Rwanda for malaria control if future similar surveys, such as the 2019/20 DHS, Malaria Indicator Surveys,

or other broad sampling efforts, are genotyped similarly. The longitudinal data across surveys can be

used to understand how malaria interventions impact both asymptomatic malaria and non-falciparum

malaria, both of which are not captured by standard DHS data but still important for malaria

elimination.

While this national molecular evaluation offers important perspectives, we are limited by a few key

points. As noted above, the cross-sectional nature of a single DHS limits inference regarding

transmission and time trends, but use of subsequent (2019-20 DHS) and future DHSs, would allow for

ongoing surveillance of these parasites, as has been done in the DRC.[48] We also could not determine

if *P. ovale* spp. and *P. vivax* infections were newly acquired or the result of relapse from hypnozoites.

Detection of these parasites is still important for malaria control programming. Additionally, the relative

lack of data for children under 15 makes this survey not representative of the full asymptomatic

reservoir and non-falciparum prevalence. We are limited by the sampling framework of the DHS. School

aged children are a particularly vulnerable group and often have the highest rate of malaria infection in

Africa, including for *P. vivax*.[49] Without this group, a true population prevalence of asymptomatic

infection is difficult to determine. However, we likely underestimate overall population prevalence given

the 2017 MIS had an overall prevalence of 14% for children aged 6 months to 14 years of age using a

rapid diagnostic test which typically detects 41% of infections compared to PCR.[9,50] In addition, males

were overrepresented in the sample. We used an assay that targets 18S ribosomal RNA genes for non-

falciparum species, thus could underreport infections that could have been detected with assays that

target genes with higher copy numbers in the parasite such as the assay used for falciparum in this

study. [51–53] Despite these limitations, the use of existing samples and individual level data from a DHS

is a highly informative method to gain insights into national malaria prevalence.

This study represents the first nationwide molecular investigation of asymptomatic *Plasmodium*

falciparum and non-falciparum malaria in Rwanda and highlights a significant parasite reservoir and the

essential role of molecular testing to appropriately detect them. First, we found that asymptomatic *P.*

falciparum infections in adults were significantly more prevalent than previously estimated by rapid

diagnostic tests (RDTs) and microscopy in children.[42] These infections are not benign: they carry

clinical consequences and serve as a hidden reservoir for continued transmission, underscoring the

importance of identifying and addressing asymptomatic carriers in control strategies.[10,12] Second,

molecular methods were necessary to detect *P. ovale spp.*, *P. malariae*, and *P. vivax*, which are better

characterized by molecular methods as they can be frequently missed by standard diagnostics. In this

study, most non-falciparum infections were low-density and often co-occurred with *P. falciparum*,

particularly *P. ovale spp.* and *P. vivax*, which would contribute to under-recognition of mixed infections.

The observed prevalence of *P. ovale spp.* and *P. malariae* was higher than expected and lacked

consistent risk factors. In order to assess if these species may be expanding into niches left by successful

P. falciparum control, continued monitoring with molecular methods is needed. Third, comprehensive

malaria elimination efforts must address all human malaria species—not just *P. falciparum*—including

those capable of relapse and silent transmission. Therefore, ongoing molecular surveillance, ideally led

by the national malaria control program, is critical to track changing epidemiology, guide targeted

interventions, and support Rwanda's path toward malaria elimination.

Funding: This work was funded by the National Institutes for Health (R01AI156267 to JAB, JBM and JJJ

and K24AI134990 to JJJ). The funding for the DHS was provided by the government of Rwanda, the

United States Agency for International Development (USAID), the One United Nations (One UN), the

Global Fund to Fight AIDS, Tuberculosis and Malaria (Global Fund), World Vision International, the Swiss

Agency for Development and Cooperation (SDC), and the Partners in Health (PIH). ICF International

provided technical assistance through The DHS Program, a USAID-funded project providing support and

technical assistance in the implementation of population and health surveys in countries worldwide. The

funders of the study had no role in study design, data collection, data analysis, data interpretation, or

writing of the report.

Author Contributions: CK, CM, JAB, JBM and JJJ conceived of the study. CG, CM, RK, SW, HMT, DG, KT,

and KB were involved in data generation and analysis. All authors were involved in writing the

manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Data Availability: Clinical and spatial data are available through the DHS MEASURE website

(https://dhsprogram.com/data/available-datasets.cfm). PCR data has been provided to DHS and is

available upon request with approval of use of DHS data from DHS MEASURE. Code used for analysis is

available at: https://github.com/claudiagaither.

Acknowledgements: The following reagent was obtained through BEI Resources, NIAID, NIH:

Plasmodium falciparum, Strain 3D7, MRA-102, contributed by Daniel J. Carucci. The following reagent

was obtained through BEI Resources, NIAID, NIH: Diagnostic Plasmid Containing the Small Subunit

Ribosomal RNA Gene (18S) from *Plasmodium vivax*, MRA-178, contributed by Peter A. Zimmerman. The

following reagent was obtained through BEI Resources, NIAID, NIH: Diagnostic Plasmid Containing the

Small Subunit Ribosomal RNA Gene (18S) from *Plasmodium malariae*, MRA-179, contributed by Peter A.

Zimmerman. The following reagent was obtained through BEI Resources, NIAID, NIH: Diagnostic Plasmid

Containing the Small Subunit Ribosomal RNA Gene (18S) from *Plasmodium ovale*, MRA-180, contributed

by Peter A. Zimmerman.

Table 1. Study Population Characteristics and Distribution of PCR Positives by Species. Weighted counts and percentages for all individuals positive for each species of malaria. Weights applied are DHS sampling weights, propensity score weights, and weights to correct for over-selection of high transmission intensity clusters (transmission intensity weights). Characteristics of the included study population are also shown.

		PCR positive (falciparum)		PCR negative (falciparum)		PCR positive (malariae)		PCR negative (malariae)		PCR positive (ovale)		PCR negative (ovale)		PCR positive (vivax)		PCR negative (vivax)		total
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n
	Unweighted counts	1468	29.07	3582	70.93	173	3.43	4877	96.57	283	5.60	4767	94.40	7	0.14	5043	99.86%	5050
	Weighted counts	831	16.30	4268	83.70	172	3.37	4927	96.63	241	4.73	4858	95.27	7.5	0.15	5092	99.85%	5099
Individual level covariates																		
Sex of respondent	Male	419.9	51%	2070.5	49%	93.7	54%	2396.7	49%	125.7	52%	2364.6	49%	5.7	76%	2484.7	49%	2490.4
	Female	411.3	49%	2197.8	51%	78.5	46%	2530.5	51%	114.8	48%	2494.2	51%	1.8	24%	2607.2	51%	2609.0
Age group (years)	0-15	14.2	2%	63.2	1%	2.2	1%	75.1	2%	0.7	0%	76.6	2%	0.0	0%	77.3	2%	77.3
	15-24	217.0	26%	506.8	12%	32.5	19%	691.4	14%	37.1	15%	686.7	14%	0.0	0%	723.9	14%	723.9
	25-34	230.7	28%	1419.9	33%	41.9	24%	1608.7	33%	97.8	41%	1552.8	32%	2.2	29%	1648.4	32%	1650.6
	35-44	198.0	24%	1251.5	29%	50.4	29%	1399.0	28%	59.6	25%	1389.9	29%	2.5	34%	1446.9	28%	1449.4
	45-54	117.2	14%	686.9	17%	27.8	16%	776.3	17%	33.5	17%	770.7	17%	0.0	0%	804.1	17%	804.1
	55+	54.0	7%	340.0	8%	17.4	10%	376.6	8%	11.9	5%	382.1	8%	2.8	37%	391.2	8%	394.0
Wealth Quintile	Poorest	196.3	24%	547.5	13%	23.1	13%	720.7	15%	34.6	14%	709.2	15%	2.3	31%	741.4	15%	743.8
	Poorer	160.0	19%	737.2	17%	34.4	20%	862.8	18%	31.4	13%	865.8	18%	0.2	2%	897.0	18%	897.2
	Middle	169.7	20%	747.2	18%	37.8	22%	879.0	18%	35.0	15%	881.8	18%	0.0	0%	916.8	18%	916.8
	Richer	155.3	19%	947.8	22%	35.3	21%	1067.7	22%	54.4	23%	1048.7	22%	4.9	66%	1098.1	22%	1103.1
	Richest	149.9	18%	1288.6	30%	41.6	24%	1397.0	28%	85.2	35%	1353.3	28%	0.0	0%	1438.5	28%	1438.5
Education	None/preschool	143.5	17%	496.6	12%	22.5	13%	617.7	13%	19.7	8%	620.5	13%	2.3	31%	637.8	13%	640.1
	Primary	553.2	67%	2637.8	62%	110.4	64%	3080.6	63%	148.8	62%	3042.2	63%	3.5	47%	3187.5	63%	3191.0

	Secondary	113.5	14%	960.0	22%	32.4	19%	1041.1	21%	65.3	27%	1008.1	21%	1.6	22%	1071.8	21%	1073.4
	Higher	20.5	2%	171.3	4%	7.0	4%	184.8	4%	6.7	3%	185.1	4%	0.0	0%	191.8	4%	191.8
	Missing data	0.5	0%	2.5	0%	0.0	0%	3.0	0%	0.0	0%	3.0	0%	0.0	0%	3.0	0%	3.0
Owns livestock, herds, or farm animals	No	349.0	42%	1850.3	43%	79.3	46%	2120.0	43%	101.4	42%	2097.8	43%	4.7	63%	2194.5	43%	2199.2
	Yes	482.2	58%	2417.9	57%	92.9	54%	2807.2	57%	139.1	58%	2761.0	57%	2.8	37%	2897.4	57%	2900.1
Source of drinking water	Unpiped	770.5	93%	3550.1	83%	155.4	90%	4165.2	85%	193.0	80%	4127.6	85%	0.0	0%	4320.6	85%	4320.6
	Piped	60.7	7%	718.1	17%	16.8	10%	762.0	15%	47.5	20%	731.2	15%	7.5	100%	771.3	15%	778.8
Household bed net	No	130.5	16%	773.1	18%	36.2	21%	867.4	18%	34.4	14%	869.2	18%	2.9	39%	900.7	18%	903.7
	Yes	700.6	84%	3495.1	82%	136.0	79%	4059.7	82%	206.1	86%	3989.6	82%	4.5	61%	4191.2	82%	4195.7
Slept under LLIN last night	No	334.5	40%	1591.9	37%	79.0	46%	1847.4	37%	79.3	33%	1847.1	38%	2.9	39%	1923.5	38%	1926.4
	Yes	496.7	60%	2676.3	63%	93.2	54%	3079.8	63%	161.2	67%	3011.7	62%	4.5	61%	3168.4	62%	3172.9
Insecticide-treated household net	No	1.5	0%	1.5	0%	0.1	0%	3.0	0%	0.2	0%	2.9	0%	0.0	0%	3.1	0%	3.1
	Yes	496.7	60%	2678.3	63%	93.2	54%	3081.8	63%	161.2	67%	3013.7	62%	4.5	61%	3170.4	62%	3174.9
	Missing data	332.9	40%	1588.4	37%	78.3	46%	1843.0	37%	79.1	33%	1842.2	38%	2.9	39%	1918.4	38%	1921.3
1 bed net per 1.8 household members	No	685.2	83%	3184.3	75%	135.4	80%	3734.1	76%	171.2	71%	3698.4	76%	7.5	100%	3862.1	76%	3869.6
	Yes	145.1	17%	1074.2	25%	34.9	20%	1184.4	24%	69.3	29%	1149.9	24%	0.0	0%	1219.2	24%	1219.2
Cluster level covariates																		
Region	Kigali City	94.1	11%	740.3	17%	16.9	10%	817.5	17%	40.1	17%	794.3	16%	1.6	22%	832.8	16%	834.4
	South	324.3	39%	792.3	19%	33.3	19%	1083.4	22%	76.1	32%	1040.6	21%	1.1	15%	1115.5	22%	1116.7
	West	95.1	11%	1204.6	28%	46.2	27%	1253.6	25%	39.2	16%	1260.6	26%	3.2	43%	1296.5	25%	1299.8

	North	58.8	7%	1007.3	24%	42.2	25%	1023.8	21%	22.9	10%	1043.2	21%	0.0	0%	1066.1	21%	1066.1
	East	258.8	31%	523.6	12%	33.6	20%	748.8	15%	62.2	26%	720.2	15%	1.5	20%	780.9	15%	782.4
Place of residence	Urban	111.0	13%	1128.3	26%	35.1	20%	1204.2	24%	60.0	25%	1179.3	24%	1.6	22%	1237.7	24%	1239.3
	Rural	720.1	87%	3139.9	74%	137.1	80%	3722.9	76%	180.5	75%	3679.6	76%	5.8	78%	3854.2	76%	3860.0
Elevation (m)	500 - 1000	3.6	0%	13.9	0%	0.6	0%	16.9	0%	0.0	0%	17.5	0%	0.0	0%	17.5	0%	17.5
	1001 - 1500	339.8	41%	894.7	21%	46.9	27%	1187.5	24%	78.3	33%	1156.1	24%	3.1	41%	1231.3	24%	1234.4
	1501 - 2000	426.8	51%	2204.6	52%	72.3	42%	2559.1	52%	119.4	50%	2512.0	52%	1.1	15%	2630.2	52%	2631.4
	2000 - 2500	55.7	7%	1052.6	25%	50.1	29%	1058.2	21%	42.8	18%	1065.4	22%	1.6	22%	1106.7	22%	1108.3
	> 2500	5.4	1%	102.4	2%	2.3	1%	105.5	2%	0.0	0%	107.8	2%	1.6	22%	106.2	2%	107.8
Month of data collection	Jan-15	191.4	23%	1185.8	28%	37.7	22%	1339.5	27%	56.5	23%	1320.6	27%	0.0	0%	1377.2	27%	1377.2
	Feb-15	196.2	24%	693.6	16%	27.5	16%	862.2	17%	44.1	18%	845.7	17%	2.2	29%	887.6	17%	889.8
	Mar-15	197.7	24%	833.5	20%	40.0	23%	991.1	20%	69.5	29%	961.6	20%	2.5	34%	1028.6	20%	1031.1
	Apr-15	6.1	1%	37.3	1%	2.5	1%	40.9	1%	2.1	1%	41.3	1%	0.0	0%	43.4	1%	43.4
	Nov-14	111.0	11%	484.3	9%	26.7	12%	568.6	9%	17.7	6%	577.6	10%	1.1	11%	594.1	9%	595.3
	Dec-14	128.8	15%	1033.8	24%	37.8	22%	1124.8	23%	50.6	21%	1112.0	23%	1.6	22%	1161.0	23%	1162.6
Landcover	Moderate forest	32.7	0.04	283.0	7%	8.4	5%	307.2	6%	21.4	9%	294.2	6%	0.0	0%	315.6	6%	315.6
	Sparse forest	11.7	0.01	32.6	1%	3.0	2%	41.2	1%	1.1	0%	43.2	1%	0.0	0%	44.2	1%	44.2
	Woodland	205.9	0.25	614.7	14%	36.7	21%	783.9	16%	35.6	15%	785.0	16%	0.9	12%	819.7	16%	820.6
	Closed grassland	538.5	0.65	2895.2	68%	113.0	66%	3320.7	67%	167.6	70%	3266.2	67%	6.6	88%	3427.2	67%	3433.7
	Perennial cropland	42.5	0.05	442.7	10%	11.1	6%	474.1	10%	14.9	6%	470.3	10%	0.0	0%	485.2	10%	485.2

404
405
406
407

408 **Figure 1. Calculated Parasitemia Estimates for Falciparum and Non-falciparum Infections.** Overall, falciparum had a higher parasite density with
409 a median of 10.90 (IQR 0.96-101.41). The median parasitemia level for *P. malariae*, *P. ovale spp.,* and *P. vivax* malaria were 0.35 (IQR: 0.07-
410 1.83), 0.48 (IQR: 0.16-3.25) and 0.94 (IQR: 0.20-3.44), respectively.

411
412
413

414 **Figure 2. District Level Weighted Overall Malaria Prevalence.** Weighted prevalence of any malaria infection, using HIV sampling, inverse

415 propensity for selection and transmission intensity weights (described above). Shapefiles for country borders were downloaded from ArcGIS

416 Hub, district boundaries and the lake shapefile were downloaded from the OCHA Regional Office for Southern and Eastern Africa database.

Figure 3. District Level Weighted Prevalence Estimates for Malaria Species. The weighted prevalence estimate for each species is shown. Panel A, B, C and D represent *P. falciparum, P. malariae, P. ovale spp*. and *P. vivax* malaria, respectively. Shapefiles for country borders were downloaded from ArcGIS Hub, district boundaries and the lake shapefile were downloaded from the OCHA Regional Office for Southern and Eastern Africa database.

428 **Figure 4. Bivariate associations and between demographic and environmental risk factors and *Plasmodium* spp. prevalence using weighted**

429 **survey data.** Models incorporate 2014-15 Rwanda Demographic and Health Survey weights, inverse probability of selection weights, and cluster

430 transmission intensity weights (described in **Table 1**). Point estimates of prevalence difference are surrounded by confidence intervals. The

431 reference is the second variable listed. Panel A, B and C represent *P. falciparum*, *P. malariae,* and *P. ovale sp.*, respectively. Note that each panel

432 has a different scale.

References

1.　World malaria report 2024. [cited 31 Mar 2025]. Available: https://www.who.int/teams/global-
malaria-programme/reports/world-malaria-report-2024

2.　Langford S, Douglas NM, Lampah DA, Simpson JA, Kenangalem E, Sugiarto P, et al. Plasmodium
malariae Infection Associated with a High Burden of Anemia: A Hospital-Based Surveillance Study.
PLoS Negl Trop Dis. 2015;9: e0004195.

3.　Mueller I, Zimmerman PA, Reeder JC. Plasmodium malariae and Plasmodium ovale--the "bashful"
malaria parasites. Trends Parasitol. 2007;23: 278–283.

4.　Twohig KA, Pfeffer DA, Baird JK, Price RN, Zimmerman PA, Hay SI, et al. Growing evidence of
Plasmodium vivax across malaria-endemic Africa. PLoS Negl Trop Dis. 2019;13: e0007140.

5.　Groger M, Fischer HS, Veletzky L, Lalremruata A, Ramharter M. A systematic review of the clinical
presentation, treatment and relapse characteristics of human Plasmodium ovale malaria. Malar J.
2017;16: 112.

6.　Yman V, Wandell G, Mutemi DD, Miglar A, Asghar M, Hammar U, et al. Persistent transmission of
Plasmodium malariae and Plasmodium ovale species in an area of declining Plasmodium falciparum
transmission in eastern Tanzania. PLoS Negl Trop Dis. 2019;13: e0007414.

7.　Graumans W, Ayo D, van Lieshout N, Lanke K, Bousema T, Arinaitwe E. Plasmodium malariae and
Plasmodium ovale-Prevalent and Relevant. J Infect Dis. 2024;229: 1239–1240.

8.　Boyce MR, O'Meara WP. Use of malaria RDTs in various health contexts across sub-Saharan Africa:
a systematic review. BMC Public Health. 2017;17: 470.

9.　Wu L, van den Hoogen LL, Slater H, Walker PGT, Ghani AC, Drakeley CJ, et al. Comparison of
diagnostics for the detection of asymptomatic Plasmodium falciparum infections to inform control
and elimination strategies. Nature. 2015;528: S86–S93.

10.　Chen I, Clarke SE, Gosling R, Hamainza B, Killeen G, Magill A, et al. "Asymptomatic" Malaria: A
Chronic and Debilitating Infection That Should Be Treated. PLoS Med. 2016;13.
doi:10.1371/journal.pmed.1001942

11.　Sumner KM, Mangeni JN, Obala AA, Freedman E, Abel L, Meshnick SR, et al. Impact of
asymptomatic Plasmodium falciparum infection on the risk of subsequent symptomatic malaria in a
longitudinal cohort in Kenya. 2021 [cited 15 Sep 2023]. doi:10.7554/eLife.68812

12.　Lindblade KA, Steinhardt L, Samuels A, Kachur SP, Slutsker L. The silent threat: asymptomatic
parasitemia and malaria transmission. Expert Rev Anti Infect Ther. 2013;11: 623–639.

13.　Rek J, Blanken SL, Okoth J, Ayo D, Onyige I, Musasizi E, et al. Asymptomatic School-Aged Children
Are Important Drivers of Malaria Transmission in a High Endemicity Setting in Uganda. J Infect Dis.
2022;226: 708–713.

14.　Biruksew A, Demeke A, Birhanu Z, Golassa L, Getnet M, Yewhalaw D. Schoolchildren with
asymptomatic malaria are potential hotspot for malaria reservoir in Ethiopia: implications for

malaria control and elimination efforts. Malar J. 2023;22: 311.

15.   Richter J, Franken G, Mehlhorn H, Labisch A, Häussinger D. What is the evidence for the existence
of Plasmodium ovale hypnozoites? Parasitol Res. 2010;107: 1285–1290.

16.   Mueller I, Galinski MR, Baird JK, Carlton JM, Kochar DK, Alonso PL, et al. Key gaps in the knowledge
of Plasmodium vivax, a neglected human malaria parasite. Lancet Infect Dis. 2009;9: 555–566.

17.   Wells TNC, Burrows JN, Baird JK. Targeting the hypnozoite reservoir of Plasmodium vivax: the
hidden obstacle to malaria elimination. Trends Parasitol. 2010;26: 145–151.

18.   Price RN, Commons RJ, Battle KE, Thriemer K, Mendis K. Plasmodium vivax in the Era of the
Shrinking P. falciparum Map. Trends Parasitol. 2020;36: 560–570.

19.   Mapping the global endemicity and clinical burden of Plasmodium vivax, 2000–17: a spatial and
temporal modelling study. Lancet. 2019;394: 332–343.

20.   Brazeau NF, Mitchell CL, Morgan AP, Deutsch-Feldman M, Watson OJ, Thwai KL, et al. The
epidemiology of Plasmodium vivax among adults in the Democratic Republic of the Congo. Nat
Commun. 2021;12: 4169.

21.   Karema C, Wen S, Sidibe A, Smith JL, Gosling R, Hakizimana E, et al. History of malaria control in
Rwanda: implications for future elimination in Rwanda and other malaria-endemic countries. Malar
J. 2020;19. doi:10.1186/s12936-020-03407-1

22.   [No title]. [cited 29 Apr 2024]. Available:
https://www.rbc.gov.rw/fileadmin/user_upload/report23/Rwanda%20Aide_Memoire_MTR_9%20
_March_2023_Final.pdf

23.   van Loon W, Oliveira R, Bergmann C, Habarugira F, Tacoli C, Jäger J, et al. Plasmodium vivax Malaria
in Duffy-Positive Patients in Rwanda. Am J Trop Med Hyg. 2023;109. doi:10.4269/ajtmh.23-0143

24.   Demographic ICF. health surveys standard recode manual for dhs7. The Demographic and Health
Surveys Program.

25.   Kirby R, Giesbrecht D, Karema C, Watson O, Lewis S, Munyaneza T, et al. Examining the Early
Distribution of the Artemisinin-Resistant R561H Mutation in Areas of Higher Transmission in
Rwanda. Open Forum Infect Dis. 2023;10: ofad149.

26.   Walsh PS, Metzger DA, Higushi R. Chelex 100 as a medium for simple extraction of DNA for PCR-
based typing from forensic material. BioTechniques 10(4): 506-13 (April 1991). Biotechniques.
2013;54: 134–139.

27.   Rougemont M, Van Saanen M, Sahli R, Hinrikson HP, Bille J, Jaton K. Detection of four Plasmodium
species in blood from humans by 18S rRNA gene subunit-based and species-specific real-time PCR
assays. J Clin Microbiol. 2004;42: 5636–5643.

28.   Mitchell CL, Brazeau NF, Keeler C, Mwandagalirwa MK, Tshefu AK, Juliano JJ, et al. Under the Radar:
Epidemiology of Plasmodium ovale in the Democratic Republic of the Congo. J Infect Dis. 2021;223.
doi:10.1093/infdis/jiaa478

29. Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-sensitive detection of
Plasmodium falciparum by amplification of multi-copy subtelomeric targets. PLoS Med. 2015;12:
e1001788.

30. Mercereau-Puijalon O, Barale J-C, Bischoff E. Three multigene families in Plasmodium parasites:
facts and questions. Int J Parasitol. 2002;32: 1323–1344.

31. Gumbo A, Topazian HM, Mwanza A, Mitchell CL, Puerto-Meredith S, Njiko R, et al. Occurrence and
Distribution of Nonfalciparum Malaria Parasite Species Among Adolescents and Adults in Malawi. J
Infect Dis. 2022;225. doi:10.1093/infdis/jiab353

32. Sendor R, Banek K, Kashamuka MM, Mvuama N, Bala JA, Nkalani M, et al. Epidemiology of
Plasmodium malariae and Plasmodium ovale spp. in Kinshasa Province, Democratic Republic of
Congo. Nat Commun. 2023;14: 6618.

33. de Sousa K, Sparks A, Ashmall W, van Etten J, Solberg S. Chirps: API client for the CHIRPS
precipitation data in R. J Open Source Softw. 2020;5: 2419.

34. [No title]. [cited 2 Nov 2023]. Available:
https://spatialdata.dhsprogram.com/references/DHS%20Covariates%20Extract%20Data%20Descri
ption%202.pdf

35. Guide to DHS Statistics. [cited 2 Nov 2023]. Available: https://dhsprogram.com/data/Guide-to-DHS-
Statistics/index.cfm

36. Rosenbaum PR, Rubin DB. The Central Role of the Propensity Score in Observational Studies for
Causal Effects. 1982.

37. Mitchell CL, Topazian HM, Brazeau NF, Deutsch-Feldman M, Muwonga J, Sompwe E, et al.
Household Prevalence of Plasmodium falciparum, Plasmodium vivax, and Plasmodium ovale in the
Democratic Republic of the Congo, 2013-2014. Clin Infect Dis. 2021;73: e3966–e3969.

38. Rwanda - Subnational Administrative Boundaries. [cited 15 Sep 2023]. Available:
https://data.humdata.org/dataset/cod-ab-rwa?

39. Africa Countries. [cited 14 Apr 2025]. Available: https://hub.arcgis.com/datasets/africa::africa-
countries/about

40. Rwanda - Water Bodies (Lakes). [cited 14 Apr 2025]. Available:
https://data.humdata.org/dataset/rwanda-water-bodies

41. Carlson CJ, Bannon E, Mendenhall E, Newfield T, Bansal S. Rapid range shifts in African Anopheles
mosquitoes over the last century. Biol Lett. 2023;19: 20220365.

42. [No title]. [cited 15 Sep 2023]. Available: https://dhsprogram.com/pubs/pdf/FR316/FR316.pdf

43. Sifft KC, Geus D, Mukampunga C, Mugisha JC, Habarugira F, Fraundorfer K, et al. Asymptomatic only
at first sight: malaria infection among schoolchildren in highland Rwanda. Malar J. 2016;15.
doi:10.1186/s12936-016-1606-x

44. McCaffery JN, Munyaneza T, Uwimana A, Nace D, Lucchi N, Halsey ES, et al. Symptomatic Infection
in Rwanda. Open Forum Infect Dis. 2022;9: ofac025.

45. Kotepui M, Kotepui KU, De Jesus Milanez G, Masangkay FR. Plasmodium spp. mixed infection
leading to severe malaria: a systematic review and meta-analysis. Sci Rep. 2020;10: 11068.

46. Kubana E, Munyaneza A, Sande S, Nduhuye F, Karangwa JB, Mwesigye D, et al. "A comparative
analysis of risk factors of malaria" case study Gisagara and Bugesera District of Rwanda. RDHS
2014/2015. A retrospective study. BMC Public Health. 2023;23: 1–9.

47. Rudasingwa G, Cho S-I. Determinants of the persistence of malaria in Rwanda. Malar J. 2020;19: 1–
9.

48. Deutsch-Feldman M, Aydemir O, Carrel M, Brazeau NF, Bhatt S, Bailey JA, et al. The changing
landscape of Plasmodium falciparum drug resistance in the Democratic Republic of Congo. BMC
Infect Dis. 2019;19: 872.

49. Sendor R, Mitchell CL, Chacky F, Mohamed A, Mhamilawa LE, Molteni F, et al. Similar Prevalence of
Plasmodium falciparum and Non-P. falciparum Malaria Infections among Schoolchildren, Tanzania.
Emerg Infect Dis. 2023;29: 1143–1153.

50. Habyarimana F, Ramroop S. Prevalence and Risk Factors Associated with Malaria among Children
Aged Six Months to 14 Years Old in Rwanda: Evidence from 2017 Rwanda Malaria Indicator Survey.
International Journal of Environmental Research and Public Health. 2020;17: 7975.

51. Hofmann NE, Gruenberg M, Nate E, Ura A, Rodriguez-Rodriguez D, Salib M, et al. Assessment of
ultra-sensitive malaria diagnosis versus standard molecular diagnostics for malaria elimination: an
in-depth molecular community cross-sectional study. Lancet Infect Dis. 2018;18: 1108–1116.

52. Gruenberg M, Moniz CA, Hofmann NE, Wampfler R, Koepfli C, Mueller I, et al. Plasmodium vivax
molecular diagnostics in community surveys: pitfalls and solutions. Malar J. 2018;17: 55.

53. Gruenberg M, Moniz CA, Hofmann NE, Koepfli C, Robinson LJ, Nate E, et al. Utility of ultra-sensitive
qPCR to detect Plasmodium falciparum and Plasmodium vivax infections under different
transmission intensities. Malar J. 2020;19: 319.