Estimating SARS-CoV-2 transmission parameters between coinciding outbreaks

in a university population and the surrounding community

Matthew S. Mietchen MPH, PhD[1], Erin Clancey PhD[1], Corrin McMichael MPH,

CPH[2], Eric T. Lofgren MSPH, PhD[1]

[1]Paul G. Allen School for Global Health, Washington State University, Pullman, WA

[2]Whitman County Public Health Department, Colfax, WA

Corresponding Author Email: mietchen@unc.edu

Keywords: SARS-CoV-2, COVID-19, university, sub-population, metapopulation, SEIR

model, parameter estimation, likelihood function, particle filter, pMCMC

Abstract

Prior studies suggest that population heterogeneity in SARS-CoV-2 (COVID-19)

transmission plays an important role in epidemic dynamics. During the fall of 2020,

many US universities and the surrounding communities experienced an increase in

reported incidence of SARS-CoV-2 infections, with a high disease burden among

students. We explore the transmission dynamics of an outbreak of SARS-CoV-2 among

university students, how it impacted the non-student population via cross-transmission,

and how it could influence pandemic planning and response.

Using surveillance data of reported SARS-CoV-2 cases, we developed a two-

population SEIR model to estimate transmission parameters and evaluate how these

subpopulations interacted during the 2020 Fall semester. We estimated the

transmission rate among the university students (β_U) and community residents (β_C), as

well as the rate of cross-transmission between the two subpopulations (β_M) using

particle Markov Chain Monte Carlo (pMCMC) simulation-based methods.

We found that both populations were more likely to interact with others in their

population and that cross-transmission was minimal. The cross-transmission estimate

(β_M) was considerably smaller [0.04×10^{-5} (95% CI: 0.00×10^{-5}, 0.15×10^{-5})] compared

to the community estimate (β_C) at 2.09×10^{-5} (95% CI: 1.12×10^{-5}, 2.90×10^{-5}) and

university estimate (β_U) at 27.92×10^{-5} (95% CI: 19.97×10^{-5}, 39.15×10^{-5}). The higher

within population transmission rates among the university and the community (698 and

52 times higher, respectively) when compared to the cross-transmission rate, suggests

that these two populations did not transmit between each other heavily, despite their

geographic overlap.

During the first wave of the pandemic, two distinct epidemics occurred among

two subpopulations within a relatively small US county population where university

students accounted for roughly 41% of the total population. Transmission parameter

estimates varied substantially with minimal or no cross-transmission between the

subpopulations. Assumptions that county-level and other small populations are well-

mixed during a respiratory viral pandemic should be reconsidered. More granular

models reflecting overlapping subpopulations may assist with better-targeted

interventions for local public health and healthcare facilities.

Introduction

SARS-CoV-2 transmission is well established to be largely airborne primarily due

to aerosolized virus, which caused a worldwide pandemic beginning in late 2019. Owing

to the urgent timing for policy decision-making to reduce hospitalizations and mortality,

many studies, including modeling studies, being utilized early in the pandemic assumed

minimal or no population-level heterogeneity, often exploring only large-scale

interactions[1–7]. Geopolitical boundaries, such as states and counties, also encourage a

tendency to focus on single population units due to policy decisions, as well as data

collection efforts, legal jurisdictions, and access to testing and treatment programs

generally governed by these same boundaries.

However, recent studies have found that various groups or subpopulations within

a county, for example, have been affected by SARS-CoV-2 differently, particularly when

there are large universities or industries that residents are associated with in some

capacity[8–10]. The study conducted by Bharti *et al*.,2022 describes how mobile device

data showed a non-student residential population restricted movement during the

pandemic and were able to show that the outbreak was primarily among students,

suggesting distinct strategies for specific sub-populations should be considered for

county-level outbreak management[8]. Additionally, the research by Arnold *et al*., 2022

studied SARS-CoV-2 seroprevalence among county residents, and a returning-student

population were different, implying that heterogeneity in SARS-CoV-2 transmission can

occur within a geographically defined population[9]. Additionally, model-based estimates

such as the time-varying effective reproduction number (R_t) have been used to help

governmental decision-making, such as considering when to relax various non-

pharmaceutical interventions (NPIs) and were often highlighted by state officials in

updates to the public. However, when a heterogenous population may be physically

located in a small spatial environment such as a rural city or county, summary estimates

for the whole population may be deceptive.

Like many universities, Washington State University (WSU), a large rural land-

grant university located in southeastern WA, made the decision to go fully remote for all

students for the Fall 2020 semester. However, many students, faculty, and staff either

remained in the city or returned before the beginning of the semester. While there was

no in-person instruction at the university and widespread lockdown measures were in

place, a large outbreak still occurred. University testing sites, in collaboration with local

public health, enabled all returning students, faculty, and staff to receive free SARS-

CoV-2 (COVID-19) tests before and after the beginning of the semester. While some

negative test requirements were in place, many of these relied on self-attestation. To

our knowledge, all positive student cases were initiated for contact tracing, attempted to

be interviewed, and then required to isolate per university guidelines.

For members of the wider community, SARS-CoV-2 (COVID-19) testing was

available throughout the city for residents at clinics, pharmacies, and non-permanent

testing sites. Additionally, masking requirements and other NPI barrier methods (e.g.,

queue spacing, plastic dividers, etc.) were implemented throughout the city in public

spaces and private businesses. Despite these mitigation efforts, the city experienced a

sharp, dramatic rise in reported cases, primarily among those associated with WSU.

Within the first three weeks of the semester, Whitman County, the county that WSU

resides within, had one of the highest rates of reported SARS-CoV-2 in Washington

State[11].

Considering a multi- or metapopulation framework allows for a higher fidelity look

at subpopulation interactions, potentially identifying opportunities for public health

interventions to suppress those interactions to decreases in incidence and shorter

epidemic durations[12]. Metapopulation models have a rich history of describing the

spatial structure of sub-populations as the driving indicator of disease dynamics and

continue to support how it influences both the rate and range of pathogen dispersal and

the final prevalence of disease within each population[13–16].

In this study, we use a two-population Susceptible-Exposed-Infected-Recovered

(SEIR) transmission model framework to study the transmission of SARS-CoV-2 in a

relatively small rural community and estimate the transmission parameters within and

between two populations defined by an association with a university (or lack thereof)

within a confined geographic area during the Fall of 2020. Doing so allows the

assessment of the extent to which one subpopulation might be driving the epidemic

dynamics of the other subpopulation, and whether or not an intervention targeting one

group (such as vaccination) may be expected to have wider population-level impacts.

Methods

Descriptive Epidemiology

Using SARS-CoV-2 infection data reported to Whitman County, Washington

State, USA as well as information from Washington State University, we performed a

time series analysis of weekly case counts in addition to a descriptive analysis of cases

by age group, birth sex, and university affiliation between July 26, 2020, and January 2,

2021 (Morbidity and Mortality Weekly Report weeks 31 – 53). The time period for this

study was chosen to capture at least a two-week timeframe before and after the

beginning of the Fall semester at a large land grant university prior to a large increase in

cases not previously observed in the county, while terminating before the widespread

availability of vaccines for COVID-19 might alter either transmission dynamics or

individual behaviors.

Due to the limitations and lack of negative test reporting, no data were available

to assess SARS-CoV-2 testing positivity rates. Therefore, we assumed testing rates

between those associated with the university and residents of the community were

similar and not significantly different. We also assumed the availability and access to

testing remained constant throughout the study period, as public health and testing

capacity remained steady and, to our knowledge, did not have major fluctuations.

Time-varying Reproduction Number (R_t) Estimation

Due to the lack of transmission chain data or age-structured matrices and the

focus on sub-population transmission without assumed heterogeneity, the case

reproduction numbers[17] were not estimated in our study; rather we used the R_t

estimation method from Cori *et al.*[18]. Using the case surveillance data, we estimated the

instantaneous reproduction number (R_t) values for the total number of cases and the

sub-population-specific cases throughout the study period from Morbidity & Mortality

Weekly Report (MMWR) Weeks 31-53, 2020. We used the *EpiEstim* package[18,19] for

the R statistical computing language[20].

For our analysis, we assumed a parametric Serial Interval (SI) distribution with a

mean of 5.19 days and a standard deviation of 1.46 days from a pooled estimate in a

systematic review of 11 studies using a random effects model[21]. We assume the serial

interval distribution remained constant throughout the study period. R_t estimates were

adjusted for a 14-day period from the beginning of the study timeframe to ensure

enough case data were available for accurate estimation. The statistical framework of

the Cori *et al.*, 2013 method has been extended to account for future incident cases

being imported rather than localized transmission[22]. However, we assume imported

cases had little to no effect due to the assumption that all university-associated

individuals had arrived in the city prior to the onset of the outbreak.

Model Structure

We model the transmission dynamics of SARS-CoV-2 in a closed population

consisting of two geographically overlapping but distinct subgroups, university-

associated individuals (denoted by the subscript U) and members of the broader

community (denoted by the subscript C) with no university affiliation using a two-

population compartmental model (Figure 1). The university-associated population is

assumed to be primarily undergraduate students, as these represented the vast bulk of

individuals tested by the university itself, while staff and faculty were primarily tested by

their individual healthcare providers.

This model makes several assumptions. The total population ($N_U + N_C$) during

the study period is assumed to remain constant, with no imported cases, births, or

deaths. Hospitalizations and deaths had no impact on transmission and thus were not

modeled directly in order to simplify the model. Once recovered, immunity to re-infection

occurred throughout the length of the study. No prior immunity is assumed due to the

unavailability of vaccination at this point in the pandemic and the relatively low incidence

of reported SARS-CoV-2 infections prior to the large outbreak. The latency period and

recovery rate were assumed to be the same for all those infected, regardless of

population affiliation.

While the more simplified SEIR framework has been widely used for SARS-CoV-

2 (COVID-19) transmission models, it is important to recognize that many models

consider the infected state as individuals experiencing symptoms (clinical) and those

who do not (subclinical) that still contribute to community transmission. While the clinical

course of these two types of infections can vary widely, the absence of symptom-level

information in publicly reported surveillance data, compounded with very mild symptoms

experienced by many college-aged students made disambiguating these two groups

untenable. We demonstrate in Supplement S1 that collapsing these two classes down

to a single infected "I" class effectively provides a weighted average of the two classes,

which accounts for the potential difference in transmission between asymptomatic and

symptomatic individuals at a population level.

We model the data collection process (i.e. weekly case counts) using the

negative binomial distribution. We assume that testing rate and overdispersion are

equivalent in both subpopulations and that testing in each subpopulation occurs

independently. By coupling our mathematical model of the transmission dynamics and

statistical model of the weekly case counts, we formulate a likelihood function (see

Supplement S2 and S3) that then maximize to estimate our parameters of interest.

Parameter Estimation

To further understand the disease dynamics between these two populations

during the COVID-19 pandemic, we used the previously described two-population

compartmental model to estimate the unknown transmission parameters to determine if

the observed epidemics are independent of one another or if there is cross-

transmission. Obtaining estimates of the transmission parameters requires simulation-

based methods which we implement using the Partially Observed Markov Process

(pomp) framework and R package[23,24]. We evaluated the model by simulating the

stochastic process of the model states to maximize the likelihood function and obtain

estimates of the unknown parameters listed in Table 1. Specifically, we used the particle

filter algorithm, as described in Algorithm 1 by King *et al., 2016,* to calculate the value of

the likelihood function for each set of randomly drawn unknown parameters (Table 1)

and the particle Markov Chain Monte Carlo algorithm (pMCMC), as described in

Algorithm 4 by King *et al.*, 2016 to maximize the likelihood estimate for the observed

data[23,25]. Particle Markov-chain Monte Carlo (pMCMC) was used due to its ability to

estimate time-dependent hidden variables[26] and handle relatively high dimensional

problems with reasonable computational demands, a challenge often found when fitting

a model to surveillance data due to testing bias and underreporting. Further specifics

and details of the particle filter and MCMC methods used to obtain our results are

described in Supplement S4.

The initial conditions and fixed parameter values for each subpopulation used for

model simulations are described in Table 1. Due to the lack of available data and small

numbers for both populations, we assumed no persons were in the exposed or latent

periods. Recovered individuals were defined as the number of reported cases between

MMWR weeks 1 and 28 in 2020 by population affiliation. In an effort to more accurately

estimate the number of infected individuals due to testing and reporting delays, we

adjusted the number of reported cases, by subpopulation, during MMWR weeks 29 and

30 using SARS-CoV-2 ascertainment rates obtained for Whitman County (0.1959) and

the statewide average for Washington (0.2458) in August 2020[5]. The Whitman County

rate was applied to the reported cases not affiliated with the university; however, the

statewide rate was used for those in the university subpopulation to account for a large

number of students moving into the city prior to the Fall semester. While we believe the

ascertainment rates are robust, we further adjusted the number of the infected

populations to include pre-symptomatic persons that would most likely not be diagnosed

or reported due to their pre-clinical stage. According to Davies *et al.,* 2020[27], the pre-

symptomatic period was estimated to have a mean duration of 2.1 days out of a 5-day

pre-clinic/clinical duration. Therefore, we calculated the additional pre-symptomatic

cases to be 42% of the adjusted number of infected cases. The rate was applied to both

subpopulations due to the assumption of equal pre-symptomatic durations.

Results

Descriptive Epidemiology

Although the first SARS-CoV-2 (COVID-19) case in the United States was

reported in the state of Washington, the pandemic did not reach many rural parts of the

state until later in 2020. Prior to the study period (MMWR weeks 31-53; July 27, 2020 –

January 2, 2021), only 72 cases were reported in Whitman County, WA (pop. 47,873),

the county where the city and university being studied reside[28]. All reported cases

among city residents, including the university cases, during the first three weeks of the

study period (MMWR weeks 31-33) totaled 24 compared to the following three weeks

(MMWR weeks 34, 35, and 36), where 80, 272, and 339 cases were reported to public

health, respectively (Figure 2a). Additionally, the data in this figure show two distinct

waves, an initial large wave followed by a smaller second wave, suggesting possible

heterogeneities within the population.

Analyzing the reported case data by demographics during the study time period

found that cases were evenly distributed among males and females, with 77% of the

cases being among college-aged (18-22 years) individuals, suggesting high

transmission among university students. Further stratification found the initial large peak

to be primarily among the university population (Table 2). Examining the characteristics

of the initial wave, the data revealed a large epidemic among university students with no

increase in cases among the broader community. Further investigation revealed a

smaller second wave of cases occurring several weeks later, primarily among the non-

university-associated population, appearing around week 40, peaking between weeks

46 and 47 (Figure 2b).

Time-varying Reproduction Number (R_t) Estimation

When assuming a homogenous population using aggregated time series data,

we observe one major peak with a maximum R_t value of 10.48 [8.64, 12.47] lasting 19

days with R_t >1, in addition to a second smaller peak with a maximum R_t value of 2.05

[1.62, 1.52] lasting 21 days with R_t >1 (Figure 3a). When estimating R_t for the two

subpopulations, assuming no interaction or cross-infectivity, not only do we observe

some of the estimates to peak at different points in time but also at very different

magnitudes. While the initial R_t peak among the university-associated cases begins

three days earlier compared to the initial peak among total cases, the peak remains

consistent with a maximum R_t value of 11.75 [9.67, 14.01] and lasting 19 days with R_t

>1, again supporting the initial peak was only among that specific population. However,

the estimates for the remaining duration of the study period show large increases at

different time points. Most notably, the prominent peak among community-associated

cases begins 24 days before the second peak observed within the total case count time

series. The magnitude of the peak is also higher, with a maximum value of 9.96 [4.96,

16.37] and lasting 17 days with R_t >1 (Figure 3b).

Parameter Estimation

We implemented the pMCMC algorithm using pomp[23,24] to estimate the unknown

model transmission parameters β_U (university-university), β_C (community-community),

and β_M (university-community), and additionally K and phi (Table 1). Results of the five-

chain random walks are presented in Supplemental S3. The posterior distributions for

the estimated model transmission parameters were: β_U: 27.92 x 10^{-5} (95% CI: 19.97 x

10^{-5}, 39.15 x 10^{-5}), β_C: 2.09 x 10^{-5} (95% CI: 1.12 x 10^{-5}, 2.90 x 10^{-5}), and β_M: 0.04 x 10^{-5}

(95% CI: 0.00 x 10^{-5}, 0.15 x 10^{-5}). The full distributions are shown in Figure 4.

Both populations were more likely to interact with others in their subgroups, while

the cross-transmission estimate was much smaller, or possibly zero as indicated by the

credible intervals. On a relative scale, the individual transmission rates were

considerably higher (β_U: 698 times and β_C: 52 times) within populations than the cross-

transmission rate β_M.

Discussion

Considering the Whitman County epidemic from several different analytical

standpoints, our results suggest both the university population and the wider

community, while geographically proximate, experienced two separate and largely

independent epidemics, rather than one epidemic simultaneously affecting both

populations or an epidemic that moved from one population to the other.

Because of the larger size of the epidemic within the university population, the

estimated R_t for the total population (Figure 3a) fails twice to detect distinct peaks for R_t

within the community, masking the epidemic in that population almost entirely. Given

the importance of R_t estimates in various governmental decisions about lifting NPIs,

missing these peaks, especially in the population more at risk for severe outcomes, has

profound policy implications.

In this study, we have estimated SARS-CoV-2 transmission parameters between

two populations using an SEIR modeling framework to capture concurrent epidemics.

The results from estimating the three salient transmission parameters paint a similar

picture as the estimates of R_t described above – namely that there appear to be two,

largely independent epidemics within the community. The university population had the

largest estimated transmission parameter value, roughly 13 times that of the

community, but both within-population transmission parameters dwarfed the cross-

population transmission parameter by 698 (University) and 52 (Community) times,

respectively. Of particular note, the credible interval for β_M contained zero. The relative

values of these parameter estimates suggest that transmission resulting from the mixing

of these two populations played a considerably smaller role in the overall dynamics of

the epidemic.

Using particle filtering MCMC, we show that two populations geographically co-

occurring in a small rural town primarily spread the virus within their own groups, while

cross-group transmission estimates were found to be orders of magnitude smaller than

within-group transmission, suggesting separate epidemics. Additionally, comparison of

the two groups, showed there was also considerable heterogeneity in both their timing

and transmission rates.

Conclusion

When modeling relatively small populations within geographically defined areas,

it is understandably tempting to assume homogenous mixing, especially for a disease

like COVID-19 where, due to airborne transmission from aerosolized virus, brief,

transient contacts in space and time between susceptible and infected individuals is

sufficient to cause infection.

However, it is well documented that the social network structures of most

communities vary in degrees of connectedness. In particular, communities with large

universities or single, dominant employers that group individuals together through social

structures and physical interactions. These groupings create highly connected

subpopulations with non-static or fluctuating connections between one another,

effectively creating metapopulations. Our findings suggest that during a respiratory viral

pandemic, such as the COVID-19 pandemic, well-mixed population assumptions at a

city or county level, even if the population is small, should be reconsidered when NPIs

and other public health policies are implemented. Simplistic models (e.g., SEIR) that

assume random interactions could be restructured to contain multiple populations that

are weakly connected by those interactions rather than by geography are useful when

social contact networks are unavailable. The computational costs of these models are

often lower than individual- or network-based models, which may make them more

accessible to a broader range of decision-makers. However, computational expenses

could be a potential limitation depending on model complexity and other resources.

While the quality of the data limits model fitting using surveillance data, model

convergence, and parameter estimation are achievable.

MCMC and other simulation-based methods are dependent upon model

specifications for accurate and valid results and should be implemented carefully.

Improperly specified, model results may be incorrect, regardless of whether

convergence is achieved. Model complexity is also considered a limitation due to

convergence issues arising as the model becomes more complex. More complex

models, initially explored in this study, produced convergence issues necessitating

model simplification. Limitations around an unrepresentative model, for example, the

inability to capture disease states such as pre-symptomatic, asymptomatic, and

symptomatic for the transmission of SARS-CoV-2, may be an oversimplification of

reality.

There are additional model-based limitations based on the realities of using case-

data from a relatively resource constrained public health department. Parameter

estimation from multiple infected classes proved difficult, as was the assessment of

reliable starting values. It has been suggested that asymptomatic or subclinical case

detection requires daily testing capacity and extensive serological data to improve the

certainty of transmission models[29]. Subramanian *et al*., 2021, struggled with parameter

identifiability however, and were leveraging data for a large city. Given the smaller,

sparser data in this study, and the focus on parameter identification, the detection and

testing rate of exposed persons is assumed to be equal regardless of symptomatic

status. While this admittedly simplifies the overall dynamics of the model, we do not

believe the lack of a clinical classification undermines the primary focus of this analysis,

which is on transmission.

Lastly, an important limitation of this study is that public health policies and

messaging, including legally mandated social distancing measures, were in place for

the entire duration of the study period, which precludes analysis of the level to which

these sub-populations mix absent preexisting interventions. We must assume that the

NPIs during this time period of the pandemic amplified the separation between

populations and reduced cross-transmission (indeed, this is their objective), but it is

impossible to assess from our data what the cross-group transmission would be had

these measures not been in place.

It is nevertheless important, anticipating future pandemics, to understand the

methodological aspects of modeling both smaller rural populations and those with

emergent population structures based on social and economic forces. This enables to

assessment of the effectiveness of broad public health interventions, as well as

estimating those that are targeted towards – or will have difficulty reaching – specific

sub-populations. There was a real and justifiable concern that widespread transmission

in university populations would amplify a local epidemic and then spill over into more

vulnerable populations – our results suggest that this did not occur. This, in turn, offers

a justification for some interventions – such as early vaccinations for the most medically

vulnerable, rather than vaccinations trying to contain transmission. At the same time, it

suggests that others, such as basing the suspension of some NPIs on county-level

estimates of transmission, may need to be revisited.

Additional Files

Supplemental Information:

Supplement S1: Averaging of Infected Classes

Supplement S2: Two-Population Ordinary Differential Equations (ODE) Compartment

Model

Supplement S3: Statistical Model and Likelihood Function

Supplement S4: Bayesian Inference

Availability of Data and Materials

Datasets and R code files are available in the **pullman_covid** GitHub repository

(https://github.com/epimodels/pullman_COVID).

Acknowledgments

Funding: ETL, MSM and EC were funded by contracts 75D30121P10551,

75D30120P07911 and 75D30122C15691 from the Centers for Disease Control and

Prevention, as well as RAPID 2110109 from the National Science Foundation. ETL was

also supported on R35GM147013 from the National Institutes of Health.

Conflict of Interest

The authors declare they have no conflicts of interest.

References

1. Kucharski AJ, Russell TW, Diamond C, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. *Lancet Infect Dis*. 2020;20(5):553-558. doi:10.1016/S1473-3099(20)30144-4

2. Ives AR, Bozzuto C. Estimating and explaining the spread of COVID-19 at the county level in the USA. *Commun Biol*. 2021;4(1):60. doi:10.1038/s42003-020-01609-6

3. Krishnamachari B, Morris A, Zastrow D, Dsida A, Harper B, Santella AJ. The role of mask mandates, stay at home orders and school closure in curbing the COVID-19 pandemic prior to vaccination. *Am J Infect Control*. 2021;49(8):1036-1042. doi:10.1016/j.ajic.2021.02.002

4. McKee KL, Crandell IC, Hanlon AL. County-Level Social Distancing and Policy Impact in the United States: A Dynamical Systems Model. *JMIR Public Health Surveill*. 2020;6(4):e23902. doi:10.2196/23902

5. Pei S, Kandula S, Shaman J. Differential effects of intervention timing on COVID-19 spread in the United States. *Sci Adv*. 2020;6(49):eabd6370. doi:10.1126/sciadv.abd6370

6. Zou Z, Fairley CK, Shen M, et al. Critical timing and extent of public health interventions to control outbreaks dominated by SARS-CoV-2 variants in Australia: a mathematical modelling study. *Int J Infect Dis*. 2022;115:154-165. doi:10.1016/j.ijid.2021.11.024

7. Wiens K. In-person schooling and associated COVID-19 risk in the United States over spring semester 2021. *Sci Adv*. Published online 2022:12.

8. Bharti N, Lambert B, Exten C, Faust C, Ferrari M, Robinson A. Large university with high COVID-19 incidence is not associated with excess cases in non-student population. *Sci Rep*. 2022;12(1):3313. doi:10.1038/s41598-022-07155-x

9. Arnold CRK, Srinivasan S, Rodriguez S, et al. A longitudinal study of the impact of university student return to campus on the SARS-CoV-2 seroprevalence among the community members. *Sci Rep*. 2022;12(1):8586. doi:10.1038/s41598-022-12499-5

10. Schwartz EJ, Morgan M, Lapin S. Pandemic 2009 H1N1 influenza in two settings in a small community: the workplace and the university campus. *Epidemiol Infect*. 2015;143(8):1606-1609. doi:10.1017/S0950268814002684

11. Painter I, Huynh G, Ferres JML, et al. SitRep 15: COVID-19 transmission across Washington State. Published online September 8, 2020.

12. Tao Y, Hite JL, Lafferty KD, Earn DJD, Bharti N. Transient disease dynamics across
ecological scales. *Theor Ecol*. 2021;14(4):625-640. doi:10.1007/s12080-021-00514-
w

13. Hess G. Disease in Metapopulation Models: Implications for Conservation. *Ecology*.
1996;77(5):1617-1632. doi:10.2307/2265556

14. North AR, Godfray HCJ. The dynamics of disease in a metapopulation: The role of
dispersal range. *J Theor Biol*. 2017;418:57-65. doi:10.1016/j.jtbi.2017.01.037

15. Chen S, Lanzas C, Lee C, Zenarosa GL, Arif AA, Dulin M. Metapopulation Model
from Pathogen's Perspective: A Versatile Framework to Quantify Pathogen Transfer
and Circulation between Environment and Hosts. *Sci Rep*. 2019;9(1):1694.
doi:10.1038/s41598-018-37938-0

16. Citron DT, Guerra CA, Dolgert AJ, et al. Comparing metapopulation dynamics of
infectious diseases under different models of human movement. *Proc Natl Acad Sci*.
2021;118(18):e2007488118. doi:10.1073/pnas.2007488118

17. Wallinga J. Different Epidemic Curves for Severe Acute Respiratory Syndrome
Reveal Similar Impacts of Control Measures. *Am J Epidemiol*. 2004;160(6):509-516.
doi:10.1093/aje/kwh255

18. Cori A, Ferguson NM, Fraser C, Cauchemez S. A New Framework and Software to
Estimate Time-Varying Reproduction Numbers During Epidemics. *Am J Epidemiol*.
2013;178(9):1505-1512. doi:10.1093/aje/kwt133

19. Cori A. *EpiEstim: Estimate Time Varying Reproduction Numbers from Epidemic
Curves*.; 2021. https://CRAN.R-project.org/package=EpiEstim

20. R Core Team. *R: A Language and Environment for Statistical Computing*. R
Foundation for Statistical Computing; 2021. https://www.R-project.org/

21. Rai B, Shukla A, Dwivedi LK. Estimates of serial interval for COVID-19: A systematic
review and meta-analysis. *Clin Epidemiol Glob Health*. 2021;9:157-161.
doi:10.1016/j.cegh.2020.08.007

22. Thompson RN, Stockwin JE, van Gaalen RD, et al. Improved inference of time-
varying reproduction numbers during infectious disease outbreaks. *Epidemics*.
2019;29:100356. doi:10.1016/j.epidem.2019.100356

23. King AA, Nguyen D, Ionides EL. Statistical Inference for Partially Observed Markov
Processes via the R Package pomp. *J Stat Softw*. 2016;69(12):1-43.
doi:10.18637/jss.v069.i12

24. King AA, Ionides EL, Bretó CM, et al. *Pomp: Statistical Inference for Partially
Observed Markov Processes*.; 2022. https://kingaa.github.io/pomp/

25. Ionides EL, Nguyen D, Atchadé Y, Stoev S, King AA. Inference for dynamic and
latent variable models via iterated, perturbed Bayes maps. *Proc Natl Acad Sci*.
2015;112(3):719-724. doi:10.1073/pnas.1410597112

26. Endo A, van Leeuwen E, Baguelin M. Introduction to particle Markov-chain Monte
Carlo for disease dynamics modellers. *Epidemics*. 2019;29:100363.
doi:10.1016/j.epidem.2019.100363

27. Davies NG, Klepac P, Liu Y, et al. Age-dependent effects in the transmission and
control of COVID-19 epidemics. *Nat Med*. 2020;26(8):1205-1211.
doi:10.1038/s41591-020-0962-9

28. *Washington State Department of Health COVID-19 Data Dashboard*. Washington
State Department of Health; 2022. https://doh.wa.gov/emergencies/covid-19/data-
dashboard

29. Subramanian R, He Q, Pascual M. Quantifying asymptomatic infection and
transmission of COVID-19 in New York City using observed cases, serology, and
testing capacity. *Proc Natl Acad Sci*. 2021;118(9):e2019716118.
doi:10.1073/pnas.2019716118

30. Hakki S, Zhou J, Jonnerby J, et al. Onset and window of SARS-CoV-2
infectiousness and temporal correlation with symptom onset: a prospective,
longitudinal, community cohort study. *Lancet Respir Med*. 2022;10(11):1061-1073.
doi:10.1016/S2213-2600(22)00226-0

31. *Total Student Enrollment - Washington State University*. Washington State
University; 2022. https://ir.wsu.edu/total-student-enrollment/

32. *Washington State University Environmental Health and Safety*.; 2022.
https://ehs.wsu.edu/public-health/contact-tracing/

33. *State of Washington 2021 Population Trends*.; 2022.
https://ofm.wa.gov/sites/default/files/public/dataresearch/pop/april1/ofm_april1__pop
trends.pdf

34. *Whitman County Public Health Department*.; 2021.
https://whitmancountypublichealth.org/community-health/disease-prevention/covid-
19/

516
517
518

Tables

Three parameters define SARS-CoV-2 viral transmission rates among and between these subgroups; β_U (university-university), β_C (community-community), and β_M (university-community). Exposed populations are within the latency period (α) or the time period between exposure and infectiousness, and those recovered are no longer infectious and leave the infectious period at a rate of γ or the recovery rate. Two additional parameters were estimated, ϕ or the testing probability and κ which is a variance constant (Table 1).

Table 1. Model Parameters and Initial Conditions

Parameter	Parameter Description	Value	Source(s)
β_U	Transmission Rate between university-associated individuals	Unknown	Estimated
β_C	Transmission Rate between individuals in the community	Unknown	Estimated
β_M	Cross-Transmission Rate between the university and community populations	Unknown	Estimated
α	Latency Period or the time between a person being exposed and becoming infectious with a pathogen	$3.59\ \text{days}^{-1}$	[5]
γ	Recovery Rate or the duration of infectiousness	$5\ \text{days}^{-1}$	[27,30]
ϕ	Testing probability or the proportion of those who are infected and eligible for testing (parameter is required by the model)	Unknown	Estimated
κ	Variance constant (parameter is required by the model)	Unknown	Estimated
Total University Population (N_U)	Estimated number of enrolled undergraduate students at WSU (Pullman Campus) for the 2020 Fall semester living on or near campus	14,254	[31]
Susceptible (S_U)	Number of susceptible university-associated individuals $[S_U = N_U - (E_U + I_U + R_U)]$	14,158	Calculated
Exposed (E_U)	Number of individuals associated with the university known to be exposed to SARS-CoV-2	0	Assumed
Infected (I_U)	Estimated number of infected individuals associated with the university in the two weeks prior to the study period (2020 MMWR Week 29 & 30)	81	[5,27,32]

Recovered (R_U)	Number of reported SARS-CoV-2 cases that were associated with the university prior to the study period, who are no longer infectious (2020 MMWR Week 1-28)	15	[32]
Total Community Population (N_C)	Estimnated number of non-student residents in the city of Pullman, WA	20,785	[33]
Susceptible (S_C)	Number of susceptible individuals not associated with the university [$S_C = N_C - (E_C + I_C + R_C)$]	20,642	Calculated
Exposed (E_C)	Number of individuals not associated with the university known to be exposed to SARS-CoV-2	0	Assumed
Infected (I_C)	Estimated number of infected individuals not-associated with the university in the two weeks prior to the study period (2020 MMWR Week 29 & 30)	124	[5,27,34]
Recovered (R_C)	Number of reported SARS-CoV-2 cases among the residents of Pullman, WA who were not university-associated prior to the study period, who are no longer infectious (2020 MMWR Week 1-28)	19	[34]

548 Table 2. Reported SARS-CoV-2 Infections from 07/26/2020 – 01/02/2021 by Age

549 Group, Sex, and Sub-population Demographic, Pullman, WA

University Sub-population

Age Group	Female (%) [†]	Male (%) [†]	Total (%) [†]
0-17	1 (0.1%)	0 (0%)	1 (0.1%)
18-22	740 (92.4%)	705 (87.8%)	1,445 (90.1%)
23-32	42 (5.2%)	73 (9.1%)	115 (7.2%)
33-39	7 (0.9%)	9 (1.1%)	16 (1.0%)
40-59	10 (1.3%)	12 (1.5%)	22 (1.4%)
60 and above	1 (0.1%)	4 (0.5%)	5 (0.3%)
Total (%) *	**801 (49.9%)**	**803 (50.1%)**	**1,604**

Community Sub-population

Age Group	Female (%) [†]	Male (%) [†]	Total (%) [†]
0-17	22 (9.5%)	14 (5.9%)	36 (7.6%)
18-22	50 (21.6%)	47 (19.7%)	97 (20.6%)
23-32	55 (23.7%)	70 (29.3%)	125 (26.5%)
33-39	22 (9.5%)	20 (8.4%)	42 (8.9%)
40-59	47 (20.3%)	51 (21.3%)	98 (20.8%)
60 and above	36 (15.5%)	37 (15.5%)	73 (15.5%)
Total (%) *	**232 (49.3%)**	**239 (50.7%)**	**471**

551 † Percentages across Age groups category

552 *Percentage across Sex category

553 Note: Percentage may not add up evenly due to rounding.

   Figures

561   Figure 1. Two-population SEIR Model Schematic of the University (U) and Community

562   (C) population interactions

579   Figure 2a. Reported SARS-CoV-2 epidemiological curve by MMWR week

583   Figure 2b. Reported SARS-CoV-2 epidemiological curve by MMWR week and

584   subpopulation

587   Figure 3a. R_t estimates of Total SARS-CoV-2 Reported Cases

588
589
590

591   Figure 3b. R_t estimates of SARS-CoV-2 Reported Cases by Sub-Population

597 Figure 4. Posterior distributions of Processed pMCMC for Parameter Estimation. The

598 red vertical dashed line represents the modal posterior estimate, with 95% credible

599 intervals shown as the solid black line on the x-axis for each estimate.

601