Unveiling the crucial neuronal role of the proteasomal ATPase subunit gene *PSMC5* in neurodevelopmental proteasomopathies

Sébastien Küry,[1,2,*] Janelle E. Stanton,[3,4] Geeske van Woerden,[5,6,7] Tzung-Chien Hsieh,[8] Cory Rosenfelt,[9] Marie Pier Scott-Boyer,[10] Victoria Most,[11] Tianyun Wang,[12,13,14] Jonas Johannes Papendorf,[15] Charlotte de Konink,[5,6] Wallid Deb,[1,2] Virginie Vignard,[2] Maja Studencka-Turski,[15] Thomas Besnard,[12] Anna Marta Hajdukowicz,[15] Franziska Thiel,[15] Sophie Möller,[15] Laëtitia Florenceau,[2] Silvestre Cuinat,[1,2] Sylvain Marsac,[1] Ingrid Wentzensen,[16] Annabelle Tuttle,[16] Cara Forster,[17] Johanna Striesow,[18] Richard Golnik,[19] Damara Ortiz,[20] Laura Jenkins,[20] Jill A. Rosenfeld,[21,22] Alban Ziegler,[23] Clara Houdayer,[24] Dominique Bonneau,[24,25] Amber Begtrup,[16] Kristin G. Monaghan,[16] Sureni V. Mullegama,[16] C.M.L. (Nienke) Volker-Touw,[26] Koen L. I. van Gassen,[26] Renske Oegema,[26] Mirjam de Pagter,[26] Katharina Steindl,[27] Anita Rauch,[27,28,29,30] Ivan Ivanovski,[27] Kimberly McDonald,[31] Emily C. Boothe,[31] Andrew Dauber,[32] Janice Baker,[33] Noelle Andrea V Fabie,[33] Raphael A. Bernier,[34] Tychele N. Turner,[35] Siddharth Srivastava,[36] Kira A. Dies,[36] Lindsay Swanson,[36] Carrie Costin,[37] Cherith Somerville,[38,39] Rebekah K. Jobling,[39] Tiffany Turner,[37] John Pappas,[40,41] Rachel Rabin,[40] Dmitriy Niyazov,[42] Anne Chun-Hui Tsai,[43] Karen Kovak,[44] David B. Beck,[45,46] MCV (May Christine) Malicdan,[47,48] David R Adams,[48] Lynne Wolfe,[48] Rebecca D. Ganetzky,[49,50] Colleen Muraresku,[49,50] Davit Babikyan,[51,52] Zdeněk Sedláček,[53] Miroslava Hančárová,[53] Andrew T. Timberlake,[54] Hind Al Saif,[55,56] Berkley Schmidt,[55] Kayla King,[55] Karen Kovak,[57,58] MJ Hajianpour,[59] Gregory Costain,[39,60,61,62] Tayyaba Khan,[60,61] D'Arcy Prendergast,[39,62] Chumei Li,[63] David Geneviève,[64] Antonio Vitobello,[65,66] Arthur Sorlin,[67,68] Christophe Philippe,[65,66] Tamar Harel,[69,70] Ori Toker,[71] Ataf Sabir,[72,73] Danielle Bogue,[74] Derek Lim,[72] Mark Hamilton,[75] Lisa Bryson,[76] Elaine Cleary,[77,78,79] Sacha Weber,[80] Trevor L. Hoffman,[81] Anna Maria Cueto-González,[82,83] Eduardo Fidel Tizzano,[82,83] David Gómez-Andrés,[84] Marta Codina-Solà,[82,83] Athina Ververi,[85] Efterpi Pavlidou,[86] Alexandros Lambropoulos,[87] Kyriakos Garganis,[88] Marlène Rio,[89] Jonathan Levy,[90,91] Sarah Jurgensmeyer,[92,93] Anne M. McRae,[92] Mathieu Kent Lessard,[94] Maria Daniela D'Agostino,[95] Isabelle De Bie,[96] Meret Wegler,[97] Rami Abou Jamra,[97] Susanne B. Kamphausen,[98] Viktoria Bothe,[97] Uwe Völker,[99] Elke Hammer,[99] Kristian Wende,[18] Benjamin Cogné,[1,2] Bertrand Isidor,[1,2] Julien Marcoux,[100,101] Marie-Pierre Bousquet,[100,101] Jeremie Poschmann,[102] Frédéric Laumonnier,[103,104] Peter W. Hildebrand,[105,106,107] Evan E. Eichler,[108,109] Kirsty McWalter,[16] Peter M. Krawitz,[8] Arnaud Droit,[110,111] Ype Elgersma,[5,6] Andreas M. Grabrucker,[3,4,112] Francois Bolduc,[9,113,114] Stéphane Bézieau,[1,2,115] Frédéric Ebstein,[2,115,*] Elke Krüger,[15,115,*]

[1]Nantes Université, CHU Nantes, Service de Génétique Médicale, F-44000 Nantes, France;
[2]Nantes Université, CHU Nantes, CNRS, INSERM, l'institut du thorax, F-44000 Nantes, France;
[3]Health Research Institute (HRI), University of Limerick, Limerick, Ireland;
[4]Department of Biological Sciences, University of Limerick, Limerick, Ireland;
[5]Department of Clinical Genetics, Erasmus University Medical Center, Rotterdam, 3015 GD, The Netherlands;
[6]ENCORE Expertise Centre for Neurodevelopmental Disorders, Erasmus Medical Center, Rotterdam, 3015 GD, The Netherlands;
[7]Department of Neuroscience, Erasmus Medical Center, Rotterdam, 3015 GD, The Netherlands;

[8]Institute for Genomic Statistics and Bioinformatics, University Hospital Bonn, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany;

[9]Department of Pediatrics, University of Alberta, Edmonton, AB, Canada;

[10]Centre de recherche du Chu de Quebec-Université Laval, 2705 boulevard Laurier Québec, Québec, G1V 4G2, Canada;

[11]Institute for Drug Discovery, Medical Faculty, Leipzig University, 04103 Leipzig, Germany;

[12]Department of Medical Genetics, Center for Medical Genetics, Peking University Health Science Center, Beijing 100191, China;

[13]Neuroscience Research Institute, Peking University; Key Laboratory for Neuroscience, Ministry of Education of China & National Health Commission of China, Beijing 100191, China;

[14]Autism Research Center, Peking University Health Science Center, Beijing 100191, China;

[15]Universitätsmedizin Greifswald, Institut für Medizinische Biochemie und Molekularbiologie, 17475 Greifswald, Germany;

[16]GeneDx, 207 Perry Parkway, Gaithersburg, MD 20877, USA;

[17]Loyola University Chicago, Chicago, IL 60660, USA;

[18]Leibniz Institute for Plasma Science and Technology (INP), 17489, Greifswald, Germany;

[19]Klinik für Pädiatrie I, Universitätsklinikum Halle (Saale), 06120 Halle (Saale), Germany;

[20]UPMC Children's Hospital of Pittsburgh, One Children's Hospital Drive, 4401 Penn Avenue, Pittsburgh, PA 15224, USA;

[21]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, 77030, USA;

[22]Baylor Genetics Laboratory, Houston, TX, 77021, USA.;

[23]Service de Génétique, CRMR AnDDI-Rares, CHU Reims, Reims, France;

[24]CHU Angers, Service de Génétique médicale, Angers, France;

[25]Mitovasc, UMR CNRS 6015, INSERM U1083, Angers University, Angers, France;

[26]Department of Genetics, University Medical Centre Utrecht, Utrecht University, 3508 AB Utrecht, The Netherlands;

[27]Institute of Medical Genetics, University of Zürich, Schlieren-Zurich, 8952, Switzerland;

[28]University Children's Hospital Zurich, Zurich, Switzerland;

[29]University of Zurich Research Priority Program ITINERARE: Innovative Therapies in Rare Diseases, Zurich, Switzerland;

[30]University of Zurich Research Priority Program AdaBD: Adaptive Brain Circuits in Development and Learning, Zurich, Switzerland;

[31]University of Mississippi Medical Center, 2500 North State Street, Jackson, MS 39216, P 601.984.1597, F 601.984.1916, USA;

[32]Division of Endocrinology, Children's National Hospital and Department of Pediatrics, The George Washington University School of Medicine and Health Sciences, Washington, DC 20010, USA;

[33]Department of Medical Genetics and Genomics, Children's Hospitals and Clinics of Minnesota, Minneapolis, MN 55404, USA;

[34]Department of Psychiatry & Behavioral Sciences, Center on Human Development and Disability, University of Washington, Box 357920 Seattle, WA 98195, USA;

[35]Department of Genetics, Washington University School of Medicine, St. Louis, MO 63110, USA;

[36]Rosamund Stone Zander Translational Neuroscience Center, Boston Children's Hospital, Boston, Massachusetts, USA;

[37]Department of Genetics, Akron Children's Hospital, One Perkins Square, Akron, Ohio 44308, USA;

[38]Ted Rogers Centre for Heart Research, Cardiac Genome Clinic, The Hospital for Sick Children, Toronto, ON M5G 0A4, Canada;

[39]Division of Clinical and Metabolic Genetics, The Hospital for Sick Children, Toronto, ON, Canada;

[40]Clinical Genetic Services, Department of Pediatrics, NYU School of Medicine, New York, NY 10016, USA;

[41]Clinical Genetics, NYU Orthopedic Hospital, New York, NY 10010, USA;

[42]Division of Medical Genetics, Department of Pediatrics, Duke University School of Medicine, Durham, NC 22710, USA;

[43]Department of Pediatrics, College of Medicine Peoria, University of Illinois, Chicago, IL 61605, USA;

[44]Department of Molecular and Medical Genetics, Oregon Health and Sciences University, OHSU, Portland, Oregon, USA;

[45]Division of Rheumatology, Department of Medicine, New York University Grossman School of Medicine, New York, NY, USA;

[46]Center for Human Genetics and Genomics, New York University Grossman School of Medicine, New York, NY, USA;

[47]Medical Genetics Branch, National Human Genome Research Institute, NIH, Bethesda, MD 20892-1851, USA;

[48]National Institutes of Health Undiagnosed Diseases Program, National Human Genome Research Institute, NIH, Bethesda, MD 20892, USA;

[49]Department of Pathology and Laboratory Medicine, Children's Hospital of Philadelphia, Philadelphia, PA, USA;

[50]Division of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA, USA;

[51]Department of Medical Genetics, Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia;

[52]Laboratory of Molecular Genetic, Center of Medical Genetics and Primary Health Care, Yerevan, Armenia;

[53]Department of Biology and Medical Genetics, Charles University 2nd Faculty of Medicine and University Hospital Motol, 150 06 Prague, Czech Republic;

[54]Wyss Department of Plastic Surgery, NYU Langone Medical Center, New York, NY, USA;

[55]Department of Human and Molecular Genetics, Division of Clinical Genetics, Virginia Commonwealth University School of Medicine, Richmond, VA, USA;

[56]Department of Pediatrics, Division of Clinical Genetics, Virginia Commonwealth University School of Medicine, Richmond, VA, USA;

[57]Oregon Health & Science University, USA;

[58]Oregon Genetics Program, Oregon Health Authority, 800 NE Oregon St, Ste. 370, Portland, OR, 97232, USA;

[59]Division of Medical Genetics and Genomics, Department of Pediatrics, Albany Medical College, Albany, NY, USA;

[60]Department of Molecular Genetics, University of Toronto, Toronto, ON M5S 1A4, Canada;

[61]Program in Genetics and Genome Biology, SickKids Research Institute, Toronto, ON, Canada;

[62]Department of Paediatrics, Temerty Faculty of Medicine, University of Toronto, Toronto, ON, Canada;

[63]McMaster University Medical Center, Hamilton, ON, Canada;

[64]Université Montpellier, Inserm U 1183, centre de référence maladies rares anomalies du développement, Service de génétique médicale, Hôpital Arnaud de Villeneuve, Montpellier, France;

[65]UMR 1231 GAD, Inserm, Université de Bourgogne Franche Comté, Dijon, France;

[66]Unité Fonctionnelle Innovation en Diagnostic Génomique des Maladies Rares, FHU-TRANSLAD, CHU Dijon Bourgogne, Dijon, France;

[67]Centre de génétique et UF6254 « Innovations en Diagnostic Génomique des Maladies Rares », CHU Dijon Bourgogne, France;

[68]Service de Pédiatrie 1 et de Génétique Médicale, Centre Hospitalier Universitaire Dijon Bourgogne, Dijon, France;

[69]Department of Genetics, Hadassah Medical Organization, Jerusalem, Israel;

[70]Faculty of Medicine, Hebrew University of Jerusalem, Jerusalem, Israel;

[71]Department of Pediatrics, Allergy and Clinical Immunology Unit, Shaare Zedek Medical Center, Faculty of Medicine, Hebrew University of Jerusalem, Jerusalem, Israel;

[72]Department of Clinical Genetics, Lavender House, Birmingham Women's and Children's Hospital NHS Foundation Trust, Birmingham, UK;

[73]Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK;

[74]Department of Paediatrics, Sandwell & West Birmingham NHS Trust, Birmingham, UK;

[75]West of Scotland Clinical Genetics Service, Queen Elizabeth University Hospital, Glasgow G51 4TF, UK;

[76]Department of Clinical Genetics, NHS Greater Glasgow and Clyde, Glasgow, UK;

[77]South East Scotland Genetics Service, Western General Hospital, Edinburgh, UK;

[78]Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh EH16 4SB, UK;

[79]UK Dementia Research Institute at University of Edinburgh, University of Edinburgh, Edinburgh, EH16 4SB, UK;

[80]CCA-AHU de génétique clinique et de neurogénétique, Service de Génétique et de Neurologie, CHU de Caen, Caen, France;

[81]Department of Genetics, Southern California Kaiser Permanente Medical Group, Anaheim, CA 92801, USA;

[82]Department of Clinical and Molecular Genetics, Vall d'Hebron Hospital Universitari, Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain;

[83]Medical Genetics Group, Vall d'Hebron Institut de Recerca (VHIR), Vall d'Hebron Hospital Universitari, Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain;

[84]Pediatric Neurology, Vall d'Hebron Institut de Recerca (VHIR), Hospital Universitari Vall d'Hebron, Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain;

[85]Department of Genetics for Rare Diseases, 'Papageorgiou' General Hospital, 56429 Thessaloniki, Greece;

[86]Department of Speech and Language Therapy, University Hospital of Ioannina, 45111 Ioannina, Greece;

[87]Genetic Unit, 1st Department of Obstetrics and Gynecology, School of Medicine, Aristotle University of Thessaloniki, 'Papageorgiou' General Hospital, 56429 Thessaloniki, Greece;

[88]Epilepsy Unit, St Luke's Hospital, Thessaloniki, Greece;

[89]Service de Médecine Génomique des Maladies Rares, Hôpital Necker-Enfants Malades, AP-HP, 75015 Paris, France;

[90]Department of Genetics, APHP-Robert Debré University Hospital, 75019 Paris, France;

[91]Multi-site medical biology laboratory SeqOIA - FMG2025, 75014 Paris, France;

[92]Division of Genetics, Genomics and Metabolism, Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago, USA;

[93]Department of Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, USA;

[94]Division of Medical Genetics, McGill University Health Centre, Montréal, Quebec, Canada;

[95]Division of Medical Genetics, Departments of Human Genetics and Medicine, McGill University, Montreal, QC, Canada;

[96]Division of Medical Genetics, Department of Specialized Medicine, McGill University Health Centre, Montréal, Quebec, Canada;

[97]Institute of Human Genetics, University of Leipzig Medical Center, 04103 Leipzig, Germany;

[98]Institute of Human Genetics, University Hospital Magdeburg, 39120 Magdeburg, Germany;

[99]Universitätsmedizin Greifswald, Interfakultäres Institut für Genetik und Funktionelle Genomforschung, Abteilung für Funktionelle Genomforschung, 17487 Greifswald, Germany;

[100]Institut de Pharmacologie et de Biologie Structurale (IPBS), Université de Toulouse, CNRS, Université Toulouse III - Paul Sabatier (UT3), Toulouse, 31077, France;

[101]Infrastructure Nationale de Protéomique, ProFI, FR 2048, Toulouse, France ;

[102]Nantes Université, CHU Nantes, INSERM, Center for Research in Transplantation and Translational Immunology, UMR 1064, 44000, Nantes, France;

[103]UMR 1253, iBrain, Université de Tours, Inserm, 37032 Tours, France;

[104]Service de Génétique, Centre Hospitalier Régional Universitaire, 37032 Tours, France;

[105]Institut für Medizinische Physik und Biophysik, Universität Leipzig, Medizinische Fakultät, Härtelstr. 16-18, 04107 Leipzig, Germany;

[106]Charité Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Institute of Medical Physics and Biophysics, Berlin, Germany;

[107]Berlin Institute of Health, 10178 Berlin, Germany;

[108]Department of Genome Sciences, University of Washington School of Medicine, Seattle,WA 98195, USA;

[109]Howard Hughes Medical Institute, Seattle, WA 98195, USA;

[110]Département de médecine moléculaire, Faculté de médecine, Université Laval, 2325 rue de l'Université, Québec, G1V 0A6, Canada;

[111]Centre de recherche du CHU de Québec-Université Laval, 2705 boulevard Laurier Québec, Québec, G1V 4G2, Canada;

[112]Bernal Institute, University of Limerick, Limerick, Ireland;

[113]Neuroscience and Mental Health Institute, University of Alberta, Edmonton, AB, Canada;

[114]Department of Medical Genetics, University of Alberta, Edmonton, AB, Canada;

[115]These authors equally contributed to this work.

*Corresponding authors: E-mails: sebastien.kury@chu-nantes.fr (S.K.); frederic.ebstein@univ-nantes.fr (F.E.) ; elke.krueger@uni-greifswald.de (E.K.)

Abstract

Neurodevelopmental proteasomopathies have recently emerged as a distinct class of neurodevelopmental disorders (NDD). These conditions involve alterations in genes of the 26S proteasome, a protein complex essential for maintaining protein homeostasis in eukaryotic cells. Our study identified 23 unique variants in *PSMC5*, which encodes the AAA-ATPase proteasome subunit PSMC5/Rpt6, causing syndromic NDD in 38 unrelated individuals. *PSMC5* loss-of-function resulted in abnormal protein aggregation, profoundly affecting innate immune signaling, mitophagy rate, and lipid metabolism in samples of affected individuals samples. Moreover, the morphology of human hippocampal neurons was altered by overexpression of *PSMC5* variants, while *PSMC5* knockdown led to impaired reversal learning in flies and loss of excitatory synapses in rat hippocampal neurons. Promisingly, targeting key integrated stress response components like PKR and GCN2 kinases alleviated immune alterations in cells of affected individuals. These findings deepen our understanding of the molecular mechanisms underlying neurodevelopmental proteasomopathies and link these disorders with neurodegenerative disease research.

Introduction

The proper structure and function of eukaryotic cells rely on the maintenance of the intracellular proteome composed of thousands of proteins. The dynamic balance between protein synthesis and degradation is orchestrated by the proteostatic network[1,2], an integral component of which is the ubiquitin-proteasome system (UPS), that selectively eliminates short- and long-lived proteins, as well as misfolded proteins modified with ubiquitin[3]. At the very heart of the UPS is the 26S proteasome, a large multi-subunit protease consisting of a 20S core particle and a 19S regulatory particle, comprising a lid and a base[3]. Lid subunits recognize and bind polyubiquitinated target proteins[1]. Subsequently, base subunits Rpt1-6 (PSMC1-6) utilize the energy released by ATP hydrolysis for deubiquitinating, unfolding and translocating the substrates into the proteolytic chamber of the 20S core particle in which they are degraded into oligopeptides. These in turn can be further hydrolyzed into amino acids by peptidases, enabling efficient recycling within the cellular protein synthesis machinery[1,3].

The remarkable dynamics and plasticity of proteasomes confer them with a significant impact on cellular physiology[3], particularly in neurons. Neuronal synaptic plasticity relies on continuous proteome renewal, necessitating robust translation at ribosomes and concurrent breakdown of defective ribosomal products (DRiPs)[2,4]. This maintenance of protein homeostasis poses a considerable challenge in cortical neurons, where synapses may contain hundreds of proteins[4]. Effective protein clearance by the proteasome is crucial for various neuronal processes, including synaptic remodeling, cell migration, neurotransmitter release, long-term potentiation, long-term depression, and memory formation[5,6]. Experimental inhibition of proteasome activity has been shown to disrupt synapse composition and promote the aggregation of polyubiquitinated proteins, ultimately causing neurodegeneration[6]. Similar observations have been made in brain tissues from individuals with neurodegenerative disorders[6] and schizophrenia[7]. Genetic investigations have further revealed that loss-of-function variants in proteasome genes contribute to the onset and

progression of a unique class of NDDs, specifically referred to as the neurodevelopmental proteasomopathies[8]. The inheritance pattern for these disorders can either be dominant, as seen with *PSMD12* variants associated with Stankiewicz-Isidor syndrome [MIM: 617516][9,10] and *PSMC3*[11], or recessive, as observed in cases involving *PSMB1* [MIM: 620038][12] and *PSMC3* [MIM: 619354][13] variants.

The proteasome subunit PSMC5/Rpt6 is of particular interest in this context due to its role in regulating proteasome activity and synaptic remodeling. This ATPase serves as a neuronal sensor that dynamically adjusts proteasome activity and localization in response to protein turnover requirements through phosphorylation by CAMKIIα[14]. The present study further confirms the pivotal role of *PSMC5* in neuronal function. It identified 23 variants in *PSMC5*, the vast majority heterozygous and *de novo*, from 38 unrelated individuals with impaired neurodevelopment. Functional analyses performed on patient cells and animal models unveiled that *PSMC5* alterations lead to a loss of proteasome function, triggering a complex cellular program involving profound remodeling of innate immunity, inflammation, and lipid metabolism. Intriguingly, the integrated stress response (ISR) mediates most of these changes, and potential molecular targets have been identified, holding promise for the development of biomarkers and therapeutic interventions.

Materials and Methods

Genetic studies and ethics statement

The 38 affected individuals of the present study were enrolled together with their healthy parents whenever possible by participating teams involved in diagnostic or research activities on developmental disorders. All 38 affected individuals were clinically assessed by at least one expert clinical geneticist from each participating center to whom they gave written informed consent for participating in the study. Routine clinical genetic and metabolic screenings, including array comparative genomic hybridization and fragile X testing, were performed during initial workup and did not permit diagnosis of a known inherited condition, which warranted further investigation on a research basis, using a genome- or exome-trio sequencing approach. Connections between the participating centers were aided by the web-based tool GeneMatcher. Consent for the publication of photographs was obtained for subjects S1, S3, S9, S10, S12, S13, S19, S30-34 and S38. The photographs are not shown in the current version of the manuscript but are deposited in GestaltMatcher database; accession links are provided in Figure 2, in place of the original pictures. The study has been approved by the CHU de Nantes-ethics committee (number CCTIRS: 14.556). This research was ethically approved by CPP Ouest V (File 06/15) on 04/08/2015 (Ref MESR DC 2017 2987).

Three-dimensional (3D) structural analysis of PSMC5/Rpt6 modeling the effects of the identified missense variants

This analysis was carried out to investigate the spatial localization within PSMC5/Rpt6 of the amino acid substitutions corresponding to the missense variants. Our objective was to offer insights into potential effects of the mutations on subunit arrangements, interactions, and related conformational changes. To achieve this, we employed two distinct structures: a Cryo-EM structure engaged with a substrate (PDB ID: 6MSK) and a substrate-free structure (PDB ID: 7W37). These structures allowed us to place the AAA+ ATPase subunit PSMC5/Rpt6 within the context of the 26S proteasome, in

particular within the 19S regulatory particle, of which it is an integral part. The resulting refined 3D map was then visualized using molecular visualization software UCSF ChimeraX.

Facial image analysis

We performed the GestaltMatcher approach[15] on *PSMC5* subjects to analyze the facial similarities among the 13 subjects (S1, S3, S9, S10, S12, S13, S19, S30-34 and S38) whose family consented to facial analysis. We first utilized the model ensemble and test-time augmentation to encode each photo into 12 512-dimensional vectors. Each image *i* was represented in the space as 12 512-dimensional vectors $x_{i,k} \in \mathbb{R}^{512} \ and \ k \in [1,12]$. The similarity between two subjects can be quantified by averaging 12 cosine distances between two subjects. When the cosine distance is smaller, the two subjects are more similar. To validate whether the subjects of the given cohort C, *PSMC5* subjects, are similar to each other, we performed the statistical analysis at the cohort level and the pairwise comparison analysis at the individual level.

We first performed a statistical analysis based on the mean pairwise cosine distance. We selected 1,555 images from different subjects with 328 different syndromes from GestaltMatcher Database (GMDB)[16] and sampled two control distributions: (1) distances between subjects of the same syndrome and (2) distances between random patients. In the end, we compared the mean pairwise distance between patients of the given cohort C to these two distributions.

For each syndrome *S* and a random cohort size *n* $(2 \leq n \leq |S|)$, we sampled two cohorts: (1) C_S consisting of *n* patients from the chosen syndrome *S*, (2) C_R consisting of *n* randomly chosen patients from GMDB. We then calculated the mean pairwise cosine distance for each of two cohorts. This was repeated 100 times per syndrome, duplicated sampled cohorts were removed, resulting in a distribution of mean pairwise cosine distances between (1) patients stemming from the same syndrome, and (2) random patients, respectively.

We conducted a Receiver Operating Characteristic (ROC) analysis on the sampled cohorts from the same syndrome ("controls") and from random patients ("cases"). We used the mean pairwise cosine distance as a measure of discrimination and performed the 5-fold cross-validation. We chose the threshold c by selecting the fold with the highest Youden index to decide whether the patients inside a given cohort are similar (mean pairwise cosine distance $< c$) or different (mean pairwise cosine distance $> c$), resulting in $c=0.915$, corresponding to a sensitivity of 0.851, a specificity of 0.862, and an Area Under the Curve (AUC) of 0.895

To examine the similarity of the *PSMC5* subjects, we calculated their mean pairwise cosine distances and compared them to the sampled distributions originating from patients from the same syndrome and random patients. Then, we sampled a cohort C_C of random sample size n ($2 \leq n \leq |C|$) consisting of n patients from the given cohort C and calculated the mean pairwise cosine distance of this cohort. This was repeated 10,000 times and duplicated sampled cohorts were removed resulting in a distribution of mean pairwise cosine distances between patients stemming from the given cohort C.

In the end, we performed the pairwise comparison analysis on 13 *PSMC5* subjects. To simulate the real-life scenario, we first compared photographs of the 13 *PSMC5* subjects to 7,459 images with 449 different disorders stored in GMDB. For each *PSMC5* subject tested, we performed a leave-one-out cross-validation by placing the other 12 *PSMC5* subjects into the space of the 7,459 images and calculating the ranks of 12 subjects relative to the *PSMC5* subject tested.

Behavioral studies in fly

Background. Rpt6 is the Drosophila ortholog of PSMC5. When using the DRSC integrative ortholog prediction tool (DIOPT), it has a rank of high homology and a weighted score of 15/16. The KD approach was substantiated by the presence of truncating

variants in the cohort (V5:p.(Lys196Argfs*29) and V23:p.(Asp394Glufs*2)) and the resultant loss of proteasomal function.

Drosophila Strains. The *Rpt6* (*Psmc5*) Drosophila RNAi stock (34712) was obtained from Bloomington Drosophila Stock Center. Wild-type and ELAVGAL4 stocks were obtained from Dr. Tim Tully (Cold Spring Harbor Laboratory). Stocks were raised on a cornmeal agar medium at 22°C and 40% humidity. University of Alberta ethics and regulations were followed.

Genetic Crosses. Virgin females of wild-type or ELAVGAL4 were collected and crossed to theUAS-RNAi males.

Olfactory Learning. Drosophila olfactory learning was conducted by training 100 flies to associate an odour with a simultaneously presented footshock. A second control odour was then presented without a footshock. Flies were then funneled into a choice point and given a choice between the two odours. Flies that had correctly learned the association would avoid the odour that had been paired with the shock. This entire process was then repeated with 100 naive flies but with the odour shock pairing reversed in order to rule out any odour bias. The combined performance between these two trials constituted the performance index and n=1.

In reversal olfactory learning, the same procedure as normal learning was conducted with the added step of an additional training cycle after, in which the stimulus pairing was reversed. For example, flies were presented with odour A paired with a footshock and then odour B without footshock for the initial training cycle, followed byan additional training cycle where now odour B was paired with a footshock and odour A was presented without. This is a more challenging task to perform as flies are required to unlearn the initial association in addition to learning the new pairing. Once again, this entire process was then repeated with 100 naive flies but with the odour pairings reversed in order to rule out any odour bias.

Experiments were conducted at 25°C and 70% humidity. Adult Drosophila were transferred to bottles the day before testing. At the time of testing, flies were ages1-3 days.

Statistical analysis. For the learning experiments, a two-tailed t-test was performed. For the reversal learning experiments, a one-way ANOVA followed by Tukey's test was performed.Prism software was used for all statistical analysis.

Analysis of *PSMC5* knockdown in rat hippocampal neurons

Cultivation of primary neurons. Rat hippocampal neurons (RN-h, P10101, Innoprot) were plated on poly-L-lysine (PLL) (0.1 mg/mL)-coated coverslips in a 24- well plate at a density of 5×10^4 on coverslips placed in a 24-well plate. RN-h were cultivated using Neurobasal medium containing 2% B27 (#17504044, Gibco), 1% Glutamax (#35050079, Gibco), 100 µg/mL penicillin and streptomycin. Cultures were incubated at 37°C with 5% CO_2 in a sterile CO_2 incubator for 14 days.

Lentiparticle-based Psmc5 knockdown. 24 hours after plating, *Psmc5* - Rat shRNA lentiviral particles (TL711271V, Origene) were introduced into the culture medium at a multiplicity of Infection rate of 5 and incubated for 18 hours before a complete medium change was carried out.

Immunocytochemistry. Neurons were first washed with 1x PBS and fixed using 4% PFA for 15 min at room temperature. Permeabilization was performed by applying 0.2% Triton-X for 5 min and washing with 1x PBS. Next, blocking solutions (BS) (10% FBS/ 1x PBS) was applied for 1 h at room temperature. After that, the primary antibody (MAP2 (1:1000, Synaptic Systems #18006); vGlut (1:1000, Synaptic Systems #135304); vGAT (1:200, Synaptic System #131003)); PSMC5 (1:400, Sigma-Aldrich)) in BS was applied, and cells were incubated at 4°C overnight. Next, 1x PBS was applied for 3x 5 min, followed by incubating the secondary antibody diluted in BS in darkness for 1 h at room temperature. Cells were washed with 1x PBS twice for 5 min. An additional wash step includes the addition of 4,6-di-amino-2-phenylindol (DAPI) diluted in 1xPBS for 5 min. Finally, the cells were washed with 1x PBS, and a wash step with ddH_2O was also carried out for 5 min. An ImageXpress Micro Confocal microscope (Molecular Devices, San Jose, USA) was used to

obtain the images at a 40X magnification. Image analysis was carried out using ImageJ v1.53.

Analysis of neuronal morphology in mouse

Cloning. Constructs containing human *PSMC5*^WT or mutated *PSMC5* cDNA sequences were isolated from the pcDNA3.1/Zeo(+) expression vector by PCR and tagged with AscI and PacI sequences, to allow ligation in the dual promotor expression vector, containing a CAGG-promotor for expression of the gene of interest and a PGK-promotor driven tdTomato gene to easily distinguish transfected cells from non-transfected cells. The following primers were used: (AscI) FW 5'-gaatccggcgcgccaccatggcgcttgacggaccagagc and (PacI) RV 5'-gaatccttaattaatcacttccataatttcttgatg. The empty vector control (EV) in the *in vitro* assays is merely the backbone plasmid, lacking any *PSMC5* sequence.

Mice. For the primary hippocampal neurons, FvB/NHanHsd females (ordered at 6-8 weeks old from Envigo) were crossed with FvB/NHanHsd females (ordered at 6-8 weeks old from Envigo). All mice were group-housed in IVC cages (Sealsage 1145T, Tecniplast) with bedding material (Lignocel BK 8/15 from Rettenmayer) on a 12/12 h light/dark cycle in 21°C (±1°C), humidity at 40-70% and with food pellets (801727CRM(P) from Special Dietary Service) and water available *ad libitum*. All animal experiments were conducted in accordance with the European Commission Council Directive 2010/63/EU (CCD project license AVD101002017893), and all described experiments and protocols were subjected to ethical review (and approved) by an independent review board (IRB) of the Erasmus MC.

Primary hippocampal cultures. To obtain primary hippocampal neurons, hippocampi were isolated from brains from E16.5 embryos and collected in ice cold Neural Basal Medium (NB; Gibco, 21103049). After two washes with ice cold NB, the hippocampi were incubated with Trypsin/EDTA (Sigma; T3924) for 20 min at 37°C. The tissue was again washed twice with warm NB and finally kept in supplemented NB (1%penicillin/streptomycin (Sigma; p4333)/1%GlutaMax (Gibco; 350500-38)/2%B27-supplement (Gibco; 17504044) for dissociation. Single cells were seeded in 12-wells plates containing Poly-D-Lysine (Sigma;

P0899) coated coverslips and 1ml supplemented NB per well. The cells were incubated at 37°C/5% CO_2 until transfection.

Neuronal transfections and immunological stainings. Primary hippocampal neurons (e16.5) were transfected on DIV3 with the following constructs: Empty expression vector; *PSMC5*^{WT} having the wild type human cDNA sequence referring to mRNA sequence NM_002805.6; PSMC5^{Arg201Trp} containing variant c.601C>T p.(Arg201Trp); *PSMC5*^{Ala202Val} containing variant c.605C>T p.(Ala202Val); *PSMC5*^{Thr207Met} containing variant c.620C>T p.(Thr207Met); *PSMC5*^{Pro320His} containing variant c.959C>A p.(Pro320His); *PSMC5*^{Pro320Arg} containing variant c.959C>G p.(Pro320Arg); *PSMC5*^{Arg325Trp} containing variant c.973C>T p.(Arg325Trp); *PSMC5*^{Δex10} having a deletion of exon 10 mimicking the likely consequence of splice site variant V20:c.970-2A>G. 2.5µg DNA was transfected of each PSMC5 construct. As control, 1.8µg DNA of the empty expression vector was transfected. Neurons were transfected by mixing the DNA with NB and Lipofectamin2000 according to the manufacturer's instructions (Invitrogen; 11668-019)

The cells were fixed 5 days post-transfection (DIV8) using 4%PFA/4%sucrose. Fixed cells were labeled for MAP2 (1:500, Synaptic System; #188004) by o/n incubation at 4°C and visualized with a conjugated secondary antibody donkey-anti-guinea pig Alexa647 (1:200; Jackson ImmunoResearch #706-605-148), incubated for 1h at RT and finally covered with Mowiol (home-made) to allow for fluorescence imaging using confocal microscopy.

Confocal microscopy. All images were acquired by using a LSM700 confocal microscope (Zeiss). For imaging overexpression in primary hippocampal neurons, images were taken from 10 transfected neurons per condition, per neuronal batch (20x objective, 0.5 zoom, 2048 x 2048).

Analysis and statistics. The total neurite length and number of branches was traced using the program ImageJ and its plug-in NeuronJ. Within every batch, the data was normalized to the PSMC5^{WT}, allowing us to pool the normalized data from different batches. For statistics, a one-way ANOVA (Dunnett's multiple comparison test) was done, using Prism GraphPad. The n represents data from an individually traced neuron. For every condition a

minimum of 8 neurons were imaged per batch, for a minimum of 2 independent primary hippocampal neuron batches.

Expansion of T cells from PBMC isolated from control and patient samples

Whole-blood samples collected from healthy individuals as well as Subjects S1, S6, S11, S12, S21, S32 and S33 and their respective parents and/or siblings whenever possible were processed using spin medium gradient centrifugations (pluriSelect) to separate and isolate peripheral blood mononuclear cells (PBMC) for cryopreservation using a standard freezing medium consisting of 90% FBS and 10 % DMSO. At a later point in time, PBMC were plated on 96-well plates together with irradiated allogeneic PBMC in the presence of IL-2 and PHA-L for T cell expansion as in [11]. Human T cells were maintained in RPMI1640 supplemented with 10 % human AB serum and 1% penicillin/streptomycin and analyzed at a resting state after 3-4 weeks of expansion.

SDS-PAGE and western-blot analysis

Resting T cells and/or SH-SY5Y neuroblastoma cells were lysed in RIPA buffer (50 mM Tris pH 7.5, 150 mM NaCl, 2 mM EDTA, 1% NP40, 0.1% SDS) and proteins were quantified using a standard BCA assay (Thermofisher). Twenty micrograms of protein lysates were loaded on 10-15% SDS-PAGE and subsequently transferred onto PVDF membranes under wet conditions (200V, 400 mA, 1 h at 4°C). Following a 20-min blocking with ROTI®Block (Carl Roth), membranes were probed with primary antibodies overnight at 4°C under permanent shaking. Antibodies used in this study were directed against HA (BioLegend, clone HA.11), PSMC5/Rpt6 (Enzo Life Sciences, clone p45-110), α-tubulin (Abcam, clone DM1A), α6 (Enzo Life Sciences, clone MCP20), ubiquitin K48-linked ubiquitin chains (Cell Signaling, clone D9D5), GAPDH (Cell Signaling, clone 14C10), GRP94 (Invitrogen, PA5-27860), ATF6 (Cell Signaling, clone D4Z8V), BNIP3L (Cell Signaling, 12396). Antibody binding was detected using HRP-conjugated anti-mouse, -rabbit or -goat

secondary antibodies and chemiluminescence (Biorad) following the manufacturer's recommendations.

RNA isolation, reverse-transcription and PCR analysis

Total RNA was extracted from snap frozen SH-SY5Y and/or T cell pellets using the innuPREP RNA Mini Kit from Analytic Jena AG, following the manufacturer's recommendations. Five hundred nanograms of the isolated RNA was reverse transcribed using the M-MLV reverse transcriptase from Promega. Real-time PCR was performed using the Premix Ex Taq™ (probe qPCR, TaKaRa) in duplicates to determine the mRNA levels of the interferon (IFN)-stimulated genes (ISG) *IFIT1*, *IFI27*, *IFI44*, *IFI44L*, *ISG15*, *MX1*, *RSAD2* as well as *GAPDH* (housekeeping) using FAM-tagged TaqMan™ Gene Expression Assays purchased from Thermo Scientific following the manufacturer's guidelines. The cycle threshold (Ct) values for target genes were converted to values of relative expression using the relative quantification (RQ) method (2-$\Delta\Delta$Ct). Target gene expression was calculated relative to Ct values for the GAPDH control housekeeping genes. In some experiments, interferon (IFN) scores were calculated as the median of the RQ of the seven ISG over a single calibrator control following the procedure of Rice et al. as in [11]. Semi-quantitative PCR was conducted on SH-SY5Y cDNA to amplify overexpressed HA-*PSMC5* using a forward (5'-ATGGCGCTTGACGGACCAGAGCAGA-3') primer binding to *PSMC5* and a reverse primer (5'-GACAGTGGGAGTGGCACCTTCCAGGGTCAAGG-3') binding to the polyadenylation signal of the pcDNA3.1/Zeo(+) expression vector. Amplified cDNA products were resolved on 1.7% agarose gels, stained with 1.0 mg/ml GelRed®, and visualized by a UV transilluminator at 312 nm.

Native-PAGE and in-gel activity assays

Resting T cells from control and NDD affected individuals with *PSMC5* variants were subjected to non-denaturing protein extraction by three cycles of freeze (liquid nitrogen) and thaw using TSDG buffer (10 mM Tris pH 7.0, 10 mM NaCl, 25 mM KCl, 1.1 mM $MgCl_2$, 0.1

mM EDTA, 1 mM DTT, 1 mM ATP, 1 mM NaN$_3$ and 20 % Glycerol). Samples were then centrifuged (14,000xg) for 15 minutes at 4°C and protein content in supernatants was determined using a standard Bradford assay. Twenty micrograms of whole-cell extract proteins were separated on 3-12% gradient Bis-Tris gels (Invitrogen) and in-gel activity assay was performed to visualize proteasome complexes by incubating gels with 0.1 mM fluorogenic peptide (LLVY-AMC, purchased from Bachem) for 30 minutes at 37°C. In addition, gels were blotted onto PVDF membranes (200V, 400 mA, 1h at 4°C under wet conditions) which were subsequently incubated with primary antibodies directed against the proteasome subunits PSMC5/Rpt6 (Enzo Life Sciences, clone p45-110) and α6 (Enzo Life Sciences, clone MCP20) overnight at 4°C. Antibody binding was then detected using an anti-mouse horseradish peroxidase (HRP)-labelled secondary antibody and chemiluminescence (Biorad).

Analysis of gene expression through NanoString.

A total of 100 nanograms of RNA was extracted from T cells sourced from both control and affected samples. The isolated RNA underwent hybridization using the NanoString nCounter® Human AutoImmune Profiling Panel. Following this, gene expression quantification adhered to the manufacturer's instructions. The acquired data were subsequently normalized to housekeeping genes in accordance with the manufacturer's guidelines.

Results

Rare pathogenic *PSMC5* variants cause a syndromic neurodevelopmental disorder

We identified 38unrelated individuals presenting similar clinical features and harboring a rare single nucleotide variant (SNV) or indel in *PSMC5*. Thirty cases were *de novo*, five assumed *de novo*, two inherited from an affected mother, and one recessive. A total of 23 distinct variants were found, four recurring (**Table 1**), mostly missense (18/23) affecting residues in regions highly conserved from human to yeast (**Fig. S1**). These variants predominantly occurred in the "ATPases Associated with diverse cellular Activities (AAA-ATPase)" domain (14/23), including the recurring variants NM_002805.6: c.959C>G p.(Pro320Arg) (7/38 subjects) and c.973C>T p.(Arg325Trp) (8/38) near the end of the domain (**Fig. 1A**).

Our three-dimensional structural analysis showed notably that 13 of the variants primarily located in the lower ATPase domain interface with neighboring base ATPase subunits PSMC1/Rpt2 and PSMC4/Rpt3, subunits of the lid complex like PSMD14/Rpn11, and the substrate in its engaged state (**Fig. 1B,C,S2A,B**). They are likely to affect proteasome function in multiple ways, by disrupting proteasomal complex formation (**Fig. S2C**), impairing the transition from substrate-free to substrate-engaged states of the 26S proteasome (**Fig. S2D**), affecting ATP binding and hydrolyzation and thereby the integrity of the tertiary proteasomal structure (**Fig. S2D, S3A**), hindering substrate processing (**Fig. 1B,S2E,S3B**), or perturbing protein folding and subunit incorporation (**Fig. 1B,S2B, S3C**). Overall, these findings, detailed in supplemental notes, align with the criteria that would classify of all 23 variants as pathogenic according to the American College of Medical Genetics and Genomics (ACMG) guidelines if *PSMC5* was an OMIM morbid gene (**Table S1**).

Abnormal neurodevelopment was evident in all subjects, with developmental delay in 35/38 (92%) cases (**Tables 2-S2b**). Neurological manifestations included speech absence or delay (33/35; 94%), abnormal muscle tone (26/34; 76%), motor delay or impairment (27/36;

75%), intellectual disability (19/26 (73%)), abnormal behavior (23/34; 68%, including autism spectrum disorder (ASD) and attention deficit hyperactivity disorder (ADHD)), and seizures (10/37; 27%) (**Tables 2-S2**). Brain magnetic resonance imaging revealed frequent but varied abnormalities in the examined subjects (14/22; 64%). The phenotype appeared syndromic (**Fig. 2A; Table S2b/S2c**), with notable non-neurological findings such as ophthalmological anomalies (24/32; 75%), skeletal malformations (21/38; 55%), feeding difficulties (20/37; 54%), cardiac abnormalities (11/38; 29%), hearing loss (9/35; 26%, conductive in four individuals, sensorineural and mixed in one patient each), kidney abnormalities (5/37; 14%) and genital abnormalities (5/38; 13%).

Subjects with *PSMC5* variants present a similar facial gestalt

Most affected individuals exhibited abnormal facial shape (28/35; 80%), often accompanied by craniofacial abnormalities, including microcephaly (12/32; 38%) or abnormality of the mandible (11/35; 31%, including micrognathia, retrognathia and prognathia) (**Fig. 2B**). However, the manifestations were heterogeneous and did not allow conclusive diagnosis of the disorder through clinical assessment alone. We therefore performed facial analysis with GestaltMatcher[15] to validate a common facial phenotype among *PSMC5* subjects. Firstly, we evaluated the similarity between 13 subjects from the PSMC5 cohort with photos provided. Ninety-five percent of subject combinations had mean pairwise distances below the threshold (*c=0.915*), indicating strong facial resemblance (**Fig. S4**). Secondly, we extended the analysis to 7,459 images from 449 disorders in GMDB. Pairwise comparison matrix (**Fig. 2C**) revealed that 10 of 13 *PSMC5* subjects (S3/9/10/12/13/19/30-33) matched at least another *PSMC5* subject with a rank below 50. Subject pairs/trios (S9 and S33), (S10, S19 and S31), and (S13 and S30) were highly similar, with ranks below five, indicating again strong resemblance. These findings collectively suggest a recognizable facial gestalt among *PSMC5* subjects.

Knockdown (KD) of *Psmc5* alters fly reversal learning and excitatory/inhibitory balance of rat hippocampal neurons

Pan-neuronal RNAi KD of *Psmc5*/Rpt6 in *Drosophila melanogaster* had no significant effect on normal learning performance (p=0.3059) (**Fig. 3A**). However, when conducting the more difficult reversal learning paradigm, KD of Rpt6 resulted in significantly impaired performance as compared to controls (p<0.0001) (**Fig. 3B**).

We studied the effects of *Psmc5* loss on primary rat hippocampal neurons by gene KD, using *Psmc5*-specific shRNA or scrambled shRNA for controls (scrambled Ctrl) along with coexpressed GFP delivered via lentiviral-particle infection of cells at day *in vitro* (DIV) 1 of development (**Fig. 3C-J**). After 14 days in culture, a significant reduction of PSMC5 levels was measured by fluorescent labelling with PSMC5-specific antibodies (**Fig. 3C-D**). Notably, under control conditions, PSMC5 labelling appeared predominantly in neurons and much less in glial cells.

To assess the general morphology of neurons developing under *PMSC5* KD, we measured their dendritic arborization at DIV 14. No significant differences were observed in *PSMC5* KD neurons compared to controls (**Fig. 3E-F**). We also measured synapse density of excitatory synapses (visualized using vGlut as excitatory presynaptic marker) and inhibitory synapses (visualized using vGat as inhibitory presynaptic marker) (**Fig. 3G-J**). After 14 days, *PSMC5* loss led to a significantly reduced number of vGlut positive puncta along primary dendrites (**Fig. 3G-I**), whereas the number of vGat positive puncta remained unchanged. (**Fig. 3H-J**). This indicates an excitation/inhibition (E/I) imbalance due to PSMC5 loss in neurons.

Overexpression of *PSMC5* affects neuronal morphology in mouse

A subset of identified missense variants was assessed for their impact on neuronal morphology through overexpression in primary mouse hippocampal neurons. Neurite length

of mouse hippocampal cells was significantly increased during the neurodevelopmental phase (DIV3) by overexpression of protein product PSMC5-WT compared to empty vector control (**Fig. 4A,B**), suggesting a potential role of *Psmc5* in promoting neurite outgrowth. Overexpression of variants p.(Arg201Trp), p.(Pro320His) and p.(Arg 325Trp) yielded the same effect as PSMC5-WT, suggesting that in this assay these variants do not alter the effect of overexpression of *PSMC5*. By contrast, overexpression of variants p.(Ala202Val), p.(Thr207Met), p.(Pro320Arg) and PSMC5-Δex10 (c.970-2A>G) showed reduced neurite length compared to PSMC5-WT (**Fig. 4A,B**); this effect, similar to the one induced by the empty vector, suggest that these variants have a loss of function (LoF) effect in the assay. Notably, no effect on arborization was induced by PSMC5-WT and all variants had no impact on arborization compared to the empty vector control (**Fig. 4C**).

The *PSMC5* variants associated with NDD do not equally impact the PSMC5/Rpt6 subunit steady-state expression and subsequent incorporation into 26S proteasomes

To analyze the ability of PSMC5/Rpt6 variant subunits to integrate 26S proteasome complexes, we ectopically expressed 13/23 identified alterations as N-terminally HA tagged-*PSMC5* versions in SHSY-5Y neuroblastoma cells, as previously described[11].

While the plasmid-driven production of *PSMC5* transcripts was similar between the wild-type and the 13 variant cell lines (**Fig. S5A**), steady-state protein expression levels of PSMC5/Rpt6 were differentially impacted by the variants: they were reduced profoundly by p.(Ala202Val), moderately by p.(Pro183Leu), p.(Arg201Trp), p.(Gly216Asp), p.(Glu250Val) and p.(Pro320Arg), and mildly by p.(Arg258Trp), p.(Pro320His) and p.(Arg325Trp), whereas they were unaltered by p.(Thr207Met), p.(Met368Thr) and p.(Asp394Glufs*2) (**Fig. S5B**). Overall, analyses of the 14 *PSMC5* variants examined revealed no significant impact on *in vitro* PSMC5/Rpt6 abundance (**Fig. S5B**, right panel), indicating that haploinsufficiency is unlikely the main driver of variant pathogenicity.

We also observed distinct effects of the 13 *PSMC5* variants on proteasome assembly in this *in vitro* assay. Notably, p.(Ala202Val) prevented PSMC5 incorporation into 19S, 26S, and 30S complexes (**Fig. S5C**), due to the instability of the HA-PSMC5/Rpt6 full-length protein (**Fig. S5B**). Conversely, despite their stable expression in SH-SY5Y cells (**Fig. S5B**), p.(Met368Thr) and p.(Asp394Glufs*2) accumulated in 19S precursors and fully assembled 19S particles without associating with 26S and/or 30S proteasome complexes (**Fig. S5C**). Unlike variants p.(Gly216Asp), p.(Gln221Arg), p.(Arg201Val) and p.(Arg258Trp) which showed unchanged incorporation efficiency compared to their wild-type counterpart, p.(Pro183Leu), p.(Glu250Val), p.(Pro320His) and p.(Pro320Arg) only minimally assembled into mature proteasomes, (**Fig. S5C**). Surprisingly, p.(Thr210Met) had a higher propensity to integrate 19S-capped proteasomes than the wild-type. Densitometric analysis showed that the 13 *PSMC5* variants studied had similar incorporation into 19S, 26S, and/or 30S complexes compared to their wild-type counterparts (**Fig. S5C**, lower panel). These findings underscore the varied effects of these variants on subunit expression, protein level stability, and incorporation into mature 26S/30S proteasome complexes.

Most *PSMC5* variants do not lead to haploinsufficiency but result in severe proteasome assembly defects

To assess the impact of *PSMC5* variants in samples from patients, we conducted experiments using T cells derived from affected subjects. Except p.(Pro320Arg), no *PSMC5* variants reduced PSMC5/Rpt6 levels in subjects' T cells compared to control samples (**Fig. 5A**). The unchanged levels of PSMC5/Rpt6 in S12 [p. (Glu250Val)] and S32's [(p. Arg325Trp)] T cells, were discordant with *in vitro* results that had hinted at variant-induced protein instability in SHSY-5Y cells (**Fig. S5B**). This observation suggests a full compensation by the wild-type allele and counters haploinsufficiency as a cause of the disorder. Notably, while the PSMC5/Rpt6 steady-state protein expression levels were comparable between control and subjects, all T cell specimens with *PSMC5* variants showed

extra lower-migrating PSMC5/Rpt6 species compared to controls (**Fig. 5A, lower panel**), possibly due to a dysregulation of *PSMC5* transcript splicing.

In line with *in vitro* findings, T cells of subjects S6 [p.(Arg201Trp)], S21 [p.(Pro320Arg)], S32 [p.(Arg325Trp)], and S33 [p.(Arg325Trp)] exhibited reduced incorporation of PSMC5/Rpt6 into 26S and 30S proteasome complexes, as evidenced by less intense staining of PSMC5/Rpt6 in the 19S-capped proteasomes (**Fig. 5B**). However, for two other *PSMC5* variants, there was a discrepancy between *in vivo* and *in vitro* results: (i) p.(Gln221Arg) was associated with decreased PSMC5/Rpt6 subunit incorporation in S11's T cells (**Fig. 5B**), potentially influenced by a second proteasomal variant, NM_002815.3:c.268C>T p.(Arg90*) in *PSMD11*, inherited from a moderately affected father. Although, we did not consider this variant in our *in vitro* assays, its effect on 26S proteasome assembly seems evident, given the reduced amounts of 30S proteasomal complexes as demonstrated by the decreased intensity of α6 staining in S11's father's T cells; (ii) p.(Glu250Val) did not impair PSMC5/Rpt6 assembly into 26S/30S proteasomal complexes in S12's T cells (**Fig. 5B**), in contrast to the significant disruption observed in SHSY-5Y cells (**Fig. S5C**). In summary, *PSMC5* variants behave diversely, some of them failing to maintain stable expression or integrate into mature proteasomes.

PSMC5 variants profoundly disrupt protein homeostasis and lipid metabolism, and activate mitophagy in T cells from NDD subjects

To determine the effect of *PSMC5* variants on protein homeostasis, we evaluated the propensity of T cells from individuals S1/6/11/12/21/32 to form protein aggregates. We observed increased aggresome formation in all subject samples with S6 and S21 exhibiting a particularly high aggresome propensity factor (APF) (**Fig. 6A**). Accordingly, this coincided with the accumulation of high molecular weight (HMW) ubiquitin-modified proteins detected in all five subjects (**Fig. 6B**). These findings demonstrate the loss-of-function nature of *PSMC5* variants, which prevent T cells from coping with proteotoxic stress.

To delve deeper into the cellular repercussions of proteasome loss of function, we conducted proteomics investigations on subject-derived T cells. Our findings reveal a strong correlation between proteasome dysfunction and increased expression levels of HLA class I molecules and immune regulatiors, (**Fig. S6, Table S3d**). Furthermore, we observed many enriched proteins involved in amino acid or protein metabolism (FKBP1A, NAPG, LXN, DDAH2, FAH, DPP4, CSTB, GGT3P, AASDHPPT), oxidative stress (GPX4, GSTM1), intracellular transport and signaling (RRAGC, AGA, VPS35L, ARL8A, TBC1D10A, NCALD), regulation of proliferation and cell death (TRIAP1, BRD8, DFFA, CASP6, TRADD), as well as apoliproteins (**Table S3d**).

The massive enrichment of apolipoproteins prompted us to investigate lipid compositions of T cell samples of patients versus healthy controls. Surprisingly, we observed a massively altered distribution of lipids in samples of patients with *PSMC5* variants (**Fig. 6C,D**). Thus, phospholipids of membranes such as phosphatidylserine (PS), phosphatidylethanolamine (PE), phosphatidylcholine (PC), as well as di- (DG) or tri-acylglycerols (TG), were significantly decreased in patients, whereas cholesterolesters were increased by 50%.

The enrichment of mitochondrial proteins (**Fig. S6B**) may point to impaired mitochondrial import or protein quality control. To investigate whether defective proteasomal degradation was accompanied by mitochondrial damage, we monitored mitophagy rates by flow cytometry in S1/6/11/12/21/33, using pH-sensitive MtPhagy dye. Indeed, T cells from individuals with *PSMC5* variants exhibited higher lysosomal-targeted mitochondria proportions than controls (**Fig. 6E**). This confirms that proteasome dysfunction is associated with increased degradation of mitochondria in these subjects, indicating mitochondrial damage.

NDD individuals harboring *PSMC5* variants display sterile type I IFN responses predominantly triggered by the ISR

So far all proteasompathy syndromes (PRAAS and NDD) were linked to dysregulated type I interferon (IFN) signaling irrespective of the gene variants. Indeed, our focused transcriptomic analysis using a NanoString encounter autoimmune gene panel revealed a strongly elevated type I IFN gene signature of five affected individuals compared to healthy controls or relatives (**Fig. 7A**).

To further confirm the association between *PSMC5* variants and a type I IFN gene signature, we calculated the IFN scores of T cells from S1/11/12/21/32/33 and unaffected controls, based on transcript expression of seven IFN-stimulated genes (ISGs: *IFIT1*, *IFI27*, *IFI44*, *IFI44L*, *ISG15*, *MX1* and *RSAD2*). All affected individuals showed elevated expression of the ISGs compared to controls (**Fig. S7**), resulting in high IFN scores (**Fig. 7B**) and confirming the above theory. Intriguingly, S11's father, bearing heterozygous *PSMD11* variant p.(Arg90*) (**Table S2**), exhibited a positive type I IFN score (**Fig. 7B**), strengthening the idea that proteasome loss-of-function variants consistently lead to type I IFN gene signatures, even in mildly affected subjects.

Considering the known role of protein kinase R (PKR) in triggering sterile autoinflammation in response to proteasome dysfunction[17], we assessed its contribution to the initiation of type I IFN responses in T cells from subjects with *PSMC5* variant, using C16, a specific PKR inhibitor. The IFN scores from T cells treated with C16 were significantly lower in T cells treated with C16 than in the same T cells exposed to DMSO (controls), confirming PKR's role in spontaneous ISG induction in the disorder (**Figs. 7C,S8**). A comparable reduction was observed upon H-151 treatment (**Figs. 7C,S8**), underscoring an additional contribution of the cGAS-STING pathway sensing host-derived DNA this process. Of note, the IFN scores dropped dramatically with JAK inhibitor baricitinib (**Figs. 7C, S8**), confirming the autocrine/paracrine nature of this phenomenon. While blocking the UPR with

the IRE1 inhibitor 4µ8C did not yield any significant effects (**Figs. S8**), GCN2 inhibition by A92 significantly lowered IFN scores for all subjects, emphasizing the key role of ISR in the generation of type I IFN.

Discussion

Rare syndromic NDDs caused by *PSMC5* loss-of-function underscore the vulnerability of genes encoding proteasome subunits to genomic lesions, leading to a wide range of severe neurological phenotypes. *PSMC5* now joins a growing list of proteasome subunit genes linked to various central nervous system (CNS) disorders[9-12]. It further highlights the crucial developmental role of the proteasome. which has already been emphasized in studies involving plants and mice, where proteasome disruption is known to cause severe organ defects or non-viability[18,19].

While *PSMC5* is expressed throughout the body, phenotyoing points to a predominant neurodevelopmental phenotype in all affected individuals. The connection between *PSMC5* and NDDs had already been strongly suggested by a large international cohort study, which had stressed a significant enrichment of *de novo* variants in this gene[20]. This aligns with the described critical role of PSMC5/Rpt6 in neuronal function, particularly in regulating synaptic function in hippocampal neurons through CAMKIIα-mediated phosphorylation[21,22]. However, *PSMC5* loss of function impacts development beyond the CNS, spanning various physiological systems (**Fig. 2A, Tables 2,S2**), albeit with variable severity and extent among individuals.

Utilizing the T-cell model, our objective was to elucidate the fundamental cellular mechanisms involved with primary cells from patients. Our *in vitro* functional investigations on this model showed that T cells with *PSMC5* variants present a characteristic cellular phenotype consisting notably of proteasome loss-of-function (**Figs. 5,6**), increased mitophagy (**Figs. 6C,S6B**)**,** and a type I IFN gene signature (**Figs. 7B,S7**). The latter feature bears resemblance to what is observed in individuals with chronic atypical neutrophilic dermatosis with lipodystrophy and elevated temperature (CANDLE, also known as proteasome-associated autoinflammatory syndromes, PRAAS), the first described monogenic disorders associated with proteasome gene variants[23]. Nevertheless, unlike

individuals with CANDLE/PRAAS, who exhibit severe and typical cutaneous and systemic inflammatory symptoms[23], individuals with NDD in the present series do not exhibit striking signs of autoinflammation, although pathogenesis mechanisms partially overlap in proteasome impairment, activation of proteotoxic stress and dysregulation of type I IFN signaling. Similar to the Aicardi-Goutières-Syndrome, a prototypic type I interferonopathy with CNS manifestation, developmental defects such as growth retardation also have been observed in patients with CANDLE/PRAAS[23].

Although it remains unclear whether dysregulated type I IFN production actively contributes to the acquisition of NDD phenotypes, it is noteworthy that two potential markers of the disorders described here, PKR and GCN2—both sensors of the integrated stress response (ISR)—belong to the eIF2α kinase (EIF2AK) family. This kinase family has already been associated with NDD via two of its members, *EIF2AK1* and *EIF2AK2*[24]. Interestingly, in mammals, GCN2, a regulator of neurogenesis known for its inhibitory effect on spontaneous neuritogenesis, also plays a significant role in behavioral control and memory consolidation[25]. In turn, PKR is involved in the pathogenesis of neurodegenerative disorders, notably Alzheimer's disease[26]. In this context, our results hold great promise, as type I IFN in T cells harboring *PSMC5* variants could be reduced using inhibitors of PKR, GCN2, and JAK (**Figs. 7C,S8**). These findings merit further investigation to clarify the role of inflammation on neurogenesis in the current disorder.

PSMC5 variants notably disrupt lipid homeostasis: (i) in S6 and S11, we noted an enrichment of apolipoproteins APOA1, APOA2, APOE, and APOC3 **(Fig. S6)**, whose degradation partially rely on proteasomes[27,28]; and (ii) S1/6/11/12/19/21/33's cells were endowed with reduced amounts of triglycerides, diglycerides and glycerophospholipids, while their pools of cholesterol ester were increased **(Fig. S6D)**. Perturbations of lipid homeostatis were noted in CANDLE/PRAAS subjects, typically suffering from lipodystrophy[23].

Furthermore, lipid remodeling has been associated with other neurodevelopmental pathologies, including ASD[29] and ADHD[30].

To emphasize the neuronal implications of *PSMC5* variants, we employed both hippocampal cell and *Drosophila melanogaster* models. Assays in primary hippocampal neurons suggested that LoF variants impair the neuritogenic role of *Psmc5**(**Fig. 4**), Additionally, the *Drosophila Psmc5*-knockdown (KD) model demonstrated the influence of *Psmc5* loss on cognitive performance. Our assays showed that *Psmc5* KD impaired reversal learning without compromising memory (**Fig. 3A,B**). These findings echo observations from the paralogous gene *PSMC3*, whose KD produces similar effects in flies[11]. Additionally, parallels emerge from studies showcasing behavioral deficits in male rats with reduced proteasomal activity during memory recovery[31]. Interestingly, mouse investigations suggested the involvement of *Psmc5* in molecular and behavioral plasticity, particularly in response to stimuli like cocaine[32]. These studies demonstrated that Psmc5/Rpt6, when forming a complex with ΔFosB and other proteins in the nucleus accumbens, regulates chromatin remodeling and gene expression. This observation gains significance considering the enriched presence of dopaminergic neurons in the midbrain region, pivotal in behavioral flexibility[33]. In addition to its connection with ΔFosB, PSMC5/Rpt6 was found to physically interact with type B GABA (GABAB) receptors, modulating their presence on cell surfaces and subsequently affecting neuronal activity[34]. This suggests a potential link between *PSMC5* variants and GABAB signaling, a key element in behavioral flexibility[35], which is impaired in ASD and ADHD[36].

In the last experimental model involving rat hippocampal cells, *Psmc5* KD induced a lowered E/I ratio (**Fig. 3G-J**). This phenomenon has been associated with modified cognitive function in humans, affecting decision-making negatively[37]. The imbalance in E/I ratio is implicated in conditions such as ASD and schizophrenia, where it disrupts brain neural circuits and leadds to a decline in cognitive abilities[38].

Strikingly, findings obtained from the various experimental models of the PSMC5-associated NDD systematically guide us to draw parallels with the processes inherent in aging and neurodegeneration. On the one hand, T cells derived from individuals with *PSMC5* variants exhibit characteristics reminiscent of senescent cells. One example is the decreased proteasomal activity in T cells indicated by the accumulation of polyubiquitinated proteins (**Fig. 6A,B**). Proteasomal activity is known to decline with age in humans[39], and its beneficial impact on lifespan has been extensively documented, exemplified by human bone marrow multipotent stromal cells[40] or *Drosophila*[41] models. Also consistent with observations in T cells (**Figs. 6C,7,S6,S7**), age-related cellular traits comprise impaired mitophagy[42], type I IFN response[43], and elevated levels of APOA2[44]. Correspondingly, the elevated prelamin A (LMNA) levels observed in affected individuals (**Fig. S6C**) mirror the excess LMNA resulting from mTOR activation reported in a rat progeria model, where it correlated with premature aging and compromised autophagy[45]. These observations could potentially clarify the usage of phosphorylated Psmc5 (Rpt6) and the accumulation of K48-linked polyubiquitinated proteins as markers to monitor proteasomal activity in the aging brain of another rat model[31]. This raises the question of whether *PSMC5* variants induce premature aging in the individuals studied and whether they play a role in neurodegenerative diseases. Indeed, reduced proteasome activity significantly contributes to the onset and progression of age-related neurodegenerative diseases, such as Alzheimer, Parkinson or Huntington disease[39]. Remarkably, the conditional inactivation of *Psmc1*, a paralog of *Psmc5*, successfully recapitulates Parkinson disease phenotype in both mice[46] and *Drosophila*[47]. Despite the average pediatric age of the *PSMC5* individuals (9.1 years) and the limited follow-up on disease progression, it is noteworthy that three cases of developmental regression have been evoked in the series. On the other hand, the findings made in neuronal or animal models of the disorder also concur with the hypothesis linking *PSMC5* variants to senescence and age-related dementias. Thus, impaired neuritogenesis was reported for instance in an iPSC-derived neuronal model of Parkinson disease[48], cognitive inflexibility was

observed in individuals with Parkinson and Alzheimer diseases[36], and imbalance in E/I ratio was similarly noticed in individuals with Alzheimer disease[49].

To conclude, the combined inputs of T cells, neuronal models, animal studies, and cognitive paradigms contribute to a multi-dimensional understanding of the pivotal role for PSMC5/Rpt6 in neurodevelopment and neuronal function. Our data reveal major overlaps in the molecular pathogenesis of NDD stemming from *PSMC5*, *PSMC3* and *PSMD12* variants which are collectively characterized by protein aggregation (**Fig. 6A, B**), increased mitophagy (**Fig. 6E**) and activation of the ISR leading to sustained type I IFN production (**Figs. S7,8**)[10,11], which, very engagingly, can be mitigated by the use of inhibitors targeting PKR, GCN2, or JAK. These findings lay the foundation for novel diagnostic procedures and forthcoming therapeutic strategies that will primarily target individuals with neurodevelopmental proteasomopathies, but could eventually benefit a broader patient cohort, considering the remarkable similarities observed between the cellular characteristics of individuals with *PSMC5* variants and those with neurodegenerative diseases.

Web resources

Combined Annotation Dependent Depletion (CADD), https://cadd.gs.washington.edu/

dbSNP, http://www.ncbi.nlm.nih.gov/projects/SNP/

DIOPT, https://www.flyrnai.org/cgi-bin/DRSC_orthologs.pl

GeneMatcher, https://genematcher.org/

gnomAD, http://gnomad.broadinstitute.org/

InterVar, https://wintervar.wglab.org/

Metadome, https://stuart.radboudumc.nl/metadome/

Missense Tolerance Ratio Gene Viewer, http://mtr-viewer.mdhs.unimelb.edu.au/

MobiDetails, https://mobidetails.iurc.montp.inserm.fr/MD/

OMIM, http://www.omim.org/

RCSB Protein Data Bank, https://www.rcsb.org/

UCSF ChimeraX, https://www.rbvi.ucsf.edu/chimerax/

Conflict of interest

Acknowledgments

The Deciphering Developmental Disorders (DDD) study presents independent research commissioned by the Health Innovation Challenge Fund (grant number HICF-1009-003). IDB, MDD'A and MKL are contributors to the Care for Rare Solve study led by the Care for Rare Canada Consortium, Children's Hospital of Eastern Ontario Research Institute, Ottawa, ON, Canada.

Funding

For this work, SK awarded research grants from the Agence Nationale de la Recherche (ANR) for the projet ANR-21-CE17-0005 (UPS-NDDecipher) and as a partner of the European Joint Programme on Rare Diseases (EJP RD) for the project ANR-22-RAR4-0001-01 (UPS-NDDiag), and from la Region des Pays de la Loire as part of the National Trajectory program for the project TN_2021_AAP_UPS-NDDECIPHER_INSERM_154550. The EJP RD initiative has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement Number 825575. EK received funding from the German Research Foundation (RTG PRO 2719) and the work was further supported by COST (European Cooperation in Science and Technology) Action ProteoCure CA20113 for EK. The authors are grateful to Robert Beyer and Anne Brandenburg for their excellent technical assistance. FE is a recipient of an I-SITE NExT Junior Talent Chair. SB received financial support from the University Hospital Center (CHU) of Nantes for the BioTND-UPS biobank (PROG/09/72-03). The TND-UPS project led by SB, SK and FE is funded for 3 years

through the patronage of the Mutuelles AXA as part of its health program dedicated to supporting innovative research projects in France.

References

1. Bhattacharyya, S., Yu, H., Mim, C. & Matouschek, A. Regulated protein turnover: snapshots of the proteasome in action. *Nat. Rev. Mol. Cell Biol.* **15**, 122-133 (2014).
2. Hipp, M.S., Kasturi, P. & Hartl, F.U. The proteostasis network and its decline in ageing. *Nat Rev Mol Cell Biol* **20**, 421-435 (2019).
3. Rousseau, A. & Bertolotti, A. Regulation of proteasome assembly and activity in health and disease. *Nat. Rev. Mol. Cell Biol.* **19**, 697-712 (2018).
4. Grochowska, K.M., Andres-Alonso, M., Karpova, A. & Kreutz, M.R. The needs of a synapse-How local organelles serve synaptic proteostasis. *EMBO J* **41**, e110057 (2022).
5. Ehlers, M.D. Activity level controls postsynaptic composition and signaling via the ubiquitin-proteasome system. *Nat Neurosci* **6**, 231-42 (2003).
6. Tai, H.-C. & Schuman, E.M. Ubiquitin, the proteasome and protein degradation in neuronal function and dysfunction. *Nat. Rev. Neurosci.* **9**, 826-838 (2008).
7. van Scheppingen, J. *et al.* Dysregulation of the (immuno)proteasome pathway in malformations of cortical development. *J Neuroinflammation* **13**, 202 (2016).
8. Cuinat, S. *et al.* Understanding neurodevelopmental proteasomopathies as new rare disease entities: a review of current concepts, molecular biomarkers and perspectives. *Genes Dis.* (in press).
9. Küry, S. *et al.* De Novo Disruption of the Proteasome Regulatory Subunit PSMD12 Causes a Syndromic Neurodevelopmental Disorder. *Am. J. Hum. Genet.* **100**, 689 (2017).
10. Isidor, B. *et al.* Stankiewicz-Isidor syndrome: expanding the clinical and molecular phenotype. *Genet Med* **24**, 179-191 (2022).
11. Ebstein, F. *et al.* PSMC3 proteasome subunit variants are associated with neurodevelopmental delay and type I interferon production. *Sci Transl Med* **15**, eabo3189 (2023).
12. Ansar, M. *et al.* Biallelic variants in PSMB1 encoding the proteasome subunit beta6 cause impairment of proteasome function, microcephaly, intellectual disability, developmental delay and short stature. *Hum Mol Genet* (2020).
13. Kroll-Hermi, A. *et al.* Proteasome subunit PSMC3 variants cause neurosensory syndrome combining deafness and cataract due to proteotoxic stress. *EMBO Mol Med* **12**, e11861 (2020).
14. Hamilton, A.M. *et al.* Activity-dependent growth of new dendritic spines is regulated by the proteasome. *Neuron* **74**, 1023-30 (2012).
15. Hsieh, T.C. *et al.* GestaltMatcher facilitates rare disease matching using facial phenotype descriptors. *Nat Genet* **54**, 349-357 (2022).
16. Lesmann, H. *et al.* GestaltMatcher Database - a FAIR database for medical imaging data of rare disorders. *medRxiv*, 2023.06.06.23290887 (2023).
17. Davidson, S. *et al.* Protein kinase R is an innate immune sensor of proteotoxic stress via accumulation of cytoplasmic IL-24. *Sci Immunol* **7**, eabi6763 (2022).
18. Book, A.J. *et al.* The RPN5 subunit of the 26s proteasome is essential for gametogenesis, sporophyte development, and complex assembly in Arabidopsis. *Plant Cell* **21**, 460-78 (2009).
19. Tanaka, K. The proteasome: overview of structure and functions. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* **85**, 12-36 (2009).
20. Kaplanis, J. *et al.* Evidence for 28 genetic disorders discovered by combining healthcare and research data. *Nature* **586**, 757-762 (2020).
21. Bingol, B. *et al.* Autophosphorylated CaMKIIalpha acts as a scaffold to recruit proteasomes to dendritic spines. *Cell* **140**, 567-78 (2010).
22. Jarome, T.J., Ferrara, N.C., Kwapis, J.L. & Helmstetter, F.J. CaMKII regulates proteasome phosphorylation and activity and promotes memory destabilization following retrieval. *Neurobiol Learn Mem* **128**, 103-9 (2016).
23. Torrelo, A. CANDLE Syndrome As a Paradigm of Proteasome-Related Autoinflammation. *Front Immunol* **8**, 927 (2017).

24. Mao, D. *et al.* De novo EIF2AK1 and EIF2AK2 Variants Are Associated with Developmental Delay, Leukoencephalopathy, and Neurologic Decompensation. *Am J Hum Genet* **106**, 570-583 (2020).
25. Roffe, M., Hajj, G.N., Azevedo, H.F., Alves, V.S. & Castilho, B.A. IMPACT is a developmentally regulated protein in neurons that opposes the eukaryotic initiation factor 2alpha kinase GCN2 in the modulation of neurite outgrowth. *J Biol Chem* **288**, 10860-9 (2013).
26. Martinez, N.W., Gomez, F.E. & Matus, S. The Potential Role of Protein Kinase R as a Regulator of Age-Related Neurodegeneration. *Front Aging Neurosci* **13**, 638208 (2021).
27. Anjum, F. *et al.* Dysregulation of ubiquitin-proteasome pathway and apolipoprotein A metabolism in sickle cell disease–related pulmonary arterial hypertension. *Pulmonary Circulation* **3**, 851-855 (2013).
28. Ginsberg, H.N. & Fisher, E.A. The ever-expanding role of degradation in the regulation of apolipoprotein B metabolism. *J Lipid Res* **50 Suppl**, S162-6 (2009).
29. Tierney, E. *et al.* Sterol and lipid analyses identifies hypolipidemia and apolipoprotein disorders in autism associated with adaptive functioning deficits. *Translational Psychiatry* **11**, 471 (2021).
30. Ugur, C. *et al.* The assessment of serum lipid profiles of children with attention deficit hyperactivity disorder. *Psychiatry Research* **264**, 231-235 (2018).
31. Dulka, B.N., Trask, S. & Helmstetter, F.J. Age-Related Memory Impairment and Sex-Specific Alterations in Phosphorylation of the Rpt6 Proteasome Subunit and Polyubiquitination in the Basolateral Amygdala and Medial Prefrontal Cortex. *Front Aging Neurosci* **13**, 656944 (2021).
32. Ohnishi, Y.H. *et al.* PSMC5, a 19S Proteasomal ATPase, Regulates Cocaine Action in the Nucleus Accumbens. *PLoS One* **10**, e0126710 (2015).
33. Aquili, L., Liu, A.W., Shindou, M., Shindou, T. & Wickens, J.R. Behavioral flexibility is increased by optogenetic inhibition of neurons in the nucleus accumbens shell during specific time segments. *Learn Mem* **21**, 223-31 (2014).
34. Zemoura, K. & Benke, D. Proteasomal degradation of gamma-aminobutyric acidB receptors is mediated by the interaction of the GABAB2 C terminus with the proteasomal ATPase Rtp6 and regulated by neuronal activity. *J Biol Chem* **289**, 7738-46 (2014).
35. Beas, B.S., McQuail, J.A., Ban Uelos, C., Setlow, B. & Bizon, J.L. Prefrontal cortical GABAergic signaling and impaired behavioral flexibility in aged F344 rats. *Neuroscience* **345**, 274-286 (2017).
36. Uddin, L.Q. Cognitive and behavioural flexibility: neural mechanisms and clinical considerations. *Nat Rev Neurosci* **22**, 167-179 (2021).
37. Lam, N.H. *et al.* Effects of Altered Excitation-Inhibition Balance on Decision Making in a Cortical Circuit Model. *J Neurosci* **42**, 1035-1053 (2022).
38. Gao, R. & Penzes, P. Common mechanisms of excitatory and inhibitory imbalance in schizophrenia and autism spectrum disorders. *Curr Mol Med* **15**, 146-67 (2015).
39. Saez, I. & Vilchez, D. The Mechanistic Links Between Proteasome Activity, Aging and Age-related Diseases. *Curr Genomics* **15**, 38-51 (2014).
40. Lu, L. *et al.* Ameliorating replicative senescence of human bone marrow stromal cells by PSMB5 overexpression. *Biochem Biophys Res Commun* **443**, 1182-8 (2014).
41. Velentzas, A.D. *et al.* Proteomic mapping of Drosophila transgenic elav.L-GAL4/+ brain as a tool to illuminate neuropathology mechanisms. *Sci Rep* **10**, 5430 (2020).
42. Chen, G., Kroemer, G. & Kepp, O. Mitophagy: An Emerging Role in Aging and Age-Associated Diseases. *Front Cell Dev Biol* **8**, 200 (2020).
43. Baruch, K. *et al.* Aging. Aging-induced type I interferon response at the choroid plexus negatively affects brain function. *Science* **346**, 89-93 (2014).
44. Ma, C., Li, J., Bao, Z., Ruan, Q. & Yu, Z. Serum Levels of ApoA1 and ApoA2 Are Associated with Cognitive Status in Older Men. *Biomed Res Int* **2015**, 481621 (2015).
45. Pan, X. *et al.* Accumulation of prelamin A induces premature aging through mTOR overactivation. *FASEB J* **34**, 7905-7914 (2020).

46.   Bedford, L. *et al.* Depletion of 26S proteasomes in mouse brain neurons causes neurodegeneration and Lewy-like inclusions resembling human pale bodies. *J Neurosci* **28**, 8189-98 (2008).

47.   Fernandez-Cruz, I., Sanchez-Diaz, I., Narvaez-Padilla, V. & Reynaud, E. Rpt2 proteasome subunit reduction causes Parkinson's disease like symptoms in Drosophila. *IBRO Rep* **9**, 65-77 (2020).

48.   Bogetofte, H. *et al.* Post-translational proteomics platform identifies neurite outgrowth impairments in Parkinson's disease GBA-N370S dopamine neurons. *Cell Rep* **42**, 112180 (2023).

49.   van Nifterick, A.M. *et al.* Resting-state oscillations reveal disturbed excitation-inhibition ratio in Alzheimer's disease patients. *Sci Rep* **13**, 7419 (2023).

Figure legends

Figure 1: Structural analysis of *PSMC5* variants reveals most missense variants are located in the AAA+ ATPase domain of PSMC5/Rpt6, and affect proteasome dynamics or subunit interactions. *A.* Schematic representation of the PSMC5 (i.e. RPT6) protein on which the 23 alterations identified in patients with NDD are designated. Approximately half of these mutations fall into three distinct hotspot regions within the AAA+ ATPase domain of PSMC5/RPT6, as indicated. Numbers refer to the sums of unrelated NDD subjects in which the highlighted variant has been identified. Missense variants are colored in blue, frameshift variants in red and splice site variants in purple. *B.* Basing on the structures 6MSK from RCSB Protein Data Bank, we localized the missense variants identified in the AAA+ ATPase subunit PSMC5/Rpt6 within the 26S proteasomal complex. The variants of interest (spheres) are primarily located in the lower ATPase domain. Part of the 26S proteasome was hidden for better visibility; Rpt6 is shown in cartoon representation.

Figure 2: Clinical phenotypes associated with *PSMC5* variants.

A. To visually represent the phenotypic data from tables S2b-2c, we organized the HPO terms into categories and ranked the top categories based on their prevalence within the cohort. Within each category, circles illustrate the number of phenotypes assessed and documented for each individual. The size of a circle corresponds to the number of phenotypes evaluated for a particular subject. If no phenotypes were assessed, there is no circle present. The color of each circle indicates the proportion of these confirmed phenotypes in a subject, with a spectrum ranging from dark blue (indicating the lowest fraction) to red (indicating the highest fraction). The top categories are arranged in descending order according to the highest average fraction observed among subjects. Notably, the most commonly observed categories among these are nervous system and head and neck anomalies. Intriguingly, there was no apparent correlation between genotype and phenotype, as significant variability in phenotypic expression was observed even among individuals with the same genetic variant. *B.* Dysmorphic facial features included notably

abnormal ear morphology as seen in subjects S3/9/10/13/19/30/32-34 (this feature is present in 19/31 (61%) subjects in the whole cohort), abnormal palpebral fissures (13/35; 37% in the whole cohort) including downslanted ones as in subjects S3/10/13/30, thin upper lip vermilion as in subjects S3/34 (7/35; 20%), abnormal palate as in subject S10 (7/35; 20%), tall or broad forehead as in S3/33/34 (9/35; 26%), epicanthus as in S4/38 (5/35; 14%), and orofacial clefts as in subject S10 (2/35; 6%). *C.* Facial image analysis using GestaltMatcher. The pairwise rank matrix and hierarchical clustering of 13 *PSMC5* subjects. Each column is the result of testing one subject in the column and ranking of the remaining 12 photos in each row. For example, by testing the similarities between S10 and the 7,459 images of affected individuals from GMDB, S31 was ranked 1st and S19 was ranked 7th as being most similar to S10. The red box is the cluster of subjects with at least one match below the rank 50, which indicates this cluster shares a similar facial phenotype.

**Figure 3: Suppression of *Psmc5* leads to changes in reversal learning in flies and disrupts the balance between excitatory and inhibitory signals in rat hippocampal neurons. *A.* Pan-neuronal knockdown of Rpt6 in *Drosophila* did not significantly affect normal olfactory learning (p=0.3059, n=4). *B.* When faced with the more difficult task of reversal olfactory learning, knocking down Rpt6 pan-neuronally in *Drosophila* resulted in a significant decrease in performance (p<0.0001, n=4). *(C-J)* Primary rat hippocampal neurons infected DIV1-DIV14 with scrambled shRNA or PMSC5 knockdown (KD) shRNA delivering lenti-particles that co-express GFP. *C.* Assessment of PMSC5 fluorescent signal intensity in neurons at DIV14 shows a significant reduction in cells infected (GFP positive) with *PMSC5* knockdown (KD) compared to cells infected with scrambled shRNA (n=10, Mann Whitney test, p = 0.0002). *D.* Exemplary images of neurons infected with scrambled shRNA (left) or PMSC5 shRNA (right) lenti-particles. PMSC5 labeling (arrow) is significantly reduced, confirming the KD of PMSC5. *E-F.* Infected neurons stained for MAP2 to visualize neuronal morphology. *E.* No significant difference in dendritic branching measured by the mean number of primary, secondary, and tertiary dendrites per neuron at DIV14 was found

comparing scrambled Ctrl and PMSC5 KD conditions (n=6). *F.* Exemplary images of neurons infected with scrambled shRNA (left) or PMSC5 shRNA (right) lenti-particles. MAP2 labeling is shown in red. *G-J.* Infected neurons stained for vGlut to visualize excitatory presynaptic compartments or vGat to visualize inhibitory presynapses. *G.* The number of vGlut positive puncta along the primary dendrites was measured at DIV14. In *Psmc5* KD conditions, a significant reduction was observed compared to scrambled Ctrls (n=5, Mann Whitney test, p = 0.0045). *H.* The number of vGat positive puncta along the primary dendrites was not significantly altered (n=5, Mann Whitney test, p = 0.2988). *I-J.* Exemplary images of primary dendrites of neurons infected with scrambled shRNA (left) or *Psmc5* shRNA (right) lenti-particles. VGlut (full arrow) positive signals per dendrite length are significantly reduced, while VGat (open arrow) positive signals per dendrite length are not altered after *Psmc5* KD.

Figure 4: Overexpression of *PSMC5* affects neuronal morphology.

A. Representative images of primary hippocampal neurons transfected with the empty vector control (EV), *PSMC5*-WT or the different *PSMC5* variants. Red indicates the tdTomato blue indicat MAP2. *B.* Analysis of the total neurite length in the different conditions reveal: (i) a significant increase induced by overexpression of PSMC5-WT compared to empty vector control (One-way ANOVA, F=3.296, p= 0.0014; empty vector versus PSMC5-WT, p=0.0046, Dunnet's multiple comparison test); (ii) a similar effect as PSMC5-WT induced by overexpression of variants p.(Arg201Trp), p.(Pro320His) and p.(Arg325Trp) (PSMC5-WT versus p.(Arg201Trp), p=0.8; PSMC5-WT versus p.(Pro320His), p=0.8; PSMC5-WT versus p.(Arg325Trp), p=0.5, Dunnet's multiple comparison test); (iii) a decrease induced by overexpression of variants p.(Ala202Val), p.(Thr207Met), p.(Pro320Arg) and PSMC5-Δex10, compared to PSMC5-WT (PSMC5-WT versus p.(Ala202Val), p=0.0031; PSMC5-WT versus p.(Thr207Met), p=0.02; PSMC5-WT versus p.(Pro320Arg), p=0.006; PSMC5-WT versus PSMC5-Δex10, p=0.03, Dunnet's multiple comparison test). The effects of these variants were indistinguishable from the empty vector control (empty vector versus p.(Ala202Val), p=0.4; empty vector versus p.(Thr207Met), p=0.9; empty vector versus p.(Pro320Arg), p=0.9;

empty vector versus PSMC5-Δex10, p=0.9, Dunnet's multiple comparison test). *C.* Analysis of the arborization showed no differences between conditions compared to the empty vector control (One-way ANOVA, F=2.179, p= 0.03; empty vector versus PSMC5-WT, p=0.9, PSMC5-WT versus p.(Arg201Trp), p=0.9; PSMC5-WT versus p.(Ala202Val), p=0.9; PSMC5-WT versus p.(Thr207Met), p=0.6; PSMC5-WT versus p.(Pro320His), p=0.3; PSMC5-WT versus p.(Pro320Arg), p=0.9; PSMC5-WT versus p.(Arg 325Trp), p=0.9; PSMC5-WT versus PSMC5-Δex10, p=0.9, Dunnet's multiple comparison test). Error bars indicate SEM, n (number of neurons traced): EV=40, PSMC5-WT=30, p.(Arg201Trp)=18 p.(Ala202Val)=20, p.(Thr207Met)=27, p.(Pro320Arg)=29, p.(Pro320Arg)=19, p.(Arg325Trp)=20 and PSMC5-Δex10=20. *p<0.05; **p<0.01. Scale bar: 100μm.

Figure 5: T cells from NDD subjects carrying *PSMC5* heterozygous variants exhibit abnormalities in their proteasome expression and/or activity profiles. *A.* T cells expanded from PBMC isolated from NDD subjects S6/11/12/21/32 as well as related controls (father and/or mother) and heathy donors (noted 'un. control' for unrelated controls) were lysed in RIPA buffer prior to SDS-PAGE/western-blotting analysis using antibodies specific PSMC5/RPT6 and GAPDH (loading control). Arrow indicates full-length endogenous PSMC5/RPT6 at the expected size of approximately 45 kDa detected in all samples. PSMC5/Rpt6 staining of T cell lysates from subjects 6/11/21/32 uncovered a further shorter PSMC5/Rpt6 immunoreactive band migrating at ~30kDa (denoted as a low-molecular mass (LMM) species by the square brackets) following prolonged exposure times, whereas T cells from subject S12 exhibited an extra PSMC5/Rpt6 signal running a little bit lighter than PSMC5/Rpt6 expected size just below 40 kDa (indicated by an asterisk). Lower panel: the RPT6 and GAPDH immunoreactive bands were quantified by densitometry. Data are presented as RPT6 full-length and low molecular weight (LMW) species on GAPDH ratio mean values ± SD for the control (n=9) and patients (PSMC5 lesions, n=5) groups, as indicated. Statistical analysis was performed using the Mann-Whitney U test where * indicates *p*<0.001. *B.* Resting T cells from NDD subjects S6/11/12/21/32/33 as well as

related (subject's father and/or mother, as indicated) and unrelated (healthy donor) control T cells were analyzed for proteasome function and abundance. They were subjected to non-denaturing protein extraction using TSDG buffer prior to native-PAGE/western-blotting analysis using monoclonal antibodies specific for the α6 and PSMC5/RPT6 subunits, as indicated. Proteasome complexes (20S and 26S) were also visualized by in-gel activity assay using the Suc-LLVY-AMC fluorogenic peptide reflecting chymotrypsin-like activity. Lower panel: the LLVY-AMC fluorescent signals as well as the α6 and RPT6 immunoreactive bands in 30S and/or 26S proteasome complexes were quantified by densitometry and presented as fold change values in patients S6, S11, S12, S21, S32 and S33 versus the control group for each gel whose densitometry measurements were set to 1 (grid line). Columns indicate foldchange mean values ± SD of the patient group (n=6) for LLVY-AMC, α6 and RPT6 in 30S and/or 26S complexes, as indicated. Statistical analysis was performed using the Mann-Whitney U test where * indicates p<0.05.

Figure 6: T cells from NDD subjects with *PSMC5* heterozygous variants are characterized by disrupted protein homeostasis, increased mitophagy rates and a specific lipid signature. *A.* T cells expanded from PBMC isolated from NDD subjects S6/11/12/21/32 were stained with 1 µM of the PROTEOSTAT® dye prior to flow cytometry analysis using the B3 (PerCP-Vio 700) channel. PROTEOSTAT®, a dye that specifically intercalates into the cross-beta spine of quaternary protein structures typically found in misfolded and aggregated proteins. All T cells with PSMC5 variants exhibited increased aggresome formation, as evidenced by increased PROTEOSTAT® fluorescence intensity detected using flow cytometry. Shown are one representative histogram overlay plot of the five T cell NDD subject samples over a control one (left panel), the individual mean fluorescence intensity (MFI) values of the patient (*PSMC5* lesions, n=5) and control (n=5) groups (middle panel) (*p<0.05, Mann-Whitney U test) and the foldchange of the. aggresome propensity factor (APF) for each NDD subject sample compared to the mean value of the five controls (right panel). ***B.*** T cells from NDD subjects S1/6/11/12/21/32, relatives (subject's

father, mother and/or brother) and unrelated controls (healthy donors) were subjected to RIPA-mediated protein extraction and subsequent SDS-PAGE/western-blotting using antibodies directed against ubiquitin, β-actin (loading control) and GAPDH (loading control), as indicated. Right panel: the ubiquitin, β-actin immunoreactive bands were quantified by densitometry and data are presented as ubiquitin/loading control mean values ±SD for both control (n=6) and patient (PSMC5 lesion, n=6) groups, as indicated. Statistical analysis was performed using the Mann-Whitney test with ** indicating p<0.01. *C.* In an attempt to deepen our understanding of the involvement of proteasomes in the regulation of lipid metabolism, we undertook an untargeted lipidomic analysis of T cells from seven unrelated *PSMC5* Subjects (S1, S6, S11, S12, S19, S21 and S33) together with their six relative controls (probands' brother, father and/or mother). Principal component analysis (PCA) plots were first generated to visualize lipid distribution and identify specific patterns across control and patient samples with positive (left panel) and negative (right panel) ion modes. The control and patient groups are presented in red and blue, respectively. *D.* Fold change analysis showing the lipid classes undergoing significant changes in T cells derived from NDD patients when compared to those from healthy donors. *E.* T cells expanded from PBMC isolated from the six NDD subjects S1/6/11/12/21/33 as well as from related and unrelated healthy donors (n=7) were incubated overnight with 100 nM of the Mtphagy dye prior to flow cytometry analysis using the B4 (PE-Vio 770) channel. Shown are the percentage values of the patient (PSMC5 lesions) and control groups (**$p<0.01$, Mann-Whitney U test).

Fig. 7: Gene expression analysis reveals a specific gene signature and a spontaneous type I IFN response in T cells from NDD subjects with *PSMC5* heterozygous variants.
A. The heat map indicates the fold-change of expression of immune-related genes between T cell samples from five subjects with NDD (S1/11/12/21/32) and five heathy individuals (probands' fathers and unrelated donors). Our profiling was based on a panel of 700 immune system genes, using the previously described NanoString nCounter® technology[11]. Differentially expressed genes are hierarchically clustered. Upregulated and downregulated

genes are represented in red and green, respectively. Forty genes were differentially expressed, almost all upregulated. The vast majority of them belonged to the large family of type I interferon (IFN)-stimulated genes (ISGs), including in particular the signal transducers STAT1 and STAT2 and members of the ISG15 conjugation machinery (e.g. UBE2L6 and HERC5). **D.** A comparative expression analysis of seven ISG (*IFIT1*, *IFI27*, *IFI44*, *IFI44L*, *ISG15*, *MX1* and *RSAD2*) was done by quantitative real-time PCR (see Materials & Methods), as previously described [11] between T cells from *PSMC5* subjects S1/11/12/21/32/33, unrelated healthy donors (HD) and subjects' parents. Type I IFN score was determined for each sample by calculating the median of the normalized fold-change values of the seven ISG relative to one control calibrator. Shown are the IFN scores of each group of donor and parental controls (n=8; left panel) and affected individuals (n=6; right panel). (*$p < 0.05$, Mann-Whitney U test). **E.** We evaluated the potential contribution of PKR to the initiation of type I IFN responses in NDD patients with *PSMC5* variants. To this end, T cell from subjects S1/11/12/21/32/33 were exposed to PKR specific inhibitor C16 for 12 h prior to quantification of ISG transcripts by qPCR. T cells expanded from PBMC isolated from NDD subjects were subjected were subjected to an 8-hr treatment with DMSO (vehicle) or the PKR specific inhibitor C16 (3 µM), 4µ8C (100 µM), H-151 (2 µM), baricitinib (1 µM) or A92 (10 µM). Then RNAs were extracted and submitted to RT-qPCR for ISG gene expression analysis. Shown are the median values of the IFN scores calculated for each T cell patient sample (n=6) under different treatment conditions (*$p < 0.05$, Mann-Whitney U test).

Table 1. Main characteristics of the *PSMC5* variants identified in the affected individuals included in the study.

Variant	cDNA change	Protein change	Inheritance	Variant databases	Intervar	Mobidetails	Altered splicing	Subjects
V1	c.166+3del	p.?	*de novo*	-	Pat	212698	+	S1
V2	c.206A>G	p.(Gln69Arg)	*de novo*		Pat	212702	-	S2
V3	c.414G>C	p.(Met138Ile)	*de novo*	-	Pat	212704	-	S3
V4	c.548C>T	p.(Pro183Leu)	assumed *de novo*	-	Pat	212706	-	S4
V5	c.587del	p.(Lys196Argfs*29)	maternally inherited*	-	Pat	306340	-	S5
V6	c.601C>T	p.(Arg201Trp)	assumed *de novo/ de novo*	-	Pat	212708	-	S6, S7 (2)
V7	c.605C>T	p.(Ala202Val)	*de novo*	-	Pat	212710	-	S8
V8	c.620C>T	p.(Thr207Met)	recessive (homozygous)	gnomAD(1); ALFA(1); rs749452987	Pat	212712	-	S9
V9	c.647G>A	p.(Gly216Asp)	*de novo*	-	Pat	212714	-	S10
V10	c.662A>G	p.(Gln221Arg)	assumed *de novo*	-	Pat	212716	+	S11
V11	c.749A>T	p.(Glu250Val)	*de novo*	-	Pat	212718	-	S12
V12	c.749A>G	p.(Glu250Gly)	*de novo*	-	Pat	212720	-	S13
V13	c.754G>A	p.(Asp252Asn)	*de novo*	-	Pat	212722	-	S14, S15 (2)
V14	c.772C>T	p.(Arg258Trp)	*de novo*	gnomAD(2); rs11543211	Pat	212724	-	S16
V15	c.773G>A	p.(Arg258Gln)	maternally inherited*	rs781522827, but no MAF	Pat	274699	-	S17
V16	c.845G>A	p.(Gly282Asp)	*de novo*	-	Pat	307153	-	S18
V17	c.959C>G	p.(Pro320Arg)	*de novo*	-	Pat	212726	-	S19-S25 (7)
V18	c.959C>A	p.(Pro320His)	assumed *de novo*	BRAVO (1) ; rs1349754006	Pat	212728	-	S26
V19	c.968_969+1del	p.(Glu323Glyfs*26)	*de novo*	-	Pat	307156	+	S27
V20	c.970-2A>G	p.?	assumed *de novo*	-	Pat	212730	+	S28
V21	c.973C>T	p.(Arg325Trp)	*de novo*	-	Pat	212732	-	S29-S36 (8)
V22	c.1103T>C	p.(Met368Thr)	*de novo*	gnomAD (1)	Pat	212734	-	S37
V23	c.1182_1183del	p.(Asp394Glufs*2)	*de novo*	-	Pat	212736	-	S38

RefSeq transcript used for PSMC5 is NM_002805.6. *inherited from an affected mother
Variant database used: gnomAD v2.1.1/v3.1.2, dbSNP v154, Clinvar, ALFA v20201027095038, BRAVO/TOPmed data freeze 8; MAF: minor allele frequency;
tools used for assessing effect on splicing: SPIP, SpliceSiteFinder, MaxEntScan, NNSplice, GeneSplicer; Intervar: Pat= pathogenic
The access to detailed predictions for variant XXXXXX (e.g. 212698) using Mobidetails is as follows: https://mobidetails.iurc.montp.inserm.fr/MD/variant/XXXXXX

Table 2. Clinical features of the subjects with PSMC5 variants and indels (1/2).

Subject	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12	S13	S14	S15	S16	S17	S18	S19	S20
Variant	V1	V2	V3	V4	V5	V6	V6	V7	V8	V9	V10	V11	V12	V13	V13	V14	V15	V16	V17	V17
Sexr	M	M	M	M	F	F	F	F	M	M	F	F	F	M	F	M	M	M	M	F
Age (in years)	0-5	0-5	0-5	0-5	0-5	6-10	0-5	16-20	11-15	0-5	11-15	6-10	16-20	11-15	0-5	6-10	26-30	11-15	0-5	0-5
Neurodevelopmental abnormality	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Speech impairment (delay, absence)	+	+	+	-	+	+	+	+	ND	+	+	+	+	+	+	-	+	+	+	+
Abnormal facial shape	+	+	+	-	+	-	+	+	+	+	-	-	+	+	+	ND	-	-	+	+
Abnormal muscle tone	+	-	-	-	+	+	+	+	-	+	-	-	+	+	+	ND	-	-	+	+
Motor delay/impairment	+	-	+	+	+	+	+	+	+	+	-	+	+	-	+	ND	-	-	+	+
Ophthalmological findings	+	-	-	+	-	+	-	+	+	+	-	+	+	+	+	ND	ND	-	+	-
Intellectual disability	-	ND	ND	+	+	+	ND	-	-	ND	+	+	+	+	+	-	-	+	ND	ND
Abnormal behavior	-	-	+	+	+	+	-	+	+	-	+	-	-	+	-	+	+	+	+	-
Brain MRI abnormalities	-	ND	-	ND	ND	-	-	+	ND	+	ND	ND	+	+	-	ND	-	-	+	ND
Abnormal ear morphology	-	+	+	-	+	-	+	ND	+	+	-	-	+	+	-	ND	-	-	+	-
Other findings	-	-	+	+	+	+	+	+	-	-	+	-	-	+	+	+	+	-	-	-
Skeletal abnormalities (not involving skull)	-	-	+	-	+	-	+	+	+	-	-	-	-	+	-	+	-	-	-	-
Feeding difficulties	-	+	-	+	+	+	+	-	-	+	-	+	-	+	-	-	-	-	+	+
Abnormal weight	+	+	-	+	+	+	+	-	+	-	+	-	+	+	+	-	-	-	-	+
Abnormal length	+	+	+	+	+	-	+	-	+	-	+	-	+	-	-	-	-	-	-	+
Abnormal OFC	+	-	-	+	+	-	+	+	-	-	+	-	+	-	-	-	-	-	-	+
Other malformations	-	-	-	-	+	+	-	-	+	-	-	-	+	+	-	ND	-	+	+	-
Retrognathia/microretrognathia	-	+	+	-	-	-	-	ND	-	+	-	-	+	+	+	ND	-	-	-	-
Cardiac abnormalities	-	+	-	-	-	-	-	-	-	-	-	+	+	-	-	-	+	-	+	-
Seizures	+	-	-	-	-	+	-	-	-	-	-	-	+	-	+	-	+	+	-	-
Hearing loss	-	-	-	-	-	-	-	+	+	-	-	ND	-	+	-	-	-	-	-	-
Renal abnormalities	-	-	-	-	-	+	-	-	-	-	-	-	-	-	+	-	-	-	-	-
Genital abnormalities	-	-	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+
Developmental regression	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+	-

ND: not determined; OFC: occipital-frontal head circumference; MRI: magnetic resonance imaging. Detailed clinical findings are reported in Table S2a.

Table 2. Clinical features of the subjects with PSMC5 variants and indels (2/2).

Subject	S21	S22	S23	S24	S25	S26	S27	S28	S29	S30	S31	S32	S33	S34	S35	S36	S37	S38	Total
Variant	V17	V17	V17	V17	V17	V18	V19	V20	V21	V21	V21	V21	V21	V21	V21	V21	V22	V23	
Sex	M	M	M	M	F	F	M	M	M	F	M	M	M	F	M	M	M	F	24M14F
Age (in years)	6-10	16-20	0-5	0-5	0-5	6-10	6-10	11-15	0-5	11-15	6-10	16-20	6-10	0-5	11-15	0-5	6-10	0-5	0-5 to 26-30
Neurodevelopmental abnormality	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	38/38 (100%)
Speech impairment (delay, absence)	+	+	+	+	+	+	+	+	ND	+	+	+	+	+	ND	+	+	+	33/35 (94%)
Abnormal facial shape	ND	+	ND	+	+	+	+	+	+	+	+	+	+	+	-	+	+	+	28/35 (80%)
Abnormal muscle tone	+	ND	ND	+	+	+	+	+	+	+	+	+	+	+	ND	+	+	+	26/34 (77%)
Motor delay/impairment	+	+	+	+	+	-	-	+	+	+	+	+	-	+	ND	+	+	-	27/36 (75%)
Ophthalmological findings	+	ND	ND	ND	+	+	+	+	+	+	-	ND	+	+	+	+	+	+	24/32 (75%)
Intellectual disability	+	+	ND	ND	ND	+	-	+	ND	+	+	+	-	ND	+	ND	+	+	19/26 (73%)
Abnormal behavior	+	ND	ND	+	-	+	+	+	-	+	+	+	+	-	ND	+	ND	+	23/34 (68%)
Brain MRI abnormalities	ND	ND	ND	ND	+	ND	+	+	ND	ND	+	+	+	+	ND	+	+	-	14/22 (63%)
Abnormal ear morphology	ND	ND	ND	+	+	ND	-	+	+	+	+	+	-	+	ND	+	+	-	19/31 (61%)
Other findings	-	-	-	+	+	+	+	-	-	-	+	+	+	+	+	+	+	-	22/38 (58%)
Skeletal abnormalities (not involving skull)	+	+	+	+	-	+	+	+	+	+	+	+	+	-	-	-	+	+	21/38 (55%)
Feeding difficulties	-	-	-	+	-	-	-	-	+	+	+	+	+	+	ND	+	+	+	20/37 (54%)
Abnormal weight	-	ND	ND	-	-	-	-	+	+	ND	ND	-	+	-	ND	-	+	-	16/33 (48%)
Abnormal length	-	ND	ND	-	-	-	-	+	-	ND	-	+	+	-	+	-	+	-	15/35 (43%)
Abnormal OFC	-	ND	ND	-	-	-	-	+	+	ND	-	-	+	+	ND	+	-	-	13/34 (38%)
Other malformations	+	-	-	+	-	-	+	-	+	-	-	-	+	-	+	+	-	-	14/37 (38%)
Retrognathia/microretrognathia	-	+	-	-	-	+	-	-	-	+	-	+	-	-	ND	-	-	+	11/35 (31%)
Cardiac abnormalities	-	-	-	+	-	-	-	+	+	-	-	+	+	+	-	-	-	-	11/38 (29%)
Seizures	-	-	-	-	-	-	-	-	-	-	+	+	+	-	ND	-	+	-	10/37 (27%)
Hearing loss	-	-	-	ND	+	-	-	-	ND	+	+	-	+	+	+	-	-	-	9/35 (26%)
Renal abnormalities	-	-	-	-	+	-	ND	-	-	+	-	-	-	-	-	-	+	-	5/37 (14%)
Genital abnormalities	-	-	-	-	-	-	-	-	-	-	+	-	+	-	-	-	-	-	5/38 (13%)
Developmental regression	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	3/38 (8%)

12   ND: not determined; OFC: occipital-frontal head circumference; MRI: magnetic resonance imaging. Detailed clinical findings are reported in Table S2a.

A

Fraction of phenotypes present in subjects

50%
33%
25%
10%
0%

Phenotypes assessed

Subject: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38

Nervous system
Head or neck
Eye
Cardiovascular system
Ear
Skeletal system
Limbs
Integument
Growth
Digestive system
Genitourinary system

B

The photos have been removed from this version of the manuscript. However, they can be viewed in the GestaltMatcher database, using the following links (requires login to GMDB database):

S1: https://db.gestaltmatcher.org/patients/9968
S3: https://db.gestaltmatcher.org/patients/9969
S9: https://db.gestaltmatcher.org/patients/9970
S10: https://db.gestaltmatcher.org/patients/9971
S12: https://db.gestaltmatcher.org/patients/9972
S13: https://db.gestaltmatcher.org/patients/9973
S19: https://db.gestaltmatcher.org/patients/9974
S30: https://db.gestaltmatcher.org/patients/9975
S31: https://db.gestaltmatcher.org/patients/9976
S32: https://db.gestaltmatcher.org/patients/9977
S33: https://db.gestaltmatcher.org/patients/9978
S34: https://db.gestaltmatcher.org/patients/9979
S38: https://db.gestaltmatcher.org/patients/9980

C

PSMC5 pairwise rank