Trajectories in county-level Low Birthweight Rates and Associated Contextual Factors

in the United States, 2016-2021

AUTHORS: Pallavi Dwivedi, MS, MPH[a], Thu T. Nguyen, ScD MSPH[a], Xin He, PhD[a], Quynh C. Nguyen, PhD[a]

AFFILIATIONS: [a] Department of Epidemiology and Biostatistics, University of Maryland School of Public Health, College Park, Maryland, United States

CORRESPONDENCE TO: Pallavi Dwivedi, Department of Epidemiology and Biostatistics, University of Maryland School of Public Health, College Park, Maryland

Email: pdwived1@umd.edu

Manuscript word count: 3082   Abstract word count: 237

Tables: 4   Figures: 1

eTables: 1   eFigures: 1

Conflict of Interest statement: The authors have no conflict of interest to disclose.

ABSTRACT

Introduction

Infants with low birthweight (less than 2500 grams) have greater risk of mortality, long-term neurologic disability and chronic diseases such as diabetes and cardiovascular disease as compared to infants with normal birthweight. This study examined the trajectories of low birthweight rate in the U.S. across the metropolitan and non-metropolitan counties over the time period of 2016-2021 and the associated contextual factors.

Methods

This longitudinal study utilized data on 21,759,834 singleton births across 3,108 counties. Data on birthweight and maternal sociodemographic and behavioral characteristics was obtained from the National Center for Health Statistics. A generalized estimating equations model was used to examine the association of county-level contextual variables with low birthweight rates.

Results

A significant increase in low birthweight rates was observed across the counties over the duration of the study. Large metro and small metro counties had significantly higher low birthweight rates as compared to non-metro counties. High percentage of Black women, underweight women, age more than 35 years, lack of prenatal care, uninsured population, and

high violent crime rate was associated with an increase in low-birth-weight rates. Other contextual characteristics (percentage of married women, American Indian/Alaskan Native women, and unemployed population) differed in their associations with low birthweight rates depending on county metropolitan status.

Conclusions

Our study findings emphasize the importance of developing interventions to address geographical heterogeneity in low birthweight burden, particularly for metropolitan areas and communities with vulnerable racial/ethnic and socioeconomic groups.

Keywords

Low Birthweight, contextual factors, heterogeneity, maternal factors

INTRODUCTION

In comparison with normal birthweight infants, neonates with low birthweight (< 2500 grams at birth) have higher risk of mortality, and chronic diseases such as cardiovascular disease and diabetes.[1,2,3,4] The determinants of low birthweight include multiple gestation or carrying more than one fetus, obstetric complications, infections, smoking, drinking alcohol, drug abuse, stress, undernutrition and chronic maternal conditions such as hypertension.[2,4,5,6] Prenatal exposure to air pollution has also been shown to be associated with low birthweight.[2] Previous studies have also suggested that racial discrimination is associated with increased risk of preterm and low birthweight.[7,8] The low birthweight rate for Black people has been more than twice that for White people from 2006-2016.[9]

The prevalence of low birthweight in the U.S. was estimated to be 8.3% in 2018 by National Center for Health Statistics.[10] The Healthy People 2020 goal was to reduce the rate of low birthweight to 7.8%.[11] While several previous studies have examined the maternal factors, which are associated with low birthweight of infants, none of these studies examined the patterns of low birthweight rates based on the metropolitan status of counties across the U.S.. In this study, we examined whether there is any variation in the rates of low birthweight rates based on the metropolitan status of U.S. counties. In addition, we examined whether the effect of certain county-level contextual factors such as socioeconomic disadvantage, and maternal factors such as marital status and race or ethnicity on low birthweight rates vary based on the metropolitan status of counties. We also examined the distribution of low birthweight risk across the counties. Examination of geographical variation in low birthweight rates and

associated contextual factors can help to inform interventions and policies and reduce cases of

low birthweight.

Study aims and hypotheses.

The aim of this study was to examine the trajectories of low birthweight rate at the county

scale based on the metropolitan status of counties and the association of county-level

contextual factors with low birthweight rates over the time period of 2016-2021. We

hypothesized that there is variation in low birthweight rates based on the metropolitan status

of counties and certain county-level contextual factors such as socioeconomic disadvantage,

violent crime rate, and maternal factors such as marital status and race or ethnicity might

influence the disparities in low birthweight rate across the counties.

METHODS

County-level health data

This longitudinal study utilized data on 21,759,834 singleton births across 3,108 counties for the

time period of 2016-2021. Data on birthweight and maternal characteristics (e.g., live birth

order, race or ethnicity, age, marital status, prenatal care, body mass index, cigarette smoking

status during first, second or third trimester of pregnancy, maternal education) were obtained

from the restricted natality files with geographical identifiers from the National Center for

Health Statistics.[12] Data on county-level time-varying contextual variables, such as percentage

of unemployed population, percentage of uninsured population, violent crime rate per 100,000

population, and primary care provider rate per 100,000 population were obtained from the

County Health Rankings data.[13]

Measures

Counts of low birthweight cases were aggregated by county in two-year study periods (2016-

2017, 2018-2019 and 2020-2021) to enable stabilization of low birthweight rates. Similarly, data

on county-level time-varying contextual variables, such as percentage of unemployed

population, percentage of uninsured population, violent crime rate per 100,000 population,

primary care provider rate per 100, 000 population were aggregated for the two-year study

periods. Data on maternal characteristics (live birth order, race or ethnicity, age, marital status,

prenatal care, body mass index, cigarette smoking status during first, second or third trimester

of pregnancy, maternal education) was also aggregated for the two-year study periods. The

National Center for Health Statistics (NCHS) Urban–Rural Classification Scheme for counties was

used to classify counties.[14] The NCHS scheme is a 6-category classification scheme based on

county population. The six categories were combined into 3 categories, and the counties were

classified as large metropolitan counties (metro areas of 250,000 population or more), small

metropolitan counties (metro areas of fewer than 250,000 population) and non-metropolitan

counties.

Statistical Analysis

The geographic distribution of low birthweight rates for each two-year study period was

examined by calculating the Standardized Incidence Ratios (SIRs). The SIRs can be calculated as

the ratio of observed to expected count of low birthweight cases for each county. The state-

level low birthweight rates were used to calculate county-level expected count of low

birthweight cases by multiplying state-level low birthweight rates with total number of births

per county. The county level observed, or raw rates of low birthweight can be misleading,

particularly if the data are sparse due to small population size. Therefore, a Bayesian

hierarchical model was used to compute smoothed SIRs and exceedance probability of SIR

being greater than 1 for each county during each study period.[15,16] The Bayesian model

comprised of a spatial random effect term, which incorporates the influence of neighboring

counties, and a non-spatial or unstructured variability term. The spatial random effect term can

be defined as a conditional autoregressive Gaussian process with mean equal to the mean of

the spatial effects of the neighboring geographic units and precision (τ_v). The non-spatial

random effect term was normally distributed with mean equal to zero and precision (τ_u).

Gamma (0.5, 0.0005) priors were assigned to the precision terms in the model. The R-INLA

package in R was utilized to conduct Bayesian inference by using the integrated nested Laplace

approximation (INLA) method.[17,18.]

A generalized estimating equations (GEE) model with a log link function was used to examine

the association of county-level contextual variables with low birthweight rates from 2016-2021.

The variances of low birthweight cases within each type of county classification (non-

metro/small-metro/large-metro) was higher than the mean low birthweight cases, which

suggested the presence of overdispersion in the data (eTable 1). Therefore, negative binomial

regression was utilized, and the log of the total number of births per county was used as an

offset variable to allow for the interpretation as low birthweight rates. A random effect term

was included to adjust for the within-county correlations among repeated measures during

three measurement periods (2016-2017, 2018-2019 and 2020-2021). We hypothesized that

metropolitan status of counties moderates or modifies the effect of socioeconomic status of

women on low birthweight rates. Therefore, interaction between metropolitan status and

county-level percentage of uninsured population and percentage of unemployed population

were examined. We also examined the interaction between metropolitan status and

measurement period, maternal race and ethnicity, and marital status of women. The variables

such as the county-level percentage of women with live birth order more than one, women

within age range of 35-54 years, women who did not receive any prenatal care, underweight

women based on their Body Mass Index (BMI), women who smoked cigarette during

pregnancy, women with education more than high school, county-level primary care provider

rate (per 100,000 population) and violent crime rate per 100,000 population were included as

control variables in the model. This study was determined exempt by the University of

Maryland College Park Institutional Review Board (1797788-1).

RESULTS

There were a total of 1,444,384 low birthweight cases, out of 21,759,834 singleton births, over

the duration of 2016-2021. The percentage of low birthweight cases was similar over the three

study periods (486,893 [6.49%] in 2016-2017; 481,152 [6.63%] in 2018-2019; 476,339 [6.8%] in

2020-2021). The median low birthweight rate was similar over the three study periods (~ 4 per

100 births). Large metropolitan counties accounted for the greatest number of low birthweight

cases (1,201,200 [83.16%]), followed by small metropolitan counties (142,938 [9.90%]) and

non-metropolitan counties (100,246 [6.94%]).

Geographic distribution of low-birth-weight rates

Figure 1 presents the smoothed Standardized Incidence Ratio (SIRs) of low birthweight and

Bayesian exceedance probability for SIR greater than 1 for the three study periods. The county-

level SIRs ranged from 0.23 to 13.68, 0.24 to 9.89 and 0.18 to 12.45 in 2016-2017, 2018-2019

and 2020-2021, respectively. The median (interquartile range) for SIR for the study periods of

2016-2017, 2018-2019 and 2020-2021 were 0.76 [0.57-0.98], 0.74 [0.56-0.98] and 0.68 [0.50-

1.01] respectively. SIRs greater than 1 indicate higher than expected risk of low birthweight,

and SIRs less than 1 represent lower than expected risk of low birthweight. We observed an

elevated risk of low birthweight in several counties in California, Arizona, Oregon, Washington,

Idaho, Montana, Wyoming, Colorado, New Mexico, Texas and in southern, midwestern and

northeastern states over the three study periods (Figure 1).

Contextual variables associated with low birthweight rates.

GEE model was used to examine the association of county-level contextual variables with low

birthweight rates (Table 1). We examined whether metropolitan status of counties moderates

the effect of socioeconomic status, marital status and racial group of women on low

birthweight rates. Therefore, we examined interactions between contextual variables such as

percentage of uninsured population, unemployed population, maternal racial group, marital

status and county type (large metropolitan, small metropolitan and non-metropolitan). We also

examined interaction between county type and study period. The interactions between county

type and county-level percentage of American Indian/Alaskan native women, percentage of

married women and unemployed population were significant. Stratified models based on the

county type or metropolitan status were used to examine the effect of county-level percentage

of married women, American Indian/Alaskan Native women and unemployed population on

low-birth-weight rates after adjusting for all the covariates (Tables 2 - 4). County-level

percentage of women with live birth order of more than one, women within age range of 35-54

years, underweight women based on BMI, women with education more than high school,

women who smoked cigarette during pregnancy, women who did not receive any prenatal care

and county-level violent crime rate per 100,000 population and primary care provider rate (per

100,000 population) were included as control variables in the model.

Table 1 presents the incidence rate ratios (IRR) for low birthweight for each predictor variable

after adjusting for all the other covariates. All predictor variables were standardized to have a

mean of 0 and standard deviation of 1. After adjusting for all the covariates, one standard

deviation increase in the county-level percentage of Black women (incidence rate ratio, IRR =

1.22, 95% CI = 1.20, 1.24) was associated with higher low birthweight rates. One standard

deviation increase in the percentage of women who did not obtain any prenatal care, women

who were underweight based on their BMI and women who smoked cigarette during

pregnancy was associated with increased low birthweight rates (IRR for no prenatal care =1.27,

95% CI = 1.23, 1.32; IRR for underweight based on BMI = 1.08, 95% CI = 1.03, 1.13; IRR for

smoking during pregnancy= 1.24 , 95% CI = 1.21, 1.28 respectively). An increase in the

percentage of women of age 35-54 years was associated with an increase in low birthweight

rates (IRR = 1.06, 95% CI = 1.01, 1.11). One standard deviation increase in the percentage of

women with live birth order more than one was associated with reduced low birthweight rates

(IRR = 0.89, 95% CI = 0.85, 0.93). One standard deviation increase in the county-level

percentage of uninsured women was associated with a significant increase in low birthweight

rates (IRR = 1.10, 95% CI = 1.09, 1.12). An increase in the county-scale violent crime rate was

associated with higher low birthweight rate (IRR =1.07, 95% CI = 1.06, 1.09). One standard

deviation increase in the county-level primary care provider rate was associated with a

significant increase in low birthweight rates (IRR = 1.09, 95% CI = 1.07, 1.11).

We did not observe any significant interaction between metropolitan status or county type and

study periods. Compared to the baseline study period of 2016-2017, there was a significant

increase in low birthweight rates in the study periods of 2018-2019 and 2020-2021 in all the

counties after adjusting for all the covariates (2018-2019: IRR = 1.11, 95% CI = 1.09, 1.13; 2020-

2021: IRR = 1.15, 95% CI = 1.13, 1.18). Large metropolitan counties and small metropolitan

counties had higher low birthweight rates as compared to non-metropolitan counties over the

study duration after adjusting for all the covariates (Large metro: IRR = 1.24, 95% CI = 1.20,

1.29; Small metro: IRR = 1.21, 95 % CI =1.16, 1.27). eFigure 1 presents the trajectory of low

birthweight rate for large metropolitan, small metropolitan and non-metro counties over the

duration of the study.

An increase in the county-level percentage of married women was associated with reduced low

birthweight rates in non-metropolitan and large metropolitan counties after adjusting for all

the covariates (Table 2: Non-metro: IRR = 0.90, 95% CI =0.86, 0.94; Large metro: IRR = 0.98, 95%

CI = 0.96, 0.99). One standard deviation increase in the county scale percentage of unemployed

population was associated with higher low birthweight rate in non-metropolitan counties

(Table 3: IRR = 1.04, 95% CI = 1.02, 1.06). Table 4 presents the incidence rate ratio of low

birthweight for one standard deviation increase in county scale percentage of American

Indian/Alaskan Native women based on the metropolitan status of counties. An increase in the

percentage of American Indian/Alaskan native women was associated with higher low

birthweight rates in small metropolitan counties (IRR = 1.08, 95% CI = 1.02, 1.15) and reduced

low birthweight rates in non-metro counties (IRR = 0.97, 95% CI = 0.95, 0.98).

DISCUSSION

This study examined county-level low birthweight rate trajectories in the U.S. during the 6-year

period from 2016 to 2021 across the metropolitan and non-metropolitan counties.

There was a significant increase in low birthweight rates in all the counties over the duration of

the study. Large metropolitan and small metropolitan counties had significantly higher low

birthweight rates as compared to non-metropolitan counties over the duration of the study.

These differences in low birthweight rates based on the metropolitan status of counties might

be attributed to demographic or racial and socioeconomic disparities across these

counties.[19,20,21] Black race was associated with high low birthweight rates. We also observed a

significant increase in low birthweight rates with an increase in county level violent crime rate

(per 100,000 population). A study conducted by Messer et al.[22] reported that Black women are

more likely to live in disadvantaged neighborhoods, or neighborhoods with high poverty,

unemployment, and violent crime rate as compared to White women. Additionally, the authors

suggested that Black women living in racially heterogenous areas with high crime rates might

be exposed to stress resulting from racial discrimination. Several other studies have suggested a

positive association between stress and adverse birth outcomes.[23,24]

An increase in the county-level percentage of married women was significantly associated with

reduced low birthweight rates in non-metro and large metropolitan counties. Several studies

have suggested that unmarried women are more likely to have poor socioeconomic

conditions.[25,26] A study conducted by Taffel[27] found that unmarried Black women are more

likely to give birth to a low birthweight infant as compared to unmarried White women. Studies

conducted by Rubin[28] and Alhusen et al. [29] found that maternal sociodemographic

characteristics, health behavior and prenatal care do not adequately account for disparities in

birth outcomes among Black and White women. An increase in the county-level percentage of

the unemployed population was significantly associated with higher low birthweight rates in

non-metropolitan counties. As study conducted by Hargraves[30] suggested that women living in

rural or non-metropolitan counties are affected by inequalities and biases based on gender,

reduced transportation and employment with lower salaries and inadequate health insurance.

The women of racial and ethnic minority groups are more likely to experience these limitations

in rural counties as compared to White women.

An increase in the county-level percentage of women who did not receive any prenatal care

and the uninsured population was significantly associated with high low birthweight rates. A

previous study, conducted by Taylor et al., [31] suggested that women with Medicaid or no

insurance are less likely to seek preventive care and have adverse birth outcomes such as

preterm birth, preeclampsia and low birthweight as compared to women with commercial

insurance. High percentage of underweight women based on their BMI and women who

smoked cigarette during first, second or third trimester of pregnancy was associated with high

low birthweight rates. These findings are consistent with the findings from the study conducted

by Gunther et al., [32] which reported that exposure to nicotine or cigarette smoking during

pregnancy are associated with adverse birth outcomes. While maternal age within the range of

35-54 years was associated with higher low birthweight rate, an increase in the percentage of

women with previous live births (live birth order >1) was associated with reduced low

birthweight rate. Few previous studies have reported similar effect of maternal age and live

birth order on birth weight of infants.[33,34.]

An increase in the county-level percentage of American Indian/Alaskan Native women was

significantly associated with an increase in low birthweight rate in small metropolitan counties

and a decrease in low birthweight rate in non-metropolitan counties. American Indian/Alaskan

Native racial group has been reported to be at increasing rates of socioeconomic disadvantage

as compared to White women. Several other studies have also reported disparities in birth

outcomes between the American Indian/Alaskan Native women and White women and other

racial groups.[35,36] A study conducted by Call et al. [37] reported that American Indian/Alaskan

Native racial group experiences barriers to healthcare due to racial discrimination and biases

based on their religious beliefs by health care providers. Though access to healthcare has

improved among this subpopulation due to Indian Health Service (IHS), disparities in birth

outcomes exist in this racial group due to reduced availability of IHS in urban areas.[38,39]

CONCLUSIONS

Low birthweight infants have a higher risk of mortality and chronic disorders as compared to

infants with normal birthweight. This study identified potential risk factors associated with low

birthweight and the variation in the low birthweight rates based on the metropolitan status of

counties. Based on our study findings, maternal factors such as racial or ethnic group, BMI, age,

cigarette smoking during pregnancy, socioeconomic status, marital status, prenatal care, health

insurance and county-scale violent crime rate are significant predictors of low birthweight rate

at the county level. We also observed a significant increase in low birthweight rates over the

duration of the study. Large metropolitan and small metropolitan counties had high rates of

low birthweight as compared to non-metropolitan counties over the duration of the study. Our

study findings emphasize the importance of developing interventions tailored specifically for

high risk or vulnerable racial or ethnic and socioeconomic groups to reduce low birthweight

rates. In addition, we examined geographical distribution of low birthweight risk across the

counties and find heterogeneity which suggests additional resources might be warranted to

further support areas at higher risk of adverse birth outcomes.

Funding

Research reported in this publication was supported by the National Institute on Minority Health and Health Disparities [R00MD012615 (TTN), R01MD015716 (TTN), R01MD016037 (QCN], the National Library of Medicine [R01LM012849 (QCN)]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

1. Wardlaw, T. M. (2004). *Low Birthweight: Country, Regional and Global Estimates*. UNICEF.

2. Cutland, C. L., Lackritz, E. M., Mallett-Moore, T., Bardají, A., Chandrasekaran, R., Lahariya, C., Nisar, M. I., Tapia, M. D., Pathirana, J., Kochhar, S., Muñoz, F. M., & Brighton Collaboration Low Birth Weight Working Group. (2017). Low birth weight: Case definition & guidelines for data collection, analysis, and presentation of maternal immunization safety data. *Vaccine*, *35*(48 Pt A), 6492–6500.

3. Malin, G. L., Morris, R. K., Riley, R., Teune, M. J., & Khan, K. S. (2014). When is birthweight at term abnormally low? A systematic review and meta-analysis of the association and predictive ability of current birthweight standards for neonatal outcomes. In *BJOG: An International Journal of Obstetrics & Gynaecology* (Vol. 121, Issue 5, pp. 515–526). https://doi.org/10.1111/1471-0528.12517

4. Blencowe, H., Krasevec, J., de Onis, M., Black, R. E., An, X., Stevens, G. A., Borghi, E., Hayashi, C., Estevez, D., Cegolon, L., Shiekh, S., Ponce Hardy, V., Lawn, J. E., & Cousens, S. (2019). National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *The Lancet. Global Health*, *7*(7), e849–e860.

5. Ratnasiri, A. W. G., Parry, S. S., Arief, V. N., DeLacy, I. H., Halliday, L. A., DiLibero, R. J., & Basford, K. E. (2018). Recent trends, risk factors, and disparities in low

birth weight in California, 2005–2014: a retrospective study. *Maternal Health,*

Neonatology and Perinatology, 4(1), 15.

6. Ashdown-Lambert, J. R. (2005). A review of low birth weight: predictors, precursors and

morbidity outcomes. *The Journal of the Royal Society for the Promotion of*

Health, 125(2), 76–83.

7. Mustillo, S., Krieger, N., Gunderson, E. P., Sidney, S., McCreath, H., & Kiefe, C. I. (2004).

Self-Reported Experiences of Racial Discrimination and Black–White Differences in

Preterm and Low-Birthweight Deliveries: The CARDIA Study. *American Journal of Public*

Health, 94(12), 2125–2131.

8. Collins, J. W., Jr, David, R. J., Handler, A., Wall, S., & Andes, S. (2004). Very

low birthweight in African American infants: the role of maternal exposure

to interpersonal racial discrimination. *American Journal of Public Health, 94*(12),

2132–2138.

9. Womack, L. S., Rossen, L. M., & Martin, J. A. (2018). *Singleton Low birthweight rates,*

by race and hispanic origin: United States, 2006--2016.

https://stacks.cdc.gov/view/cdc/52611

10. Martin, J. A., Hamilton, B. E., Osterman, M. J., & Driscoll, A. K. (2019). Births: Final

data for 2018.

11. US Department of Health and Human Services. (2012). Maternal, infant, and child

health. Healthy People 2020.

12. National Center for Health Statistics. Restricted-Use Vital Statistics Data. (2021).

13. County Health Rankings data. (2016-2021).

https://www.countyhealthrankings.org/explore-health-rankings/rankings-data-

documentation

14. National Center for Health Statistics (NCHS) Urban-Rural Classification Scheme for

counties. (2013). https://www.cdc.gov/nchs/data_access/urban_rural.htm

15. Besag, J., York, J., & Mollié, A. (1991). Bayesian image restoration, with two applications

in spatial statistics. *Annals of the Institute of Statistical Mathematics*, *43*(1), 1–20.

16. Richardson, S., Thomson, A., Best, N., & Elliott, P. (2004). Interpreting Posterior Relative

Risk Estimates in Disease-Mapping Studies. *Environmental Health Perspectives* (Vol. 112,

Issue 9, pp. 1016–1025). https://doi.org/10.1289/ehp.6740

17. Rue, H., Martino, S., Lindgren, F., Simpson, D., Riebler, A., & Krainski, E. T. (2014).

INLA: Functions which allow to perform full Bayesian analysis of latent

Gaussian models using Integrated Nested Laplace Approximaxion. *R Package Version 0. 0-*

1404466487, http://www. R-INLA.org.

18. Blangiardo, M., Cameletti, M., Baio, G., & Rue, H. (2013). Spatial and

spatio-temporal models with R-INLA. *Spatial and Spatio-temporal Epidemiology*

(Vol. 4, pp. 33–49). https://doi.org/10.1016/j.sste.2012.12.001

19. Wartko, P. D., Wong, E. Y., & Enquobahrie, D. A. (2017). Maternal birthplace is associated

with low birth weight within racial/ethnic groups. *Maternal and child health journal*, *21*(6),

1358-1366.

20. Clay, S. L., Woodson, M. J., & Kersh, R. (2021). An Exploration of Socio-

demographic, Economic, and Environmental Factors in Black/White Disparities in

Low Birth Weight Outcomes: A Cross-Sectional Study. *Journal of Research in*

Health Sciences, *21*(2), e00511.

21. Zeng, N., Erwin, E., Wen, W., Corsi, D. J., Wen, S. W., & Guo, Y. (2021). Comparison

of adverse perinatal outcomes between Asians and Caucasians: a population-

based retrospective cohort study in Ontario. BMC Pregnancy and Childbirth, 21(1), 1-9

22. Messer, L. C., Kaufman, J. S., Dole, N., Herring, A., & Laraia, B. A. (2006). Violent

crime exposure classification and adverse birth outcomes: a geographically-

defined cohort study. International journal of health geographics, 5(1), 1-12.

23. Lobel, M., Dunkel-Schetter, C., & Scrimshaw, S. C. (1992). Prenatal maternal stress

and prematurity: a prospective study of socioeconomically disadvantaged women.

Health psychology, 11(1), 32.

24. Dole, N., Savitz, D. A., Hertz-Picciotto, I., Siega-Riz, A. M., McMahon, M. J.,

& Buekens, P. (2003). Maternal stress and preterm birth. American journal

of epidemiology, 157(1), 14-24.

25. Masho, S. W., Chapman, D., & Ashby, M. (2010). The Impact of Paternity and

Marital Status on Low Birth Weight and Preterm Births. *Marriage & Family Review* (Vol.

46, Issue 4, pp. 243–256). https://doi.org/10.1080/01494929.2010.499319

26. Wood, C. (1997). The Association of Marital Status with Low Birthweight:

North Carolina, 1994-1995. State Center for Health Statistics.

27. National Center for Health Statistics (US). (1989). Trends in low birth weight:

United States, 1975-85 (No. 48). US Department of Health and Human Services,

Public Health Service, Centers for Disease Control, National Center for Health Statistics.

28. Rubin, L. P. (2016). Maternal and pediatric health and disease: integrating biopsychosocial

models and epigenetics. *Pediatric Research* (Vol. 79, Issues 1-2, pp. 127–135).

https://doi.org/10.1038/pr.2015.203

29. Alhusen, J. L., Bower, K. M., Epstein, E., & Sharps, P. (2016). Racial discrimination and

adverse birth outcomes: an integrative review. *Journal of midwifery & women's*

health, *61*(6), 707-720.

30. Hargraves, M. (2002). Elevating the voices of rural minority women. American Journal of

Public Health, 92(4), 514-515.

31. Taylor J.Y., Liu, T., Howell, E. (2020). Insurance Differences in Preventive Care Use

and Adverse Birth Outcomes Among Pregnant Women in a Medicaid Nonexpansion State:

A Retrospective Cohort Study. *Journal of Women's Health,* 29(1).

32. Günther, V., Alkatout, I., Vollmer, C., Maass, N., Strauss, A., & Voigt, M. (2021). Impact

of nicotine and maternal BMI on fetal birth weight. BMC pregnancy and childbirth,

373   21, 1-6.

33. Bohn, C., Vogel, M., Poulain, T., Spielau, U., Hilbert, C., Kiess, W., & Körner, A. (2021).

Birth weight increases with birth order despite decreasing maternal pregnancy

weight gain. Acta Paediatrica, 110(4), 1218-1224.

34. Gantt, A., Metz, T. D., Kuller, J. A., Louis, J. M., Cahill, A. G., & Turrentine, M. A.

(2022). Pregnancy at Age 35 Years or Older ACOG Obstetric Care Consensus No.

11. OBSTETRICS AND GYNECOLOGY, 140(2), 348-366.

35. Grossman, D. C., Baldwin, L. M., Casey, S., Nixon, B., Hollow, W., & Hart, L. G.

(2002). Disparities in infant health among American Indians and Alaska Natives in

US metropolitan areas. Pediatrics, 109(4), 627-633.

36. Morisaki, N., Kawachi, I., Oken, E., & Fujiwara, T. (2017). Social and anthropometric

factors explaining racial/ethnical differences in birth weight in the United States. *Scientific*

Reports, *7*, 46657.

37. Call, K. T., McAlpine, D. D., Johnson, P. J., Beebe, T. J., McRae, J. A., &

Song, Y. (2006). Barriers to care among American Indians in public health

care programs. Medical care, 595-600.

38. Gwin, K. M., Schrader, R., Peters, K., Moreno, A., Thiel, K. W., & Leslie, K. K. (2012).

An exploratory study of the variables impacting preterm birth rates in New Mexico.

BMC pregnancy and childbirth, 12, 1-8.

39. Raglan, G. B., Lannon, S. M., Jones, K. M., & Schulkin, J. (2016). Racial and ethnic

disparities in preterm birth among American Indian and Alaska Native women.

Maternal and child health journal, 20, 16-24.

401 **TABLES**

Table 1: Association of county-level contextual variables with low birthweight rates during the study period (2016-2021).

County-level contextual variables[a]	Incidence rate ratio of low birthweight (95% CI)	p-value
County type (Reference = Non-metro; N = 1836)		
Small metro (N = 347)	1.21 (1.16, 1.27)	< 0.0001
Large metro (N = 804)	1.24 (1.20, 1.29)	< 0.0001
Time period (Reference = 2016-2017)		
2018-2019	1.11 (1.09, 1.13)	< 0.0001
2020-2021	1.15 (1.13, 1.18)	< 0.0001
% Black	1.22 (1.20, 1.24)	< 0.0001
% Asian	1.01 (0.99, 1.03)	0.22
% Native Hawaiian/Pacific Islander	0.99 (0.97, 1.01)	0.27
% BMI underweight	1.08 (1.03, 1.13)	< 0.001
% Cigarette smoking during pregnancy	1.24 (1.21, 1.28)	< 0.0001
% Live Birth order > 1	0.89 (0.85, 0.93)	< 0.0001
% Mother's age 35-54 years	1.06 (1.01, 1.11)	0.02
% No prenatal visits	1.27 (1.23, 1.32)	< 0.0001
% Maternal education more than high school	1.02 (0.98, 1.05)	0.34
% Uninsured population	1.10 (1.09, 1.12)	< 0.0001
Violent crime rate (per 100,000)	1.07 (1.06, 1.09)	< 0.0001
Primary care provider rate (per 100,000)	1.09 (1.07, 1.11)	< 0.0001

N= 3,108 counties
[a]All predictor variables were standardized to have a mean of 0 and standard deviation of 1.

Table 2: Incidence rate ratios (IRR*) of low birthweight for 1-SD increase in the county-level percentage of married women.

County type	Incidence rate ratio of low birthweight (95% CI)	p-value
Non metro	0.90 (0.86, 0.94)	< 0.0001
Small metro	0.97 (0.94, 1.01)	0.16
Large metro	0.98 (0.96, 0.99)	0.01

N= 3,108 counties
*IRR estimates were generated after adjusting for all the county-level contextual variables

Table 3: Incidence rate ratios (IRR*) of low birthweight for 1-SD increase in the county-level percentage of unemployed population.

County type	Incidence rate ratio of low birthweight (95% CI)	p-value
Non metro	1.04 (1.02, 1.06)	< 0.001
Small metro	0.99 (0.97, 1.01)	0.16
Large metro	1.00 (0.98, 1.02)	0.99

N= 3,108 counties
*IRR estimates were generated after adjusting for all the county-level contextual variables

Table 4: Incidence rate ratios (IRR*) of low birthweight for 1-SD increase in the county-level percentage of American Indian/Alaskan Native women.

County type	Incidence rate ratio of low birthweight (95% CI)	p-value
Non metro	0.97 (0.95, 0.98)	< 0.001
Small metro	1.08 (1.02, 1.15)	0.01
Large metro	1.03 (0.94, 1.13)	0.56

N = 3,108 counties
*IRR estimates were generated after adjusting for all the county-level contextual variables

Figure 1: Smoothed Standardized Incidence Ratios (SIRs) of low birthweight and exceedance probability of low birthweight SIR > 1 for counties across the U.S. for 3 study periods (2016-2017, 2018-2019 and 2020-2021)