An ecological study of COVID-19 outcomes among Florida counties

Sobur Ali[1], Taj Azarian[1*]

[1] Burnett School of Biomedical Sciences, University of Central Florida, Orlando, Florida, USA.

*Corresponding author: Taj Azarian, Email: taj.azarian@ucf.edu

Abstract

During the COVID-19 pandemic, Florida reported some of the highest cases and deaths in the US. County-level variation in COVID-19 outcomes has yet to be comprehensively investigated. Here, we assess corelates of COVID-19 outcomes among Florida counties that explain variation in case, mortality, and case fatality rates (CFR) across pandemic waves. County-level administrative data and COVID-19 case reports was obtained from public repositories. We tested spatial autocorrelation to assess geographic clustering of case rate, mortality rate, and CFR. Stepwise linear regression was employed to investigate the association between COVID-19 outcomes and 18 demographic, socioeconomic, and health-related county-level predictors. We found mortality rate and CFR were significantly higher in rural counties compared to urban counties, among which significant differences in vaccination coverage were also observed. Multivariate analysis found that the percentage of the population aged over 65 years, the percentage of obese people, and the percentage of the rural population were significant predictors of COVID-19 case rate. Median age, vaccination coverage, percentage of people who smoke, and percentage of the population with diabetes were significant influencing factors for CFR. Importantly, higher vaccination coverage was significantly associated with a reduction in case rate (R = -0.26, p = 0.03) and mortality (R = -0.51, p < 0.001). Last, we found that spatial dependencies play a role in explaining variations in COVID-19 CFR among Florida counties. Our findings emphasize the need for targeted, equitable public health strategies to reduce disparities and enhance population resilience during public health crises.

Keywords: COVID-19, Florida, Urban rural, Socioeconomic, Regression analysis, CFR

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), responsible for the pandemic of the coronavirus disease 2019 (COVID-19), has posed an unprecedented crisis to public health worldwide. As of December 2023, COVID-19 has resulted in 103 million cases and 1.14 million deaths in the US. As the pandemic unfolded, it became increasingly evident that demographic, socioeconomic, and health-related risk factors play crucial roles in influencing the COVID-19 outcome among communities in the United States (US) [1,2].

Socioeconomic predictors, including income, education, employment, and access to healthcare services, have long been recognized as critical determinants of health outcomes [3]. These factors may be associated with an individual's capacity to adhere to public health guidelines, access healthcare resources, and implement preventive measures, which can impact the spread and severity of infectious diseases such as COVID-19. Furthermore, individuals residing in rural counties, characterized by socioeconomic challenges such as higher poverty rates, limited access to healthcare services, and lower educational attainment, have been found to be particularly vulnerable to COVID-19 [4,5]. However, variations in study time periods, population demographics, risk factors assessed, and endpoints have resulted in mixed findings among studies investigating the influence of demographic and socioeconomic risk factors on COVID-19 incidence rates [6].

Florida, one of the most populous states in the US, is notable for its unique demographic composition, high rates of immigration, recognition as a popular tourist destination, and climate. The diverse population, ranging from densely populated urban centers to sparsely populated rural counties, contributes to a complex web of potential influences on disease transmission and outcomes [7,8]. Considering the socioeconomic variation among counties in Florida, it is crucial to examine how these factors may have influenced COVID-19 outcomes.

In this study, we first investigated geographic clustering of COVID-19 outcomes. We then analyzed variation in case rate, mortality rate, and CFR across epidemic waves

60   and assessed the association of demographic, socioeconomic, and health-related

61   factors, and COVID-19 outcomes. Our results show that demographic, socioeconomic,

62   and health-related differences explain a significant proportion of the variation in COVID-

63   19 outcomes and that those inequalities disproportionately affected rural counties in

64   Florida.

Methods

Map data

Florida Rural counties designations were obtained from the Florida Department of Health (FDOH) [9]. FDOH defines rural as (i) a county with a population of 75,000 or less, (ii) a county with a population of 125,000 or less which is contiguous to a county of 75,000 or less, (iii) any municipality within a county as described above [9]. County-level cartographic boundary Shapefiles were downloaded from the US Census Bureau TIGER Geodatabase [10].

COVID-19 Outcome Data

County-level confirmed COVID-19 case and death data were obtained from The New York Times GitHub repository (https://github.com/nytimes/covid-19-data), based on reports from state and local health agencies [11]. Population statistics were obtained from the US Census Bureau (https://www2.census.gov/geo/docs/reference/ua/). Case rates represent the number of cumulative cases per 100,000 population and the mortality rate the number reported cumulated deaths from COVID-19 per 100,000 population as of December 2022. Each county's CFR was determined by dividing the cumulative deaths associated with COVID-19 by cumulative COVID-19 cases [12].

We categorized the pandemic into five epidemic waves, delineated based on the epidemic curve and coinciding with the emerge of notable variants of concern during specific date ranges. The initial wave occurred from June 1, 2020, to September 30, 2020, corresponding to the onset of the pandemic and the original strain. Subsequent waves included the second wave from October 1, 2020 to May 30, 2021, associated with the Alpha variant; the third wave from July 1, 2021, to October 31, 2021, marked by the Delta variant; the fourth wave from December 1, 2021 to February 28, 2022, characterized by the emergence of the Omicron variant (mainly BA.1, BA.2 subvariant); and the fifth wave from April 1, 2022 to October 31, 2022, comprised largely of Omicron sub-variants BA.4 and BA.5.

Covariates

Eighteen demographics, socioeconomic, and health-related factors were included in the study (**Table 1**). Data were collected from the Florida Community Health Assessment Resource Tool Set (CHARTS, www.flhealthcharts.gov), maintained by FDOH. Data for the percent of the population in the county within rural blocks in Florida was obtained from the US Census Bureau and COVID-19 vaccination data for Florida counties from the Center for Disease Control and Prevention (CDC) [13]. Age-adjusted deaths from all causes per 100,000 population were collected from Florida CHARTS. Age-adjusted rates, which accounts to variation in county age structure, were calculated as described here (https://www.flhealthcharts.gov/charts/OpenPage.aspx?tn=665). All data used in this study were retrieved from publicly available databases.

Statistical Analysis

All variables were reported as mean and standard deviation, and the coefficient of variation (CV) was calculated to assess the relative dispersion. Spread and collinearity of the predictor variables were assessed through histograms, bivariate scatterplots, and Spearman correlation coefficients using R-package psych v2.3.6 [14]. Correlation coefficients were calculated for each predictor with the outcome using Spearman or Pearson correlation according to data type and distribution. Florida county-level case rates, mortality rates, and CFR were visualized in the map using ggplot2 v3.4.3 and sf v1.0.14 package in R v4.3.1.

We used multivariate ordinary least squares (OLS) regression model to examine the relationship of predictor variables and COVID-19 outcomes. Then, we built multivariate models for each outcome using a stepwise backward procedure, starting with the fully saturated model, and then iteratively dropping variables. Akaike Information Criteria (AIC) and Variance Inflation Factor (VIF) were used to assess model fit and the degree of collinearity among covariates, respectively. R^2 was calculated for the final model [15]. All statistical analysis and visualization were performed using R v4.3.1 [16] with Rstudio v2023.06.1 [17] statistical software.

Spatial autocorrelation

We calculated the global Moran I statistic to examine whether county COVID-19 case rate, mortality rate, and CFR were spatially autocorrelated [18]. First-order Queen contiguity binary neighbor list and spatial weights matrix were constructed using R package spdep v1.2.8. Counties were considered neighbors when they shared one common point or vertex [19]. Local spatial autocorrelation identified four distinctive categories: High-High, Low-Low, High-Low, and Low-High. Local spatial clusters of COVID-19 outcome were measured using GeoDa v1.22 software [20].

We compared the final selected ordinary least squares (OLS) model and two spatial regression models: spatial lag model (SLM) and spatial error model (SEM). SLM examines how the adjacent counties influence an index county's outcome variable. The spatial lag parameter (r) estimates the association between the average logged period rate in nearby counties and the logged period rate in the index county. In SEM, the parameter (l) determines how well a county's ordinary least squares (OLS) residual correlates with the residuals of its neighbors. This spatial error parameter (l) represents the strength of the association between the average residuals or errors in nearby counties and the residual or error of the current county [21]. We used the R "lagsarlm" and "errorsarlm" functions from spatialreg package to perform spatial regression analysis [22].

Results

Descriptive mapping of COVID-19 outcomes in Florida

Florida comprises 67 counties (**Figure 1A**) with a population of 21 million, where a higher percentage of the state's population (87.7%) lives in urbanized areas. By December 2022, Florida reported around seven million confirmed cases and around 84,000 COVID-19 associated deaths. Visualization of cumulative rates across the five waves showed geographical heterogenicity in COVID-19 case rate, mortality rate, and CFR across Florida counties. Seminole, Miami-Dade, and Broward Counties had the highest case rate, while Glades, St. Johns, and Sumter Counties exhibited lower rates (**Figure 1B**). Union, Suwannee, Citrus Counties had the highest death rate, whereas Orange, Monroe, and St. Johns Counties had the lowest mortality rate (**Figure 1C**). Citrus, Highlands, and Union counties had the highest CFR and Leon, Orange, Monroe counties had lowest CFR (**Figure 1D**).

Table 1 presents the summary and descriptive statistics of COVID-19 outcome variables and independent variables, and CV, which informs dispersion of the data. Mortality rate and CFR were more dispersed than case incidence rate. Demographic, socioeconomic, and health-related risk factors vary widely among counties. The percentage of people in the rural areas, hospital bed rate, Hispanic population, and poverty rate are the most heterogeneous factors.

Spatial autocorrelation of COVID-19 outcome between Florida counties

Moran's I statistics were calculated to variables among Florida counties. Assessment of spatial autocorrelation in COVID-19 outcomes indicated significant but weak positive spatial autocorrelation for case rate (I = 0.12, p = 0.037), mortality Rate (I = 0.22, p =0.001), and CFR (I = 0.3, p < 0.00). The Local Moran test was conducted to examine spatial clustering in more detail. The "High-High" category signifies areas with high values neighboring other counties with high values, "High-Low" counties with high values neighboring counties with low values, and "Low-Low" counties with low values neighboring other counties with low values. Orange County (Orlando, FL) belonged to a High-High case rate cluster, and   Flagler, Putnam, Marion, Citrus and Hernando

County a Low-Low cluster (**Figure 2A**). Madison, Suwannee, Lafayette, Gilchrist, and

Columbia Counties belonged to a High-High mortality rate cluster, and Okaloosa,

Collier, and Miami-Dade Counties a Low-Low cluster (**Figure 2B**). For CFR, we

identified one High-High cluster consisting of Marion, Citrus, Sumter, and Hernando

Counties and a Low-Low cluster consisting of four additional counties (**Figure 2C**).

Trends in COVID-19 case and mortality rates

Our analysis revealed higher case rates in the first and third waves in rural counties

compared to urban counties. However, no significant difference in case rates was

observed for the second and fourth waves (**Figure 3A**). Furthermore, we observed a

noteworthy temporal difference. Over the five pandemic waves, rural counties appeared

to experience a more acute increase in COVID-19 case incidence, while urban areas

described by a more protracted epidemic curve (**Figure 4**). Our analysis indicates that,

despite the differences in epidemic trajectories, cumulative COVID-19 case rates did not

differ significantly between urban and rural counties (**Figure 3B**).

We further assessed mortality rates in urban and rural counties across five pandemic

waves in the context of SARS-CoV-2 variants (**Figure 5**). We found that the mortality

rates in rural counties was significantly higher than in urban counties during the second,

third, and fifth waves **(Figure 3C)**. While the difference in mortality rate between urban

and rural counties during the first and fourth waves was not significant, the overall

mortality rate was significantly higher among rural counties (**Figure 3D**). Variation in the

case and mortality rate trends across the five waves of the pandemic was also

observed. In particular, the third wave (Delta variant) exhibited the highest mortality

rate, while the fourth wave (Omicron variant) showed a significantly higher case rate.

CFR was significantly higher (p = 0.02) in rural counties than in urban **(Figure 3F)**.

However, we observed inconsistent patterns among urban and rural counties when we

assessed CFR differences during the five waves. In the first wave, CFR was

significantly higher in urban counties, whereas in rural counties CFR was higher in the

second and fifth waves (**Figure 3E**). As decreased access to SARS-CoV-2 testing

among rural counties may have resulted in an under estimation of case rates, we

conducted a sensitivity analysis finding that even after accounting for a 2.6%

underreporting of cases among rural counties, the CFR remained significantly higher (p

= 0.04).

Lastly, we extended our investigation to include age-adjusted all-cause mortality rates,

comparing urban and rural Florida counties before and during the pandemic. We

observed higher all-cause mortality among rural counties than urban counties before

and during the pandemic (**Figure 6A**). Our analysis also indicated that the difference in

death rates between urban and rural counties increased during the pandemic (**Figure**

6A). By calculating the percent change in the age-adjusted all-cause death rate from

2019 (pre-pandemic) to subsequent pandemic years (2020, 2021 and 2022), we found

a consistent and statistically significant trend: rural counties experienced a higher mean

percentage change in age-adjusted all-cause mortality than urban counties across the

analyzed years (**Figure 6B**).

Vaccination disparities between urban and rural counties in Florida

To investigate the role of vaccination on the trajectory of the pandemic, we analyzed

differences in vaccination coverage, finding significantly lower vaccination coverage in

rural counties compared to urban counties (**Figure 7A**). Further, when considering

vaccination trends, a persistent pattern became evident. Urban counties (Mean: 69.48,

95% CI: 66.55-72.41) consistently had ~ 20% higher vaccination coverage than rural

counties (Mean: 45.94, 95% CI: 42.39-49.50) (**Figure 7B**). However, several rural

counties were outliers, exhibiting higher vaccination rates than certain urban counties.

Correlation analysis

We evaluated the correlation between selected influencing factors and COVID-19 case

rate, mortality rate, and CFR. A significant positive correlation was observed between

case rate and individuals below poverty (r = 0.36, P = 0.003) and percent of the

population of black race (r = 55, P < 0.001); significant negative correlation was

observed between case rate and median age (r = -0.58, P < 0.001), percent of

population insured (r = -0.32, p=0.008) and COVID-19 vaccination coverage (r = -0.26,

P = 0.03). We also found a significant positive correlation between mortality rate and

percent of rural population (r = 0.53, P = <0.001), percent of population with obesity

(BMI>30) (r = 0.34, P = 0.006), percent of population with diabetes (r = 0.43, P < 0.001),

and percent of population who are smokers (r = 0.48, P < 0.001). In contrast, a

significant negative correlation was observed between the mortality rate and median

household income (r = -0.52, P <0.001) and vaccination coverage (r = -0.51, P < 0.001).

The CFR is significantly correlated with median age (r = 0.43, P <0.001), rural

population (r = 0.41, P <0.001), diabetes (r = 0.50, P <0.001), and smoking (r = 0.39, P

= 0.001) (**Table 2**).

Multivariate analysis

The spread and collinearity of covariates were assessed (**Supplementary Figure S1**).

COVID-19 vaccination with one dose and fully vaccinated was positively correlated, and

as a result, we retained fully vaccinated in the final analysis. For all other highly

correlated variables, such as hospital and acute care beds, which are both proxies for

healthcare access, we retained only one variable for the multivariate analysis.

In the final model for case rate, the percentage of the population in counties with rural

blocks, percent of black population, percent of obese population, Hispanic population,

and percent of people aged over 65 years remained in the final stepwise model

(R^2=0.42, P < 0.01). A low value of VIF (< 3) suggested that the covariates in the final

model did not have significant multi-collinearity and have substantial power to explain

maximum model variation (**Table 3**).

For the mortality rate, the final selected model included median age, percent of people

in the rural area, percent of black population, vaccination coverage, percent of

population who smokes, and percentage of Hispanic population, which explained 37%

of variation of mortality rate in Florida. The result of collinearity diagnostics (VIF < 5)

indicated that the collinearity between predictor variables was acceptable in this multiple

linear regression model (**Table 3**).

Based on the stepwise multiple linear regression analysis, median age, acute care beds

rate vaccination coverage, percent of the population who smokes, percent of the

diabetes population were found to explain 65% of variation in the CFR in Florida (**Table**

3). The collinearity diagnostics (VIF < 3) suggested that the predictor variables are
relatively independent of each other in the final model.

Spatial regression

We compared the final OLS models for case rate, mortality rate, and CFR with SLM and
SEM to assess spatial association between the predictor covariates and outcome
variables. For the case rate and mortality rate, coefficients of the OLS and SEM models
are closer than SLM (**Table 3**). However, AIC values were lowest for the OLS model
compared to the spatial models, indicating that case and mortality rate can be explained
better by OLS (**Table 4**). We then explored SLM and SEM models to account for spatial
dependencies in CFR. Notably, the estimates and significance of the covariates in the
SLM model remained consistent with the OLS model, indicating that the spatial lag
effects did not significantly alter the relationships between predictors and CFR. The
SLM also estimated a spatial lag coefficient (rho) of 0.26, informing the extent to which
neighboring counties' CFR values influence each other. As with SLM models, the
predictor variables of SEM exhibit estimates and significance similar to the OLS model.
Even after accounting for these predictor variables, the SEM estimated a spatial error
coefficient (lambda) of 0.31, indicating persistent spatial variation in CFR (**Table 3**). The
AIC values further informed our model selection, with the SLM having the lowest AIC,
indicating the best trade-off between model fit and complexity. The SEM followed
closely, while the OLS model had the highest AIC (**Table 4**). Therefore, spatial
dependencies play a role in explaining variations in COVID-19 CFR among Florida
counties.

Discussion

Socioeconomically disadvantaged communities often bear a greater burden during epidemics [23,24], and disparities were evident during the COVID-19 pandemic [25,26]. In Florida, links between geographic factors and COVID-19 outcomes have previously been identified [27,28]. Here, we further explore these associations by comprehensively investigating the interplay between demographic, socioeconomic, and public health factors. Through our analysis, we observed significant temporospatial variations in county-level COVID-19 outcomes, which are explained in part by variation in socioeconomic and health-related risk factors as well as vaccination coverage.

While Florida's epidemic from the onset of the pandemic until December 2022 followed the US trend, subsequent county-level analysis revealed several notable trends. Rural counties experienced higher case rates in the first and third waves than their urban counterparts. Moreover, during the pandemic's second and fourth wave, rural counties experienced significantly higher mortality than urban counties as well as significantly higher cumulative CFRs, a finding consistent with other US-based studies [29,30]. Variation in CFR was also observed across the five waves, highlighting differences in epidemic trajectories between rural and urban counties, which corresponded with differences in vaccine uptake. Specifically, CFR remained elevated among rural counties during the fourth and fifth waves after the introduction of COVID-19 vaccines, which are highly efficacious in reducing mortality [31]. Mirroring the trend in CFR, all-cause mortality, which is robust to potential variation in COVID-19 mortality reporting among rural and urban counties, followed the same pattern. Together, these observations elucidate a broader and potentially longer-lasting impact of the pandemic. Detrimental collateral effects of the pandemic on the diagnosis and management of chronic health conditions have been described [32], and these effects often disproportionally impact rural communities.

Higher rates of SARS-CoV-2 transmission, limited access to tertiary healthcare, high prevalence of commodities, and decreased vaccine uptake may contribute to observed differences in COVID-19 outcomes. In general, rural populations tend to be un- or under-insured, older, more likely to live farther away from tertiary medical facilities and

have underlying health conditions, which increase their likelihood for adverse outcomes

[33]. Indeed, here we find that median income, adult obesity, smoking rates, and

diabetes were associated with higher county-level mortality rates. Obesity, diabetes,

and smoking are well-recognized risk factors for COVID-19 severity and mortality

[34,35]. In univariate analysis, number of acute care and hospital beds and the

proportion uninsured in a county were inversely associated with all COVID-19

outcomes, suggesting that access to care contributed to the observed disparities.

The relationship between race and ethnicity, socioeconomic status, and health is well

established. Higher rates of crowding and increased household size are often observed

among African American and Latinx communities [36], which in turn may result in more

frequent person-to-person contact and increased SARS-CoV-2 transmission [37].

Furthermore, African Americans have higher prevalence of cardiovascular disease,

obesity, diabetes and hypertension, which may contribute to COVID-19 severity [38,39].

Throughout the pandemic, race has been a consistent factor associated COVID-19

outcomes [40,41]. Here, we found a significant positive association between the

percentage of Black persons in a county with case rate and CFR. Interestingly, while

previous studies have reported increased COVID-19 cases and mortality rates among

Hispanic/Latinx population in US [42,43], we did not observe a significant association

between the percent Hispanic population in a county and increased morbidity and

mortality in our study. However, we found that both the percentage of Black and

Hispanic populations were significantly and positively associated with county-level

immunization rates. As higher immunization rates were protective in multivariable

models of mortality and CFR, this may partially explain the lack of association between

Hispanic percentage and COVID-19 outcomes, while raising questions about why the

percentage Black population remained associated with increased morbidity and

mortality. Further research is required to better understand the factors of the observed

variation in COVID-19 outcomes in Latinx and Black populations. Overall, pandemic

preparedness should recognize the substantial impact of systemic flaws within the

structural framework of American society, encompassing issues related to healthcare

accessibility and the availability of community-level resources [44].

Our analysis also indicates that rural counties had slower vaccine uptake and

significantly lower vaccination coverage than urban counties, which may have resulted

in excess mortality during later pandemic waves. Indeed, county vaccination rates

remained significantly associated with lower mortality rates when controlling for social

and demographic variables. These findings align with recent studies conducted in the

US [45,46]. Several factors may contribute to variation in vaccination coverage. First,

healthcare access in rural counties remains a challenge, with residents of rural areas

more likely to report decreased satisfaction with healthcare facilities than those in urban

areas. Decrease healthcare access may have limited accessibility of COVID-19

vaccination [33]. Second, there are potential differences in knowledge and attitudes

between urban and rural areas regarding the severity of COVID-19 and the

implementation of mitigation strategies, which are influenced by different political

ideologies and sociocultural identities [33]. Third, rural areas have historically higher

vaccine hesitancy for routine vaccines, which likely contributed to reduced COVID-19

vaccination coverage. For example, during the pandemic, rural adults in the US were

reported to be nearly three times more likely than urban adults to respond that they

"definitely won't" get vaccinated against COVID-19 [47]. Our multivariate analysis of

county level predictors of vaccination coverage indicates that all these factors likely

contributed to low vaccination rates (Supplementary Table S1). Further, our finding that

increased county-level vaccination coverage was associated with decreased mortality

rate and CFR indicates that a targeted approach to increasing vaccine confidence is

needed to close the gap in vaccination coverage between rural and urban communities.

While this study provides valuable insights, several limitations warrant consideration.

The use of cross-sectional data limits the ability to infer causal relationships.

Additionally, other unmeasured factors, such as variation in county-level interventions,

population mobility, and cultural practices, may also shape COVID-19 outcomes. In

Florida in particular, there was significant variation in county-level mitigation strategies

such as non-pharmaceutical interventions due to the absence of state-level policies.

Therefore, our analysis did not account for differences in mask mandates, stay-at-home

orders, or school-based interventions. Last, an underlying assumption of our analysis is

that case, death, and vaccine reporting was consistent among Florida counties. In

particular, differences in testing capacity between rural and urban counties may have

led to underreporting of cases among rural counties and a subsequent over-estimation

of the CFR [48]. However, our sensitivity analysis suggests that our results are robust to

modest differences in testing and case reporting, and our assessment of all-cause

mortality in addition to COVID-19 mortality supports the observed disparities.

377   Conclusion

In conclusion, this study contributes to our understanding of the dynamics of the

COVID-19 pandemic in Florida and accentuates the vulnerability of rural communities to

health crises. Spatial variation and associations between demographic, socioeconomic,

and health factors emphasize the need for tailored and equitable public health

strategies addressing infectious diseases and the broader determinants of health. The

insights gained from this study have the potential to inform evidence-based

interventions aimed at reducing health disparities and enhancing overall population

resilience in the face of public health crises.

Authorship contribution

Sobur Ali: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing – original draft, Writing – review & editing

Taj Azarian: Conceptualization, Project administration, Resources, Supervision, Writing – review & editing

Funding

There was no funding associated with this study.

Competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1.   **Onder G, Rezza G, Brusaferro S**. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. Jama American Medical Association, 2020; **323**: 1775–1776.

2.   **Grasselli G, et al.** Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy Region, Italy. Jama American Medical Association, 2020; **323**: 1574–1581.

3.   **Braveman P, Gottlieb L**. The social determinants of health: it's time to consider the causes of the causes. Public health reports SAGE Publications Sage CA: Los Angeles, CA, 2014; **129**: 19–31.

4.   **Khanijahani A, et al.** A systematic review of racial/ethnic and socioeconomic disparities in COVID-19. International journal for equity in health BioMed Central, 2021; **20**: 1–30.

5.   **Marmot M, Allen J**. COVID-19: exposing and amplifying inequalities. J Epidemiol Community Health BMJ Publishing Group Ltd, 2020; **74**: 681–682.

6.   **Martins-Filho PR, et al.** Socio-economic inequalities and COVID-19 incidence and mortality in Brazilian children: a nationwide register-based study. Public Health Elsevier, 2021; **190**: 4–6.

7.   **Neiderud C-J**. How urbanization affects the epidemiology of emerging infectious diseases. Infection ecology & epidemiology Taylor & Francis, 2015; **5**: 27060.

8.   **Reyes R, et al.** Urbanization and infectious diseases: general principles, historical perspectives, and contemporary challenges. Challenges in infectious diseases Springer, 2013; : 123–146.

9.   **FloridaHealth**. Florida Rural Counties (Last Upadate: February 2023). 2023(chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www.floridahealth.gov/programs-and-services/community-health/_documents/Rural_Counties_February_2023.pdf). Accessed 1 August 2023.

10. **USCensus**. United States Census Bureau, Cartographic Boundary Files. 2022(https://www.census.gov/geographies/mapping-files/time-series/geo/cartographic-boundary.html). Accessed 1 August 2023.

11. **The New York Times**. Coronavirus (Covid-19) Data in the United States. The New York Times. 2021(https://github.com/nytimes/covid-19-data). Accessed 10 July 2023.

12. **Cao Y, Hiyoshi A, Montgomery S**. COVID-19 case-fatality rate and demographic and socioeconomic influencers: worldwide spatial regression analysis based on country-level data. BMJ open British Medical Journal Publishing Group, 2020; **10**: e043560.

13. **CDC**. COVID-19 Vaccinations in the United States,County. Centers for Disease Control and Prevention. 2023(https://data.cdc.gov/Vaccinations/COVID-19-Vaccinations-in-the-United-States-County/8xkx-amqh). Accessed 1 August 2023.

14.   **Revelle W**. Psych: procedures for psychological, psychometric, and personality research.
Northwestern University (2020). Evanston, Illinois, R package 2023; Published online:
2023.

15.   **Middya AI, Roy S**. Geographically varying relationships of COVID-19 mortality with
different factors in India. Scientific reports Nature Publishing Group UK London, 2021; **11**:
7890.

16.   **Team RC**. R: a language and environment for statistical computing. Vienna: R
Foundation for Statistical Computing. (No Title) 2021; Published online: 2021.

17.   **RStudio T**. RStudio: integrated development for R. Rstudio Team, PBC, Boston, MA
URL http://www. rstudio. com 2020; Published online: 2020.

18.   **Getis A**. Spatial autocorrelation. Handbook of applied spatial analysis: Software tools,
methods and applications. Springer, 2009, p. 255–278.

19.   **Ramírez-Aldana R, Gomez-Verjan JC, Bello-Chavolla OY**. Spatial analysis of COVID-
19 spread in Iran: Insights into geographical and structural transmission determinants at a
province level. PLoS neglected tropical diseases Public Library of Science San
Francisco, CA USA, 2020; **14**: e0008875.

20.   **Anselin L, Syabri I, Kho Y**. GeoDa: an introduction to spatial data analysis. Handbook of
applied spatial analysis: Software tools, methods and applications. Springer, 2009, p. 73–
89.

21.   **Sun F, et al.** A spatial analysis of the COVID-19 period prevalence in US counties
through June 28, 2020: where geography matters? Annals of epidemiology Elsevier,
2020; **52**: 54–59.

22.   **Bivand R, Hauke J, Kossowski T**. Computing the J acobian in G aussian Spatial
Autoregressive Models: An Illustrated Comparison of Available Methods. Geographical
Analysis Wiley Online Library, 2013; **45**: 150–179.

23.   **Meara ER, Richards S, Cutler DM**. The gap gets bigger: changes in mortality and life
expectancy, by education, 1981–2000. Health affairs Project HOPE-The People-to-
People Health Foundation, Inc., 2008; **27**: 350–360.

24.   **Ottersen OP, et al.** The political origins of health inequity: prospects for change. The
Lancet Elsevier, 2014; **383**: 630–667.

25.   **Karmakar M, Lantz PM, Tipirneni R**. Association of social and demographic factors with
COVID-19 incidence and death rates in the US. JAMA network open American Medical
Association, 2021; **4**: e2036462–e2036462.

26.   **Bollyky TJ, et al.** Assessing COVID-19 pandemic policies and behaviours and their
economic and educational trade-offs across US states from Jan 1, 2020, to July 31, 2022:
an observational analysis. The Lancet Elsevier, 2023; **401**: 1341–1360.

27.   **Backer S, et al.** Socioeconomic determinants of COVID-19 incidence and mortality in

Florida. Cureus Cureus, 2022; **14**Published online: 2022.

28.   **Khan MM, Odoi EW**. Geographic disparities in COVID-19 testing and outcomes in
474   Florida. BMC Public Health BioMed Central, 2023; **23**: 1–13.

475   29.   **Pro G, et al.** Differences in US COVID-19 case rates and case fatality rates across the
476   urban-rural continuum. Rural and remote health NIH Public Access, 2020; **20**: 6074.

477   30.   **Iyanda AE, et al.** Racial/ethnic heterogeneity and rural-urban disparity of COVID-19 case
478   fatality ratio in the USA: a negative binomial and GIS-based analysis. Journal of racial
479   and ethnic health disparities Springer, 2022; **9**: 708–721.

480   31.   **Mohammed I, et al.** The efficacy and effectiveness of the COVID-19 vaccines in
481   reducing infection, severity, hospitalization, and mortality: a systematic review. Human
482   vaccines & immunotherapeutics Taylor & Francis, 2022; **18**: 2027160.

483   32.   **Hacker KA, et al.** Peer reviewed: COVID-19 and chronic disease: the impact now and in
484   the future. Preventing chronic disease Centers for Disease Control and Prevention, 2021;
485   **18**Published online: 2021.

486   33.   **Ullrich F, Mueller K**. COVID-19 cases and deaths, metropolitan and nonmetropolitan
487   counties over time (update). Policy File Rural Health Research Gateway, 2021;
488   **2020**Published online: 2021.

489   34.   **Kim L, et al.** Risk factors for intensive care unit admission and in-hospital mortality
490   among hospitalized adults identified through the US coronavirus disease 2019 (COVID-
491   19)-associated hospitalization surveillance network (COVID-NET). Clinical infectious
492   diseases Oxford University Press US, 2021; **72**: e206–e214.

493   35.   **Shang J, et al.** Cell entry mechanisms of SARS-CoV-2. Proceedings of the National
494   Academy of Sciences National Acad Sciences, 2020; **117**: 11727–11734.

495   36.   **U.S. Census Bureau**. America's families and living arrangements: 2020 – Table AVG1.
496   2020(https://www.census.gov/data/tables/2020/demo/families/cps-2020.html). Accessed
497   19 February 2024.

498   37.   **Shah M, Sachdeva M, Dodiuk-Gad RP**. COVID-19 and racial disparities. Journal of the
499   American Academy of Dermatology Elsevier, 2020; **83**: e35.

500   38.   **Reyes MV**. The disproportional impact of COVID-19 on African Americans. Health and
501   human rights Harvard University Press & François-Xavier Bagnoud Center for Health
502   and …, 2020; **22**: 299.

503   39.   **Yancy CW**. COVID-19 and african americans. Jama American Medical Association,
504   2020; **323**: 1891–1892.

505   40.   **Hooper MW, Nápoles AM, Pérez-Stable EJ**. COVID-19 and racial/ethnic disparities.
506   Jama American Medical Association, 2020; **323**: 2466–2467.

41.   **Golestaneh L, et al.** The association of race and COVID-19 mortality. EClinicalMedicine

Elsevier, 2020; **25**Published online: 2020.

42.   **Khanijahani A**. Racial, ethnic, and socioeconomic disparities in confirmed COVID-19
cases and deaths in the United States: a county-level analysis as of November 2020.
Ethnicity & health Taylor & Francis, 2021; **26**: 22–35.

43.   **Glance LG, Thirukumaran CP, Dick AW**. The unequal burden of COVID-19 deaths in
counties with high proportions of Black and Hispanic residents. Medical care Wolters
Kluwer Health, 2021; **59**: 470.

44.   **Hawkins RB, Charles EJ, Mehaffey JH**. Socio-economic status and COVID-19–related
cases and fatalities. Public health Elsevier, 2020; **189**: 129–134.

45.   **Saelee R, et al.** Disparities in COVID-19 vaccination coverage between urban and rural
counties—United States, December 14, 2020–January 31, 2022. Morbidity and Mortality
Weekly Report Centers for Disease Control and Prevention, 2022; **71**: 335.

46.   **Hernandez I, et al.** Disparities in distribution of COVID-19 vaccines across US counties:
A geographic information system–based cross-sectional study. PLoS Medicine Public
Library of Science San Francisco, CA USA, 2022; **19**: e1004069.

47.   **Sparks G, et al.** COVID-19 Vaccine Monitor: Differences in Vaccine Attitudes Between
Rural, Suburban, and Urban Areas. Kaiser Family Foundation, 2022; Published online:
2022.

48.   **Souch JM, Cossman JS**. A commentary on rural-urban disparities in COVID-19 testing
rates per 100,000 and risk factors. The Journal of Rural Health Wiley-Blackwell, 2021; **37**:
188.

Figures

Figure 1: (A) Florida map showing geographic distribution of urban and rural counties.
Statewide distribution of county-level COVID-19 confirmed cases per 100,000
population (B), mortality per 100,000 population (C), case fatality rate (D) in Florida from
March 2020 to December 2022.

Figure 2: Spatial autocorrelation of COVID-19 outcomes by Florida county considering
queen contiguity weights. Significant spatial clustering for case rate (B), mortality rate
(C), and CFR (D) was observed. The High-High cluster (red) indicates counties with
high values of a variable that are significantly surrounded by regions with similarly high
values. The "Low-Low" cluster (blue) refers to counties where low COVID-19 outcome is
surrounded by other counties with low COVID-19 outcome.

Figure 3: Variation in COVID-19 outcomes in Florida. (A) Variation of case rate in urban and rural counties in five pandemic waves and (B) difference in cumulative case rate (B) in urban and rural counties. (C) Variation of mortality rate in urban and rural counties in five pandemic waves and (D) difference in cumulative mortality rate. (E) Difference in CFR in urban and rural counties in five pandemic waves and (F) difference in overall CFR in Florida. Significance was assessed using a t-test. Data distributions are presented as box plots. Each dot point indicates the data of each county.

Figure 4: Trend in COVID-19 case rate among Florida counties. The heat map
illustrates daily COVID-19 cases per 100,000 population with red coloration indicating
higher values, and blue coloration indicating lower values. Counties are sorted by rural
population percentage, with higher percentages at the top and lower percentages at the
bottom. County names are color-coded, with orange indicating urban counties and blue
representing rural counties, as defined by the Florida Health Department. The bar plot
on the right side of the figure shows each county's COVID-19 case rate per 100,000
population from the pandemic's beginning to December 2022. At the top, a trend line
displays the 7-day moving average of daily COVID-19 cases in Florida, with annotations
marking five significant pandemic waves as of December 2022.

Figure 5: Trend in COVID-19 mortality rates among Florida counties. The heat map illustrates daily COVID-19 deaths per 100,000 population with red coloration indicating higher values, and blue coloration indicating lower values. Counties are sorted by population percentage within rural blocks, with the county having the highest percentage at the top. County names are color-coded, with orange indicating urban counties and sky blue representing rural counties, as defined by the Florida Health Department. At the top, a trend line displays the 7-day moving average of daily COVID-19 associated deaths in Florida, with annotations marking five significant pandemic waves. The right side presents the county-wise mortality rate as of December 2022.

Figure 6: Age-adjusted all-cause death rate (per 100k population) in Florida during pre-
pandemic and pandemic times. (A) Violin plots and boxplots of the age-adjusted all-
causes death rates in urban (orange) and rural (sky-blue) counties in pre-pandemic
(2018-19) and pandemic years (2020-22). (B) Boxplot presents the age-adjusted all-
cause death rate change from pre-pandemic (2019) to pandemic times (2020-22) in
urban and rural counties. For the t-test between urban and rural counties, significant
differences are represented with asterisks ($*$, $p < 0.05$; $**$, $p < 0.01$; $***$, $p < 0.001$).

586   **Figure 7:** The variation of vaccination coverage in Florida. (A) Boxplot shows the

587   vaccination coverage in urban counties(orange) and rural counties (sky-blue) in Florida.

588   (B) The trend of COVID-19 vaccination coverage in urban and rural counties from

589   December 2020 to December 2022.

Supplementary Figure S1: Correlation matrix of covariates. Spearman's correlation coefficient was used to measure the correlation between predictor variables. The diagonal shows the distribution of each variable in the histogram. On the top are the values of the correlations with significance measures (asterisks), and the bivariate scatter plots are shown at the bottom. p-value: *** < 0.001; ** <0.01; * < 0.05.

Table 1: Definition and Summary statistics of different variables and COVID-19 outcomes in 67
countries in Florida. Mean, standard deviation (SD) and Coefficient of Variation (CV) of
variables are listed.

Variable	Description	Mean	SD	CV
Response Variables (COVID-19 outcomes)				
Case rate	Confirmed cases per 100,000 population	30368	7163	23.59
Mortality rate	Reported deaths per 100,000 population	460.30	147.47	32.04
Case fatality rate	Case Fatality Rate in percentage	1.55	0.49	31.79
Demographic and socioeconomic factors				
Median age	Population median age	44.03	6.54	14.85
Aged	% of people aged over 65 years	22.96	7.93	34.54
Income	Median household income in Doller	51290	10300	20.08
Diploma	% Individuals with a high school diploma (aged 25 years and older)	33.25	7.20	21.65
Rural population	% of the population in the county within rural blocks	37.50	32.26	86.02
Unemployment rate	Unemployment rate, % of labor force	6.64	1.53	23
Poverty rate	% of total population, poverty rate	15.18	5.10	33.6
Below poverty level	% of total population, individuals below poverty level	14.96	5.03	33.59
Health and behavioral factors				
Hospital beds	Hospital beds, rate per 100,000 population	209.14	141.73	67.77
Acute care beds	Acute care beds, rate per 100,000 population	173.37	111.53	64.33
Health Insurance	% of population, with any type of health care insurance coverage	82.61	4.31	5.22
Vaccinated one dose	% of population who received one dose of vaccine	68.45	17.37	25.38
Fully vaccinated	% of population fully vaccinated	58.24	14.95	25.67
Obesity	% of total population, who are obese	32.46	6.07	18.69
Diabetes	% of total population, who have ever been told they had diabetes	13.37	3.09	23.11
Smoking	% of total population, who are current smokers	19.14	5.08	26.55
Race and Ethnicity				
Black	% of Population Black (of total population)	14.86	9.38	48.24
Hispanic	% of Population Hispanic (of total population)	14.59	13.09	89.75

602 **Table 2:** Univariate correlation analysis with predictor variables for case rate, mortality rate, and

603 CFR.

Variable	Case Rate		Mortality Rate		Case Fatality Rate	
	Correlation Coefficient	P value	Correlation Coefficient	P value	Correlation Coefficient	P value
Median age	-0.58	<0.001***	0.11	0.367	0.43	<0.001***
Aged	-0.62	<0.001***	0.12	0.332	0.44	<0.001***
Median income	-0.2	0.11	-0.52	<0.001***	-0.47	<0.001***
High school diploma	0.19	0.123	0.55	<0.001***	0.49	<0.001***
Rural population	0.24	0.056	0.53	<0.001***	0.41	<0.001***
Unemployment Rate	-0.15	0.221	-0.18	0.143	-0.12	0.303
Poverty rate	0.36	0.003**	0.29	0.017*	0.15	0.237
Below poverty level	0.34	0.004**	0.26	0.032*	0.13	0.282
Hospital beds	-0.17	0.179	-0.24	0.049*	-0.13	0.282
Acute care beds	-0.13	0.289	-0.17	0.178	-0.09	0.471
Health insurance	-0.32	0.008**	-0.24	0.0042*	0.02	0.819
Vaccination one dose	-0.26	0.035*	-0.51	<0.001***	-0.39	0.001**
Fully vaccinated	-0.26	0.035*	-0.51	<0.001***	-0.38	0.001**
Adult obesity	0.29	0.019*	0.34	0.006**	0.23	0.067
Diabetes	0.03	0.807	0.43	<0.001***	0.5	<0.001***
Smoking	0.1	0.423	0.48	<0.001***	0.39	0.001**
Black	0.55	<0.001***	-0.07	0.591	-0.24	0.047*
Hispanic	0.03	0.833	-0.29	0.017*	-0.33	0.006**

604 Notes: * p < 0.05 significant at 0.05 level

605 　　　　** p < 0.01 significant at 0.01 level

606 　　　　*** p < 0.001 significant at 0.001 level

 Table 3: Multivariable regression models of the association between socioeconomic characteristics and risk factors with the COVID-19

608 outcome variables. The three outcome variables included are case rate, death rate, and case fatality rate. All are calculated per 100,000

609 people, and data was included up to December 2022.

Dependent Variable	Variables	OLS				Spatial lag model			Spatial Error model		
		Adjusted R square	Estimate	Std. Error	(VIF)	Rho	Estimate	Std. Error	Lambda	Estimate	Std. Error
Case Rate	(intercept)	0.42	44912***	6163		0.04	43657***	7560	-0.02	44861***	5870.21
	Rural Population		96.05**	30.53	2.15		94.27***	29.82		96.73***	29.02
	Black		199.70*	82.88	1.34		195.17*	80.69		200.80*	78.80
	Obese		-486.22**	168.63	2.32		-482.07**	161.49		-485.75**	160.94
	Age over 65 years		-372.39***	101.07	1.42		-370.53***	96.66		-372.62***	95.99
	Hispanic		220.59***	58.65	1.3		219.72***	55.99		220.28***	55.55
Mortality rate	(intercept)	0.37	33.51	213.04		0.13	-25.39	209.78	0.21	17.37	201.87
	Median Age		10.95***	2.97	1.82		10.86***	2.81		11.61***	2.96
	Rural Population		-0.34	0.83	3.48		-0.50	0.79		-0.64	0.78
	Black		2.90	1.90	1.54		2.85	1.79		3.09	1.80
	Vaccination		-5.40**	1.97	4.18		-5.30**	1.86		-5.75**	1.89
	Smoking		10.03*	4.43	2.44		10.06*	4.16		10.46*	4.25
	Hispanic		2.53	1.39	1.59		2.66*	1.30		3.07*	1.40
Case Fatality Rate	(intercept)	0.65	-0.81	0.45		0.26	-1.03*	0.43	0.31	-0.56	0.45
	Median Age		0.05***	0.01	1.27		0.04***	0.01		0.05***	0.01
	Acute Care Beds		0.001***	< 0.001	1.49		0.001***	< 0.001		0.001***	<0.001
	Vaccination		-0.02***	0.004	2.96		-0.01***	<0.001		-0.02***	<0.001
	Smoking		0.02*	0.01	2.02		0.02*	0.01		0.02*	0.01
	Diabetes		0.04**	0.01	1.37		0.03**	0.01		0.03*	0.01

610 Notes: * p < 0.05 significant at 0.05 level
611   ** p < 0.01 significant at 0.01 level
612   *** p < 0.001 significant at 0.001 level

613 Table 4: Comparison of AIC values in OLS, SLM and SEM models for case rate, mortality rate and

614 CFR

Outcome variables	OLS	SLM	SEM
Case Rate	1350.8	1352.7	1352.8
Mortality Rate	836.8	838.2	837.8
Case Fatality Rate	32.3	29.9	31.2

617 **Supplementary Table S1.** Multivariable regression models of the association between

618 socioeconomic characteristics and risk factors with the vaccination coverage for COVID-19.

619 Vaccination coverage is measured per 100 people, and data was included from December 2020 to

620 December 2022.

Variables	Adjusted R square	Estimate	Std. Error	t value	Pr(>|t|)	VIF
Intercept		-5.56	9.14	-0.61	0.545	
Median Income		< 0.001	< 0.001	6.57	< 0.001***	1.91
Unemployment Rate		2.03	0.69	2.93	0.004**	2.07
Rural population (%)	0.85	-0.15	0.04	-4.07	< 0.001***	2.69
% of black population		0.32	0.09	3.55	< 0.001***	1.37
Aged over 65 years		0.64	0.11	5.81	< 0.001***	1.43
% of Hispanic population		0.23	0.07	3.10	0.003**	1.79