Time to Eat - A Personalized Circadian Eating Schedule Leads to Weight Loss Without Calorie Restriction: A Randomized Controlled Trial

Isabell Wilming[a], Jana Tuschewski[a], Jessie M Osterhaus[a], Theresa JG Bringmann[a], Anisja Hühne-Landgraf[a,b] & Dominic Landgraf[a,✉]

[a]Circadian Biology Group, Department of Molecular Neurobiology, Clinic of Psychiatry and Psychotherapy, University Hospital, Ludwig Maximilian University, Munich, Germany

[b]Munich Medical Research School, Ludwig Maximilian University, Munich, Germany

✉Dr. Dominic Landgraf
Clinic of the University Munich
Clinic for Psychiatry and Psychotherapy
Nussbaumstr. 7
80336 Munich
Germany
Tel.: +49 (0)89 4400 52753
Fax: +49 (0)89 4400 54741
E-Mail: dominic.landgraf@med.uni-muenchen.de

Highlights:

- Individual optimal times for meals are determined via an app-based meal diary.
- Generation of a structure plan for mealtimes are adjusted to individual circadian clocks.
- Following this plan, participants lost an average body weight of 2.6 kg over six weeks.
- Weight loss is achieved without changes in self-reported food quantity or composition.
- Regular mealtimes contribute to the improvement of the general well-being.

Abstract

Objective: Many weight loss strategies are based on the restriction of calories or certain foods. Here, we tested a weight loss intervention based solely on increasing the regularity of meals, presuming that this allows the circadian system to optimally prepare the food metabolism for these times.

Participants & Methods: In a two-group, single center randomized-controlled single-blind pilot study (pre-registration DRKS00021419) with participants aged 18-65 years and BMI $\geq$ 22 kg/m², we used a smartphone application to identify the times at which each participant eats particularly frequently and asked participants of the experimental group to restrict their meals to only these times for six weeks. Control participants received sham treatment. Primary outcome was self-reported body weight/BMI and secondary outcome the well-being of participants.

Results: Of 148 participants entering the pilot study, 121 were randomized and of these 100 (control: 33, experimental: 67) finished the study. Our results show that the more regular the meals of participants of the experimental group became, the more weight/BMI they lost, averaging 2.62 kg (0.87 kg/m²); $p < 0.0001$ (BMI: $p < 0.0001$) compared to an insignificant weight loss of 0.56 kg (0.20 kg/m²) in the control group; $p = 0.0918$ (BMI: $p = 0.0658$). Strikingly, weight loss was not related to changes in self-reported calories, food composition, and other food-related factors. Additionally, physical and mental well-being improved significantly.

Conclusion: In summary, increasing the regularity of meals causes participants to lose excess body weight and improves overall well-being.

Key words:

Circadian Clock, Obesity, Overweight, Personalized Therapy, Mealtime Pattern, Mental Health, Weight Loss, Diet

Abbreviations:

CG = Control group; EG = Experimental group; IDS-SR = Self-Assessment Inventory of Depressive Symptoms; MCTQ = Munich Chronotype Questionnaire; MTVS = Meal time variability score; PSQI = Pittsburgh Sleep Quality Index; SF-36 = 36-Item Short Form Health Survey; SWE = Scale of General Expectations of Self-Efficacy; TRE/TRF = time-restricted eating/feeding

1. Introduction:

Circadian clocks induce endogenous 24-hour rhythms in the expression of more than half of all genes across all tissues [1, 2], allowing the body to anticipate daily changes between day and night and to prepare and harmonize physiological processes accordingly. However, this requires exposure to very regular environmental time cues, so-called *Zeitgebers*, such as food. Thus, on the one hand, the circadian system contributes to the optimization of food metabolism; on the other hand, the intake of food serves as a potent *Zeitgeber*. Both mouse and human studies have shown the negative consequences of circadian disturbances on metabolism, body weight [3-11], and also mental health [12-14] and demonstrated that restriction of meals to certain time spans of the day can counteract these harmful effects [15-19]. In the recent past, some human studies have shown that restricting food intake to certain times of the day (time-restricted eating, TRE / time-restricted feeding, TRF) can indeed lead to weight loss. In most cases, the interventions were successful, and participants lost ~0.9 - 3.5 kg body weight within 12 weeks when they reduced their food intake to 8–10-hour intervals per day [19-22]. Furthermore, they suggest that TRE is more successful when the meal window is earlier in the day rather than later in the day [23]. However, previous human studies merely established rough periods during which participants could still eat irregularly.

It has previously been shown that the success of a hypocaloric diet can be increased if the distribution of daily calories consumed is adapted to the individual chronotypes in such a way that morning types consume many calories in the first half of the day and evening types consume many calories in the second half of the day [24]. We expect that the success of previous chrononutrition interventions can be further improved by determining each participant's eating time profile and creating a personalized meal plan accordingly, even if participants continue to consume their usual amounts and foods. We are testing this in the present pilot study, which is intended to examine in principle whether this assumption and possibly a more in-depth continuation of the study is justified.

Therefore, the primary aim is to individually determine and set optimal times for food intake and to correlate the reduction of mealtime variability with alterations in body weight. We include participants of Body Mass Index (BMI) classifications normal to extremely obese to test whether participants who exceed normal BMIs particularly benefit from the intervention. Because of the impact of circadian clocks on other body functions, we further hypothesize that regular food intake contributes to an overall improvement in well-being, such as the subjective feeling of general health, sleep quality, affective state, and self-efficacy. After screening, recruitment, a session to introduce the content and technical aspects of the study, and the completion of questionnaires on well-being, the eating times and food composition of all participants were recorded for two weeks using a smartphone application (Fig. S1A). Based on this data, a meal schedule was designed for each participant individually, which the

experimental participants were asked to follow for the next six weeks while data on eating times and food composition continued to be collected. For control subjects, we specified a long 18-h window during which they could continue to eat irregularly. The same questionnaires as at the beginning were then collected again. The participants then had the opportunity to end the intervention or to continue without further data collection. In both cases, however, an attempt was made to reach them for a follow-up assessment. Since the program focuses on eating times and, unlike many other dietary programs, explicitly does not aim to limit calories or certain types of food, we have named it "Time to Eat" (Fig. S1B).

2. Results

2.1. Participants

For this pilot study, a total of 148 participants were recruited. Of these, 121 were randomly allocated to the experimental group (EG) (n=79) and the control group (CG) (n=42) (Fig. S1C), which had non-significantly different mean BMIs of 25.9 (SD: ± 4.138) (CG) and 27.5 (SD: ± 4.627) (EG) (Tab. S1). At the end of the intervention (T2), data were available from 100 participants (EG: n=67, CG: n=33). Their baseline data indicate that participants in both groups were, on average, living with overweight according to BMI classification, with the included CG participants having a BMI 25.5 (SD: ± 3.827) and being slightly less overweight, on average, than the analyzed EG participants with a BMI of 27.4 (SD: ± 4.616) (Tab. S1). There were no significant BMI differences between completers and drop-outs in either group (Fig. S2A, Tab. S2). Reasons for drop-outs were, according to self-report, illness/accident during the study and unwillingness or failure to adhere to fixed eating times.

2.2. Improvement of eating time regularity

All study participants were asked to record each occasion when anything containing calories, such as meals, snacks, sugared or milk-containing drinks, was consumed during a 14-day exploration and a six-week intervention phase (Fig. 1A, B, S1C). During the exploration and intervention phases, a total of 13,838 and 34,564 calorie intake occasions were recorded from completers, respectively. Most EG participants were assessed as having three eating times (n=53), some reported eating four meals (n=13), and one had two main meals per day. The average MTVS for breakfast, lunch, dinner, and total meals of the exploration phase of both groups was around 4, which corresponds to a daily deviation of ~120 min for each meal. In the EG group, mean scores improved to less than to 2 during the intervention phase, corresponding to a mean deviation of less than ± 30min (Fig. 1C, D, Tab. S3). In contrast, in the CG group, there was no improvement but some significant increase of the MTVS.

2.3. Weight change

During the exploration phase, when no instructions on eating times were given, CG and EG participants had a statistically non-significant average loss of 0.17 kg (0.06 kg/m²) and 0.21 kg (0.06 kg/m²), respectively (Figs. 1E, S2B-F, Tabs. S4, S5). However, during the intervention phase, when EG participants took their meals more regularly, they lost an average of about 2.6 kg (95% CI [-2.906, -1.915]) and 0.81 kg/m² (95% CI [-0.9729, -0.6420]) (Figs. 1E, S2B-E, Tabs. S4, S5), which translates to an average weight loss of 0.40 kg (0.145 kg/m²) per week during the intervention. The weight loss observed in this group is independent of the age of the participants and their sex (Fig. S2G, Tabs. S6, S2). In contrast, weight and BMI changes of -0.39 kg (95% CI [-0.9125, 0.1307]) and -0.14 kg/m² (95% CI [-0-3152, 0.0304]), respectively, were not significant in the CG (Figs. 1E, S2B, C, F, Tabs. S4, S5), resulting in a significant difference in weight loss between groups at the end of the intervention (Figs. 1F, S2C, Tab. S5). The inclusion of the last available weight and BMI data of the drop-outs in an end-point analysis did not lead to any notable change in this result (Fig. S2H, Tab. S5).

Four weeks after the intervention ended, all participants were contacted again for a follow-up assessment. 34 of the 67 EG participants reported to have maintained the regularity of their meals and continued to lose body weight significantly, on average another 1.12 kg (95% CI [-1.479, -0.7626]) and 0.36 kg/m² (95% CI [-0.4907, -0.2457]), respectively (Figs. 1E, S2B, D, E, Tab. S4). In contrast, participants who reported to have resumed irregular eating gained a significant average of 0.48 kg (95% CI [0.1984, 0.7616]) and 0.17 kg/m² (95% CI [0.07434, 0.2603]), respectively (Fig. 1E, S2B, D, E, Tab. S4).

2.4.   Relation between weight change and improvement in the regularity of meals

To examine the extent to which EG participants' weight loss was effectively related to improvements in meal regularity, we correlated changes in body weight and BMI with individual MTVS changes. Consistent with our hypothesis, a significant relationship exists between weight loss and improvement in regularity in the EG (Fig. 1G, Tab. S6). A breakdown of meals shows that increasing the regularity of lunch and dinner has the greatest effects, with no significant relationship between increasing the regularity of breakfast and weight loss (Fig. 1H, Tab. S6). In contrast, in the CG, there were no or little improvements in regularity and, accordingly, no correlation with weight/BMI changes.

Further analyses revealed that all MTVS up to 3, which corresponds to a time window for meals of up to ±45 min, resulted in comparably effective weight loss, whereas the efficiency of weight loss decreases significantly with an MTVS greater than 3 (Tab. 1).

***Table 1:** Association of weekly MTVS with weekly weight loss. The table shows the weekly weight change of all participants during the exploration and the intervention phase while*

Weekly MTVS	Corresponding time window (min)	Weekly weight change (kg)	SD	n-value	Sig. difference from MTVS =1
= 1	≤ 30	-0.47	0.1995	50	
≤ 2	≤ 60	-0.39	0.5144	318	n.s.
≤ 3	≤ 90	-0.38	0.5881	426	n.s.
> 3	> 90	-0.12	0.5415	280	*
> 4	> 120	-0.10	0.4035	137	*

2.5. Relation between weight change and baseline BMI

A weight loss intervention which is particularly based on metabolic optimization is likely to be most effective for individuals with exceeded body weight. In fact, the higher the baseline BMI of EG participants at T_1, the greater the weight loss (Fig. S2I, Tab. S6). Stratifying the subjects by baseline BMI, including the control group, additionally shows that in all strata the BMI of EG subjects decreased more than that of CG subjects, even if they had the same initial BMI (Fig. S2J, Tab. S7). Multiple regression shows that the reduction in MTVS and baseline BMI together predict the weight loss of the participants (Fig. 1I, Tab. S8). These data show that regularity in meal intake can lead to a significant loss of body weight and suggest that this effect is especially pronounced in individuals with a higher BMI.

Fig. 1: Improvement in regularity of meals, body weight changes, and correlation between loss of body weight and regularity of eating meals. (A, B) Times of day of calorie intake occasions of 15 representative participants of the EG (A) and the CG (B) during the two-week exploration phase (left) and the six-week intervention phase (right). Each point represents a calorie intake occasion. (C, D) Change in MTVS of participants in the CG (gray) and EG (cyan) of breakfast, lunch, and dinner (C) and of all meals combined (D), including snacks, if snacks were taken. 2-way repeated-measures ANOVA with Bonferroni post hoc test comparing CG and EG during the exploration or intervention phase and T0-T1 and T1-T2 within groups, * p ≤ 0.05, ** p ≤ 0.01, **** p ≤ 0.0001. CG: n=33; EG: n=64-67, missing n-values in the EG are due to the different number of meals for each participant. Detailed statistics in Tables S3. (E) Within group differences of body weight during the study. Body weight was collected at different time points, at the beginning of the study (T0), after the exploration phase (T1), after the intervention phase (T2), and after the follow-up phase (T3). During the follow-up phase, 34 EG participants voluntarily continued the intervention (continuing; black line and open circle) while 33 participants no longer followed the intervention (discontinuing; gray, dashed line and open circle). Only three control subjects continued their intervention in the follow-up phase, which is why a division into continuing and discontinuing was omitted in this group. Data are normalized so that all baseline data at T0 equals 0 to represent change. Differences of all CG and EG participants between T0, T1, and T2 were calculated with a 1-way ANOVA, as were differences between T2 and T3 of CG participants (Bonferroni post-hoc test, **** p ≤ 0.0001). Differences between T2 and T3 of continuing and discontinuing EG participants were calculated with a paired t-test °° p ≤ 0.01, °°°° p ≤ 0.0001. Detailed statistics in Table S4. (F) Between group differences of BMI from T0-T3. Mixed-effect model with Bonferroni post-hoc test comparing CG and EG at different phases during the study, **** p ≤ 0.0001. Detailed statistics in Table S5. (G) Significant correlation between body weight/BMI and MTVS. The more participants improved their regularity of meals (decreasing MTVS), the more weight/BMI they lost. Linear regression, CG: n=33; EG: n=67, ** p ≤ 0.01. Detailed statistics in Tab. S6. (H) Significant correlation between body weight/BMI and MTVS of lunch and dinner. The more participants improved their regularity of lunch and dinner (decreasing MTVS), the more weight/BMI they lost. Linear regression, CG: n=33; EG: n=67, ** p ≤ 0.01. Detailed statistics in Tab. S6. (I) Significant correlation between change in BMI, baseline BMI, and improvement in MTVS. Individuals with higher BMI benefit more from similar improvements in MTVS (shades of data points) than individuals with lower BMI in EG participants. Multiple regression, n=67, * p ≤ 0.05, ** p ≤ 0.01. Detailed statistics in Tab. S8.

2.6.   *Relation between weight change and self-reported food quantity and composition*

By restricting meals to specific times and shortening the time window in which food is eaten, it is feasible that EG participants ate less or changed food composition during the intervention phase and therefore lost weight. However, on average, the reported food energy ingested by CG and EG participants did not substantially change during the study (Fig. 2A, S2K, L, Tab. S3, 6), apart from a minor average reduction of ~350 kJ (95% CI [-627.6, -67.67]) per day in the EG (Fig. S2K, Tab. S3), and did not differ between CG and the EG. As they are rather small, these differences do not add up to a significant reduction in the total average energy consumed over the course of the study. Accordingly, the determined cumulative average energy intake during the intervention phase did not differ on any day from the estimated energy intake based on an extrapolation of the cumulative food consumption during the exploration phase, neither in the CG nor in the EC group (Fig. 2B). Nevertheless, there were some participants who reported having consumed fewer calories during the intervention phase, while some also reported having ingested more calories. However, consistent with our hypothesis that high regularity of mealtimes optimizes metabolism of similar amounts of food, there is no relationship between reported changes in average daily and cumulative energy intake and change in body weight/BMI of either CG or EG participants (Fig. 2C, D, Tab. S6). Likewise, the composition of the ingested foods did not change considerably in either group, and participants reported a normal distribution of the macronutrients of fat, carbohydrate, and protein in both phases of the study (Fig. 2E, S2M, Tab. S3). Also, there was no correlation between weight change and change in reported average daily and cumulative macronutrient intake (Fig. 2F, S2N, Tab. S6).

2.7. Relation between weight change and other meal timing-related factors

Besides increasing regularity, restricting meals to certain times of the day can also shorten the daily window in which meals are eaten and shift meals to other times of the day. In fact, our intervention caused many EG participants to avoid occasional meals at extremely early or late times, limiting their meals to much shorter intervals of 11 h (95% CI [10.468, 11.520]) (Fig. 3A, B, Tab. S3), while those of the CG remained long with 13.7 h (95% CI [12.902, 14.562]). Additionally, compared with before and with the CG, the EG intervention resulted in the first meal of the day being delayed to a later time for many subjects (EG: 09:21 (95% CI [08:58, 09:45]) vs. CG: 08:03 (95% CI [07:32, 08:33])) and the last meal of the day being advanced to an earlier time (EG: 20:17 (95% CI [19:59, 20:35]) vs. CG: 21:31 (95% CI [21:05, 21:57])) (Fig. 3C, D, Tab. S3). However, weight changes of participants do not correlate with these changes in eating characteristics (Fig. 3B-D, Tab. S6).

2.8. Effect of meal schedule personalization on weight loss

A key part of our hypothesis involves the personalization of meal schedules. Consistent with the concept of social jet lag (the discrepancy between socially imposed and biological times), our data show that during the exploration phase, eating times on workdays differ from those on work-free days. Because social constraints usually matter less on non-working days, we expect eating times on these days to correspond more closely to individual circadian characteristics, such as the chronotype. The MCTQ questionnaire measures the chronotype based on the mid-sleep phase (mSP), i.e., the midpoint between sleep onset and waking up

[25]. Corresponding to the mSP, we determined the mid-eat phase (mEP), which we define as
the midpoint between the first and the last meal of the day. Indeed, our data show that the
later the mSP of participants, the later their mEP on weekends (Fig. 3E, Tab. S6), with the
mSP and mEP being almost antiphasic on average, i.e. 12 hours apart (Fig. 3F, Tab. S9). The
EG intervention maintains most of this relationship. However, the residuals of the regression
become smaller, resulting in a model fitting the data better (Fig. 3E, Tab. S6). Interestingly,
the closer the intervention brings an observed value to the predicted value, i.e., the smaller
the residual becomes, the more weight the participant loses (Fig. 3G, Tab. S6). This indicates
that the personalized adjustment of eating times to the individual chronotype, i.e., the
establishment of a highly accurate phase relation between mEP and mSP, leads to a
particularly effective reduction of excess weight.

Another part of the hypothesis is that the adaptation of eating times to the times of hunger
presumably determined by the circadian system contributes to weight loss. The calculation of
individual eating times was based on the frequency of the hours at which meals were eaten on
all days of the exploration phase. These days typically included more workdays than non-
workdays (Fig. 3H). Consequently, for some subjects, the EG intervention created a
discrepancy between what may have been physiologically optimal eating times, presumably
reflected better by data from non-workdays, and the calculated eating times. As the EG
participants were instructed to adhere to the calculated eating times even on non-working
days, their usual eating times at the weekend were sometimes greatly shifted (Fig. 3I, Tab.
S2). However, the magnitude of these shifts does not correlate with weight changes (Fig. S2O,
Tab. S6). On the contrary, if the calculated eating times resulted in large shifts of the weekend
meals, this also means that previously particularly large "meal jet lags" between working and
non-working days were eliminated. Indeed, the more the calculated eating times led to a shift
of late weekend breakfasts to earlier times and early weekend dinners to later times, the more
the participants lost weight (Fig. 3J, Tab. S6).

The independent variables used above for simple regressions may influence each other and
therefore their influence on BMI may be confounded. Therefore, a multiple regression with
the most important and least multicollinear variables (ΔMTVS, ΔkJ, Δcalorie intake interval,
ΔmSp-mEP relation, Δweekend breakfast, Δweekend dinner) was done to study their
individual influences on the BMI of EG participants while keeping the other variables
constant. According to this analysis, only the change in MTVS and also the resulting
elimination of "dinner jet lags" have significant influence on the BMI (Fig. 3K, Tab. S8). The
elimination of "breakfast jet lags" and the improvement of the mSP-mEP ratio did not reach
significance but showed a strong tendency to influence BMI. In contrast, changes in caloric
intake or the time interval during which food is consumed had no effect on the BMI. A

concurrent analysis of covariance shows that MTVS is most strongly intertwined with BMI, while other variables play a role to a much lesser extent (Fig. 3L).

Fig. 3: Influence of other changes in eating times and of personalization of meal schedules on weight loss. (A) 95th percentile of the time period during which food was ingested by 15 example participants during the exploration phase (yellow) and the intervention phase (purple). (B) Change in length of this period and its relation to weight/BMI changes. (C, D) Change in time of first (C) and last (D) caloric intake and their association with weight/BMI changes. (B-D) 2-way repeated-measures ANOVA with Bonferroni post hoc test comparing CG and EG during the exploration or intervention phase and T0-T1 and T1-

T2 within groups, *** p ≤ 0.001, **** p ≤ 0.0001. Linear regression. EG: n=67, varying n-
numbers result from incomplete entries in the diaries. Detailed statistics in Tables S3 and
S6. (E) The relation between mid-Sleep phase and mid-Eat phase (brown) on the weekends
of the exploration phase. The goodness of fit of this association improves during through EG
intervention and affects both workdays and weekend days (purple). Linear regression,
n=65, **** p ≤ 0.0001. Detailed statistics in Table S6. (F) Relation of the mid-Sleep phase
(mSP) and mid-Eat phase (mEP) during exploration and intervention phase. The circles
represent 24 h, the gray area represents the estimated sleep time, and the black arrow
represents the mid-sleep phase, which is normalized for all subjects. The colored triangles
represent the mid-Eat phase of each subject in relation to their mid-Sleep phase. The red
arrow shows the average mid-Eat phase across all subjects. n=65. Detailed statistics in
Table S9. (G) The smaller the residuals, i.e., the vertical distances of observation points
from the regression line, became, the more BMI the EG subjects lost. Linear regression,
n=65, ** p ≤ 0.01. Detailed statistics in Table S6. (H) Representative example of a meal
diary showing a shift from meals on weekends. The purple arrows represent the calculated
times, based on the events of the more frequent workdays, for the meal schedule of the
intervention phase. (I) Shifts from weekend meals caused by the EG intervention. Paired t-
test, n=65, *** p ≤ 0.001, **** p ≤ 0.0001. Data are normalized so that all baseline data at
T0-T1 equal 0 to represent the change in hours. Detailed statistics in Table S2. (J) Large
phase shifts of breakfast and dinner, equivalent to a correction of the social jetlag of meals
on weekends, are related to greater weight loss. Linear regression, breakfast: n=55, lunch:
n=64, dinner: n=63, ** p ≤ 0.01. Missing n-numbers are due to incomplete diary entries on
weekends during the exploratory phase or because participants usually did not have a
particular meal. Detailed statistics in Table S6. (K) Results of a multiple regression to
investigate the influence of individual variables while holding the others constant on the
BMI of EG participants. The Quantile-Quantile (QQ) plot shows the data are normally
distributed (left). The multiple regression plot and R2 value shows that the multiple
regression model explains the data well (middle). Additionally, the assignment of each
regression beta coefficient to the independent variables is shown along with the p-values
showing the influence on BMI. The data used is the same as those used in the individual
regressions shown previously. Detailed statistics in Tab. S8. (L) Covariance matrix with
normalized covariances for each pair of parameters. Values close to 1 (blue) or -1 (red)
indicate high intertwining of the data, values close to 0 (white) indicate independence or
orthogonality of the data. The assignment of the coefficients to the independent variables is
the same as in (K).

2.9.   General well-being outcomes

Since stable circadian rhythms are generally associated with many aspects of health, regular
entrainment of a variety of peripheral and brain clocks by the *Zeitgeber* food may enhance
overall well-being. To test this, participants were asked to complete clinical questionnaires
measuring subjective health status, sleep quality, depressed mood, and self-efficacy at T0 and
at T2. In addition, chronotype was also determined. Many aspects of subjective physical well-
being (SF-36 - physical health items) improved significantly during the intervention in the
EG (Fig. 4A, Tab. S3), reversing previous differences between CG and EG. In contrast, there
was only a slight improvement in the general health item within the CG. In addition, sleep
quality (PSQI) of the EG group improved significantly during the intervention, but not of the
CG group, again eliminating prior differences between CG and EG. Furthermore, mental
health, including vitality, depressed mood, and self-efficacy (SF-36 mental health items, IDS-
SR, SWE) also improved significantly in EG participants, but not in the CG (Fig. 4A, Tab. S3).
Consistent with our aim to optimize rather than to change individual rhythms, chronotype
did not change during the study (Fig. 4B, Tab. S3).

382
Fig. 4: Improvement in overall well-being over the course of the study. (A) Significant
improvements are observed in various aspects of physical and mental health (SF-36), sleep
quality (PSQI), depressed mood (IDS-SR), and self-efficacy (SWE) in participants of the EG
(colored columns), but hardly among CG participants (gray columns). Scales are either
upward or downward, depending on whether low or high scores on the questionnaires
mean subjectively good or poor well-being. (B) The chronotype of EG and CG participants
did not change during the study. 2-way repeated-measures ANOVA with Bonferroni post
hoc test comparing CG and EG during the exploration or intervention phase and T0-T1 and
*T1-T2 within groups, * p ≤ 0.05, ** p ≤ 0.01, *** p ≤ 0.001, **** p ≤ 0.0001. CG: n=33; EG:*
n=67. Detailed statistics in Tab. S3.

3. Discussion

In our pilot study, we show that increased regularity of the *Zeitgeber* food can lead to a significant reduction in body weight. Previously published data show that increased regularity of breakfast in particular, and not so much that of dinner, was identified as a predictor of weight loss [26]. In contrast, in our study, an increase in the regularity of lunch, and somewhat more strongly that of dinner, is related to weight loss. Although the diary entries of some participants show that they changed their energy intake and also the food composition of fat, carbohydrates, and proteins during the study, the observed weight loss is not related to these changes.

Related results have already been shown in mice, although here with a restriction to the entire active phase rather than specific eating times. Not surprisingly, mice gain weight and develop diabetes-like states under a high-fat diet [17]. It is surprising, however, that the negative physiological consequences of the high-fat diet can be almost completely prevented if the mice consume the equivalent amounts of high-fat food exclusively during their activity phase and do not receive any food during their sleep phase [17, 27].

There are also similar studies with humans in which the time window of food intake is limited to a certain length. This is, for example, 16:8 intermitted fasting or TRE/TRF, in which meals are taken only during specific time window, yet at unspecified times during that period. The effects of these protocols is mixed and often less pronounced than in our study, as less weight is lost [19-23, 28] or a longer period is required to achieve similar results [29] than in our study - with the caveat that participants with more overweight and older participants participated in these studies than in ours. Besides increased regularity, our intervention has similar effects on eating behavior, such as shortening the hours in which food is eaten. Interestingly, in our study these parameters are not related to the observed weight loss. However, other studies have also explored the effect of restricting individual meals to specific times. In mice, restricting the same amount of high-fat chow to two times per day reduces the development of obesity [30]. And in humans, restricting meals to three fixed, uniform times leads to similar weight loss like we observed in our study - but in combination with calorie reduction and standardized food [31].

The fact that our participants lost weight to a similar extent despite free choice of meals might be due to the individual tailoring of meal schedules. It is true that our data show that scheduling single meals to supposedly circadian optimal eating times of the participants is not essential for successful weight loss. Circadian clocks can adjust flexibly to *Zeitgebers* , and it is apparently more important to choose times for meals when people (have to) eat particularly frequently than the time that is probably genetically determined. That is, if the intervention leads to eating at times that are far from those normally chosen on non-working

days, this has no negative impact on weight loss efficiency. Our data suggest that the
regularity of eating times seems to be more important than the times themselves.

However, our pilot data still show that personalization of meal schedules is nevertheless
useful. First, depending on the individual chronotype of the participants, their mealtime
windows were earlier or later in the day. And those for whom the intervention optimized this
relationship were particularly effective at losing excess weight. Second, weight loss is more
successful if the eating time schedule is such that an individual who usually eats dinner at an
early time on non-working days, but has dinner later on working days, consistently eats at the
later time. This also tends to apply to breakfast if an individual eats breakfast earlier on
working days than on non-working days and the meal plan is aligned with the early time of
the working days.

Since food serves as a *Zeitgeber* for a large number of tissues, including several brain regions,
it can be hypothesized that regular eating not only has a positive effect on body weight, but
also on other aspects of well-being. A previous study by Panda et al. [18] has already shown
that shortening the eating interval to 10-12 hours per day has a positive effect on the general
well-being of the participants. Similarly, in our study, we also found significant
improvements in different aspects of subjective well-being over the course of the study,
specifically in physical and mental health, sleep quality, and self-efficacy. This shows that
adhering to daily structured mealtimes can improve not only metabolic but also other
physical and also psychological levels of health.

There are several explanatory approaches for why increased meal regularity can lead to
weight loss despite equal caloric intake. For example, circadian clocks influence the
composition of the microbiome, which regulates metabolic balance and body weight [32-37].
Additionally, circadian clocks help to anticipate mealtimes and provide the necessary
digestive components in advance of food intake in advance [38-41]. Accordingly, the number
of rhythmic genes in the liver of mice increases from ~350 to ~3,000 when the animals can
follow their natural feeding rhythm and even to ~5,000 when the feeding time is restricted to
a certain period of time [42]. There could therefore be a change in metabolic programming
that leads to a change in energy expenditure, for example.

3.1. Limitations
However, in our pilot study, we did not collect metabolic parameters that could provide
mechanistic explanations for the observed weight loss. Therefore, the assumption of an
optimized metabolism remains speculation and should be investigated in future follow-up
studies, for example by occasional indirect calorimetry measurements that can provide
information about changes in energy expenditure and the actual amount of food consumed in
different phases of the study.

Furthermore, we cannot conclude whether the circadian rhythms in the digestive tract of the
participants were indeed changed by the intervention. At the behavioral level, we see that the
eating rhythms show significantly less variability during the intervention than before or in
the CG. However, whether this has an effect on other rhythms, such as sleep-wake rhythms or
physiological rhythms, remains unknown. It is known from animal studies that precise eating
rhythms have a positive effect on endogenous rhythms and it is also known from human
studies that regular meals help the body to adapt endogenous rhythms to meals and
anticipate them [23, 38-41]. It can be assumed that this also applies to the participants in this
study. Nevertheless, measurements must be integrated in future studies to confirm this
assumption.

Another limitation of the pilot study is the self-assessment of key data on body weight, eating
times, and food composition. Self-reported data from participants who are not subject to
objective monitoring are prone to unreliability and errors. However, we have undertaken
measures and analyses to reduce this uncertainty. Body weight could not be collected in our
clinic under supervision because of the Covid-19 pandemic but was measured at T0, T1, and
T2 in the virtual presence of study staff (by camera) in order to be able to objectively check
this data for reliability. Regarding food intake, some participants underreported during the
study, as indicated by unrealistically low kJ intakes. However, our analyses suggest that
weight changes of individual participants are not related to their reliability of data recording
as they show that the individual reporting behavior of each participant remained largely the
same during the exploration and intervention phases, so underreporting is not indicative of a
change in eating behavior and, conversely, the observed kJ differences are likely due to actual
changes in eating behavior. However, when continuing the study, the transfer of photos of all
meals consumed should be implemented to objectively assess the composition and quantity
of food.

It was important for the design of this study that the participants were able to carry it out in
their natural environment and not in a laboratory setting, in order to be able to convincingly
demonstrate its applicability in everyday life. Instead of entries in the smartphone
application, there would have been the option of questionnaires or 24-hour dietary recall
outside the laboratory setting. These methods, for their part, harbor risks of misreporting due
to subsequent misjudgments of food intake or possible forgetting of snacks or caloric drinks
taken between meals [43, 44]. For the present pilot study, the food was to be recorded in real
time in order to avoid these possible sources of error. In our view, in the natural
environment, the limitation of self-reporting is unavoidable and must be accepted. The
above-mentioned inclusion of indirect calorimetry in the study protocol can reveal errors in
the self-report but will not be able to replace daily data collection by means of self-report.

However, the fact that the participants were only supervised online during the pilot study due to the pandemic may have had an influence. It cannot be ruled out that an occasional meeting in person would have improved compliance with data collection, as the participants might have felt more monitored by the study staff.

Although our evaluations suggest that this is the case, our data cannot clearly clarify whether the personalization of meal plans actually increases the success of weight loss. For this purpose, another control group is missing, in which the participants should have followed either a standardized meal plan or individual meal plans with specifically unsuitable eating times. This could be part of a future follow-up study, although it is difficult to predict how body weight would change in such a control group, as circadian clocks in the digestive organs can adapt to unusual eating times and the misalignment would disappear over time. However, compliance could suffer under such meal schedules, as it is more difficult for the control subjects to adhere to the eating times. This could lead to less successful weight loss.

In addition, after exclusion and drop-out of some participants, in the remaining evaluable 100 participants, there was a significant difference in baseline body weight and BMI between CG and EG. However, even if CG participants had been heavier from the start, it cannot be assumed that the applied sham treatment with an eating window of 18 hours during wake time would have had a stronger effect. Second, those subjects in the EG group whose baseline BMI was similar to those of the CG subjects lost significant weight during the intervention, while CG participants did not. Taking these two arguments into account, it cannot be assumed that a floor effect, in which the CG participants cannot lose any more weight due to a lower baseline weight, is the reason for the lack of weight loss in the CG. Lasty, studies show that even much larger baseline BMI differences do hardly affect the success of weight loss programs [45, 46], so we assume that this also applies to our study. During the course of the study, there were some drop-outs in both groups. Importantly, these did not differ in baseline weight from the completers, which is why it can be ruled out that the intervention was not effective in a particular BMI group. Rather, the subjects who dropped out were those who either could not continue due to illness or did not adhere to the meal plan. However, the inclusion of drop-outs in the analysis of weight progression does not fundamentally change the results.

3.2.   Conclusion

The present study can be regarded as a pilot study, the results of which we believe justify a continuation with adjustment of the above-mentioned limitations. In summary, our data show that the increase of regularity of meals to a period of 90 min for each meal promotes a significant reduction of body weight and a significant increase of well-being within short time. Importantly, the successes achieved do not seem to be related to a reduction in calories,

a change in food composition, or other eating behaviors. Additionally, we believe that this intervention poses hardly any health risks since it is most likely solely based on the optimization of metabolic processes. On the contrary, the strong increase in general well-being during the study rather indicates an improvement in health on many different levels.

4. Methods

4.1. Experimental approach

The study was a randomized controlled intervention trial with repeated measures investigating the effect of regularity of mealtimes on body weight and parameters related to overall well-being. Participants were blinded to the group assignment in that both control and experimental subjects were equally informed to participate in an intervention. The study was carried out at the Clinic for Psychiatry and Psychotherapy of the Ludwig Maximilian University (LMU), Munich, Germany. The study is registered at the German Clinical Trial Register with the trial number DRKS00021419 and was approved by the Ethics Committee of the LMU.

4.2. Participants

The participants were recruited across Germany via flyers at universities, fitness studios, adult education centers, pharmacies, grocery stores, and via Facebook® between September 2020 and August 2021. Those interested were able to contact the study staff by phone, email, or by making an entry in an online calendar tool. Participants had to be between 18-65 years old and have a BMI $\geq$ 22 kg/m². A requirement for participation in the study was that no other diet was concurrently followed, and that no medication was being taken regularly that could influence appetite or weight. No participants were included who knowingly suffer from a metabolic, mental, or addictive disorder. Further exclusion criteria were pregnancy, blindness, bedriddenness, and dependence on assistance with eating. Special emphasis was placed on the participants' ability to understand the details of the study and to give written consent to participate.

4.3. Study Protocol

Each participant completed a 12-week program, which consisted of the following steps (Fig. S1A):

4.3.1. Introductory Session and Initial Questionnaire Assessment

At the introductory session, the eligibility for possible participation in the study was determined and demographic data was collected. Then, suitable participants received a general introduction to the study, which included the theoretical background of the study and

a detailed explanation of the study procedure. Subsequently, the participants were informed about their rights, the voluntary nature of their participation in the study, and data protection regulations. All participants had to sign an informed consent form for study participation and for the recording and use of their data. Participants weighed and measured their height themselves while study staff connected with them online via camera. Afterwards, the participant received questionnaires to fill out (Tab. S10). This initial data assessment is referred to as T0 in the analysis.

4.3.2. Two-Week Exploration Phase

The introduction was followed by a two-week exploration phase in which the participants were asked to follow their usual eating habits and to document all calorie intake occasions (main meals, snacks, caloric drinks) using the smartphone application "FDDB" as accurately as possible. Based on this data, the personalized nutritional schedule was then developed. Data from the FDDB smartphone application also allowed us to calculate individual caloric intake along with the ratio of consumed macronutrients for each day during the study. Additionally, the participants weight was measured again as described above. The end of the exploration phase is called T1 in the analysis.

4.3.3. Six-Week Intervention Phase

Experimental Group (EG): Following the exploration phase, the participants received their personalized mealtime schedule, which was prepared beforehand by the study staff based on the meal diary of the exploration phase. Together with the participants, the nutritional behavior of the last two weeks was reviewed and analyzed. Special attention was paid to irregularities in mealtimes and to times of day when food was eaten particularly frequently. In this way, participants were made aware that they eat irregularly, but that there are nevertheless times when their internal clock frequently triggers hunger. This was followed by a six-week intervention phase, in which the participants were instructed to have their meals only at times according to the mealtime schedule, for each meal at least within a time window of ±30 min from the specified time. Participants were explicitly told that they were not restricted in their choice and quantity of food. During the intervention phase, participants should continue to document all calorie intake occasions using the FDDB App. In addition, participants were asked to weigh themselves weekly on the same scales they used at the start of the study and report their weight to the study staff to measure their progress. Halfway through the intervention phase, participants were contacted at least once by telephone to clarify possible questions and maintain compliance. Further, in case participants stopped making entries in their FDDB diary for two or more days, they were contacted to ask for reasons and to reinstate compliance again.

Control Group (CG): In principle, participants in the CG followed the same procedure. However, they were given a sham treatment, for which they were asked to restrict their meals to a freely chosen 18-hour time window per day, during which they could eat what they wanted and, above all, when they wanted. It can be assumed that restricting the eating time window to 18 hours should have no effect on weight, since such a long eating time window is hardly ever exceeded by most people anyway [18] and therefore hardly any change in eating behavior is to be expected.

4.3.4. Final Questionnaire Assessment

At the end of the intervention phase, the participants were asked to fill out the same questionnaires as at the introduction session (Tab. S10). Additionally, weight was measured again while study staff connected with them online via camera. This data assessment is called T2 in the analysis.

4.3.5. Follow-up Assessment

Four weeks after completion of the study, the study staff contacted the participants again by telephone to inquire about their current weight and to obtain information about whether the participants had voluntarily continued the program after the end of the intervention phase. If the participants indicated that they did not continue the program, possible reasons were evaluated, and suggestions were received on how the current protocol could be improved to make it more attractive for permanent implementation in everyday life. Participants of the control group were informed at this point about their allocation and the actual question of the study. The final data assessment is called T3.

4.3.6. Restrictions due to the Covid-19 pandemic

Due to restrictions imposed by Covid-19, the study was carried out online. Data collection and communication was exclusively done via email, Zoom Meetings (Zoom Video Communications, Inc.), Facebook (Meta Platforms), or telephone.

4.4. Assessment of General Well-Being

To measure whether restricting meals to times when we believe the circadian system triggers hunger leads to an increase in general well-being, questionnaires were collected to measure sleep quality, physical well-being, self-efficacy, and depressive mood. Additionally, the chronotype was assessed. The questionnaires used are described in Tab. S10.

4.5. Mealtime Diary with FDDB Application

The nutrition diary of the smartphone application FDDB was used to document all calorie intake occasions. FDDB is freely available in the usual application stores and is operated by the independent food industry company FDDB Internetportale GmbH, Berlin. Users can

record each calorie intake occasion with detailed information about the amount, type, and method of preparation of each food, which can be selected from a large database of different foods, which also contains data about the energy value of each product. The app automatically records the date and time of each calorie intake occasion. In case a certain food is not listed in the database, placeholder events with estimated energy values can be recorded. In addition to the participants, the study staff also had access to each account to check compliance and regularly download diary data.

4.6.   Creation of Personal Meal Schedules

At the end of the exploration phase, FDDB data were downloaded, and clusters of times of meals were identified. First, for each participant the optimal number of mealtime clusters was determined by the heuristic technique *elbow method*. With the number of meals calculated in this way, the optimal times for each of these meals were then calculated using the *k-means* algorithm with the scikit-learn machine learning package for Python. The variable minutes was created to adjust the format of the clock times in the data. The *k-means* algorithm finds a clustering structure, that minimizes the sum squared error which measures the distance of each datapoint to its representative value. The center of each cluster was then calculated as the mean of all observations in the cluster, which was extracted in hours and minutes. Calculated times were rounded to quarter, half, three-quarter, or whole hours and visually displayed in scatterplots for discussion with participants of the EG. If the calculated times were incompatible with, for example, participants' work schedules, they had the opportunity to adjust the times of meals accordingly. Then, an agreement was reached to eat only at these times for the next six weeks. For each meal, a time window of plus/minus 30 minutes around the specified time was granted. The participants were instructed not to advance missed meals or to make up for them at times that did not correspond to the established times, but to wait for the next established mealtime if possible. A scatterplot was also created for participants of the CG, but they were not provided with meal timings; instead, an 18-hour time window for meals was established with them.

4.7.   Statistical Analyses

Statistical analyses were performed with SPSS 24, Python, GraphPad Prism 9.2.0, and R. Details about statistical tests used for specific experiments are indicated in the main text and the corresponding figure legends.

4.7.1.   Mealtime Variability Score (MTVS)

A Python script was written to generate the MTVS, which is calculated from the deviation of each calorie intake occasion from the specified mealtime closest to the event. Similar to a previous study, each meal was given a score value based on the time deviation of the specified meal in minutes: 1 = +/- 0-15 min, 2 = +/- 16-30 min, 3 = +/- 31-45 min, 4 = +/- 46-60 min,

5 = +/- 61-75 min, 6 = +/- 76-90 min, 7 = +/- 91-105 min, 8 = +/- 106-120 min, 9 = +/- 2-3 h,

10 = +/- 3-4 h, 11 = over +/- 4 h [47]. Since the participants were given a 60-minute (±30

min) time slot for each meal, a score of up to 2 is considered very regular. Based on individual

score values of each meal, an average score value can be calculated for each individual day,

for each individual week or for the entire period, i.e., a daily, a weekly, or a total score.

Similarly, daily, weekly, and total scores can be calculated for each type of meal, i.e.,

breakfast, lunch, dinner.

4.7.2.   Sample Size and randomization

For the sample size calculation, we assumed an average of 2.5 kg of body weight loss, based

on TRE studies [18, 19, 23, 28, 29], as we are not aware of any other study in which eating

times were personalized and scheduled. The reported standard deviation for body weight loss

in such studies is 3.2 [19]. Randomization was to be 2:1 (EG:CG) because no significant

change in body weight was expected from the control treatment. The randomized assignment

was carried out using a continuous list prepared by the study supervisor with the Excel®

randomization function. Participant enrollment and assignment was carried out by study

staff. On this basis a large effect size $d = 0.8$ was assumed for the calculation of the power

analysis in G*Power and transformed into the effect size $f = 0.4$. The sample size calculation

was performed a priori for two groups (EG and CG) and the main three measuring time

points (T0, T1, T2) resulting in a sample size of N = 100 participants.

4.7.3.   Within- and Between Group Comparisons across T0-T1 and T1-T2

Two-way repeated-measures ANOVA was used to compare group and time effects in the CG

and EG and in the exploration (T0-T1) and intervention (T1-T2) phases, as well as effects of

the interaction of group and time. Additionally, Bonferroni's multiple comparison tests were

used to compare effects of time within groups (T0-T1 vs. T1-T2 within CG or EG) and effects

between groups at either the exploration or intervention phase (CG vs. EG at T0-T1 or T1-T2).

This form of analysis was applied to data in Figures 1C & D, 2A & E, 3B-D, 4A & B, and S2K.

If missing data points prevented a repeated-measure analysis, a mixed-effects model with

Bonferroni's multiple comparison test was applied instead. This applies to the data in Figures

1F, S2C, H & M. When comparisons were made only between exploration and intervention

phases within a group, a paired t-test was applied, which is the case for data shown in Figures

3I and S2G. When comparisons were made with groups with very different n-numbers,

Welch's test was used instead of the t-test. This applied to data shown in Figure S2A.

4.7.4.   Body Weight Development over Time

710 To analyze the change in body weight within the CG and EG groups over the T0-T2 period,

711 one-way repeated measures ANOVA with Bonferroni's multiple comparison test was applied.

712 The division of CG and EG into continuing and discontinuing did not allow inclusion of T3

data in the repeated measures analysis. Therefore, the comparison between T2 and T3 was completed with a paired t-test. This type of analysis was applied in data of Figures 1E and S2A.

4.7.5.   *Relationships between parameters*

Simple linear regression was used to determine whether two different parameters were related. This analysis was performed with data from Figures 1G & H, 2C, D & F, 3B-E, G & J, and S2G,, I, L, N & O. Relationships of more than two parameters were analyzed with multiple regression, which applies to data from Figures 1H and 3K.

4.7.6.   *Circadian Phase Distribution*

The mSP was evaluated according to the MCTQ [25]. The mEP was calculated in the same way, but instead of using the time of falling asleep and the time of waking up as in the calculation of the mSP, the first and the last meal of the day were used for the calculation. The distribution of each calculated phase was analyzed using Rayleigh's uniformity test. This analysis concerns the data shown in Figure 3F.

References

730   1.   Mure, L.S., et al., *Diurnal transcriptome atlas of a primate across major neural and peripheral*
731   *tissues.* Science, 2018.

732   2.   Ruben, M.D., et al., *A database of tissue-specific rhythmically expressed human genes has*
733   *potential applications in circadian medicine.* Sci Transl Med, 2018. **10**(458).

734   3.   Di Lorenzo, L., et al., *Effect of shift work on body mass index: results of a study performed in*
735   *319 glucose-tolerant men working in a Southern Italian industry.* Int J Obes Relat Metab
736   Disord, 2003. **27**(11): p. 1353-8.

737   4.   Antunes, L.C., et al., *Obesity and shift work: chronobiological aspects.* Nutr Res Rev, 2010.
738   **23**(1): p. 155-68.

739   5.   Vetter, C., et al., *Association Between Rotating Night Shift Work and Risk of Coronary Heart*
740   *Disease Among Women.* JAMA, 2016. **315**(16): p. 1726-34.

741   6.   Sun, M., et al., *Meta-analysis on shift work and risks of specific obesity types.* Obes Rev, 2018.
742   **19**(1): p. 28-40.

743   7.   Morris, C.J., et al., *Effects of the Internal Circadian System and Circadian Misalignment on*
744   *Glucose Tolerance in Chronic Shift Workers.* J Clin Endocrinol Metab, 2016. **101**(3): p. 1066-
745   74.

746   8.   Turek, F.W., et al., *Obesity and metabolic syndrome in circadian Clock mutant mice.* Science,
747   2005. **308**(5724): p. 1043-5.

748   9.   Barclay, J.L., et al., *High-fat diet-induced hyperinsulinemia and tissue-specific insulin*
749   *resistance in Cry-deficient mice.* Am J Physiol Endocrinol Metab, 2013. **304**(10): p. E1053-63.

750   10.   Shi, S.Q., et al., *Circadian disruption leads to insulin resistance and obesity.* Curr Biol, 2013.
751   **23**(5): p. 372-81.

752   11.   Zarrinpar, A., A. Chaix, and S. Panda, *Daily Eating Patterns and Their Impact on Health and*
753   *Disease.* Trends Endocrinol Metab, 2016. **27**(2): p. 69-83.

754   12.   Lee, A., et al., *Night Shift Work and Risk of Depression: Meta-analysis of Observational*
755   *Studies.* J Korean Med Sci, 2017. **32**(7): p. 1091-1096.

756   13.   Torquati, L., et al., *Shift Work and Poor Mental Health: A Meta-Analysis of Longitudinal*
757   *Studies.* Am J Public Health, 2019. **109**(11): p. e13-e20.

758   14.   Landgraf, D., et al., *Genetic Disruption of Circadian Rhythms in the Suprachiasmatic Nucleus*
759   *Causes Helplessness, Behavioral Despair, and Anxiety-like Behavior in Mice.* Biological
760   Psychiatry, 2016. **80**: p. 827-835.

761   15.   Kolbe, I., et al., *Circadian clock network desynchrony promotes weight gain and alters glucose*
762   *homeostasis in mice.* Mol Metab, 2019. **30**: p. 140-151.

763   16.   Chaix, A., et al., *Time-restricted feeding is a preventative and therapeutic intervention against*
764   *diverse nutritional challenges.* Cell Metab, 2014. **20**(6): p. 991-1005.

765   17.   Hatori, M., et al., *Time-restricted feeding without reducing caloric intake prevents metabolic*
766   *diseases in mice fed a high-fat diet.* Cell Metab, 2012. **15**(6): p. 848-60.

767   18.   Gill, S. and S. Panda, *A Smartphone App Reveals Erratic Diurnal Eating Patterns in Humans*
768   *that Can Be Modulated for Health Benefits.* Cell Metab, 2015. **22**(5): p. 789-98.

769   19.   Wilkinson, M.J., et al., *Ten-Hour Time-Restricted Eating Reduces Weight, Blood Pressure, and*
770   *Atherogenic Lipids in Patients with Metabolic Syndrome.* Cell Metab, 2020. **31**(1): p. 92-104
771   e5.

772   20.   Schroder, J.D., et al., *Effects of time-restricted feeding in weight loss, metabolic syndrome*
773   *and cardiovascular risk in obese women.* J Transl Med, 2021. **19**(1): p. 3.

774   21.   Gabel, K., et al., *Effect of time restricted feeding on the gut microbiome in adults with obesity:*
775   *A pilot study.* Nutr Health, 2020. **26**(2): p. 79-85.

776   22.   Lowe, D.A., et al., *Effects of Time-Restricted Eating on Weight Loss and Other Metabolic*
777   *Parameters in Women and Men With Overweight and Obesity: The TREAT Randomized*
778   *Clinical Trial.* JAMA Intern Med, 2020.

779   23.   Xie, Z., et al., *Randomized controlled trial for time-restricted eating in healthy volunteers*
780   *without obesity.* Nat Commun, 2022. **13**(1): p. 1003.

24. Galindo Munoz, J.S., et al., *Effect of a chronotype-adjusted diet on weight loss effectiveness:*
A randomized clinical trial. Clin Nutr, 2020. **39**(4): p. 1041-1048.

25. Roenneberg, T., A. Wirz-Justice, and M. Merrow, *Life between clocks: daily temporal patterns*
of human chronotypes. J Biol Rhythms, 2003. **18**(1): p. 80-90.

26. Fleischer, J.G., et al., *Associations between the timing of eating and weight-loss in calorically*
restricted healthy adults: Findings from the CALERIE study. Exp Gerontol, 2022. **165**: p.
111837.

27. Chaix, A., et al., *Time-Restricted Feeding Prevents Obesity and Metabolic Syndrome in Mice*
Lacking a Circadian Clock. Cell Metab, 2019. **29**(2): p. 303-319 e4.

28. Domaszewski, P., et al., *Effect of a Six-Week Intermittent Fasting Intervention Program on the*
Composition of the Human Body in Women over 60 Years of Age. Int J Environ Res Public
Health, 2020. **17**(11).

29. Gabel, K., et al., *Effects of 8-hour time restricted feeding on body weight and metabolic*
disease risk factors in obese adults: A pilot study. Nutr Healthy Aging, 2018. **4**(4): p. 345-353.

30. Martinez-Lopez, N., et al., *System-wide Benefits of Intermeal Fasting by Autophagy.* Cell
Metab, 2017. **26**(6): p. 856-871 e5.

31. Jakubowicz, D., et al., *Reduction in Glycated Hemoglobin and Daily Insulin Dose Alongside*
Circadian Clock Upregulation in Patients With Type 2 Diabetes Consuming a Three-Meal Diet:
A Randomized Clinical Trial. Diabetes Care, 2019. **42**(12): p. 2171-2180.

32. Thaiss, C.A., et al., *Transkingdom control of microbiota diurnal oscillations promotes*
metabolic homeostasis. Cell, 2014. **159**(3): p. 514-29.

33. Mashaqi, S. and D. Gozal, *"Circadian misalignment and the gut microbiome. A bidirectional*
relationship triggering inflammation and metabolic disorders"- a literature review. Sleep
Med, 2020. **72**: p. 93-108.

34. Voigt, R.M., et al., *Circadian disorganization alters intestinal microbiota.* PLoS One, 2014.
9(5): p. e97500.

35. Kuang, Z., et al., *The intestinal microbiota programs diurnal rhythms in host metabolism*
through histone deacetylase 3. Science, 2019. **365**(6460): p. 1428-1434.

36. Wang, Y., et al., *The intestinal microbiota regulates body composition through NFIL3 and the*
circadian clock. Science, 2017. **357**(6354): p. 912-916.

37. Heddes, M., et al., *The intestinal clock drives the microbiome to maintain gastrointestinal*
homeostasis. Nat Commun, 2022. **13**(1): p. 6068.

38. Scheving, L.A., *Biological clocks and the digestive system.* Gastroenterology, 2000. **119**(2): p.
536-49.

39. Cummings, D.E., et al., *A preprandial rise in plasma ghrelin levels suggests a role in meal*
initiation in humans. Diabetes, 2001. **50**(8): p. 1714-9.

40. Natalucci, G., et al., *Spontaneous 24-h ghrelin secretion pattern in fasting subjects:*
maintenance of a meal-related pattern. Eur J Endocrinol, 2005. **152**(6): p. 845-50.

41. Landgraf, D., A.M. Neumann, and H. Oster, *Circadian clock-gastrointestinal peptide*
interaction in peripheral tissues and the brain. Best Pract Res Clin Endocrinol Metab, 2017.
31(6): p. 561-571.

42. Vollmers, C., et al., *Time of feeding and the intrinsic circadian clock drive rhythms in hepatic*
gene expression. Proc Natl Acad Sci U S A, 2009. **106**(50): p. 21453-8.

43. Bailey, R.L., *Overview of dietary assessment methods for measuring intakes of foods,*
beverages, and dietary supplements in research studies. Curr Opin Biotechnol, 2021. **70**: p.
91-96.

44. Lichtman, S.W., et al., *Discrepancy between self-reported and actual caloric intake and*
exercise in obese subjects. N Engl J Med, 1992. **327**(27): p. 1893-8.

45. Unick, J.L., et al., *The long-term effectiveness of a lifestyle intervention in severely obese*
individuals. Am J Med, 2013. **126**(3): p. 236-42, 242 e1-2.

46. Barte, J.C., et al., *Differences in weight loss across different BMI classes: a meta-analysis of*
the effects of interventions with diet and exercise. Int J Behav Med, 2014. **21**(5): p. 784-93.

47.   Soehner, A.M., K.S. Kennedy, and T.H. Monk, *Circadian preference and sleep-wake regularity:*
associations with self-report sleep parameters in daytime-working adults. Chronobiol Int,
2011. **28**(9): p. 802-9.

Data availability

All data produced in the present study are available upon reasonable request to the authors.

Code availability

The codes used to calculate MTVS, individual optimal number of meals, and personalized optimal eating times will be available at https://github.com/dolandgraf/Time-To-Eat.git as of the date of publication of this paper.

Acknowledgments

We would like to thank all participants for their participation in this study. We also gratefully acknowledge Thomas Schneider-Axmann (Ludwig Maximilians Universität Munich, Germany) and Elisabeth Paul (Linköping University, Sweden) for their valuable contribution to the statistical analysis.

Funding

This work was supported by an Emmy Noether fellowship: LA4126/1-1 of the Deutsche Forschungsgemeinschaft to DL. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

DL, AHL, IW: Conceptualization

IW, JT, JMO, TJGB, DL: Data curation

DL: Formal Analysis

DL: Funding acquisition

IW, JT, JMO, TJGB: Investigation

IW, DL, AHL: Methodology

DL: Project administration

DL, IW: Resources

IW, JT: Software

DL: Supervision

DL, IW, AHL: Validation

DL: Visualization

870 DL: Writing – original draft

871 DL, AHL, IW: Writing review & editing

Ethics declaration

874 The conduct of the study was evaluated and approved by the Ethics Committee of Ludwig
875 Maximilian University under file number 19-975.

Competing interests

The authors declare no competing interest.

Fig. S1: (A) Timeline of the study. INTRODUCTORY SESSION: Test participants are
screened, recruited by a member of the study staff, and sign the consent form. This is
followed by an introductory session where they are informed about the contents of the
study and familiarized with the FDDB smartphone application. In addition, questionnaires
are filled out and weight and abdominal girth are measured. EXPLORATION PHASE:
During the following two weeks, the participants are asked to record each calorie intake
occasion with the FDDB app. Afterwards, a mealtime schedule is developed for each

participant. INTERVENTION PHASE: During the following six weeks, the participants are instructed to take their meals only at times according to the mealtime schedule and to report their weight weekly. In the middle of the intervention phase, a telephone appointment is made to maintain compliance. FINAL QUESTIONNAIRE ASSESSMENT: At the end of the intervention phase, the participants are asked to fill out the same questionnaires as at the introduction session. FOLLOW-UP ASSESSMENT: After another four weeks, the test participants are contacted again to explore whether they have continued the intervention voluntarily and to ask again for their body weight. (B) Logo of the "Time To Eat" study. The logo represents a clock in which the letters act as hands. (C) Consort diagram through the phases of the trial. A total of 148 initial participants were recruited. Of these, 48 were excluded or withdrew from the study in the course of the trial. Thus, the final study sample consisted of 100 participants of whom 67 were allocated to the EG and 33 to the CG. Until the follow-up appointment, 34 EG participants voluntarily continued the intervention. 30 EG participants discontinued the intervention but were available for further data collection, while three EG participants were no longer available and thus no analyzable data were available from them at T3. Of the 33 CG participants, only three voluntarily continued the sham intervention until T3.

Fig. S2: (A) BMI differences between study completers and drop-outs. Differences were calculated with a Welch's test (detailed statistics in Tab. S2). (B) Within group differences of BMI during the study. BMI was collected at different time points, at the beginning of the study (T_0), after the exploration phase (T_1), after the intervention phase (T_2), and after the follow-up phase (T_3). During the follow-up phase, 34 EG participants voluntarily continued the intervention (continuing; black line and open circle) while 33 participants dropped out (discontinuing; gray, dashed line and open circle). Only three control subjects continued their intervention in the follow-up phase, which is why a division into continuing and discontinuing was omitted in this group. Data are normalized so that all baseline data at T_0 equals 0 to represent change. Differences of all CG and EG participants between T_0, T_1, and T_2 were calculated with a 1-way ANOVA, as were differences between T_2 and T_3 of CG participants (Bonferroni post-hoc test, **** $p \leq 0.0001$). Differences between T_2 and T_3 of continuing and discontinuing EG participants were calculated with a paired t-test °° $p \leq 0.01$, °°°° $p \leq 0.0001$ (detailed statistics in Tab. S4). (C) Between group differences of BMI from T_0-T_3. Mixed-effect model with Bonferroni post-hoc test comparing CG and EG at different phases during the study, **** $p \leq 0.0001$ (detailed statistics in Tab. S5). Individual body weight (D) and BMI (E) changes of all 34 continuing and all 33 discontinuing participants of the EG group and of CG participants (F) during the study. Data is normalized so that all baseline data at T_0 equals 0 to represent change. (G) No correlation between age and weight change (top) and no difference in weight change between female and male participants (bottom). Linear regression, n=67 (detailed statistics in Tab. S6) and paired t-test, n=41/26 (f/m) (detailed statistics in Tab. S2) (H) Between group differences of body weight and BMI from T_0-T_3 including the last available values of study drop-outs which were carried forward. Mixed-effect model with Bonferroni post-hoc test comparing CG and EG at different phases during the study, **** $p \leq 0.0001$ (detailed statistics in Tab. S5). (I) Correlation between baseline BMI at T_1 and change in BMI in EG participants. Linear regression, n=67 (detailed statistics in Tab. S6). (J) Between group differences after stratification according to different baseline BMIs. 2-way repeated-measure ANOVA with Bonferroni post-hoc test, *** $p \leq 0.001$ (detailed statistics in Tab. S7). (K) Raw data of change in caloric intake during intervention. 2-way repeated-measure ANOVA with Bonferroni post-hoc test, CG: n=30; EG: n=67, * $p \leq 0.05$, (detailed statistics in Tab. S3). (L) Correlation between reported ingested energy during the exploration phase vs. the intervention phase. Linear regression, CG: n=30; EG: n=67 (some participants were excluded from this analysis because of several erroneous diary entries about unrealistic amounts of food), (detailed statistics in Tab. S6). (M) Determined and estimated cumulative intake of fat, carbohydrates, and proteins during the study. Mixed-effects model with Bonferroni post hoc test comparing determined and estimated cumulative energy intake. The estimated macronutrient intake is formed as an assumed continuation of the average daily intake during the exploration phase. The Bonferroni post hoc test showed no significant differences on any study day. (N) No significant correlation between body weight/BMI and changes of cumulative macronutrient intake. Linear regression, CG: n=30; EG: n=67, (detailed statistics in Tab. S6). (K, L, M, N) Some participants were excluded from this analysis because of several erroneous diary entries about unrealistic amounts of food. (O) The absolute deviation between the times of weekend meals and the times of food intake calculated for the EG intervention did not affect the subjects' BMI change. Linear regression, n=67 (detailed statistics in Tab. S6).

Table S1: Baseline characteristics of final 100 participants.

Characteristic	CG (n = 33; male: 10. female: 23)			EG (n = 67; male: 26, female: 41)			Difference CG - EG	
	Mean	Range	SD	Mean	Range	SD	p-value	Summary
BMI (kg/m²) incl. drop-outs	25.9	20.3 / 35.7	4.138	27.5	21.7 / 43.8	4.627	.0597	n.s.
Age (years)	33	20-60	12.08	38	19 - 64	14.49	.0795	n.s.
BMI (kg/m²) (T0/T1)	25.5 / 25.4	20.4 − 35.8 / 20.3 − 35.7	3.83 / 3.88	27.4 / 27.3	22.0 − 43.6 / 21.7 − 43.8	4.62 / 4.59	.0420 / .0420	* / *
BMI (T2)	25.3	20.4 − 34,6	3.77	26.5	21.2 − 42.5	4.47	.1680	n.s.
Δ BMI (T0-T2)	-.20	-1.17 − 1.52	.55	-.87	-2.8 − 1.0	.72	.0420	*
Body weight (kg) (T0/T1)	75.04 / 74.87	54.2 − 134.6 / 54.0 − 134.2	16.94 / 17.07	82.73 / 82.52	58.5 − 129.0 / 58.1 − 129.5	16.72 / 16.59	.0338 / .0342	* / *
Body weight (T2)	74.48	53.5 − 130.2	16.84	80.11	56.3 − 125.8	16.30	.1116	n.s.
Δ Body weight (T0-T2)	-.56	-4.4 − 5.1	1.72	-2.62	-8.5 − 3.5	2.23	.0338	*
MTVS exploration (total)	3.35	1.32 − 4.79	.8081	3.88	1.11 − 5.76	.9086	.0058	**
MTVS intervention (total)	3.94	2.45 − 5.13	.7203	1.72	1.02 − 3.84	.6156	<.0001	****
ΔMTVS (total)	.583	-.940 − 3.47	.9051	-2.16	-4.69 -.420	.9571	<.0001	****
KJ intake exploration	7,211	3,350 − 1.537	1,596	7,071	2,505 − 15,185	1,854	.7197	n.s.
KJ intake intervention	7,248	3,324 − 11,829	1,766	6,723	3,496 − 12,932	1,661	.1618	n.s.
% Fat exploration	20.73	11.59 − 26.12	3.639	21.85	10.86 − 32.30	3.811	.1773	n.s.
% Fat intervention	22.06	12.54 − 28.98	3.282	21.40	11.70 − 28.91	3.158	.3477	n.s.
% Carbohydrate exploration	57.51	47.63 − 71.40	6.536	56.23	42.69 − 71.11	6.247	.3607	n.s.
% Carbohydrate intervention	56.54	45.29 − 70.90	5.712	55.90	35.20 − 68.59	5.282	.5969	n.s.
% Protein exploration	21.77	12.56 − 31.56	4.456	21.92	13.29 − 35.51	4.520	.8747	n.s.
% Protein intervention	21.41	15.07 − 30.13	3.688	22.70	16.80 -.39.78	40.70	.1398	n.s.

Table S2: Statistical data on the difference between two groups. Missing n-numbers in 3I are due to incomplete diary entries on weekends during the exploratory phase or because participants usually did not have a particular meal. The data in the table refers to the data shown in Fig. S2A, S2G, and 3I.

Fig.	Group	Variable value (mean ± SD)				Statistical test	d.f.	t-value	Significance	
		Completers	n-value	Drop-outs	n-value				p-value	Summary
S2A	CG	25.40 ± 3.882	33	27.65 ± 4.793	9	Welch's test	11.03	1.298	.2207	n.s.
	EG	27.32 ± 4.586	67	28.53 ± 4.932	12		14.61	.7873	.4437	n.s.
		female		male						
S2G	EG		41		26	t-test	65	1.238	.2203	n.s
		Exploration phase T0-T1		Intervention phase T1-T2						
3I	Breakfast	.0 ± .0	55	-.6477 ± 1.120	55	paired t-test	54	4.288	**<.0001**	****
	Lunch	.0 ± .0	64	-.6384 ± 1.346	64		63	3.796	**.0003**	***
	Dinner	.0 ± .0	63	-.0002648 ± 1.013	63		62	.0002648	.9984	n.s.

SD = Standard deviation

d.f. = degree of freedom

n.s. = not significant

Table S3: Statistical data from all Two-way-RM ANOVAs analyzing time, group, as well as interaction effects between time x group. In data for Figure 1, missing *p*-values in the and EG are due to the different number of meals for each participant. In the data for Figure 2, some participants were excluded from this analysis because of several probably erroneous diary entries about unrealistic amounts of food. In data for Figure 3, varying n-numbers result from incomplete entries in the diaries. In data for Figure 4, missing data are due to incomplete questionnaires.

Fig.	Meal	CG Exploration phase	CG Intervention phase	n-value	EG Exploration phase	EG Intervention phase	n-value	Statistical test	ANOVA results	Time	Group	Time x Group	CG p-val.	Sum.	EG p-val.	Sum.	T0-T1 p-val.	Sum.	T1-T2 p-val.	Sum.
1C	Breakfast	3.257; (2.742, 3.773)	4.352; (3.755, 4.948)	33	3.893; (3.508, 4.278)	1.722; (1.543, 1.902)	66		d.f.	1, 97	1, 97	1, 97	**.0012**	**	**<.0001**	****	.0564	n.s.	**<.0001**	****
									F-value	21.20	8.111	74.63								
									p-value	**<.0001**	**.0054**	**<.0001**								
									Summary	****	**	****								
	Lunch	3.331; (2.93. 3.731)	3.743; (3.352, 4.135)	33	4.021; (3.75. 4.292)	1.668; (1.484, 1.852)	67	Two-way RM ANOVA	d.f.	1, 98	1, 98	1, 98	.0543	n.s.	**<.0001**	****	**.0030**	**	**<.0001**	****
									F-value	14.45	74.51	151.4								
									p-value	**.0003**	**<.0001**	**<.0001**								
									Summary	***	****	****								
	Dinner	3.358; (3.033, 3.633)	3.807; (3.527, 4.087)	33	3.740; (3.477, 4.002)	1.778; (1.619, 1.937)	64		d.f.	1, 95	1, 95	1, 95	**.0384**	*	**<.0001**	****	.0814	n.s.	**<.0001**	****
									F-value	32.57	42.20	107.8								
									p-value	**<.0001**	**<.0001**	**<.0001**								
									Summary	****	****	****								
1D	Total	3.353; (3.066, 3.640)	3.936; (3.681, 4.192)	33	3.879; (3.658, 4.101)	1.723; (1.573, 1.873)	67	Two-way RM ANOVA	d.f.	1, 98	1, 98	1, 98	**.0011**	**	**<.0001**	****	**.0032**	**	**<.0001**	****
									F-value	41.87	61.87	187.7								
									p-value	**<.0001**	**<.0001**	**<.0001**								
									Summary	****	****	****								
2A	Daily Kj intake (x-fold change)	1.0; (….0)	1.011; (.9599, 1.062)	30	1.0; (….0)	.9718; (.927. 1.017)	67	Two-way RM ANOVA	d.f.	1, 95	1, 95	1, 95	>.9999	n.s.	.3604	n.s.	>.9999	n.s.	.2877	n.s.
									F-value	.2152	1.077	1.077								
									p-value	.6438	.3020	.3020								
									Summary	n.s.	n.s.	n.s.								
S2K	Average daily Kj intake	7212; (6615, 7808)	7248; (6588, 7907)	30	7071; (6619, 7523)	6723; (6318, 7128)	67	Two-way RM ANOVA	d.f.	1, 95	1, 95	1, 95	>.9999	n.s.	**.0271**	*	>.9999	n.s.	.3416	n.s.
									F-value	1.571	.8500	.1256								
									p-value	.2131	.3589	.1256								
									Summary	n.s.	n.s.	n.s.								
2E	% Fat	20.73; (19.37, 22.09)	22.06; (20.83, 23.28)	30	21.85; (20.92, 22.78)	21.40; (20.63, 22.17)	67		d.f.	1, 95	1, 95	1, 95	**.0394**	*	.4577	n.s.	.2895	n.s.	.7767	n.s.
									F-value	1.686	.1115	6.998								
									p-value	.1973	.7391	**.0095**								
									Summary	n.s.	n.s.	**								
	% Carbo-hydrates	57.51; (55.07, 59.95)	56.54; (54.4. 58.67)	30	56.23; (54.7. 57.75)	55.90; (54.62, 57.19)	67	Two-way RM ANOVA	d.f.	1, 95	1, 95	1, 95	.4796	n.s.	>.9999	n.s.	.6494	n.s.	>.9999	n.s.
									F-value	1.719	.6365	.4293								
									p-value	.1930	.4270	.5139								
									Summary	n.s.	n.s.	n.s.								
	% Pro-teins	21.77; (20.1. 23.43)	21.41; (20.03, 22.78)	30	21.82; (20.82, 23.02)	22.70; (21.71, 23.69)	67		d.f.	1, 95	1, 95	1, 95	.9850	n.s.	.0564	n.s.	>.9999	n.s.	.3322	n.s.
									F-value	.4446	.6848	3.284								
									p-value	.5065	.4100	.0731								
									Summary	n.s.	n.s.	n.s.								
3B	95%ile calorie intake interval	13.29; (12.24, 14.34)	13.73; (12.9. 14.56)	32	13.00; (12.45, 13.56)	10.99; (10.47, 11.52)	67	Two-way RM ANOVA	d.f.	1, 97	1, 97	1, 97	0.4686	n.s.	**<.0001**	****	>.9999	n.s.	**<.0001**	****
									F-value	12.31	11.15	30.00								
									p-value	**.0007**	**.0012**	**<.0001**								
									Summary	***	**	****								
3C	Breakfast time	8.39 ± 1.685	8.06 ± 1.434	33	8.24 ± 1.516	9.37 ± 1.633	67	Two-way RM ANOVA	d.f.	1, 98	1, 98	1, 98	0.2939	n.s.	**<.0001**	****	>.9999	n.s.	**.0002**	***
									F-value	8.122	3.637	27.49								
									p-value	**.0053**	.0594	**<.0001**								
									Summary	**	n.s.	****								
3D	Dinner time	21.26± 1.139	21.53± 1.217	32	20.96± 1.143	20.3 ±.6156	67	Two-way RM ANOVA	d.f.	1, 97	1, 97	1, 97	0.2969	n.s.	**<.0001**	****	.4725	n.s.	**<.0001**	****
									F-value	3.199	11.06	17.51								
									p-value	.0768	.0012	**<.0001**								
									Summary	n.s.	**	****								

Fig	Measure	Col A	Col B	n	Col C	Col D	n	Test	Stat	(1)	(2)	(3)								
4A	SF-36: PF	98.03; (96.95, 99.11)	98.33; (97.38, 99.29)	33	93.73; (91.45, 96.01)	96.27; (94.79, 97.75)	67	Two-way RM ANOVA	d.f.	1, 98	1, 98	1, 98	>.9999	n.s.	**.0020**	**	**.0055**	**	.2940	n.s.
									F-value	4.769	6.375	2.951								
									p-value	**.0314**	**.0132**	.0890								
									Summary	*	*	n.s.								
	SF-36: RLPH	98.49; (96.34, 100.63)	94.70; (88.14, 101.26)	33	90.30; (84.89, 95.71)	97.76; (94.63, 100.90)	67		d.f.	1, 98	1, 98	1, 98	0.7540	n.s.	**.0288**	*	**.0464**	*	.7856	n.s.
									F-value	.4966	1.092	4.655								
									p-value	.4827	.2985	**.0334**								
									Summary	n.s.	n.s.	*								
	SF-36: BP	93.64; (89.86, 97.42)	92.73; (87.2, 98.25)	33	86.57; (82.22, 90.91)	91.57; (88.05, 95.08)	67		d.f.	1, 98	1, 98	1, 98	>.9999	n.s.	.0779	n.s.	.0626	n.s.	>.9999	n.s.
									F-value	.9674	2.688	2.017								
									p-value	.3278	.1043	.1586								
									Summary	n.s.	n.s.	n.s.								
	SF-36: GH	76.06; (69.58, 82.54)	82.27; (77.05, 87.50)	33	72.01; (68.64, 75.39)	81.79; (78.78, 84.81)	67		d.f.	1, 98	1, 98	1, 98	**0.0263**	*	**<.0001**	****	.3732	n.s.	>.9999	n.s.
									F-value	28.30	.7258	1.406								
									p-value	**<.0001**	.3963	.2386								
									Summary	****	n.s.	n.s.								
	SF-36: Vit	66.06; (61.07, 71.05)	65.91; (60.16, 71.66)	33	55.97; (51.97, 59.97)	68.62; (64.37, 72.87)	67		d.f.	1, 98	1, 98	1, 98	>.9999	n.s.	**<.0001**	****	**.0084**	**	.8745	n.s.
									F-value	11.66	1.552	12.23								
									p-value	**.0009**	.2158	**.0007**								
									Summary	***	n.s.	***								
	SF-36: SF	91.29; (86.65, 95.92)	89.02; (82.48, 95.55)	33	87.68; (83.233, 92.14)	90.63; (85.97, 95.28)	67		d.f.	1, 98	1, 98	1, 98	0.9033	n.s	.3271	n.s	.6901	n.s.	>.9999	n.s.
									F-value	.03326	.08925	2.023								
									p-value	.8557	.7658	.1581								
									Summary	n.s.	n.s.	n.s.								
	SF-36: RLEP	91.92; (85.99, 97.85)	86.87; (76.24, 97.50)	33	83.58; (75.41, 91.76)	93.04; (87.64, 98.43)	67		d.f.	1, 98	1, 98	1, 98	0.7519	n.s	**.0393**	*	.2999	n.s	.5728	n.s.
									F-value	.4026	.05550	4.371								
									p-value	.5272	.8143	**.0392**								
									Summary	n.s.	n.s.	*								
	SF-36: EWB	77.94; (74.36, 81.52)	76.85; (72.24, 81.46)	33	72.93; (68.74, 77.11)	81.07; (77.08, 85.07)	67		d.f.	1, 98	1, 98	1, 98	>.9999	n.s.	**.0002**	***	.2491	n.s.	.3901	n.s.
									F-value	4.207	.02041	7.210								
									p-value	**.0429**	.8867	**.0085**								
									Summary	*	n.s.	**								
	PSQI	4.032; (3.177, 4.887)	4.097; (3.325, 4.868)	31	5.921; (5.219, 6.622)	3.365; (2.894, 3.836)	63	Two-way RM ANOVA	d.f.	1, 92	1, 92	1, 92	>.9999	n.s.	**<.0001**	****	**.0006**	***	.3055	n.s.
									F-value	21.75	1.776	24.06								
									p-value	**<.0001**	.1860	**<.0001**								
									Summary	****	n.s.	****								
	IDS-SR	8.818; (6.621, 11.016)	7.121; (5.312, 8.931)	33	12.94; (10.86, 15.02)	7.662; (5.887, 9.436)	66	Two-way RM ANOVA	d.f.	1, 96	1, 96	1, 96	0.3229	n.s	**<.0001**	****	**.0155**	*	>.9999	n.s.
									F-value	22.30	3.019	5.877								
									p-value	**<.0001**	.0855	**.0172**								
									Summary	****	n.s.	*								
	SWE	32.70; (31.28, 34.12)	33.21; (31.99, 34.43)	33	30.87; (29.79, 31.94)	32.46; (31.36, 33.57)	67	Two-way RM ANOVA	d.f.	1, 98	1, 98	1, 98	.7093	n.s.	**.0002**	***	.0873	n.s.	.8140	n.s.
									F-value	9.741	2.382	2.556								
									p-value	**.0024**	.1259	.1131								
									Summary	**	n.s.	n.s.								
4B	MCTQ	3.802; (3.37, 4.234)	3.720; (3.223, 4.217)	32	3.746; (3.465, 4.026)	3.828; (3.542, 4.114)	66	Two-way RM ANOVA	d.f.	1, 98	1, 98	1, 98	>.9999	n.s.	.8189	n.s.	>.9999	n.s.	>.9999	n.s.
									F-value	$1.85e^{-6}$	.01149	.8940								
									p-value	.9989	.9149	.3468								
									Summary	n.s.	n.s.	n.s.								

SD = Standard deviation
RM = Repeated Measure
n.s. = not significant
PF = Physical functioning
BP = Bodily pain
VT = Vitality
RE = Role of limitation emotional Problems
PSQI = Pittsburgh Sleep Quality Index
SWE = Scale of General Expectations of self-efficacy

CG = Control Group, EG = Experimental Group
d.f. = degree of freedom
SF-36 = Short Form 36 Health Survey Questionnaire
RP = Role of limitation physical functioning
GH = General Health
SF = Social functioning
EWB = Emotional well-being
IDS-SR = Self-Assessment Inventory of Depressive Symptoms
MCTQ = Munich Chronotype Questionnaire

MTVS = Mealtime Variability Score

Table S4: Statistical data from the development of normalized body weight and BMI of participants during the exploration and the intervention phase, and of follow-up data of continuing and discontinuing participants. At T3, three participants could no longer be reached, which is why the n-number deviates here. The data in the table refers to the data shown in Figures 1E and S2A.

Fig.	Variable	Group	Baseline T0	Explor. end T1	Interv. end T2	Follow-up T3	n-value	Statistical test	d.f.	F/t-value	p-value	Summary	T0 vs. T1 p-val.	Sum.	T1 vs. T2 p-val.	Sum.	T2 vs. T3 p-val.	Sum.
1E	BW	CG	.0 ± .0	-.1697 ± .9777	-.5606 ± 1.720		33	RM 1-way-ANOVA	2, 32	2.69	.0918	n.s.	.6524	n.s.	.2734	n.s.	.9787	n.s.
		EG	.0 ± .0	-.2104 ± 1.048	-2.621 ± 2.226		67	RM 1-way-ANOVA	2, 66	83.76	**<.0001**	****	.2099	n.s.	**<.0001**	****		
	BW_c	EG			-3.379 ± 2.100	-4.500 ± 2.696	34	paired t-test	33	6.369	**<.0001**	°°°°						
	BW_d				-2.077 ± 2.110	-1.597 ± 2.152	30		29	3.487	**.0016**	°°						
S2B	BMI	CG	.0 ± .0	-.06364 ± .3219	-.1970 ± .5537		33	RM 1-way-ANOVA	2, 32	3.12	.0658	n.s.	.5292	n.s.	.2418	n.s.	.9524	n.s.
		EG	.0 ± .0	-.0620 ± .3446	-.8663 ± .7376		67	RM 1-way-ANOVA	2, 66	83.75	**<.0001**	****	.2913	n.s.	**<.0001**	****		
	BMI_c	EG			-1.244 ± .6465	-1.728 ± .7676	34	paired t-test	33	6.072	**<.0001**	°°°°						
	BMI_d				-.7033 ± .7289	-.5392 ± .7147	30		29	3.433	**.0018**	°°						

SD = Standard deviation

CG = Control Group, EG = Experimental Group

BW = body weight

BMI = Body mass index

RM = Repeated Measure

_c = continuing

_d = discontinuing

d.f. = degree of freedom

n.s. = not significant

Table S5: Statistical data from the development of normalized body weight and BMI of participants at the different phases of the study. At T3 only continuing participants were included, which is why the n-number deviates here from the other time points. The data in the table refers to the data shown in Figures 1F, S2C, and S2H.

Fig.	Body weight change (mean ± SD)																Statistical test	ANOVA results	Source of Variation			Bonferroni post-hoc					
	T0				T1				T2				T3									T1		T2		T3	
	CG	n-value	EG	n-value	CG	n-value	EG	n-value	CG	n-value	EG	n-value	CG	n-value	EG	n-value			Time	Group	Time x Group	p-val.	Sum.	p-val.	Sum.	p-val.	Sum.
1F	.00 ±.00	33	.00 ±.00	67	-.170 ±.978	33	-.210 ±1.048	67	-.561 ±1.720	33	-2.621 ±2.226	67	-.796 ±1.992	26	-4.500 ±2.696	34	Mixed-effects model	d.f.	1.6, 131.6	1, 98	3, 254	>.9999	n.s.	<.0001	****	<.0001	****
																		F-value	61.18	35.51	29.56						
																		p-value	<.0001	<.0001	<.0001						
																		Summary	****	****	****						
S2C	.00 ±.00	33	.00 ±.00	67	-.064 ±.322	33	-.060 ±.346	67	-.197 ±.554	33	-.875 ±.735	67	-.273 ±.640	26	-1.479 ±.865	34	Mixed-effects model	d.f.	1.7, 140.4	1, 98	3, 254	>.9999	n.s.	<.0001	****	<.0001	****
																		F-value	62.88	35.95	29.72						
																		p-value	<.0001	<.0001	<.0001						
																		Summary	****	****	****						
S2H	.00 ±.00	33	.00 ±.00	67	-.170 ±.978	33	-.210 ±1.048	67	-.367 ±1.58	42	-2.270 ±2.22	79					Mixed-effects model	d.f.	1.4, 137.9	1, 119	2, 196	>.9999	n.s.	<.0001	****		
																		F-value	35.35	11.14	19.62						
																		p-value	<.0001	.0011	<.0001						
																		Summary	****	**	****						
	.00 ±.00	33	.00 ±.00	67	-.064 ±.322	33	-.060 ±.346	67	-.13 ±.513	42	-.759 ±.735	79					Mixed-effects model	d.f.	1.4, 138	1, 119	2, 196	>.9999	n.s.	<.0001	****		
																		F-value	36.63	10.88	20.17						
																		p-value	<.0001	.0013	<.0001						
																		Summary	****	**	****						

SD = Standard deviation

CG = Control Group, EG = Experimental Group

d.f. = degree of freedom

n.s. = not significant

Table S6: Statistical data from all simple regression. The data in the table refers to the data shown in Figs. 3A, B, and C. In data for Figures 2B, D, and S2H, Some participants were excluded from this analysis because of several probably erroneous diary entries about unrealistic amounts of food. In data for Figures 3B-D, varying n-numbers result from incomplete entries in the diaries. In date for Figures 3E, G, J, and S2I, missing n-numbers are due to incomplete diary entries on weekends during the exploratory phase or because participants usually did not have a particular meal.

Fig.	Group		Variables	n-value	Statistical test	Slope ± SD	R²	F-value	d.f.	p-value	Summary
S2G	EG		Age	67	Linear Regression	.02203 ± .01896	.02033	1.349	1, 65	.2497	n.s.
1G	CG		Δ MTVS vs. Δ body weight	33		-.03350 ± .3929	.0002345	.007273	1, 31	.9326	n.s.
			Δ MTVS vs. Δ BMI	33		-.06239 ± .1099	.01029	.5743	1, 31	.5743	n.s.
	EG		Δ MTVS vs. Δ body weight	67	Linear Regression	.8064 ± .2802	.1130	8.284	1, 65	**.0054**	**
			Δ MTVS vs. Δ BMI	67		.2763 ± .09323	.1190	8.782	1, 65	**.0042**	**
1H	EG	B	Δ MTVS Breakfast vs. body weight	67		-.1995 ± .1439	.02874	1.923	1, 65	.1702	n.s.
			Δ MTVS Breakfast vs. BMI	67		-.07272 ± .04790	.03425	2.305	1, 65	.1338	n.s.
		L	Δ MTVS Lunch vs. body weight	67	Linear Regression	.6103 ± .2122	.1192	8.273	1, 65	**.0054**	**
			Δ MTVS Lunch vs. BMI	67		.2050 ± .07080	.1142	8.380	1, 65	**.0052**	**
		D	Δ MTVS Dinner vs. body weight	67		.5944 ± .2036	.1159	8.520	1, 65	**.0048**	**
			Δ MTVS Dinner vs. BMI	67		.2067 ± .06762	.1257	9.342	1, 65	**.0032**	**
S2I	EG		Baseline BMI vs. Δ BMI	67	Linear Regression	-.03515 ± .01779	.05974	4.130	1, 65	**.0465**	*
2C	CG		Δ KJ intake vs. Δ body weight	30		$6.526e^{-5} \pm .0003174$	.0001508	.04228	1, 28	.8386	n.s.
			Δ KJ intake vs. Δ BMI	30		$1.302e^{-5} \pm .0001046$	.0005529	.01549	1, 28	.9018	n.s.
	EG		Δ KJ intake vs. Δ body weight	67	Linear Regression	$-4.326e^{-5} \pm .00022$	.0005975	.03886	1, 65	.8443	n.s.
			Δ KJ intake vs. Δ BMI	67		$-3.172e^{-5} \pm 7.319^{e-5}$	.002881	.1878	1, 65	.6662	n.s.
S2L	CG		KJ Exploration vs. KJ Intervention	30	Linear Regression	.8787 ± .1271	.6305	47,78	1, 28	**<.0001**	****
	EG			67		.7094 ± .06780±	.6275	109,5	1, 65	**<.0001**	****
2D	CG		Δ cumulative KJ intake vs. Δ body weight	30		$1.205e^{-6} \pm 7.270^{e-6}$	.0009794	.02745	1, 28	.8696	n.s.
			Δ cumulative KJ intake vs. Δ BMI	30		$2.604e^{-7} \pm 2.395^{e-6}$	.0004219	.01182	1, 28	.9142	n.s.
	EG		Δ cumulative KJ intake vs. Δ body weight	67	Linear Regression	$-1.627e^{-6} \pm 5.237^{e-6}$	.001483	.09655	1, 65	.7570	n.s.
			Δ cumulative KJ intake vs. Δ BMI	67		$-9.296e^{-7} \pm 1.746^{e-5}$	.004341	.2834	1, 65	.5963	n.s.
2F	CG	Fat	Δ intake vs. Δ body weight	30		-.02193 ± .2513	.0002719	.007615	1, 28	.9311	n.s.
			Δ intake vs. Δ BMI	30		-.001989± .08280	2.062^{e-5}	.0005774	1, 28	.9810	n.s.
	EG		Δ intake vs. Δ body weight	67		.07281 ± .1600	.003176	.2071	1, 65	.6506	n.s.
			Δ intake vs. Δ BMI	67		.02129 ± .05344	.002436	.1587	1, 65	.6916	n.s.
	CG	Carb	Δ intake vs. Δ body weight	30		-.03933 ± .1090	.004631	.1303	1, 28	.7209	n.s.
			Δ intake vs. Δ BMI	30		-.01866 ± .03580	.009609	.2717	1, 28	.6063	n.s.
	EG		Δ intake vs. Δ body weight	67	Linear Regression	-.1421 ± .08956	.03728	2,517	1, 65	.1175	n.s.
			Δ intake vs. Δ BMI	67		-.04338 ± .03000	.03116	2,091	1, 65	.1530	n.s.
	CG	Prot	Δ intake vs. Δ body weight	30		.02385 ± .2127	.0004489	.01257	1, 28	.9115	n.s.
			Δ intake vs. Δ BMI	30		.02433 ± .06992	.004307	.1211	1, 28	.7304	n.s.
	EG		Δ intake vs. Δ body weight	67		-.3055 ± .1843	.04054	2,747	1, 65	.1023	n.s.
			Δ intake vs. Δ BMI	67		-.1050 ± .06147	.04293	2,916	1, 65	.0925	n.s.

					N	Test		R²	F	d.f.	p	sig
S2N	CG	Fat	Δ intake vs. Δ body weight		30		-.000138 ± .000476	.002939	.08254	1, 28	.7760	n.s.
			Δ intake vs. Δ BMI		30		-3.246e^{-5} ± .000157	.001526	.04279	1, 28	.8376	n.s.
	EG		Δ intake vs. Δ body weight		67		-8.216e^{-5} ± .000319	.001021	.06640	1, 65	.7975	n.s.
			Δ intake vs. Δ BMI		67		-4.923e^{-5} ± .000106	.003286	.2143	1, 65	.6450	n.s.
	CG	Carb	Δ intake vs. Δ body weight		30		9.359e^{-5} ± .000181	.008762	.2475	1, 28	.6227	n.s.
			Δ intake vs. Δ BMI		30		1.814e^{-5} ± 6.214^{e-5}	.003034	.08522	1, 28	.7725	n.s.
	EG		Δ intake vs. Δ body weight		67	Linear Regression	-.000212 ± .000141	.03345	2.249	1, 65	.1385	n.s.
			Δ intake vs. Δ BMI		67		-7.545e^{-5} ± 4.712^{e-5}	.03796	2.565	1, 65	.1141	n.s.
	CG	Prot	Δ intake vs. Δ body weight		30		-.000119 ± .000666	.001131	.03170	1, 28	.8600	n.s.
			Δ intake vs. Δ BMI		30		3.299e^{-5} ± .000219	.0008070	.02261	1, 28	.8815	n.s.
	EG		Δ intake vs. Δ body weight		67		-.000806 ± .000449	.04714	3.216	1, 65	.0776	n.s.
			Δ intake vs. Δ BMI		67		-.000281 ± .000150	.05154	3.532	1, 65	.0647	n.s.
3B	CG		Δ 95%ile interval vs. Δ body weight		32		.2213 ± .1476	.06972	2.248	1, 30	.1442	n.s.
			Δ 95%ile interval vs. Δ BMI		32	Linear Regression	.04259 ± .04231	.03268	1.013	1, 30	.3221	n.s.
	EG		Δ 95%ile interval vs. Δ body weight		67		-.04842 ± .1218	.002426	.1581	1, 65	.6922	n.s.
			Δ 95%ile interval vs. Δ BMI		67		-.02160 ± .04063	.004328	.2825	1, 65	.5968	n.s.
3C	CG		Δ Breakfast time vs. Δ body weight		33		.05080 ± .2608	.001223	.03794	1, 31	.8468	n.s.
			Δ Breakfast time vs. Δ BMI		33	Linear Regression	-.007265 ± .07335	.0003164	.009811	1, 31	.9217	n.s.
	EG		Δ Breakfast time vs. Δ body weight		67		.1142 ± .1833	.005937	.3882	1, 65	.5354	n.s.
			Δ Breakfast time vs. Δ BMI		67		.03573 ± .06123	.005210	.3404	1, 65	.5616	n.s.
3D	CG		Δ Dinner time vs. Δ body weight		32		.3149 ± .3682	.02380	.7313	1, 31	.3992	n.s.
			Δ Dinner time vs. Δ BMI		32	Linear Regression	-.01494 ± .1047	.0006776	.02034	1, 31	.8875	n.s.
	EG		Δ Dinner time vs. Δ body weight		67		-.1096 ± .2361	.003308	.2157	1, 65	.6439	n.s.
			Δ Dinner time vs. Δ BMI		67		-.05984 ± .07860	.008838	.5796	1, 65	.4492	n.s.
3E	EG		midsleep Phase vs. midEat Phase	Exploration weekend	65	Linear Regression	.5902 ± .1449	.2084	16.58	1,63	**.0001**	****
				Intervention	65		.6532 ± .1176	.3286	30.83	1, 63	**<.0001**	****
3G	EG		Δ Residuals vs. Δ BMI	mSP-mEP relation	65	Linear Regression	-.2522 ± .09116	.1084	7.656	1, 63	**.0074**	**
3J	EG		Correction of weekend meals vs. Δ BMI	Breakfast	55		.1427 ± .08443	.05114	2.856	1, 53	.0969	n.s.
				Lunch	64	Linear Regression	.0460 ± .06415	.008225	.5142	1, 62	.4760	n.s.
				Dinner	63		-.2490 ± .08055	.1355	9.558	1, 61	**.0030**	**
S2O	EG		Dev. weekend and scheduled meals	Breakfast	55		-.1174 ± .1031	.02391	1.298	1, 53	.2597	n.s.
				Lunch	64	Linear Regression	-.1409 ± .08067	.04690	3.051	1, 62	.0857	n.s.
				Dinner	63		-.03707 ± .1313	.001305	.07969	1, 61	.7787	n.s.
S2L	CG		KJ Exploration vs. KJ Intervention		30	Linear Regression	.8787 ± .1271	.6305	47,78	1, 28	**<.0001**	****
	EG				67		.7094 ± .06780±	.6275	109,5	1, 65	**<.0001**	****

BMI = Body mass index

CG = Control Group, EG = Experimental Group

MTVS = Mealtime Variability Score

d.f. = degree of freedom

n.s. = not significant

Table S7: Statistical data from BMI changes depending of the baseline BMI of each participant. The data in the table refers to the data shown in Figure S2J.

Fig.	BMI change (mean ± SD)												Statistical test	ANOVA results	Source of Variation			Bonferroni post-hoc					
	20.0-24.9				25.0-29.9				>30									20.0-24.9		25.0-29.9		>30	
	CG	n-value	EG	n-value	CG	n-value	EG	n-value	CG	n-value	EG	n-value			Weight	Group	Weight x Group	p-val.	Sum.	p-val.	Sum.	p-val.	Sum.
S2J	-.018 ±.396	17	-.742 ±.546	26	-.255 ±.468	11	-.776 ±.709	24	-.304 ±.728	5	-.951 ±.804	17	2-way ANOVA	d.f.	2, 94	1, 94	2, 94	<.0001	***	.0694	n.s.	.1295	n.s.
														F-value	1.012	19.03	.2299						
														p-value	.3673	<.0001	.7950						
														Summary	n.s.	****	n.s.						

SD = Standard deviation

CG = Control Group, EG = Experimental Group

d.f. = degree of freedom

n.s. = not significant

Table S8: Statistical data from multiple regression analysis testing the effects of different variables while keeping the others constant. The data in the table refers to the data shown in Figures 1H and 3K.

Fig.	Model summary		Model		Coefficients						
			Analysis of Variance	F (DFn, DFd)	Parameter estimates	Variable	Estimate	Stand. Err.	|t|	P value	Summary
1I			Regression	F (2, 64) = 5.729	β0	Intercept	.7183	.5139	1,398	.1670	n.s.
	R²	.1518	D_MTVS	F (1, 64) = 6.949	β1	D_MTVS	.2438	.09248	2,636	**.0105**	*
			Pre_BMI	F (1, 64) = 4.439	β2	Pre_BMI	-.03588	.01703	2,107	**.0391**	*
3K			Regression	F (6, 45) = 4.939	β0	Intercept	-.1691	.2434	.6947	.4908	n.s.
			D_MTVS	F (1, 45) = 5.977	β1	D_MTVS	.2582	.1056	2,445	**.0185**	*
			D_KJ	F (1, 45) = .1534	β2	D_KJ	$2,783^{e-5}$	7,105e-005	.3917	.6971	n.s.
	R²	.3970	Delta Intake Interval	F (1, 45) = 0.6980	β3	Delta Intake Interval	-.04606	.05513	.8355	.4079	n.s.
			Delta Residuals	F (1, 45) = 3.232	β4	Delta Residuals	-.2217	.1233	1,798	.0789	n.s.
			Breakfast Jetlag	F (1, 45) = 3.012	β5	Breakfast Jetlag	.1457	.08395	1,736	.0895	n.s.
			Dinner Jetlag	F (1, 45) = 13.01	β6	Dinner Jetlag	-.3071	.08513	3,608	**.0008**	***

BMI = Body mass index
MTVS = Mealtime variability score
d.f. = degree of freedom
Reg. Coeff = Regression coefficient
Std. Err. = Standard error

Table S9: Statistical data from Rayleigh's uniformity test analyzing phase distribution from the relation between midSleep Phase and midEat Phase. Missing n-numbers are due to incomplete diary entries on weekends during the exploratory phase. The data in the table refers to the data shown in Fig. 3F.

Fig.	Data set	Normalized midSleep Phase	Normalized midEat Phase					n-value
			Relative Phase	SD	Vector length	Rayleigh Z-value	p-value	
3F	Exploration weekend	.00	11.33	20.653°	.937	57.093	$<1e^{-12}$	65
	Intervention	.00	10.57	17.221°	.956	60.299	$<1e^{-12}$	65

SD = Standard deviation

Table S10: Questionnaires collected at T0 and T2 to assess general well-being of participants.

Construct	Questionnaire	Components
Sleep Quality	Pittsburgh Sleep Quality Index (PSQI)	Subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, daytime impairment.
Physical Well-Being	36-Item Short Form Health Survey (SF-36)	Vitality, physical functionality, physical pain, general health perception, physical role function, emotional role function, social function, mental well-being.
Self-Efficacy	Scale of General Expectations of Self-Efficacy (SWE)	Optimistic competence expectation, daring to solve life problems independently, attribution of success to own competences.
Depressive Mood	Self-Assessment Inventory of Depressive Symptoms (IDS-SR)	Sleep quality, mood, appetite, weight changes, concentration/decision making, self-esteem, future expectations, suicidal thoughts, activities, energy levels, pleasure/joy, sexual interest, slowing down, restlessness, physical/vegetative complaints, panic/anxiety, digestive problems, interpersonal sensitivity, physical heaviness.
Chronotype	Munich Chronotype Questionnaire (MCTQ)	Bedtimes, sleep times, energy levels, exposure to natural light, differences between weekdays and weekends.

A 95%ile calorie intake interval during exploration and intervention phase

Local time of day / Participants

exploration phase T0-T1
intervention phase T1-T2

B 95%ile calorie intake interval

Δ Interval of food intake (h) / Time of measurement (CG, EG)

Δ Body Weight (kg) / Δ Interval of calorie intake — n.s. / n.s. / Δ BMI

C Breakfast time

Δ Average breakfast time (local time) / Time of measurement (CG, EG)

Δ Body Weight (kg) / Shift of breakfast time (h) — n.s. / n.s. / Δ BMI

D Dinner time

Δ Average dinner time (local time) / Time of measurement (CG, EG)

Δ Body Weight (kg) / Shift dinner time (h) — n.s. / n.s. / Δ BMI

○ Exploration Phase T0-T1 — Delta Body Weight
● Intervention Phase T1-T2 — Delta BMI

E mid-Sleep Phase vs. mid-Eat Phase

Exploration Weekend: **** $p = 0.0001$ $r^2 = 0.2084$

Intervention: **** $p < 0.0001$ $r^2 = 0.3286$

mid-Eat Phase / mid-Sleep Phase

F Phase relation mid-Sleep Phase and mid-Eat Phase

Exploration Weekend — mEP / mSP

Intervention — mEP / mSP

G Improvement of mSP-mEP relation vs. weight change

Δ BMI / Change in residuals (h) ** $p = 0.0074$ $r^2 = 0.1084$

H Example of mealtime shift on weekend

Time of day / Wed Thu Fri Sat Sun Mon Tue

I Shift of weekend meal times to calculated meal times

Breakfast **** Lunch *** Dinner n.s.

Meal Shift (h) / T0-T1 W/E T1-T2

J Shifts of weekend meals vs. weight loss

Breakfast: n.s. $p = 0.097$ $r^2 = 0.051$

Lunch: n.s. $p = 0.4760$ $r^2 = 0.008$

Dinner: ** $p = 0.003$ $R^2 = 0.1355$

Δ BMI / Correction of weekend/workday meal time discrepancy (h)

K Multiple Regression QQ plot

Predicted residual / Actual residual

Multiple Regression Actual vs. Predicted plot

Predicted Y / Actual Y

$R^2 = 0.3970$

	Estimate	P-value
β0 - Intercept	-0.1691	0.4908
β1 - ΔMTVS	0.2582	0.0185 *
β2 - ΔkJ intake	0.00003	0.6971
β3 - ΔCalorie intake interval	-0.04606	0.4079
β4 - ΔmSp-mEP relation	-0.2271	0.0789
β5 - Δweekend breakfast	0.1457	0.0895
β6 - Δweekend dinner	-0.3071	0.0008 ***

L Covariance matrix

	β0	β1	β2	β3	β4	β5	β6
β0	1.00	0.75	0.12	0.21	-0.10	0.35	-0.10
β1	0.75	1.00	-0.02	-0.36	0.14	-0.08	0.01
β2	0.12	-0.02	1.00	0.10	-0.01	-0.03	-0.17
β3	0.21	-0.36	0.10	1.00	-0.27	0.42	-0.12
β4	-0.10	0.14	-0.01	-0.27	1.00	-0.18	-0.11
β5	0.35	-0.08	-0.03	0.42	-0.18	1.00	-0.12
β6	-0.10	0.01	-0.17	-0.12	-0.11	-0.12	1.00

SF-36 - Physical and Mental Health

○ T0
● T2