NOTCH1 fusion genes in pediatric T-cell lymphoblastic lymphoma hallmark a common high-risk subgroup with blood TARC levels as possible diagnostic biomarker

Emma Kroeze[1], Michelle M Kleisman[1], Lennart A Kester[1], Marijn A Scheijde-Vermeulen[1], Edwin Sonneveld[1], Jessica GC Buijs-Gladdines[1], Melanie M Hagleitner[1], Friederike AG Meyer-Wentrup[1], Margreet A Veening [1], Auke Beishuizen [1], Jules PP Meijerink [1,†], Jan LC Loeffen[1*], Roland P Kuiper[1,2]*

[1]*Princess Máxima Center for pediatric oncology, Utrecht, the Netherlands*

[2]*Department of Genetics, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands*

[†]*Presently working at Acerta-Pharma (AstraZeneca), Oss, the Netherlands*

*JLCL and RPK are considered co-last authors

Corresponding author: Prof. dr. Roland P Kuiper, Princess Maxima Center for Pediatric Oncology, Heidelberglaan 25, 3584 CS Utrecht, Netherlands, r.kuiper@prinsesmaximacentrum.nl

Abstract

Background: Twenty percent of children with T-cell lymphoblastic lymphoma (T-LBL) will relapse and have an extremely poor outcome. Currently, we can identify a genetically low-risk subgroup in pediatric T-LBL, yet these high-risk patients who need intensified or alternative treatment options remain undetected. Therefore, there is an urgent need to proactively recognize these high-risk T-LBL patients through identification of specific molecular characteristics and generic biomarkers.

Patients and Methods: In 29/43 unselected pediatric T-LBL patients diagnosed in the Princess Maxima Center for Pediatric Oncology between 2018-2023, RNA sequencing was performed at diagnosis, and blood *thymus and activation-regulated chemokine* (TARC) levels were measured at diagnosis in 22/29 patients.

Results: We discovered a previously unknown high-risk biological subgroup of children with T-LBL, characterized by *NOTCH1* gene fusions, which was found in 21% of our T-LBL cohort (6/29). All patients presented uniformly with a large mediastinal mass, pleural/pericardial effusions, and absence of blasts in the bone marrow, blood, and central nervous system. Moreover, all six patients exclusively expressed highly elevated blood TARC levels, defining a novel and previously not known clinically relevant blood biomarker for T-cell lymphoblastic lymphoma. Three out of these six patients relapsed during therapy, while a fourth developed a therapy related acute myeloid leukemia during maintenance therapy.

Conclusion: We identified a frequently occurring T-LBL entity with high risk of relapse which can be easily identified using a blood TARC screening at diagnosis. Further molecular characterization through *NOTCH1* gene fusion analysis offers these patients the opportunity for treatment intensification or new treatment strategies.

Introduction

T-cell lymphoblastic lymphoma (T-LBL) is a common pediatric malignancy accounting for approximately 20% of the non-Hodgkin lymphomas during childhood[1]. Survival rates of T-LBL are ~80%, but outcome after relapse is dismal, with salvage rates reaching only ~15%[2,3]. Considering the extremely poor prognosis after relapse and absence of clinically relevant high-risk genetics, there is an urgent need for the identification of molecular risk factors and new prognostic biomarkers in T-LBL, as well as identification of new therapeutic strategies.

Pediatric T-LBL is typically characterized by infiltration of blasts in the mediastinum (thymus) and lymph nodes. Approximately half of the cases present with pleural effusion at diagnosis and, by definition, T-LBL patients have less than 25% blasts in the bone marrow (BM)[4,5]. Based on morphology and immunophenotype, T-LBL is indistinguishable from its leukemic counterpart, T-cell acute lymphoblastic leukemia (T-ALL). T-ALL presents as leukemic disease with ≥25% blasts in the BM and presence of blasts in the peripheral blood (PB), but usually with mild mediastinal enlargement. Even though the clinical presentation of T-LBL and T-ALL largely differs, there has been no evidence so far that there are major differences in molecular genetics of T-LBL and T-ALL[6,7]. Therefore, T-LBL and T-ALL are regarded as different manifestations of the same disease[8]. Pediatric T-ALL has been extensively studied and great advances in treatment protocols have been made, including minimal residual disease (MRD) measurements as a useable stratification strategy. In contrast, molecular genetics underlying T-LBL are still poorly understood and T-LBL patients are currently mainly treated according to ALL-based protocols. Thus far, MRD assessment has not proven to be usable in T-LBL and diagnostic biomarkers to identify high-risk patients are lacking.

Diagnostic biomarkers have been successfully introduced in other types of lymphomas. For

example, blood *thymus and activation-regulated chemokine* (TARC, CCL17) has proven to

be a useful diagnostic biomarker in children with classical Hodgkin lymphoma[9,10]. It has

never been studied whether TARC can serve as a biomarker in non-Hodgkin lymphoma and

specifically T-LBL as well. TARC is constitutively produced in the thymus, acting as a

powerful T-cell chemoattractant. In classical Hodgkin lymphoma patients, TARC production

can be induced by NOTCH1 and TARC is highly expressed by the Reed-Sternberg cells,

thereby creating a specific supporting tumor microenvironment that recruits T-cells[10].

Considering the importance of NOTCH1 and strong preference of malignant T-cells in T-LBL

for the thymus, we hypothesize that TARC is of importance in the pathophysiology of T-LBL

and creating a thymic microenvironment that favors the T-LBL cells.

Recent studies show that both T-LBL and T-ALL patients with *NOTCH1* and/or *FBXW7*

DNA mutations have a better outcome compared to *NOTCH1* and *FBXW7* wildtype

patients[11,12]. The pathogenic molecular mechanisms of T-LBL patients without *NOTCH1*

and/or *FBXW7* mutations remain largely unknown, and this group probably contains both

high-risk and low-risk patients. Considering the extremely poor prognosis after relapse, it is

essential to identify these high-risk patients. Additionally, there is an urgent need for the

identification of new prognostic biomarkers in T-LBL. In this study we present a novel entity

of pediatric T-LBL patients characterized by previously unknown *NOTCH1* gene fusions and

highly elevated blood TARC levels

Methods

Patients

Pediatric T-LBL patients (n=43) were diagnosed in the Princess Máxima Center for Pediatric

Oncology between 2018-2023. RNA sequencing data at diagnosis were available for 29/43

patients and at relapse for two additional patients. Clinical information and hematological

values at diagnosis were retrieved from patient files. All patients were treated according to

the EURO-LB02 protocol[3] or its successor, the LBL2018 protocol (NCT04043494).

NOTCH1/FBXW7 mutational status was determined for most patients and retrieved from

patient files. Pediatric T-ALL patients (n=39) diagnosed between 2019-2022 at the Princess

Máxima Center for Pediatric Oncology were included as reference cohort. All sample IDs are

completely anonymized. Written informed consent was obtained from all patients and/or their

legal guardians. The study was performed in accordance with the Declaration of Helsinki.

The Medical Research Ethical Committee Utrecht declared that the Medical Research

Involving Human Subjects Act (WMO) does not apply, and approved the study (19-140/C).

RNA sequencing analysis

RNA sequencing data was obtained from the in-house diagnostics department in the Princess

Máxima Center. Pre-processing of the data was done with standardized and in-house

pipelines and guidelines[13]. Fusion detection was performed using STAR fusion[14]. In addition,

we analyzed the entire T-LBL and T-ALL cohort for exon-specific *NOTCH1* coverage as an

indication for fusions, by using DepthOfCoverage of GATK v3.8.0. Differential expression

analysis was performed with DESeq2 v1.36.0. Gene expression alterations were assessed

with log2 transformed gene length normalized read counts (transcript per million mapped

reads, TPM). Whole genome sequencing data were also obtained from the in-house

diagnostics department and used for validation of genomic breakpoints.

TARC measurements and immunohistochemistry

TARC measurements at diagnosis were performed in serum or plasma of 22 out of the 29

patients for whom RNAseq was available. Measurements were performed by

standard enzyme-linked immunosorbent assay (ELISA) using the DuoSet ELISA kit (cat. no.

DY364; R&D Systems, Inc.). Immunohistochemistry staining for TARC was performed on

the BOND-III fully automated staining system (Leica, IL, USA) using CCL17 rabbit

polyclonal antibody (ProteinTech group, Chicago IL, USA).

Results

Patient characteristics

Pediatric T-LBL patients (n=43) were diagnosed in the Princess Máxima Center for Pediatric

Oncology between 2018-2023. T-LBL diagnoses were based on histopathological

classification according to the revised world health organization for hematological

malignancies and/or flow cytometry according to the criteria of the European Group for

Immunophenotyping of Leukemias[8,15]. The median age at diagnosis for these 43 patients was

ten years, and our cohort contained more males than females (59%), in line with previously

described large T-LBL cohorts[3]. Eighty-eight percent of the patients presented with a large

mediastinal mass, which was in 53% of these cases accompanied by pleural effusion. Five out

of 43 patients (12%) had a relapse. RNAseq was performed for 29/43 patients and their

clinical characteristics are described in **Supplementary Table 1**.

Detection of NOTCH1 rearrangements

Transcriptome sequencing was performed at diagnosis from 2019 onwards when possible,

and was available for 29 patients in our cohort. This technique allows for the identification

and quantification of fusion transcripts and gene expression levels. We identified 12 fusions

transcripts (41%), including six *NOTCH1* gene fusions, with the microRNA gene *miR142HG*

on chromosome 17q22 (TLBL042 and TLBL058), *IKZF2* on chromosome 2q34 (TLBL050)

and *TRBJ* on chromosome 7q34 (TLBL033, TLBL049, TLBL050) (**Supplementary Table**

2). The fusion transcripts with *miR142HG* and *IKZF2* demonstrated correct splicing to exon

27 or 28 of *NOTCH1* and genomic breakpoints were identified using available whole genome

sequencing (WGS) data (**Supplementary Table 2**). *TRBJ::NOTCH1* fusion genes were

previously shown to express a truncated, membrane-bound form of NOTCH1[16] (**Figure 1A**).

None of the six samples with a *NOTCH1* gene fusion exhibited mutations in

NOTCH1/*FBXW7*, demonstrating the mutual exclusivity of *NOTCH1* mutations and

NOTCH1-rearrangements. Furthermore, apart from homozygous loss of the *CDKN2A/2B*

locus in two cases (TLBL042 and TLBL050), no other driver events were found in these

NOTCH1-rearranged T-LBLs. *NOTCH1* gene fusions are almost never found in T-ALL, but

considering the difficulties in detecting TR-rearranged fusions with conventional pipelines,

we also reanalyzed exon-specific *NOTCH1* expression in 39 T-ALL samples that were

diagnosed in the Princess Máxima Center for Pediatric Oncology. In line with previous

studies[17,18], none of the T-ALL samples carried gene fusions involving *NOTCH1*, thus

suggesting that the frequent occurrence of these fusions represents an important molecular

genetic discriminator between T-LBL and T-ALL.

All NOTCH1 gene fusions result in expression of intracellular NOTCH1

NOTCH1 is a transmembrane protein that is activated through ligand-receptor interaction, which induces a conformational change that results in dissociation of the heterodimerization (HD) subunits. This is followed by exposure of a cleavage site in the C-terminal part of the HD domain, resulting in the release of the intracellular domain of NOTCH1 (NICD). NICD subsequently translocates to the nucleus where it acts as a transcriptional regulator[19].

To determine whether a truncated C-terminal version of NOTCH1 was indeed expressed in the T-LBL samples with a *NOTCH1*-rearrangement, we performed Western blotting using protein lysates of four *NOTCH1*-rearranged cases and two *NOTCH1* wildtype (*NOTCH1*-WT) cases. All *NOTCH1*-rearrangements lead to expression of NICD. The *IKZF2::NOTCH1*-positive T-LBL (TLBL050) showed expression of an NICD protein co-migrating with the ~130-kDa wildtype NICD, suggesting cleavage of the chimeric protein at the □-secretase cleavage side (Val1744). In contrast, the *TRBJ::NOTCH1*-positive (TLBL033 and TLBL052) and *miR-142HG::NOTCH1*-positive (TLBL042) cases expressed a slightly larger NICD protein, which is in line with translation initiation at methionine residue 1727 (Met1727) encoded in exon 28 of *NOTCH1*, as previously reported[16,20], and lack of □-secretase cleavage (**Figure 1B**).

To explore downstream characteristics of T-LBL cases with *NOTCH1* gene fusions, we performed differential expression analysis between *NOTCH1*-rearranged and *NOTCH1*-WT T-LBLs. Two *NOTCH1*-rearranged cases (TLBL049 and TLBL058) had to be excluded from this analysis because of lower quality of the expression data. A total of 1,621 genes were found to be significantly differentially expressed, compared to only 65 genes in a comparison between *NOTCH1*-mutated and *NOTCH1*-WT T-LBLs (**Supplementary Figure 1A,B** and **Supplementary Tables 3 and 4**). Subsequently, the 100 most significantly differentially expressed genes between *NOTCH1*-rearranged and *NOTCH1*-WT T-LBLs, were selected for supervised clustering with *NOTCH1*-WT, *NOTCH1*-mutated, and all six *NOTCH1*-rearranged

samples. This analysis revealed that *NOTCH1*-rearranged samples formed a separate cluster

(**Figure 1C**). To confirm these findings, we performed clustering analysis with the 200 most

variable genes in the dataset (**Supplementary Table 5**), which showed that *NOTCH1*-

rearranged samples mostly clustered separately from the entire T-LBL cohort

(**Supplementary Figure 1C**). These findings suggest that the downstream effects and

mechanisms of action between *NOTCH1*-mutated and *NOTCH1*-rearranged T-LBLs is

different.

Clinical presentation of patients with NOTCH1 gene fusions

All six patients with *NOTCH1* gene fusions presented with a massively enlarged

mediastinum, combined with pleural/pericardial effusion. Moreover, all of the *NOTCH1* gene

fusion-positive patients were bone marrow negative, peripheral blood negative and cerebral

spinal fluid negative. These patients therefore had a uniform clinical presentation of disease,

although not differentiating between this group and the rest of the T-LBL cohort.

Flowcytometry performed at diagnosis revealed positivity for cytoplasmatic CD3 (cyCD3) in

all cases, as well as positivity for other T-cell markers. Precursor-marker Terminal

Deoxynucleotidyl Transferase (TdT) was expressed in 50% of the *NOTCH1*-rearranged

cases, which was lower than expected (~90%)[21] (**Supplementary Table 6**). Next, we

analyzed blood TARC (CCL17) levels, which were highly elevated in all patients with a

NOTCH1 gene fusion (range from 2345 to >10000 pg/ml), compared to 31-638 pg/ml in 16

patients who did not have a *NOTCH1* gene fusion (p<0.0001) (**Figure 2A**). Follow-up TARC

levels during first remission were available for 3/6 patients and revealed normalized values

(range 57-152 pg/ml) (**Figure 2B**). Two of the patients with a clinical relapse (TLBL042 and

TLBL050) also showed substantially elevated TARC levels at relapse (TLBL042:4613 pg/ml,

TLBL050:8654 pg/ml), which could be an indication that TARC levels in blood might also

increase upon progression of relapse. One patient, whose relapse was discovered with routine

imaging, presented with relatively little tumor load and low LDH levels, and did not have

increased TARC levels (TLBL033). TARC levels normalized again in second remission

(range 69-1331 pg/ml) (**Figure 2C**). Our data therefore strongly indicate that TARC can serve

as a high-risk biomarker at diagnosis. Immunohistochemistry revealed that the T-LBL cells

were not positive for TARC, in line with the finding that *CCL17* was not significantly

upregulated in *NOTCH1*-rearranged cases. This observation suggests that TARC is produced

by the tumor microenvironment in the thymus, rather than the tumor cells (**Figure 2D**).

NOTCH1 gene fusions as poor prognostic marker in T-LBL

Finally, we aimed to determine the prognostic relevance of *NOTCH1* gene fusions in T-LBL.

We found that four out of six *NOTCH1* fusion-positive patients had an event. Three patients

had a relapse during therapy (TLBL033, TLBL042, TLBL050), none of them were rescued.

Additionally, one patient had a therapy-related acute myeloid leukemia (t-AML) during

maintenance therapy of T-LBL (TLBL049). The t-AML carried the typical *KMT2A*::*MLLT3*

fusion. This patient was rescued with AML induction chemotherapy followed by allogeneic

stem cell transplantation. The other two patients did not have an event, yet both patients are

still under treatment.

The unselected 5-year T-LBL cohort (n=43) contained five patients who relapsed, of whom

three had a *NOTCH1* gene fusion. For the other two patients, only RNAseq data at relapse

was available, revealing a *DDX3X*::*MLLT10* fusion and a *JAKMIP2*::*PDGFRB* fusion at

relapse, respectively, established high-risk ALL aberrations[17,22]. Thus, *NOTCH1*-rearranged

T-LBL caused the majority of relapses in our cohort, suggesting that they cause an aggressive

T-LBL phenotype, with a significantly higher prevalence of relapse (p=0·02) compared to the

NOTCH1-fusion negative patients.

Discussion

To date, mainly clinically applicable low-risk genetics have been described for T-LBL. This

implicates that molecular genetic high-risk patients who need intensified or alternative

treatment strategies are undetected. Moreover, patients with unknown low-risk molecular

genetic profiles could also be overtreated with current treatment strategies.

We discovered a biological high-risk subgroup of T-LBL, characterized by *NOTCH1* gene

fusions. This subgroup represents 21% (6/29 patients) of our T-LBL cohort. All patients had a

similar although not unique presentation of disease predominantly consisting of a large

mediastinal mass, pleural/pericardial effusion and highly elevated TARC levels in blood.

Moreover, three out of six patients with a *NOTCH1* gene fusion had a relapse and did not

survive, indicating that *NOTCH1* fusions lead to an aggressive T-LBL phenotype. Fifty

percent of the *NOTCH1*-rearranged cases exhibited expression of TdT, which is lower than

the expected 90%. It has been described before that the TdT-negative subset often represent

diagnostically challenging cases with phenotypic features that are consistent with a late

cortical subtype, coinciding with what we found in our cohort[21].

The presence and frequency of *NOTCH1* gene fusions can currently be regarded as a major

molecular genetic difference between T-LBL an T-ALL, since fusions involving *NOTCH1* are

only extremely rarely described in T-ALL (<0·1%)[18,23-28]. The uniform absence of BM and

PB involvement in T-LBL patients with a *NOTCH1* gene fusion coincides with the fact that

these rearrangements have almost never been detected in T-ALL. *NOTCH1* gene fusions have

been described in T-LBL before[25], but given the small number of samples in these studies, as

well as difficulties in detecting TR-rearranged fusions with conventional pipelines, these

fusions might have been missed explaining the lower contribution of *NOTCH1* gene fusions

in these studies.

NOTCH1 gene fusions appear to have a bigger impact on T-LBL cells compared to *NOTCH1*

mutations, with more and larger changes in the downstream gene expression, independent of

the type of *NOTCH1*-fusion. Furthermore, whereas patients with *NOTCH1* and/or *FBXW7*

mutations are considered low risk and have a better outcome compared to *NOTCH1* and/or

FBXW7 WT patients[11,12], we demonstrate that the outcome of these recurrent *NOTCH1*-

rearranged T-LBLs is poor. It has recently been described that *NOTCH1* intronic single

nucleotide variants (SNVs) and *NOTCH1* intragenic losses were also associated with an

inferior event-free and overall survival[28], further substantiating that distinct genetic

aberrations in *NOTCH1* have a different impact on outcome. The T-LBL patients with high-

risk *NOTCH1* aberrations will probably need intensified or alternative treatment strategies. A

second consequence of our findings may be that when *NOTCH1* gene fusions are recognized

as a separate high-risk group, the survival characteristics of the remaining group of T-LBL

patients with unknown molecular genetics improves and may benefit from less intensified

treatment.

The highly elevated blood TARC levels were exclusively observed in all T-LBL patients with

a *NOTCH1*-rearrangement, even though almost all T-LBL patients had an enlarged

mediastinum. Elevated TARC levels may therefore serve as an important biomarker to assist

in the diagnosis of this high-risk group at diagnosis. Moreover, there might be a possibility

that TARC levels could be used during follow-up as well, but these findings need to be

validated in larger cohorts. It has been suggested that inhibiting TARC, produced by the

Reed-Sternberg cells, may have therapeutic consequences in classical Hodgkin lymphoma as

inhibiting TARC can decrease the recruitment of T-cells, thereby affecting the supporting

microenvironment[29]. Although our data suggests that TARC expression in *NOTCH1*-

rearranged T-LBL originates from the T-LBL microenvironment instead of the tumor cells,

this increased blood biomarker likely point towards active crosstalk between the tumor cells

and the microenvironment[30]. It is therefore intriguing to further explore whether *NOTCH1*-

rearranged T-LBLs are dependent on TARC expression and whether this would provide

opportunities for targeted treatment in a potentially high-risk subgroup of T-LBL. Measuring

blood TARC levels could also serve as an easily applicable technique to identify high-risk T-

LBL patients in low- and middle-income countries with restricted access to next generation

sequencing techniques.

In conclusion, we discovered that, in contrast to T-ALL, *NOTCH1* gene fusions are common

in T-LBL and represent a high-risk subtype with an easily applicable biomarker. The

discovery of this clinically relevant high-risk T-LBL subgroup offers opportunities to develop

intensified and targeted treatment strategies for this subgroup and decrease overtreatment in

the remaining group of T-LBL patients.

Author contributions

298 RK, JLCL, AB designed the study. EK wrote the manuscript and analyzed the data. MMK

299 and LAK performed the bioinformatic analyses. JGCB performed the wet lab analyses. ES

300 provided laboratory supervision. MSV performed the histological analyses. MMH, FAGM,

301 MAV were involved in data curation. JPPM acquired the funding.

Declaration of interests

304 None of the authors declare a conflict of interest.

References

1. Minard-Colin V, Brugieres L, Reiter A, et al. Non-Hodgkin Lymphoma in Children and Adolescents: Progress Through Effective Collaboration, Current Knowledge, and Challenges Ahead. J Clin Oncol 2015;33(27):2963-74. DOI: 10.1200/JCO.2014.59.5827.

2. Burkhardt B, Reiter A, Landmann E, et al. Poor outcome for children and adolescents with progressive disease or relapse of lymphoblastic lymphoma: a report from the berlin-frankfurt-muenster group. J Clin Oncol 2009;27(20):3363-9. DOI: 10.1200/JCO.2008.19.3367.

3. Landmann E, Burkhardt B, Zimmermann M, et al. Results and conclusions of the European Intergroup EURO-LB02 trial in children and adolescents with lymphoblastic lymphoma. Haematologica 2017;102(12):2086-2096. DOI: 10.3324/haematol.2015.139162.

4. Swerdlow SH, Campo E, Lee Harris N, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues (Revised 4th edition). World Health Organization Classification of Tumours. Lyon, France: International Agency for Research on Cancer (IARC); 2017.

5. Rosolen A, Perkins SL, Pinkerton CR, et al. Revised International Pediatric Non-Hodgkin Lymphoma Staging System. J Clin Oncol 2015;33(18):2112-8. DOI: 10.1200/JCO.2014.59.7203.

6. Khanam T, Sandmann S, Seggewiss J, et al. Integrative genomic analysis of pediatric T- cell lymphoblastic lymphoma reveals candidates of clinical significance. Blood 2020. DOI: 10.1182/blood.2020005381.

7.   Bontoux C, Simonin M, Garnier N, et al. Oncogenetic landscape of T-cell

lymphoblastic lymphomas compared to T-cell acute lymphoblastic leukemia. Mod

Pathol 2022;35(9):1227-1235. DOI: 10.1038/s41379-022-01085-9.

8.   Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health

Organization Classification of Haematolymphoid Tumours: Lymphoid Neoplasms.

Leukemia 2022;36(7):1720-1748. DOI: 10.1038/s41375-022-01620-2.

9.   Zijtregtop EAM, Diez C, Zwaan CM, Veening MA, Beishuizen A, Meyer-Wentrup

FAG. Thymus and activation-regulated chemokine (TARC) as treatment response

marker for paediatric Hodgkin lymphoma: A pilot study. Br J Haematol

2023;200(1):70-78. DOI: 10.1111/bjh.18473.

10.   Jundt F, Anagnostopoulos I, Forster R, Mathas S, Stein H, Dorken B. Activated

Notch1 signaling promotes tumor cell proliferation and survival in Hodgkin and

anaplastic large cell lymphoma. Blood 2002;99(9):3398-403. DOI:

10.1182/blood.v99.9.3398.

11.   Bonn BR, Rohde M, Zimmermann M, et al. Incidence and prognostic relevance of

genetic variations in T-cell lymphoblastic lymphoma in childhood and adolescence.

Blood 2013;121(16):3153-60. DOI: 10.1182/blood-2012-12-474148.

12.   Callens C, Baleydier F, Lengline E, et al. Clinical impact of NOTCH1 and/or FBXW7

mutations, FLASH deletion, and TCR status in pediatric T-cell lymphoblastic

lymphoma. J Clin Oncol 2012;30(16):1966-73. DOI: 10.1200/JCO.2011.39.7661.

13.   Hehir-Kwa JY, Koudijs MJ, Verwiel ETP, et al. Improved Gene Fusion Detection in

Childhood Cancer Diagnostics Using RNA Sequencing. JCO Precis Oncol

2022;6:e2000504. DOI: 10.1200/PO.20.00504.

14.   Haas BJ, Dobin A, Stransky N, et al. STAR-Fusion: Fast and Accurate Fusion

Transcript Detection from RNA-Seq. BioRxiv 2017.

15.   Bene MC, Castoldi G, Knapp W, et al. Proposals for the immunological classification

355   of acute leukemias. European Group for the Immunological Characterization of

356   Leukemias (EGIL). Leukemia 1995;9(10):1783-6.

357   (https://www.ncbi.nlm.nih.gov/pubmed/7564526).

358   16.   Palomero T, Barnes KC, Real PJ, et al. CUTLL1, a novel human T-cell lymphoma cell

359   line with t(7;9) rearrangement, aberrant NOTCH1 activation and high sensitivity to

360   gamma-secretase inhibitors. Leukemia 2006;20(7):1279-87. DOI:

361   10.1038/sj.leu.2404258.

362   17.   Steimle T, Dourthe ME, Alcantara M, et al. Clinico-biological features of T-cell acute

363   lymphoblastic leukemia with fusion proteins. Blood Cancer J 2022;12(1):14. DOI:

364   10.1038/s41408-022-00613-9.

365   18.   Brady SW, Roberts KG, Gu Z, et al. The genomic landscape of pediatric acute

366   lymphoblastic leukemia. Nat Genet 2022;54(9):1376-1389. DOI: 10.1038/s41588-

367   022-01159-z.

368   19.   Kopan R, Ilagan MX. The canonical Notch signaling pathway: unfolding the

369   activation mechanism. Cell 2009;137(2):216-33. DOI: 10.1016/j.cell.2009.03.045.

370   20.   Ashworth TD, Pear WS, Chiang MY, et al. Deletion-based mechanisms of Notch1

371   activation in T-ALL: key roles for RAG recombinase and a conserved internal

372   translational start site in Notch1. Blood 2010;116(25):5455-64. DOI: 10.1182/blood-

373   2010-05-286328.

374   21.   Patel JL, Smith LM, Anderson J, et al. The immunophenotype of T-lymphoblastic

375   lymphoma in children and adolescents: a Children's Oncology Group report. Br J

376   Haematol 2012;159(4):454-61. DOI: 10.1111/bjh.12042.

377   22.   Tasian SK, Loh ML, Hunger SP. Philadelphia chromosome-like acute lymphoblastic

378   leukemia. Blood 2017;130(19):2064-2072. DOI: 10.1182/blood-2017-06-743252.

23.   Ellisen LW, Bird J, West DC, et al. TAN-1, the human homolog of the Drosophila

notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms.

Cell 1991;66(4):649-61. DOI: 10.1016/0092-8674(91)90111-b.

24.   Speleman F, Cauwelier B, Dastugue N, et al. A new recurrent inversion,

inv(7)(p15q34), leads to transcriptional activation of HOXA10 and HOXA11 in a

subset of T-cell acute lymphoblastic leukemias. Leukemia 2005;19(3):358-66. DOI:

10.1038/sj.leu.2403657.

25.   Salmeron-Villalobos J, Ramis-Zaldivar JE, Balague O, et al. Diverse mutations and

structural variations contribute to Notch signaling deregulation in paediatric T-cell

lymphoblastic lymphoma. Pediatr Blood Cancer 2022;69(11):e29926. DOI:

10.1002/pbc.29926.

26.   Yamamoto K, Nakamachi Y, Yakushijin K, et al. A novel TRB@/NOTCH1 fusion

gene in T-cell lymphoblastic lymphoma with t(7;9)(q34;q34). Eur J Haematol

2013;90(1):68-75. DOI: 10.1111/ejh.12019.

27.   Liu Y, Easton J, Shao Y, et al. The genomic landscape of pediatric and young adult T-

lineage acute lymphoblastic leukemia. Nat Genet 2017;49(8):1211-1218. DOI:

10.1038/ng.3909.

28.   Mullighan C, Pölönen P, Giacomo DD, et al. [preprint] The genomic basis of

childhood T-lineage acute lymphoblastic leukemia. Research Square.

29.   Zijtregtop EAM, van der Strate I, Beishuizen A, et al. Biology and Clinical

Applicability of Plasma Thymus and Activation-Regulated Chemokine (TARC) in

Classical Hodgkin Lymphoma. Cancers (Basel) 2021;13(4). DOI:

10.3390/cancers13040884.

30.   Skinnider BF, Mak TW. The role of cytokines in classical Hodgkin lymphoma. Blood

2002;99(12):4283-97. DOI: 10.1182/blood-2002-01-0099.

Figure 1: NOTCH1-rearrangements in T-LBL. A) Schematic representation of three different *NOTCH1* fusions with different fusion partners. The in-frame *IKZF2::NOTCH1* fusion generates a chimeric protein in which the N-terminal DNA binding domain of IKZF2 is fused to the C-terminal intracellular domains of NOTCH1. Fusions transcripts with *miR142HG* and *TRBJ* use an alternative translation start site in exon 28 (Met1727). B) Western blot analysis using Val1744 antibody (Cell Signaling Inc.) shows that *miR142HG::NOTCH1* and the *TRBJ::NOTCH1* fusions lead to a larger NICD protein, likely representing uncleaved NICD with translation initiation at Met1727. Simultaneous Beta-actin staining was performed for loading comparisons. C) The 100 most significantly up- and

downregulated genes (from total 1,621 genes) in *NOTCH1*-rearranged compared to

NOTCH1-WT samples, revealed that *NOTCH1*-rearranged samples formed a separate cluster.

The two *NOTCH1*-rearranged cases excluded in the differential expression analysis because

of quality issues (TLBL049, TLBL058) cluster with the other four *NOTCH1*-rearranged

cases. The relapse sample of TLBL042 was used because of better quality. Range of 0-10

showing the log2 transcript per millions of normalized expression values. Significance was

determined using false discovery rate (FDR)-adjusted p-values.

Figure 2: TARC in *NOTCH1*-rearranged patients. A) TARC levels in pg/ml per patient,

showing highly elevated TARC in blood of *NOTCH1*-rearranged T-LBL patients but none of

the other patients. 10000 pg/ml is the maximum measurable TARC level with used assay.

Orange line in figure A, B and C represents maximum normal TARC level (1300 pg/ml)

Patients that had a relapse are indicated with an asterisk. Patient TLBL049 developed a

therapy-related AML (double asterisk). B) For three *NOTCH1*-rearranged patients, blood

TARC levels could be determined for a time point of remission after diagnosis, revealing

normalized TARC levels in all three cases. C) For three *NOTCH1*-rearranged patients blood

TARC levels were determined at time point of relapse and remission after relapse (second

remission), revealing increased levels in two relapses that again normalized in second

remission. D) Staining for TARC using anti-CCL17 antibody for four *NOTCH1*-rearranged

patients showing that T-LBL cells do not express TARC.