Characteristics and completeness of reporting of systematic reviews of prevalence studies

in adult populations: a meta-epidemiological study.

Diana Buitrago-Garcia [a, b], William Gildardo Robles-Rodriguez [c], Javier Eslava-Smalbach [d],

Georgia Salanti [a] *, Nicola Low [a] * [†]

[a.] Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse 43, 3012

Bern, Switzerland.

[b.] Graduate School of Health Sciences, University of Bern, Mittelstrasse 43, 3012 Bern,

Switzerland.

[c.] Department of Epidemiology, Fundación Universitaria de Ciencias de la Salud-FUCS, Cr

49 No 95 – 79, Bogotá, Colombia.

[d.] School of Medicine, Universidad Nacional de Colombia, Cr 30 No 45-03, Bogotá,

Colombia.

* GS and NL are joint senior authors

[†] Corresponding author: Nicola Low. Email: nicola.low@unibe.ch. Tel: +41 31 684 3092.

Postal address: Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse

43, Bern, CH-3012, Switzerland.

Abstract

Objective: The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement, first published in 2009, has been widely endorsed and compliance is high in systematic reviews of intervention studies. Systematic reviews of prevalence studies are increasing in frequency, but their characteristics and reporting quality have not been examined in large studies. Our objectives were to describe the characteristics of systematic reviews of prevalence studies in adults, evaluate the completeness of reporting and explore study-level characteristics associated with the completeness of reporting.

Study design and setting: We did a meta-epidemiological study. We searched 5 databases from January 2010 to December 2020 to identify systematic reviews of prevalence studies in adult populations. We used the PRISMA 2009 checklist to assess completeness of reporting and recorded additional characteristics. We conducted a descriptive analysis of review characteristics and linear regression to assess the relationship between compliance with PRISMA and publication characteristics.

Results: We included 1172 systematic reviews of prevalence studies. The number of reviews increased from 25 in 2010 to 273 in 2020. The median PRISMA score for systematic reviews without meta-analysis was 17.5 out of a maximum of 23 and, for systematic reviews with meta-analysis, 22 out of a maximum of 25. Completeness of reporting, particularly for key items in the methods section was suboptimal. Systematic reviews that included a meta-analysis or reported using a reporting or conduct guideline were the factors most strongly associated with increased compliance with PRISMA 2009.

Conclusion: Reporting of systematic reviews of prevalence was adequate for many PRISMA items. Nonetheless, this study highlights aspects for which special attention is needed. Development of a specific tool to assess the risk of bias in prevalence studies and an extension

to the PRISMA statement could improve the conduct and reporting of systematic reviews of

prevalence studies.

Keywords: Systematic reviews; prevalence, adults; reporting; meta-epidemiology; risk of bias

Running title: Systematic reviews of prevalence studies: meta-epidemiological study

Word count: main text, 3060

Abbreviations

IQR   Interquartile ranges

MOOSE   Reporting Guidelines for Meta-analyses of Observational Studies

PRISMA   Preferred Reporting Items for Systematic Reviews and Meta-Analyses

SR   Systematic review

What is new?

54 • Systematic reviews of prevalence increased from 25 in 2010 to 2020, to 273 in 2020.

55 • Reporting of systematic reviews of prevalence has improved but is still suboptimal.

56 • Reporting was better in reviews with a meta-analysis and which followed a guideline.

57 • Journals should encourage adherence to PRISMA for systematic reviews of prevalence.

58 • A risk of bias tool and a PRISMA extension for prevalence reviews should be developed.

1   Introduction

Prevalence studies quantify the occurrence of a disease and can be used to contribute to estimation of the burden of disease and as a measure to evaluate healthcare interventions [1, 2]. Systematic reviews (SRs) of prevalence studies allow the synthesis of evidence about prevalence, which also informs burden of disease estimates and provides a resource for policymakers to help set priorities [1]. The volume of SRs of prevalence studies is increasing, but the methods used to conduct them have been reported to be variable and suboptimal [3, 4].

The usefulness of any systematic review depends on the completeness of reporting and the information provided in the included publications. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement was first published in 2009 to help with transparent and complete reporting of systematic reviews that assess the benefits or harms of interventions [5]. Since then, extensions to the PRISMA statement have covered other study designs, including diagnostic test accuracy [6], protocols [7] and network meta-analysis [8]. An update to the statement, PRISMA 2020, included items that are also applicable to systematic reviews of aetiology, prognosis, and prevalence studies, whilst still being designed primarily for reviews of studies of health interventions [9]. Some studies have shown that the PRISMA statement and extensions have enhanced the reporting of systematic reviews [10-12], although others show that improvement is still needed [11, 13]. The completeness of reporting of SRs of prevalence is, to our knowledge, unknown. The objectives of this study were to describe the characteristics of SRs of prevalence studies in adults, the completeness of reporting, and to explore study level characteristics associated with the quality and completeness of reporting.

2   Methods

We conducted a meta-epidemiological study of SRs of prevalence studies. The protocol was registered in the PROSPERO register (CRD42020151625). We report our findings according to

the Guidelines for Reporting Meta-Epidemiological Research(Appendix A) [14]. Differences

between the methods in the protocol and the study reported here are in Appendix B.

2.1   Search methods

We searched MEDLINE-Ovid, Embase-Ovid, CINAHL and LILACS from January 2010 to

December 2020 without language restrictions. We also searched grey literature in

opengrey.com [15] (Appendix C).

2.2   Eligibility criteria

We included SRs of studies conducted in adults (individuals aged ≥18 years) in any setting that

assessed the prevalence of a disease, symptom, risk factor or behaviour as their primary aim.

We excluded SRs of diagnostic test accuracy and of incidence studies, unless prevalence

estimates were also presented separately. We also excluded overviews of SRs, studies that

conducted a meta-analysis or pooled prevalence data without conducting a SR and conference

abstracts, since it was not possible fully assess the completeness of reporting.

2.3   Study selection and data extraction

One author screened titles, abstracts and reviewed full-text reports for potential eligibility

(D.B.G.) and a second author (W.R.) verified 20% of studies, using the online tool Rayyan [16]

(96% agreement for title and abstract screening). One reviewer extracted data using a pre-

piloted form (D.B.G.) and a second author verified the extraction in 20% of included studies

(W.R.). We resolved disagreements through discussion. We extracted the following information:

publication year, journal and Journal Impact Factor (Web of Science Journal Citation Reports

2022 or, if not available, the impact factor reported on the journal's website) country of the first

author (using the first affiliation, if more than one was listed), number of authors, medical

speciality and targeted condition, population, primary objective, design of the included studies,

geographic coverage, type of numerical data extracted from the included studies, number of

studies included, tool reported to have been used to assess the risk of bias or quality in included

studies, statistical methods, and approaches used to assess heterogeneity. In addition, if

authors reported the use of guidelines or recommendations for SRs, such as the PRISMA

checklist [5], the Reporting Guidelines for Meta-analyses of Observational Studies (MOOSE)

[17], or for conduct, such as the Cochrane Handbook for Systematic Reviews of Interventions

[18].

2.4   Assessment of the completeness of reporting

The completeness of reporting of each SR was assessed using the PRISMA checklist published

in 2009 [5], which was appropriate for the publication dates of the included studies. The

PRISMA 2009 checklist has 27 items. We decided to exclude two items related to reporting

biases across studies, e.g., publication bias and other biases due to missing studies or missing

results within studies (item 15 in the methods and item 22 in the results). The statistical methods

used to assess these biases were developed for comparative studies, but their relevance and

interpretation in evidence from prevalence studies is less clear and needs to be further

investigated [19]. We assigned each of the 25 items one point if the item was adequately

reported or no points if the item was not reported. For some items, we awarded half a point if

the information was partially reported (Appendix D). The maximum score for SRs with a meta-

analysis was 25 points, and 23 points without a meta-analysis (items 14 and 21 were not

applicable).

2.5   Data analysis

We summarised the study characteristics (discipline, number of studies etc.) using proportions

or medians with interquartile ranges (IQR). The completeness of reporting for each review was

calculated as a) the achieved PRISMA reporting score, and b) the scaled reporting score, which

was the achieved score divided by the maximum possible value; the scaled reporting score

takes values between 0 and 100%. Completeness of reporting was summarised as, a) the

median (IQR) PRISMA scores, and b) the proportion of SRs that completely reported, partially

reported, or did not report, each PRISMA item. Suboptimal reporting for an item was defined as

less than 70%, based on a previous study [10].

We conducted univariable and multivariable linear regression analyses to assess the

relationship between the scaled reporting score and the year of publication, the Journal Impact

Factor, the journal's publishing model (open access or not), the number of co-authors, the

number of studies included in the review, the use of a guideline to report or conduct the

systematic review, the medical specialty and the type of review (SRs with or without a meta-

analysis). All analyses were performed using R version 4.3.1 [20].

3   Results

144   We screened 9580 references and included 1172 systematic reviews, which fulfilled our inclusion

145   criteria. The main reason for exclusion of potentially eligible reviews was because the primary aim

146   of the review was not to assess prevalence. (Figure 1, Appendix E).

163   *Figure 1. PRISMA flowchart of the selection of the inclusion of systematic reviews of prevalence*

164   *studies.*

 3.1   Characteristics of the reviews

The number of SRs of prevalence increased from 25 in 2010 to 273 in 2020 (Figure 2). There

were 387 SRs without meta-analysis and 785 SRs with a meta-analysis. The median number of

studies included in all SRs was 25 (IQR 14, 46). The SRs were published across 645 different

journals, with PLOS ONE being the most frequent (n=46), followed by BMC Public Health

(n=21), and BMC Infectious Diseases (n=20) (Table 1, Table S1). First authors were affiliated

with institutions in 65 countries, amongst whom half were in five countries: the United Kingdom

(n=155), the United States (n=120), Iran (n=107), Brazil (n=105), and China (n=98) (Figure S1,

Table S2). Most SRs evaluated the prevalence of a medical condition or risk factor (n=1,036,

88%) and extracted worldwide prevalence data (765, 65%). About half of the SRs (n=567, 48%)

were conducted to assess the prevalence of psychiatric conditions, infectious diseases,

cardiology, and neurology (Figure S2, Table S3). Prevalence data were extracted from diverse

populations, most commonly from general adult populations (n=436, 37%), adults with a specific

condition (n=382, 33%), older populations (n=100, 9%), women (n=89, 8%) or workers (n=34,

3%) (Table 1, Table S4).

Table 1. Characteristics of systematic reviews of prevalence studies in adults, published
between 2010-2020

Variable	Systematic reviews without meta-analysis (n=387)	Systematic reviews with meta-analysis (n=785)	Total (n=1172)
Journal Impact Factor			
Median (IQR)	3.6 (2.2, 5.4)	4.1 (2.8, 6.7)	3.9 (2.6, 6.5)
Number of co-authors			
Median (IQR)	4 (3, 6)	6 (4, 7)	5 (4, 7)
Number of studies included in the review			
Median (IQR)	22 (13, 39)	24 (14, 47)	25 (14, 46)
Top five journals, n (%)			
PLOS ONE	11 (2.8)	35 (4.5)	46 (3.9)
BMC Public Health	5 (1.3)	16 (2.0)	21 (1.8)
BMC Infectious Diseases	4 (1.0)	16 (2.0)	20 (1.7)
Journal of Affective Disorders	4 (1.0)	13 (1.7)	17 (1.5)

BMJ Open	1 (0.3)	10 (1.3)	11 (0.9)
Top five country affiliations of first author, n (%)			
United Kingdom	55 (14.0)	100 (12.7)	155 (13.2)
United States	38 (9.8)	82 (10.4)	120 (10.1)
Iran	30 (7.8)	77 (9.8)	107 (9.1)
Brazil	47 (12.0)	58 (7.4)	105 (9.0)
China	26 (6.7)	72 (9.2)	98 (8.4)
Top five study populations, n (%)			
Adults in general	146 (37.7)	290 (36.9)	436 (37.2)
Adults with a specific condition or characteristic	101 (26.1)	281 (35.8)	382 (32.6)
Older adults	47 (12.1)	53 (6.8)	100 (8.5)
Women (including pregnant women)	29 (7.5)	60 (7.6)	89 (7.6)
Workers	16 (4.1)	18 (2.3)	34 (2.9)
Geographic scope, n (%)			
Worldwide	254 (65.6)	511 (65.1)	765 (65.3)
Region	60 (15.5	87 (11.1)	147 (12.5)
Country	73 (18.9)	187 (23.8)	260 (22.2)
Top five specialties, n (%)			
Infectious diseases	38 (9.8)	168 (21.4)	206 (17.6
Psychiatry	68 (17.6)	118 (15.0)	186 (15.9)
Cardiology	26 (6.7)	68 (8.7)	94 (8.0)
Neurology	28 (7.2)	51 (6.5)	79 (6.7)
Endocrinology	23 (5.9)	52 (6.6)	75 (6.4)
Aim of the review, n (%)			
Estimate prevalence	353 (91.2)	683 (87.0)	1,036 (88.4)
Estimate prevalence, compare prevalence estimates and/or evaluate associations	34 (8.8)	102 (13.0)	136 (11.6)
Data extracted, n (%)			
Prevalence estimate	273 (70.5)	318 (40.5)	591 (50.4)
Numerator and denominator	89 (23.0)	339 (43.2)	428 (36.5)
Both	25 (6.5)	128 (16.3)	153 (13.1)
Authors reported using the PRISMA statement, n (%)			
Used	163 (42.1)	472 (60.1)	635 (54.2)
Not used	224 (57.9)	313 (39.9)	537 (45.8)
Authors reported using of MOOSE guidelines, n (%)			
Used	22 (5.7)	125 (15.9)	147 (12.5)
Not used	365 (94.3)	660 (84.1)	1,025 (87.5)
Authors reported using other guidelines (e.g., Cochrane Handbook), n (%)			
Used	17 (4.4)	48 (6.1)	65 (5.5)
Not used	370 (95.6)	737 (93.9)	1,107 (94.5)

In total, 62% (727/1172) of SRs described using one or more reporting or conducting guidelines.

The PRISMA checklist was cited most often, followed by the MOOSE guideline [17]. Most

authors included studies that answered their review questions without restrictions on study

design (n=623, 53%) (Table 1, Table S5-S6). Among SRs with meta-analysis, most (n=703,

90%) used a random-effects model (Table S6). Only 21% (n=163) of SRs reported a method

used for statistical transformation of prevalence values, such as the Freeman-Tukey double

arcsine, log, or logit functions. Heterogeneity between studies was assessed in most SRs,

mainly using the I^2 statistic (n=720; 92%).

3.2   Completeness of reporting

3.2.1   PRISMA scores

The median PRISMA score for SRs without meta-analyisis was 17.5 (IQR 15.0,19.0) out of a

possible 23, and for SRs with meta-analysis, it was 22.0 (IQR 20.5, 23.5) out of a possible 25

(Table S7). Figure 2 shows an increasing reporting score over the years. The median scaled

reporting score for all included SRs was 84. 8% (IQR 76.0, 92.0).

Figure 2. PRISMA score and number of publications (n) by year of publication and type of

systematic review.

3.2.2 Reporting of selected PRISMA items

Over 80% of the included SRs complied with more than 70% of the PRISMA checklist items.

Two of 387 SRs without meta-analysis (0.5%) and 66 of785 SRs with meta-analysis (8%) were

entirely compliant with PRISMA 2009. Completeness of reporting was below 70% for reporting

the existence of a protocol for the review, the search strategy, additional analyses and

assessment of the risk of bias (Figure 3). Findings for these items are reported below.

Protocol: Only 296/1172 (25%) of all SRs reported the existence of a protocol, which could be

accessed, while 62/1172 (6%) mentioned a protocol without information on access details. For

SRs without meta-analysis, the number with any protocol was 1/15 in 2010 and 9/47 in 2020; for

SRs with meta-analysis, 0/10 reviews in 2010 and 91/226 in 2020 had a protocol (Table S8).

Search strategy: In 607/1172 (52%) SRs, authors adequately reported information about the

search strategy for at least one database. In 498/1172 (42%) reviews, authors only reported the

keywords used and 67/1172 (6%) did not provide any information.

Assessment of risk of bias: In the *methods section*, 798/1172 (68%) review authors reported the

use of any tool to assess the risk of bias in included studies (Table S9). For 12 (1%), authors

mentioned using a tool but did not report the items assessed or the tool. The most frequently

reported tool was the Newcastle-Ottawa Scale (153/1172, 13.1%), which is for assessment of

the quality of non-randomised studies [21]. In 213/1172 (18%) reviews, authors reported the use

of a tool designed explicitly for assessment of quality or risk of bias in prevalence studies. The

most frequent was the tool developed by the JBI (formerly Joanna Briggs Institute) [22]. The risk

of bias in the included studies was adequately reported in the *results section* in 675/1172 (58%)

reviews. Completeness of reporting of the study-level risk of bias assessment in the results was

lower than in the methods section (68%). In 56/1172 reviews (5%) authors reported assessing

the risk of bias but there was no description of this in the results. In 25/1172 reviews (2%),

authors reported using the assessment of the quality of the studies to exclude studies from the

review.

Additional analyses: 526/785 (67%) SRs with meta-analysis reported in the *methods section*

additional analyses such as sensitivity, subgroup analysis, or meta-regression. We observed an

increase over the years, from 8/10 (80%) reviews in 2010 to 161/226 (71%) in 2020. Additional

analysis results were presented adequately in the *results section* in 576/785 SRs with meta-

analysis (73%).

Reporting of funding sources: 764/1172 (65%) SRs reported their source of funding. The

reporting of this item improved from 12/25 (48%) in 2010 to 202/273 (74%) in 2020.

Figure 3. Percentage of adequate reporting of PRISMA items in 2009 in 387 systematic reviews

without meta-analysis (SR-M) and 785 systematic reviews with meta-analysis (SR+M).

[a] (M), item in methods section; (R), item in results section.

3.3   Factors associated with the completeness of reporting

Inclusion of a meta-analysis in the SRs and citing the use of a reporting or methodological

guideline were the factors most strongly associated with a higher scaled reporting score (Table

2). Publishing in an open access journal, the year of publication and the Journal Impact Factor

were also positively associated with higher scaled reporting scores.

Table 2. Univariable and multivariable linear regression between characteristics of the

published systematic reviews of prevalence studies and scaled reporting score according to the

PRISMA guidelines

Variable	Univariable analysis			Multivariable analysis[*]		
	Coefficient	(95% CI)	P value	Coefficient	(95% CI)	P value
Year						
	1.20	(0.96,1.40)	<0.001	0.29	(0.06, 0.52)	<0.001
Journal Impact Factor						
	0.05	(0.01, 0.10)	0.019	0.06	(0.02, 0.10)	0.003
Open access journal						
No	Reference		0.063	Reference		0.030
Yes	1.40	(-0.08, 2.90)		1.40	(0.13, 2.70)	
Number of authors						
	0.67	(0.43, 0.91)	<0.001	0.16	(-0.05, 0.37)	0.13
Number of studies included in the review						
	2.60	(1.00, 4.20)	0.001	0.70	(-0.67, 2.10)	0.30
Report the use of guidelines to report or conduct systematic reviews[†]						
No	Reference		<0.001	Reference		<0.001
Yes	8.90	(7.50, 10.25)		5.80	(4.50, 7.04)	
Systematic review includes a meta-analysis						
No	Reference		<0.001	Reference		<0.001
Yes	13.47	(12.17, 14.76)		11.69	(10.36, 13.01)	
Medical field						
Other	Reference		>0.9	Reference		0.20
Psychiatry	1.20	(-0.95, 3.30)		0.85	(-0.90, 2.6)	
Infectious diseases	0.45	(-1.60, 2.50)		-2.10	(-3.9, -0.42)	
Neurology	0.42	(-2.60, 3.40)		0.27	(-2.20, 2.70)	
Cardiology	0.73	(-2.00, 3.50)		-0.40	(-2.70, 1.90)	
Endocrinology	-0.30	(-3.40, 2.70)		-1.20	(-3.70,1.30)	
Surgery	-1.20	(-4.30, 1.90)		-0.27	(-2.80, 2.30)	
Lifestyle characteristics	-0.45	(-4.40, 3.50)		0.84	(-2.40, 4.10)	

246 *Abbreviations:* CI, confidence intervals

247 [*] Model includes all variables reported in the table.

248 [†]Guidelines: PRISMA 2009 statement, Reporting Guidelines for Meta-analyses of Observational Studies
249 (MOOSE), Cochrane Handbook.

4 Discussion

This meta-epidemiological study found an 11-fold increase in the number of SRs of prevalence

in adult populations published from 25 in 2010 to 273 in 2020. The median PRISMA 2009 score

for SRs without meta-analysis was 17.5 (IQR 15.0, 19.0), and for SRs with meta-analysis was

22.0, (IQR 20.5, 23.5). The items with the lowest compliance (<70%) were the availability of a

protocol, search methods, assessment of the risk of bias in methods and results, additional

analyses, and sources of funding. In multivariable analysis, SRs that included a meta-analysis,

reported using a reporting or conduct guideline, and publications in more recent years, in an

open access journal, or in journals with a higher Journal Impact Factor were on average more

compliant with the PRISMA 2009 checklist.

4.1 Strengths and limitations

Strengths of this study include the detailed assessment of characteristics of SRs of prevalence,

including 11 years of publications in 1172 SRs. In addition to recording whether PRISMA

checklist items were reported, we extracted additional information for several items and

conducted a multivariable regression analysis, which allowed more detailed interpretation of the

findings than simple descriptive statistics. Our study also has limitations. First, we did not

include SRs of prevalence published before the launch of the PRISMA statement in 2009 [5],

which does not allow us to assess if there was improvement after the checklist was published.

Second, we did not extend our search after 2020, so the end date of the search means that our

findings correspond to the items and scope of the PRISMA 2009 statement [5]. The COVID-19

pandemic interrupted work on this study from 2021 and when we returned to it, the PRISMA

2020 checklist had been published [9]. Our study therefore provides an initial assessment of the

completeness of reporting of SRs of prevalence and a future assessment will help to understand

whether the extended scope of PRISMA 2020 is associated with further changes in the

completeness of reporting. Third, we did not use PRISMA extensions, such as the PRISMA

checklist for abstracts [23] or the extension for reporting literature searches in SRs [24], which

might change the results of the items assessed with PRISMA 2009. Fourth, we limited the

scope of topics to reviews conducted in adult populations, but we believe that reviews

conducted in children would yield similar methodological findings. Fifth, we acknowledge that

the PRISMA 2009 checklist was not designed to give a score. This method has been used

previously [12] and, for our objectives, provided a pragmatic, if simplified, way to highlight

aspects of reporting of SRs that could be improved.

4.2   Interpretation and comparison with other studies

Incomplete reporting of systematic reviews of prevalence should be seen in the context of

published guidance for the conduct and reporting of SRs, most of which has been developed for

randomised or non-randomised intervention studies. Whilst reported use of a guideline for

reporting or conduct of SRs was associated with more complete reporting, the content of some

items may indicate a lack of specific methodological guidance for prevalence studies. In

particular, 30% of authors did not report the use of a tool for assessment of the risk of bias in

individual studies and, amongst those that did, more than 30 different tools were used. In a

systematic search, we identified 10 tools for assessing the risk of bias in prevalence studies

[25], but only 284 (24%) of reviews in our study used one of these tools. Most of the tools listed

were not designed for use with prevalence studies, such as the STROBE checklist for reporting

of cross-sectional studies, which does not allow explicit assessment of risk of bias [26].

Completeness of reporting of SRs was associated with inclusion of a meta-analysis, publication

in open access journals and publication in journals with a higher impact factor. These

characteristics could be related to the level of experience and recognition of the methodological

requirements of reporting of a systematic review team or with the expectations and

requirements of journals.

We found two smaller studies, which assessed the characteristics of SRs of prevalence but did

not use the PRISMA checklist to quantify completeness of reporting. Borges Migliavaca et al. [3]

evaluated 235 SRs of prevalence published in 2017 and 2018 and found substantial differences

in terms of conduct, reporting, risk of bias assessment and data synthesis. Whilst we decided

not to assess the reporting of publication bias because of doubts about its relevance to

prevalence studies, Borges Migliavaca et al. extracted this information. They found that 48/235

SRs examined publication bias either graphically or using a statistical test [3]. The authors also

found that some reviews used the GRADE approach, despite the absence of GRADE guidance

on assessing the quality of the body of evidence in a SR of prevalence. Hoffmann et al. [4]

reported on 215 SRs of prevalence and incidence, identified from a random sample of

publications up to 2018. The authors did not report on their findings for SRs of prevalence and

incidence separately, but concluded that heterogeneity in characteristics, reporting, and

methodology of these SRs might be due to the absence of specific guidance.

Reporting for some items in SRs of prevalence was consistent with other study designs. Page et

al. [10] summarised meta-research studies of adherence to the PRISMA 2009 statement

published up to mid-2017. They also found that items such as protocol registration and

assessment of the risk of bias of individual studies were likely to be incomplete. Veroniki et al.

[12] assessed the reporting in 1144 SRs with network meta-analysis and also found that the

items least likely to be adequately reported were publication of a protocol (25%), and of a full

search strategy (48%). Wasiak et al. [11] assessed 50 SRs in burn care management and

concluded that methodology was the section most in need of improvement. They also found an

improvement in the PRISMA score when the systematic review incorporated a meta-analysis.

5   Conclusions and recommendations

Reporting of SRs of prevalence was adequate for many PRISMA items. The completeness of

reporting has also improved but there is room for improvement. There are items that authors

who conduct any type of SR can improve without further guidance, such as the publication of a

protocol. To improve the consistency and utility of SRs of prevalence more specific guidance

about reporting of certain methodological features is required. Development of a specific tool to

assess the risk of bias in prevalence studies and an extension to the PRISMA statement could

improve the conduct and reporting SRs of prevalence studies.

Declarations of interest

None

Declarations of interest of generative AI in scientific writing

The authors declare that no AI was used in the scientific writing of the manuscript.

Author contributions

Study design: DBG, JES, NL, GS.

Data collection: DBG, WR.

Methodology DBG, JES, NL, GS

Writing original draft: DBG

Approval

All authors approved the final version of the manuscript.

Funding

This work received support from the Swiss government excellence scholarship (grant number
2019.0774), the SSPH+ Global PhD Fellowship Programme in Public Health Sciences of the
Swiss School of Public Health.

Georgia Salanti acknowledges the Swiss National Science Foundation for funding of her work
through the NRP 78 initiative (Project 198418)

Availability of data and materials

Raw data and bibliographic details of the included studies are published on Open Science
Framework (https://osf.io/8792v)

6 REFERENCES

[1] Harder T. Some notes on critical appraisal of prevalence studies: Comment on: "The development of a critical appraisal tool for use in systematic reviews addressing questions of prevalence". Int J Health Policy Manag. 2014;3:289-90 https://doi.org/10.15171%2Fijhpm.2014.99.

[2] Denton FT, Spencer BG. Chronic Health Conditions: Changing Prevalence in an Aging Population and Some Implications for the Delivery of Health Care Services. Can J Aging. 2010;29:11-21 https://doi.org/10.1017/s0714980809990390.

[3] Borges Migliavaca C, Stein C, Colpani V, Barker TH, Munn Z, Falavigna M. How are systematic reviews of prevalence conducted? A methodological study. BMC Med Res Methodol. 2020;20:96 https://doi.org/10.1186/s12874-020-00975-3.

[4] Hoffmann F, Eggers D, Pieper D, Zeeb H, Allers K. An observational study found large methodological heterogeneity in systematic reviews addressing prevalence and cumulative incidence. J Clin Epidemiol. 2020;119:92-9 https://doi.org/10.1016/j.jclinepi.2019.12.003.

[5] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. BMJ. 2009;339:b2700 https://doi.org/10.1136/bmj.b2700.

[6] McInnes MDF, Moher D, Thombs BD, McGrath TA, Bossuyt PM, Group atP-D. Preferred Reporting Items for a Systematic Review and Meta-analysis of Diagnostic Test Accuracy Studies: The PRISMA-DTA Statement. JAMA. 2018;319:388-96 https://doi.org/10.1001/jama.2017.19163.

[7] Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;349:g7647 https://doi.org/10.1136/bmj.g7647.

[8] Hutton B, Salanti G, Caldwell DM, Chaimani A, Schmid CH, Cameron C, et al. The PRISMA Extension Statement for Reporting of Systematic Reviews Incorporating Network Meta-analyses of Health Care Interventions: Checklist and Explanations. Ann Intern Med. 2015;162:777-84 https://doi.org/10.7326/m14-2385.

[9] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The

PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ.

2021;372:n71 https://doi.org/10.1136/bmj.n71.

[10] Page MJ, Moher D. Evaluations of the uptake and impact of the Preferred Reporting Items

for Systematic reviews and Meta-Analyses (PRISMA) Statement and extensions: a scoping

review. Syst Rev. 2017;6:263 https://doi.org/10.1186/s13643-017-0663-8.

[11] Wasiak J, Tyack Z, Ware R, Goodwin N, Faggion CM, Jr. Poor methodological quality and

reporting standards of systematic reviews in burn care management. Int Wound J. 2017;14:754-

63 https://doi.org/10.1111%2Fiwj.12692.

[12] Veroniki AA, Tsokani S, Zevgiti S, Pagkalidou I, Kontouli KM, Ambarcioglu P, et al. Do

reporting guidelines have an impact? Empirical assessment of changes in reporting before and

after the PRISMA extension statement for network meta-analysis. Syst Rev. 2021;10:246

https://doi.org/10.1186/s13643-021-01780-9.

[13] Maticic K, Krnic Martinic M, Puljak L. Assessment of reporting quality of abstracts of

systematic reviews with meta-analysis using PRISMA-A and discordance in assessments

between raters without prior experience. BMC Med Res Methodol. 2019;19:32

https://doi.org/10.1186%2Fs12874-019-0675-2.

[14] Murad MH, Wang Z. Guidelines for reporting meta-epidemiological methodology research.

Evid Based Med. 2017;22:139-42 https://doi.org/10.1136/ebmed-2017-110713.

[15] OpenGrey. System for Information on Grey Literature in Europe Available:

http://www.opengrey.eu/ [Accessed 28/10/2021].

[16] Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for

systematic reviews. Syst Rev. 2016;5:210 https://doi.org/10.1186/s13643-016-0384-4.

[17] Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of

observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational

Studies in Epidemiology (MOOSE) group. JAMA. 2000;283:2008-12

https://doi.org/10.1001/jama.283.15.2008.

[18] Higgins JPT, Green S. Cochrane Handbook for Systematic Reviews of Interventions: Wiley;
2011.

[19] Page MJ, Sterne JAC, Higgins JPT, Egger M. Investigating and dealing with publication
bias and other reporting biases in meta-analyses of health research: A review. Res Synth
Methods. 2021;12:248-59 https://doi.org/10.1002/jrsm.1468.

[20] R Core Team. R: A language and environment for statistical computing. Vienna, Austria
2023. Available: https://www.R-project.org/

[21] Wells G, Shea B, O'Connell D, Peterson j, Welch V, Losos M, et al. The Newcastle–Ottawa
Scale (NOS) for Assessing the Quality of Non-Randomized Studies in Meta-Analysis. 2000.
Available: https://www.ohri.ca/programs/clinical_epidemiology/oxford.asp

[22] Joanna Briggs Institute. The Joanna Briggs Institute Critical Appraisal tools for use in JBI
Systematic Reviews Checklist for Prevalence Studies. Joanna Briggs Institute; 2017. Available:
https://jbi.global/sites/default/files/2019-05/JBI_Critical_Appraisal-
Checklist_for_Prevalence_Studies2017_0.pdf

[23] Beller EM, Glasziou PP, Altman DG, Hopewell S, Bastian H, Chalmers I, et al. PRISMA for
Abstracts: Reporting Systematic Reviews in Journal and Conference Abstracts. PLoS Med.
2013;10:e1001419 https://doi.org/10.1371%2Fjournal.pmed.1001419.

[24] Rethlefsen ML, Kirtley S, Waffenschmidt S, Ayala AP, Moher D, Page MJ, et al. PRISMA-S:
an extension to the PRISMA Statement for Reporting Literature Searches in Systematic
Reviews. Syst Rev. 2021;10:39 https://doi.org/10.1186/s13643-020-01542-z.

[25] Tonia T, Buitrago-Garcia D, Peter NL, Mesa-Vieira C, Li T, Furukawa TA, et al. Tool to
assess risk of bias in studies estimating the prevalence of mental health disorders (RoB-
PrevMH). BMJ Ment Health. 2023;26 https://doi.org/10.1101/2023.02.01.23285335.

[26] von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP.
Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement:
guidelines for reporting observational studies. BMJ. 2007;335:806-8
https://doi.org/10.1136/bmj.39335.541782.ad.