Plasma proteomic evidence for increased Alzheimer's disease-related brain pathology after SARS-CoV-2 infection

1
2
3
4
5   Eugene P Duff*[1,2], Henrik Zetterberg[3,4,5,6,7,8], Amanda Heslegrave[5], Abbas
6   Dehghan[9,1,10], Paul Elliot[9,1,10,11,12,13], Naomi Allen[14,15], Heiko Runz[16], Rhiannon Laban[6], Elena
7   Veleva[6], Christopher D Whelan[17], Benjamin B Sun[16], Paul M Matthews [§1, 2,18]
8
[1]UK Dementia Research Institute Centre at Imperial College London, London,
UK, [2]Department of Brain Sciences, Faculty of Medicine, Imperial College London, London,
UK, [3]Department of Psychiatry and Neurochemistry, Institute of Neuroscience and
Physiology, the Sahlgrenska Academy, University of Gothenburg, Mölndal,
Sweden, [4]Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal,
Sweden, [5]Department of Neurodegenerative Disease, UCL Institute of Neurology, University
College London, London, UK, [6]UK Dementia Research Institute Centre at UCL, London,
UK, [7]Hong Kong Center for Neurodegenerative Diseases, Hong Kong, China, [8]Wisconsin
Alzheimer's Disease Research Center, University of Wisconsin School of Medicine and
Public Health, Madison, WI, USA, [9]Department of Epidemiology and Biostatistics, Faculty of
Medicine, Imperial College London, London, UK, [10]MRC Centre for Environment and Health,
Imperial College London, London, UK, [11]British Heart Foundation Centre of Research
Excellence, Imperial College London, London, [12]National Institute for Health Research
Biomedical Research Centre, Imperial College London, London, [13]Health Data Research UK
at Imperial College London, Imperial College London, London, [14]Nuffield Department of
Population Health, University of Oxford, Oxford, UK, [15]UK Biobank, Stockport,
UK, [16]Translational Sciences, Biogen, Cambridge, MA, USA, [17]Neuroscience Data Science,
Janssen Research & Development, Cambridge, MA, USA, [18] The Rosalind Franklin Institute,
Harwell Science and Innovation Campus, Didcot, UK
28
29 *Communicating author: Eugene Duff e.duff@imperial.ac.uk
30 [§] Senior author: Paul Matthews p.matthews@imperial.ac.uk
31
32

Abstract

Prior studies have suggested that systemic viral infections may increase risks of dementia. Whether this holds true for SARS-CoV-2 virus infections remains uncertain but is of great consequence for predicting future dementia rates. We examined this by comparing changes in plasma biomarkers in UK Biobank participants before and after serology confirmed SARS-CoV-2 infections. We discovered biomarker changes associated with increased AD risk within this population. SARS-CoV-2 infection was associated with reduced plasma Aβ42:Aβ40 concentration ratios, and in more vulnerable participants, lower plasma Aβ42 and higher plasma pTau-181. These biomarker changes, which have been associated with brain beta-amyloid accumulation in prodromal AD, were associated here with increased brain imaging signatures of AD, poorer cognitive scores, and worse assessments of overall health. Changes were greater in participants who had been hospitalised with COVID-19 or had previously reported hypertension. Our data provide evidence for the hypothesis that SARS-CoV-2 can be associated with accelerating brain pathology related to prodromal AD.

Introduction

Exposures to infectious diseases in early life and adulthood have been linked to increased risk for neurodegenerative and other systemic disease [1–4]. Viral encephalitis or meningitis, influenza, pneumonia and viral intestinal infections all may increase risks of Alzheimer's disease (AD) [1], Vaccinations to prevent these diseases may also have protective effects [5]. A variety of central and peripheral mechanisms could underly these effects, including tissue injury during the acute phase of infections or chronic secondary inflammatory processes [6–8]. However, uncertainty remains. The epidemiological observations are confounded by heterogeneity of infections, and mechanistic hypotheses have been limited by the lack of large scale, harmonised, prospective observations in human populations that allow common factors that predispose both to infection and to dementia to be discriminated from any potential causal relationships of infection to dementia.

The recent COVID-19 pandemic provided a common viral exposure to large populations over well-defined periods. SARS-CoV-2 initiates a systemic inflammatory response, which persists in many patients beyond the acute phase and increases susceptibility to several systemic diseases. Although the virus is not neurotrophic [9–11], impairment of cognition, and brain structural changes have been reported as sequelae of SARS-CoV-2 infections, even amongst people who did not require hospitalisation or experience long-COVID [10,12]. Initial evidence for raised dementia rates in vulnerable populations following serious COVID have been reported [13]. Preclinical studies have identified potential mechanisms for increased brain inflammatory pathology with SARS-CoV-2 that could account for these observations [7,14,15]. These reports suggested to us that the COVID pandemic could provide a "natural experiment" for testing whether infection-associated systemic infection can initiate or potentiate brain pathology associated with AD. The urgency of the question seemed clear: the global scale of the pandemic and aging populations make a better understanding of the potential impact of COVID-19 and other infections on the prevalence of future dementias a public health priority [10,16].

Recently developed high-throughput plasma proteomic biomarkers that provide biologically-based markers of risk or the onset of dementia years in advance of the expression of symptoms or of a clinical diagnosis can provide sensitive outcome measures for a study of early dementia pathology [17–24]. Amongst the best characterised plasma proteomic markers for AD are lowered beta amyloid-(1-42) ($A\beta_{42}$), its ratio with beta amyloid-1-40 ($A\beta_{42}:A\beta_{40}$) and raised phosphorylated tau (pTau), all of which accompany the accumulation of pathological proteins in the brain and clinical progression of AD [17,25–27] Concentrations of neurofilament light (NFL) (a non-specific marker of neuronal injury) and glial fibrillary acidic protein (GFAP) (which is associated with astrocyte activation) also are found to increase in plasma in early AD [28–30]. Studies have identified changes in these plasma proteins prior to dementia diagnoses and brain amyloid positivity by PET [17,23,25,30]. Decreases in plasma $A\beta_{42}:A\beta_{40}$ may precede the diagnosis of AD by up to 20 years, total Tau by 15 years, with increases in NfL and GFAP detected closer to diagnosis [24]. These biomarkers provide the opportunity to assess the effect of mild to moderate SARS-CoV-2 in healthy populations on Alzheimer's prodromal pathology and risk.

Raised levels of these dementia-related proteins have been reported in patients in the
context of acute COVID-19[19,31,32]. However, interpretation of these studies is confounded by
the potential for "reverse causation", as many risk factors for severe COVID-19 are also risk
factors for dementia. Complications of severe infections also can independently increase
biomarker levels. To robustly test for relationships between SARS-CoV-2 infection and brain
pathology for AD, longitudinal study designs that define changes from before to after
infection are needed. Symptomatically mild-to-moderate infections are an ideal study target
as they are less confounded by other factors associated with severe illness and prolonged
intensive care.   Furthermore, given that dementias also are associated with genetic
background, aging, and health status, analyses need to be able to take these factors into
account for elucidation of the independent effects of infection.

We were able to apply a design incorporating these principles to address the question of
whether SARS-CoV-2 infection initiates or potentiates brain pathology associated with
prodromal AD. We took advantage of the longitudinal blood sampling and clinical data
acquisition in UK Biobank in combination with that study's surveillance for SARS-CoV-2
exposure serology during the early stage of the UK COVID-19 pandemic (2020-21). Here
we report results from over 1200 UK Biobank participants tested longitudinally for a range of
plasma proteomic biomarkers. Our primary outcome measures were plasma levels of $A\beta_{42}$
$A\beta_{40}$, the $A\beta_{42}$:$A\beta_{40}$ ratio, pTau-181, NfL and GFAP (Quanterix Simoa assays). We were
able to test for changes in these biomarkers with infection and compare these changes to
those found in a demographically matched group who were sampled prospectively over the
same interval but who were not infected with SARS-CoV-2. This longitudinal, prospective,
matched case-control design permitted the modelling of changes in AD biomarkers with
control for a variety of potential confounders including aging and baseline differences in
genetic backgrounds, sex and health status between cases and controls. The extensive
genetic, imaging and health data of the UK Biobank dataset allowed detailed assessment of
confounds and relationships.   In a subset of participants, we also were able to assess
plasma levels of a selection of over 1400 proteins obtained from a high-throughput Olink
Explore general proteomic platform, which provided the opportunity to assess the impact of
SARS-CoV-2 infection on inflammatory proteins and proteins associated with risks of other
chronic diseases [20,33,34].

Results

Study Population

We studied plasma proteomic biomarkers of AD and dementia risk, as well as proteins from
a general proteomic panel, brain images and other health data that were made available
from 1252 UK Biobank participants who had been recruited for the UK Biobank COVID-19
study (626 matched case-control pairs, 331 female)[10]. All participants had taken part in an
imaging assessment session before the pandemic onset (baseline, May 2014 - March 2020).
Participants were 46-80 years of age at this session (Fig. 1b). Individuals from these
assessments who were determined to have previously had SARS-CoV-2 infection by at least
one of: home-based lateral flow SARS-CoV-2 antibody test, PCR antigen (swab) test, GP

records or public health records were invited for the COVID-19 imaging study (see Methods).
Symptoms related to COVID-19 in cases were mostly mild, but 20 participants were
hospitalised for COVID-19 or its complications. Of 289 cases for whom vaccination status
could be determined, 40 had data indicating they had been vaccinated before their first
positive COVID test. Cases were matched individually to eligible controls (defined based on
age [+/- 6 months], sex, ethnicity [white/non-white], and location and date of initial imaging
assessment [+/- 6 months]) who had no record of confirmed or suspected SARS-CoV-2
infection at the time of the repeat assessment. The COVID imaging assessments occurred
during the UK COVID-19 pandemic (February 2021 - February 2022). Intervals between the
two assessment sessions ranged from 12-82 months, matched for case-control pairs (Fig. 1
d,e,f). Blood samples for plasma proteomics were obtained at both imaging assessment
sessions.
153
We characterised cases and controls for potential comorbidities and other factors (Table 1).
Some characteristics differed between case and control groups, reflecting lifestyle factors
that could increase the likelihood of early infection: more were employed prior to the
pandemic (395 vs 364); more cases (n=18) than controls (n=6) identified as "key workers"
during the pandemic (p=0.023); household sizes were larger on average (p=0.007*) and
cases were more active on average (p=0.002*). Cases were slightly heavier (1.4kg) than
controls (p=0.028) (although hip/waist ratio and obesity rates did not differ). Other AD
comorbidities did not show statistically significant differences across cases and controls.
Cases had slightly higher numbers in some factors including smoking and reported prior
hypertension.
164
Single molecule array (Simoa) ultrasensitive measures of plasma amyloid-β_{42} (Aβ42),
amyloid-β_{40} (Aβ42), pTau-181, glial fibrillary acidic protein (GFAP) and neurofilament light
(NfL) were available on plasma collected before and after onset of the COVID pandemic for
cases and for controls (see Online Methods). A complete set of longitudinal data was
available for 600 cases and for 600 matched controls; the remaining cases and controls
were missing data from one or more assay. Olink antibody-based proximity extension assay
proteomic measurements of concentrations of 1452 proteins (including additional assays for
NFL and GFAP) in the same plasma samples were available for 277 of the case-control
pairs[35].
174
175

176
177

Fig. 1. a. Experimental design. Protein concentrations were assayed from plasma samples acquired from the UK Biobank imaging assessment visits, the second of which was specifically recruited for the study of COVID-19. b. Distribution of participant ages at the pandemic assessment. c. Sources of evidence for case selection. Antibody – home-based lateral flow SARS-CoV-2 antibody test; Antigen – PCR antigen (swab) test; Health records: GP and/or hospital records d. Distribution of pre-pandemic assessment visit dates e. Distribution of pandemic assessment visit dates f. Distribution of intervals between assessments. g. Estimated dates of COVID symptoms (from participants with antigen test results).

186
187

Fig. 2. Schematic of data availability for the study. Primary analyses focus on 624 individually matched cases and controls from the COVID19 imaging repeat study with SIMOA plasma proteomic data. Full details of participant selection and control matching for the COVID study can be found at biobank.ndph.ox.ac.uk/ukb/ukb/docs/casecontrol_covidimaging.pdf.

192
193
194
195

		CASES	MATCHED CONTROLS	P-VALUE	N
DEMOGRAPHICS	Age (baseline)	60.21 (7.41)	60.23 (7.41)	0.324	624
	Age (pandemic)	63.45 (7.02)	63.46 (7.02)	0.456	624
	Sex	0.47 (0.50)	0.47 (0.50)	1.000	624
	Height	170.34 (8.82)	169.89 (9.08)	0.222	622
	Weight	77.06 (15.02)	75.63 (14.54)	0.028	593
	Ethnicity (white)	0.93 (0.26)	0.92 (0.28)	0.437	624
GENETICS	*APOE-ε3e4*	123	110	0.869	423
	APOE-ε4e4	11	9	0.788	423
	APOE-ε3e2	73	62	0.547	423
HEALTH MEASURE	Hip/Waist Ratio	1.16 (0.12)	1.16 (0.12)	0.831	595
	BMI	26.56 (4.21)	26.23 (4.17)	0.110	593
	BP (sys)	137.23 (19.22)	137.26 (18.58)	0.984	424
	BP (dia)	79.26 (11.04)	79.33 (10.80)	0.625	424
	Hand Grip	32.01 (10.26)	31.75 (10.42)	0.691	587
	Chest wheeze	90	94	0.872	607
	Health self-rating	3.02 (0.62)	3.09 (0.64)	0.075	618
	GFR (pre)	88.12 (10.52)	86.62 (10.87)	0.071	225
LIFE STATUS	Alcohol Intake Freq	3.19 (1.34)	3.18 (1.32)	0.966	618
	Smoker	26	18	0.283	619
	Deprivation	16.50 (11.79)	15.56 (12.14)	0.230	590
	Income	2.83 (1.70)	2.82 (1.75)	0.893	620
	Number in Household	2.46 (1.19)	2.30 (1.09)	0.007	620
	Key Worker	18	6	0.023	624
	Employed	395	364	0.061	614
	Social Isolation	34	48	0.137	624
	Moderate activity (min/d)	63.25 (79.19)	52.29 (55.34)	0.002	503
	Vigorous activity (min/d)	42.92 (41.70)	42.36 (39.26)	0.576	293
COMORBIDITY	Type II Diabetes	17	15	0.858	624
	Heart Condition	108	100	0.595	624
	Obesity	113	107	0.710	624
	Hypertension	145	127	0.244	624
	Depression	26	20	0.453	624
	IBS	39	34	0.629	624
	COPD	4	3	1.000	624
	Emphysema	3	4	1.000	624
MEDICATIONS	Blood pressure medication	53	46	0.530	624
	Cholesterol medication	46	44	0.913	624
	Diabetes medications	3	3	1.000	624
	COVID vaccinated	384	388	0.351	286
COGNITION	Cognitive Ability Score	32.84 (7.27)	32.94 (7.59)	0.967	407
NEUROIMAGING	AD Neuroimaging Phenotype	-2.29 (0.66)	-2.30 (0.66)	0.799	590

Table 1. Comparisons of baseline characteristics of cases and controls. P-values reflect paired t-tests across matched case and control pairs (uncorrected). BMI: Body Mass Index, BP: blood pressure, BP medication:

204 The plasma Aβ42:Aβ40 ratio is reduced after SARS-CoV-2
205 infection

To test for effects of SARS-CoV-2 infection on preclinical AD brain pathology, Twhether
infection status affected longitudinal changes in concentrations of plasma Aβ40, Aβ42,
pTau-181, NFL and GFAP, with the baseline assessment completed before the COVID-19
pandemic and the repeat assessment made during the pandemic, after cases had
developed serological evidence for SARS-CoV-2 infection[17,36]. In addition to the plasma
concentrations, we also assessed the Aβ42:Aβ40 concentration ratio[26,27].

We fit linear models describing each participant's pandemic-session biomarker levels in
terms of their baseline, pre-pandemic levels, time interval between the two sample
acquisitions, age, sex, and their individual case (serology positive, post SARS-CoV-2
infection) or control (serology negative, no evidence of prior SARS-CoV-2 infection) status
(see **Error! Reference source not found.**). These models therefore modelled changes in
the plasma biomarker levels between assessments, with the case-control regressor
identifying differences in these changes associated specifically with SARS-CoV-2 infection.
We fit further models assessing the associations of a variety of comorbidities and other
factors with biomarker levels at baseline and longitudinally (Online Methods, Supplementary
table 1).

Average participant plasma concentrations of Aβ42, the Aβ42:Aβ40 ratio, and pTau-181
decreased between pre-pandemic and pandemic assessments for all participants (cases
and controls), while concentrations of Aβ40, NfL and GFAP increased (all p<0.0001, paired
t-tests). Statistical testing of the case-control model parameters found SARS-CoV-2
infection was associated with a significantly greater reduction in the Aβ42:Aβ40 ratio (2%
drop from baseline, False Discovery Rate [FDR] significant **p-value, p=0.001) (Fig. 3,
Supplementary table 2). These results were maintained when an extended model including
potential confounders was used (Supplementary table 3). The estimated effect of SARS-
CoV-2 on the Aβ42:Aβ40 ratio was comparable to the effect of 4 years of aging (-0.5%
change in Aβ42:Aβ40 ratio per year of age, estimated at baseline), and around half the
average effect of heterozygosity for *APOE-ε4* (*APOE-ε4* heterozygous participants showed a
3.9% lower Aβ42:Aβ40 ratio relative to *APOE-ε3* homozygous in the baseline assessment
sessions). Other comparable comorbidities of AD did not show a statistically significant
associations with the Aβ42:Aβ40 ratio in this dataset. Greater reductions in the plasma
Aβ42:Aβ40 ratio amongst cases were associated with greater severity of symptomatic
infections: cases who were hospitalised with COVID-19 showed over twice the magnitude of
reduction relative to non-hospitalised cases (5.3% vs 2.1%).

241
242
243

244
Fig. 3. Standardised model parameters for Case/Control term in each of the AD-protein change models.
Parameters represent the SARS-CoV-2 infection associated effect on change in protein levels between pre-
pandemic and pandemic assessment sessions. Aβ42:Aβ40 ratios showed significant relative reductions in
SARS-CoV-2 individuals. P-values correspond to one-sided tests in the directory of previously reported
biomarker associations with AD. Full model fits are in Supplementary tables 2 and 3 (basic and extended
models).

Greater changes in both plasma Aβ42 and pTau-181 after SARS-CoV-2 infections were found in older participants

254
Participants who were older or had prior lifestyle or health conditions that are risk factors for
neurodegenerative disease are likely to be more vulnerable to infection-related
neurodegenerative pathology. Age is a key risk factor both for AD and for the severity of
COVID-19 but the relationships between age and risk are non-linear as older people become
more vulnerable to severe disease symptoms in later years[37]. To test for specific effects of
SARS-CoV-2 on plasma protein level changes in older, more vulnerable study participants,
we used regressors reflecting an age-related vulnerability score derived from observed
associations between age and severe neurological outcomes for COVID-19 (Fig. 4c)[37]. This
score has previously been used to analyse SARS-CoV-2 changes in MRI measures[10]. We
assessed models that included this score and its interaction with case-control status to
assess whether age-related-vulnerability predicted changes in AD-proteins between
assessments with or without dependence on SARS-CoV-2 infection status (Online Methods).
267
Across both cases and controls, a higher age-related-vulnerability score was associated with
greater decreases in Aβ42:Aβ40 ratio and greater increases in pTau, NfL and GFAP
concentrations between the pre-pandemic and pandemic assessment sessions (Fig. 4a,
Supplementary table 4). Assessing the interaction of age-related-vulnerability score and
case-control status found age-related-vulnerability score dependent increases in pTau-181
(p=0.016**) and decreases in Aβ42 (p=0.029*) in those exposed to SARS-CoV-2 relative to
controls (Fig. 4b, Supplementary table 4). These effects were strengthened when an
extended set of confounds were modelled (Supplementary table 5). Age-resolved plots
indicate that SARS-CoV-2 cases begin to show greater changes in AD proteins from around
70 years of age (Fig. 5). For an average 75-year-old participant, this model estimates an

additional 4% increase in pTau-181, a 3% decrease in Aβ40 in SARS-CoV-2 positive
participants, alongside a 5.5% change in Aβ42:Aβ40 ratio (Online Methods).
280

281
282
283 *Fig. 4. Standardised model parameters for models including age-dependent vulnerability terms in each of the*
284 *AD-protein change models. a. parameters for age-dependent vulnerability term. b. parameters for age-dependent*
285 *vulnerability term interaction with case-control status. c. Age-dependent vulnerability term weightings P-values*
286 *correspond to one-sided tests corresponding to previously reported associations with AD. Full model fits are in*
287 *Supplementary tables 4 and 5 (basic and extended models).*

AD risk factors associated with accelerated changes in plasma protein biomarker levels in cases and controls.

We assessed similar models to examine the influence of a variety of further comorbidities for
AD and other factors on the protein biomarkers (Supplementary table 1). Within assessment
sessions, AD proteins showed rich set of associations with comorbidities and other factors
(Fig. 6a, Supplementary table 6). A lower Aβ42:Aβ40 ratio was associated individuals with
an *APOE-ε4* variant (p=0.001**) and current smokers (p=0.015*). While age was matched
across cases and controls, rates of these variates varied slightly across cases and controls
(Table 1) and contributed to a difference in pre-pandemic Aβ42:Aβ40 ratios between cases
and controls (0.0747 vs 0.0765, p=0.0145) (Supplementary table 7). These differences were
attenuated by regressing *APOE* and Smoking status. While differences in baseline would

not directly affect the longitudinal analysis of SARS-CoV-2 associated differences, including
these factors in our primary models indicated that these effects were not confounding
observed SARS-CoV-2 differences (Supplementary table 8).
305
Certain prior factors were associated with the extent of longitudinal changes in the AD
proteins (Fig. 6b, Supplementary table 9, Online Methods). Independent of case-control
status, male sex, higher hip/waist ratio, and poorer health self-rating were associated with
greater increases in pTau-181 across assessments (p<0.05**). Type II Diabetes appeared
to be associated with reduced changes. Analysis with a model accounting for these
potential confounds indicated that these differences could not account for the differences in
protein levels associated with SARS-CoV-2 infection seen above (Supplementary table 10).
313
A low Glomerular Filtration Rate (GFR) indicates poor kidney function and is known to
influence plasma protein levels[39] and alter Alzheimer's disease biomarker protein levels
without increasing disease risk[38].   Here, lower estimated GFR (available for n=225 cases)
was associated with significantly increased plasma concentrations of protein levels at
baseline (Fig. 6a). Estimated GFR did not show a significant difference between cases and
controls (Table 1). Higher GFR was associated with greater reductions in Aβ42 and Aβ40
protein concentrations between assessments, but not with changes in Aβ42:Aβ40 ratio (Fig.
6b). When GFR was included as a confounding covariate in models of AD biomarker
change, it had negligible impact on effect sizes for SARS-CoV-2-associated differences in
Aβ42:Aβ40 (Supplementary table 11).
324

325
326 *Fig. 6. Associations between AD protein levels and comorbidities and other factors, as determined from*
327 *regression models across all cases and controls. Shading shows the magnitudes of standardised betas, where*
328 *model parameter fits in models controlling for age, sex and time interval between assessments. a. Associations*
329 *within baseline assessment data only. Red colours represent positive associations. b. Associations of factors*
330 *with the change in AD protein levels across assessments. Red colours represent a positive association of factors*

336 # Prior hypertension, brain structural features associated with AD
337 # and plasma inflammatory protein levels were associated with
338 # greater SARS-CoV-2 infection-related changes in plasma AD
339 # biomarkers

340
Several factors showed specific associations with changes in AD-biomarker levels between
assessment sessions in SARS-CoV-2 positive participants (Fig. 6c, Supplementary table
12). Hypertension was associated with greater reductions in amyloid markers of AD
pathology in cases positive for SARS-CoV-2 infection (diastolic blood pressure: reduced
Aβ42:Aβ40 ratio p=0.002**; hypertension: reduced Aβ42:Aβ40 ratio p=0.0018*; reduced
Aβ42 p=0.004**). Hypertension has been linked to both more severe COVID-19
symptoms[39], and the progression of AD fluid biomarkers[40]. While *APOE* genotype has been
associated with COVID-19 severity previously [44,45] our analyses showed no clear effects of
prior vaccination (p=0.44) or *APOE* status on plasma protein changes from pre- to post-
pandemic in cases compared to controls. Sex also did not have a significant effect on the
AD plasma proteomic changes.
352
A brain imaging signature was developed previously from MRI volumetric and cortical
thickness measures in the Alzheimer's Disease Neuroimaging Initiative (ADNI) database [41],
and shown to be associated with higher rates of AD and lower memory and cognitive scores
among UK Biobank participants[41]. The AD signature values were standardised for sex, age,
and intracranial volume, and showed no case-control differences at baseline (p=0.779) or
significant associations with AD protein levels prior to SARS-CoV-2 infection (all p>0.1).
However, we found that higher values of this signature at baseline were associated with
greater increases in pTau-181 (p=0.012**) and greater reductions in Aβ42 (p=0.015**) after
SARS-CoV-2 infection relative to the matched controls (Fig. 6c, Supplementary table 13).
362
We also assessed whether baseline levels of a set of Olink inflammatory proteins associated
with COVID or AD were associated with neurodegenerative biomarker changes
(Supplementary table 14). None of these protein levels differed between cases and controls
at the pre-pandemic session (Supplementary table 15). Higher baseline levels of
Macrophage Inhibitory Factor (MIF) predicted greater reductions in Aβ protein levels across
both cases and controls (Supplementary table 16). Higher baseline levels of YKL-40 (or
chitinase-3 like-protein-1) were associated with greater Aβ42:Aβ40 reductions specifically in
the SARS-CoV-2 group (standardised beta = -0.08, p=0.006**) (Supplementary table 17).
YKL-40 is highly expressed in astrocytes during neuroinflammation and is associated with
increased brain amyloid and AD symptoms[42,43].
373

Associations between changes in plasma AD biomarkers and Olink protein levels

Peripheral inflammation is a probable component of a pathway for SARS-CoV-2 induced amyloid pathology. While in mild cases of COVID-19 inflammatory protein levels have been reported to normalise in the time frame of the present study (median time since last SARS-CoV-2 positive test = 68 days)[46], it is possible that some persistent inflammation, such as that seen in long-COVID, plays a role in the AD-related pathology we observe. Of 13 proteins assessed, TRAIL protein (TNFSF10), an apoptotic TNF cytokine was significantly reduced after SARS-CoV-2 infection (p=0.002**) (Supplementary table 18). This protein has been reported to be lower in patients with severe COVID[47] and lower levels have been identified as a biomarker associated with AD risk in separate UK Biobank plasma samples and disease records[20]. Inflammation would be expected to be higher in participants with severe symptoms. Those participants who had been hospitalised with COVID-19 showed increased levels of Interleukin 6 (p=0.01*) and the long pentraxin PTX3 (p=0.012*) in the pandemic assessment relative to controls.

Using disease reports from the health records of 47,600 of the UK biobank participants[20], the Olink proteins have previously been used to estimate weightings providing individual disease risk scores (ProteinScores) (online Methods). We were able to estimate 19 of these scores for the pre-pandemic and post-pandemic assessment sessions to assess whether they were associated with changes in AD-biomarkers. There were no differences in disease risk scores between cases and controls at baseline, and the incidence of all disease risks (except endometriosis) increased with age. SARS-CoV-2 was associated with increases in disease risk scores for several co-morbidities of Alzheimer's disease: type 2 diabetes (p=0.008**), chronic obstructive pulmonary disease (COPD) (p=0.001*), ischaemic stroke (p=0.0001*) and heart disease (p=0.011*) (Supplementary table 19). These changes are consistent with recent epidemiological reports of increased incidence of some of these diseases after the pandemic [48]. The AD risk score, weighted heavily to NfL and GFAP levels, did not show a significant increase (p=0.108).

Associations between AD biomarker changes and changes in neuroimaging, cognitive performance, and general health measures

Previous work using the UK Biobank brain imaging data identified changes in brain structure following SARS-CoV-2 infection, linked to age-related-vulnerability[10]. We assessed whether there was evidence that these changes could be associated specifically with the AD-related changes we observe. An interaction analysis with the age-dependent vulnerability score showed evidence (p=0.027) for increases in the Alzheimer's Disease Neuroimaging Initiative (ADNI)-derived AD structural brain phenotype score after SARS-CoV-2 infection in older participants. Increased AD brain signature scores were associated with increased pTau-181 and NfL between visits (p=0.015* and p=0.031, respectively).

Recent work has highlighted that COVID-19 is associated with long-lasting impairments in
multiple cognitive domains[12]. To test for relationships between cognitive performance and
SARS-CoV-2 infection in this population, we evaluated a measure of general cognitive ability
derived from UK Biobank cognitive tests (see Methods)[49]. SARS-CoV-2 infection was
associated with an additional 2.05% reduction in this measure over controls, equivalent to
almost two years of age-related decline (1.16%/year, p=0.029). However, these changes
were not strongly correlated with the AD biomarkers, which can change years prior to
cognitive changes. Reductions in cognitive scores were associated with reductions in Aβ42
(p=0.037) and increases in plasma Tau-181 (p=0.011*).
425
Finally, participants were also asked to rate their overall health between poor and excellent.
There were no significant differences at baseline between cases and controls. However,
cases showed greater deterioration in their overall health ratings (2.3% p=0.007*) between
the assessment visits than did the controls. While prior health self-ratings affected pTau-181
(above), changes in general health self-ratings across were not associated with AD protein
levels.

Discussion

433
SARS-CoV-2 infections can have a wide range of profound and long-lasting effects on
human health. There have been persistent raised levels of illness and mortality since the
pandemic, even in people who experienced only clinically mild cases of COVID-19[12,48,50,51].
However, evidence for the effect of COVID-19 on risk for late-life dementia has been lacking.
Here, using longitudinal ultrasensitive Simoa platform plasma Aβ and pTau measures that
are relatively specific and sensitive for AD and a demographically matched case-control
design, we discovered plasma proteomic biomarker evidence for prodromal AD pathology
after SARS-CoV-2 infection amongst UK Biobank participants. These people largely
experienced mild to moderate symptoms. Despite this, SARS-CoV-2 infection was
associated with significant reductions in Aβ42:Aβ40 between baseline and pandemic
assessment visits. Moreover, decreases in plasma Aβ42 and increases in pTau-181 were
seen in older participants and those with MRI brain signatures of AD. These protein
biomarker changes have all previously been associated with increased AD risk[17,26,36,52].
While some of these changes have been associated previously with severe COVID-19 in
acute settings[19,53], to our knowledge they have not been assessed in the post-acute setting
for largely mild cases. Together, our results provide evidence that SARS-CoV-2 infection
initiates or accelerates brain pathology associated with AD risk and suggests increased risk
of AD after infection.
452
Changes in plasma Aβ42:Aβ40 have been found to be the earliest detectable plasma
biomarker change in preclinical stages of AD[24] and are associated with brain amyloid
accumulation even in cognitively normal participants[27]. The observed mean magnitude of
the reduction in plasma Aβ42:Aβ40 after SARS-CoV-2 infection was substantial: equivalent
to 4 years of aging (based on the baseline pre-pandemic data for this participant group) and
60% of the effect size of inheriting a single *APOE-ε4* allele. Effect sizes were larger in the
subgroup from our study who were hospitalised for COVID-19, and for those reporting prior

hypertension, suggesting that the impact on brain pathology is likely to increase with more
severe infection.
462
We observed decreases in plasma Aβ42 and increases in pTau-181 associated with SARS-
CoV-2 in participants with higher age-dependent vulnerability scores and in those with MRI
brain signatures of AD prior to infection. In prior studies, concordant changes in Aβ42 and
pTau-181 follow Aβ42:Aβ40 in the progression of plasma biomarkers in pre-AD [24,25] and
have been associated with increasing brain Aβ load [17,26,36,52,54]. These changes in more
vulnerable participants were associated with brain structural changes, increased cognitive
deficits, and decrements in measures of general health quality.
470
Increases in NfL and GFAP in CSF and plasma have been reported previously with
Alzheimer's disease, MCI, and acute COVID [17,26,53,55,56]. Here we did not find increased
levels in either following SARS-CoV-2 infection. NfL and GFAP are markers of neural injury
and astroglial activation, respectively, which are more predominant in later stages of AD
progression. While the plasma Aβ42:Aβ40 ratio have been found to decrease from control
levels up to 18 years prior to an AD diagnosis, increases in plasma NfL and decreases in
hippocampal volume were not detectable until at least 9 years later[24]. We would expect few
participants to have such progressed AD pathology. The absence of changes in these
proteins also aligns with observations made during and after COVID. In hospitalised COVID
patients, plasma NFL and GFAP concentrations have been observed to be raised, but they
returned to control levels within 3 months for the majority of patients [57].
482
There are a variety of mechanisms by which mild to moderate SARS-CoV-2 infection may
increase the brain beta-amyloid load. Brain beta-amyloid production and aggregation can be
potentiated by stress, infection and pathological conditions including diabetes, stroke and
vascular disease [58,59]. Epidemiological studies have suggested that a wide range of
systemic infections may be associated with increased risks of AD [60,61]. SARS-CoV-2 is not
neurotrophic[62], but infection is associated with systemic inflammatory activation[63].   With
associations between peripheral inflammatory events and dementia identified in numerous
studies[64], possible neuroimmune mechanisms of AD pathogenesis are increasingly being
studied [65–67]. Pre-clinical studies of peripheral inflammatory challenges have found evidence
both for potentiation of brain microgliosis and for impairment of microglial clearance of beta-
amyloid and the progression of tau pathology [7,68] Levels of midlife systemic inflammatory
markers have been associated directly with late-life relative brain atrophy comparable to that
associated with the *APOE4* allele[69], an effect size mirroring that observed here.
496
Peripheral inflammation may influence CNS inflammatory processes in several ways. For
example, systemic infection can enhance CNS inflammation directly by trafficking of
activated immune cells or indirectly with increases in systemic cytokines or inflammatory
activation of endothelial cells and breakdown of the blood-brain-barrier to allow leak of
chronically pro-inflammatory plasma proteins into the CNS[14,15]. Direct infection of vascular
endothelial cells by SARS-CoV-2 would enhance this [70]. Long-lasting immunological
changes in patients could maintain these effects for prolonged periods even in mild-to-
moderate cases[71]. One specific mechanism is increased brain interferon-induced
transmembrane protein (IFITM1-3) expression triggered by viral infection. Increased
incorporation of IFITM3 into the g-secretase complex potentiates its activity, increasing Aβ
production [72,73].

508
The AD biomarker proteins studied here are being increasingly assessed at scale[24,74], but
their patterns of their variation in the general population are not fully characterised. Aβ and
pTau are strongly linked to AD and neurodegeneration, but they can vary with specific
conditions other than AD[87]. In the present dataset, concentrations of these proteins showed
associations with a variety of traits including hypertension, weight, GFR and Type II Diabetes
diagnosis. However, the observed SARS-CoV-2 infection-related effects remained when
these factors were included in the model as potential confounders.
516
In addition to highlighting increased plasma biomarkers of prodromal AD with SARS-CoV-2
infection, application of the more general Olink plasma protein-based risk scores developed
in an earlier UK Biobank study also provides evidence for greater predicted risks for
diabetes, ischaemic stroke and heart disease, COPD, lung cancer and endometriosis in the
subset of participants under the age of 73[20]. This is consistent with emerging post-COVID-
19 epidemiology data [76–79] and proteomic studies of samples from COVID-19 patients
hospitalised with acute symptoms[19]. While we found limited evidence for comorbidities
directly confounding our AD biomarker observations, the raised risk of many - albeit not all –
diseases with SARS-CoV-2 infection raises the possibility that SARS-CoV-2 (and other
infections) may accelerate both body and brain aging [82]. The changes in these risk-markers
across diseases may be due to common (e.g., inflammation) processes initiated with SARS-
CoV-2 infection, a hypothesis supported by findings of associations of Alzheimer's disease
with multiple co-morbidities[88]. It is likely that infections can have both general and specific
impacts on disease risks, and will affect different organ systems[1,23].
531
Patients with hepatic dysfunction have raised levels of plasma beta-amyloids and increased
Aβ42/40 ratios[82,83]. However, this is opposite to the direction of change observed here. We
also did not find evidence for increases in Olink protein signatures associated with hepatic
dysfunction after SARS-CoV-2. Kidney function can also affect plasma protein levels. We
identified associations between eGFR and protein levels, but these did not confound our
primary results. Increased plasma pTau-181 is associated with combined peripheral and
central nerve pathology in ALS and with spinal muscular atrophy[76]. While further research is
necessary[77], we believe that the association of both increased plasma pTau-181 and
decreased plasma beta-amyloid 42/40 ratios with SARS-CoV-2 infection most likely reflects
brain pathology.
542
While our study was able to combine fluid biomarkers, imaging data, cognitive performance,
and other covariates in a longitudinal case-control design, it has limitations. We relied on
data linkage to hospital records and retrospective reports for information concerning
symptoms associated with SARS-CoV-2 infections, limiting our ability to investigate detailed
relationships between the clinical severity of infections and plasma biomarker changes.
While we used prospectively acquired samples from before and after SARS-CoV-2 infection,
the pragmatic, observational nature of our analyses precludes making strong causal
inferences. While case and control groups were well matched by most measures (including
cognitive scores, inflammation markers and disease risks), there were some differences
variables in body mass, activity levels, and diabetes mellitus and hypertension rates. While
analyses controlled for these and other known risk factors, it is possible there are additional,
uncontrolled susceptibility factors that distinguished the SARS-CoV-2 infection and control
groups. Asymptomatic cases may be under-represented in the sample. This does not

invalidate the assessment of accelerated changes in biomarkers in the current sample
relative to baseline levels, but it means that these changes may be lower in the broader
population. Generalisation of our results to the wider population must be done with care: the
UK Biobank sample population is not representative of the general population by many
measures[83].
561
Recent studies have compared various blood phosphorylated tau immunoassays, notably
those measuring pTau-217 and -231, for their utility as biomarkers for AD[84] . In head-to-
head assays, all were able to identify AD and brain Aβ pathology[85].   pTau-217 has shown a
greater sensitivity to later-stage disease progression, but pTau-231 and pTau-181 may show
greater sensitivity to early stage amyloid accumulation, the primary disease phase of interest
here [87,88].
568
In conclusion, our study, along with earlier studies suggesting increased rates of dementia
diagnosis following COVID-19[89], suggests that public health planning should take into
account the possibility of increased rates of AD (and other systemic diseases) in the coming
years due to the COVID-19 pandemic. Moreover, the relationships described here are
unlikely to be specific for SARS-CoV-2 and may reflect more general effects of systemic viral
or other infections on the aging brain. This highlights the need to prioritise pandemic
preparedness as part of a strategy for reducing the future long-term burden of dementia and
other chronic diseases, as well as for the direct impact of the acute infections.
577

Online Methods

Ethics

The UK Biobank is approved by the North West Multi-Centre Research Ethics Committee
(MREC) to obtain and share data and samples from volunteer participants. Written informed
consent was obtained from all participants (http://www.ukbiobank.ac.uk/ethics/).

Design

584
This study analyses Simoa ultra-sensitive assays of dementia-related proteins taken from
blood-plasma obtained from participants in the UK Biobank COVID-19 repeat imaging
study)[10]. This study identified case and control participants from those who had taken part in
the UK Biobank imaging enhancement[90], which comprises a comprehensive imaging
assessment at one of four dedicated sites. Over 40,000 participants had been assessed
prior to the COVID-19 pandemic. Inclusion criteria and other details are described in
https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/casecontrol_covidimaging.pdf.
592
Participants were identified as being SARS-CoV-2-positive by the UK Biobank if (a) there
were positive diagnostic antigen tests identified in the linkage to health-related records, (b)
COVID-19 was reported in their primary care data or hospital records, or (c) two home-
based lateral-flow antibody tests provided by the UK Biobank were positive (Fortress Fast

COVID-19 Home test, Fortress Diagnostics and ABC-19TM Rapid Test, Abingdon Health).
Both diagnostic antigen test and GP/hospital records were accompanied by data permitting
estimation of date of COVID-19 infection.
600

In all, Simoa biomarker measurements from 626 cases and their matched controls were
available. Each participant had biomarker measurements from blood samples taken
longitudinally from the two imaging assessment sessions (pre- and post-pandemic).
Complete pairs of these data were available from 600 cases and from the 600 their matched
controls had complete data. For the 626 cases, SARS-CoV-2 infection was determined from
mixed sources: 358 had a general practitioner's diagnosis of COVID-19 , 438 had record of a
positive diagnostic antigen test prior to the assessment, and 467 returned positive on home-
based antibody lateral flow test kits (Fig 1b). 24 participants had SARS-CoV-2 infection
determined during a period of hospitalisation, for which 20 had COVID-19 listed as a primary
cause. Two participants were excluded for having a prior dementia diagnosis (vascular
dementia), identified from hospital inpatient records.
612

Control participants were selected as those who had negative antibody test results, and / or
no other record of confirmed or suspected COVID-19 from available data. Control
participants were matched to positive cases based on five characteristics: sex, ethnicity
(non-white/white), date of birth (+/- 6 months), the location of first imaging assessment, and
the date of this assessment (+/- 6 months). Further details of case and control identification
are provided in the above link and [10].
619

Proteomic assay acquisition and pre-processing

Aβ40, Aβ42, NfL and GFAP concentrations were measured using the Simoa Human
Neurology 4-Plex E assay (Quanterix). The Simoa P-tau181 Advantage Kit was used to
measure P-tau181 concentration. All measurements were made on an HD-X instrument
(Quanterix) with one round of experiments and a single batch of reagents. Technicians were
blinded to participant case-control status and from which assessment visit a sample was
taken from. Measurements were all above limit of detection and coefficients of variation
within analytes were below 10%.
628

Anonymised data from 1256 participants were integrated with phenotype data via a UK
Biobank key. Four participants were excluded for data being only available for one of the
two imaging sessions. P-tau181, NfL, and GFAP markers showed positively skewed
distributions and were log-normalised. For each biomarker, outliers were identified as
measurements more than 8 times the Median Absolute Deviation (MAD) and excluded from
the analysis. PCR plate IDs were also regressed out from the data. There was no
significant imbalance of outliers or plate IDs across cases and controls. Cases and controls
were matched according to UK Biobank field 4100 with 609 case-control pairs identified.
1220 participants and 583 matched case-control pairs had complete proteomic data. All
analyses were performed on matched data, except (as noted) when key covariates of
interest were missing from a set of participants. The Simoa Neurology assays showed
patterns of covariance across participants closely matching previous reports (Fig 5) [91].
641

Olink Proteomics

We used Olink Explore proteomic data from the Pharma Proteomics Project (PPP) which assayed samples from 54,219 UK Biobank participants across the different assessment sessions [35]. Available data from the COVID-19 imaging assessment sessions for our case/control matched pairs comprised 436 matched participant pairs for which 1474 proteins from Cardiometabolic, Inflammation, Neurology and Oncology Olink panels were assayed. The Olink 3072 Explore platform extension data were not available for this COVID-19 case-control matched assessment visits. Available Olink proteins included NFL and GFAP which were also profiled in the Simoa assays. These markers were positively correlated to their Simoa counterparts (r=0.70 and 0.53 respectively).

The Olink panel included Cystatin-C, which permitted estimation of the Glomerular Filtration Rate (eGFR) in the imaging assessment sessions using the KD-EPI Cystatin C Equation[92]. We first assessed how well protein levels measured by OLINK related to serum Cystatin-C measurements made separately and included in the UK Biobank core blood biochemistry panel data. The latter were derived from the Siemens Advia 1800 platform protein and available only for the initial UK Biobank assessment session. OLINK Cystatin-C measures in our study baseline plasma samples and those from earlier plasma samples acquired on the same participants showed a correlation of 0.84, which was used for appropriate rescaling of the OLINK data to enable eGFR estimation from the OLINK data.

We assessed 13 inflammatory proteins identified from a literature search of associations of inflammatory proteins in blood or CSF with COVID and/or Alzheimer's disease (Supplementary table 14). The inflammatory proteins and risk scores were analysed for baseline case-control differences and associations with AD proteins change using the statistical models outlined below.

We also investigated whether recently defined multivariate plasma protein concentration changes associated with disease risk (ProteinScores) derived from Olink Explore plasma protein biomarkers in 47,600 UK biobank participants[20] provided evidence for increased systemic chronic disease risks following SARS-CoV-2 infection. These scores were derived using penalised Cox elastic net regression which linked protein levels some samples in the initial UK Biobank assessment session to over 16 years of primary and secondary care NHS records disclosing 21 incident outcomes. Being penalised regression, each score comprises weights from 10-20 Olink proteins, selection of which was validated using hold-out data from the UK Biobank. As these risk scores were derived and validated on participants below the age of 73 years, we limited our analysis to this subset of our case-control cohort (i.e., the 353 cases and 353 matched controls under the age of 73).

We generated these risk scores for protein profile data from the COVID-19 imaging assessment sessions. The data were rank-base inverse normal transformed and rescaled to match the processing applied in the generation of the ProteinScores. Disease weight scores were downloaded from the publication website and applied to the data[20].

Phenotypic data processing

Primary data for the study comes from the two imaging assessment sessions. Genetic and phenotypic data, including imaging and health records were available for all participants who took part in the COVID-19 reimaging project. Symptomology analyses used data from the UK Biobank COVID-19 Serology Study[93]. Data were parsed and cleaned using the FMRIB UK Biobank Normalisation, Parsing and Cleaning Kit, and integrated with other data modalities into a combined Python pandas data frame.

Genetics

APOE variant status was extracted from UK Biobank subject genotyping using PLINK-2.0[94]. We extracted variants at SNPs *rs429358* and *rs7412* to identify participants' *APOE* status. Data was available for 1027 participants, with five genotypes: ε3ε3 (n=630), ε3ε4 (235), ε2ε3 (135), ε4ε4 (20), and ε2ε2(7). We used ε3ε3 genotype as reference when one was required. Additional AD and COVID-19 -associated single nucleotide polymorphisms in the UK Biobank dataset were sequenced directly or imputed.

Imaging data and neuroimaging AD phenotype score

To avoid the multiple testing challenge of large numbers of neuroimaging measures and provide a measure recognised to be relevant to AD, we estimated a structural brain data AD phenotype score for each participant at each assessment session using structural brain imaging-derived measures made available through the UK Biobank Showcase[95,96]. The AD phenotype score, developed on the Alzheimer's Disease Neuroimaging Initiative (ADNI) dataset, reflects the prediction of a Bayesian machine learning neural that a participant has AD from 155 FreeSurfer cortical volume and thickness measures[41]. Monte Carlo dropout was used to approximate Bayesian inference within a two-layer neural network. This phenotype model was trained on 736 individuals (331 AD) from the ADNI database, and validated on 5209 participants from the National Alzheimer's Coordinating Centre and 37,104 participants in the UK Biobank [41], where predicted at-risk participants showed cognitive profiles representative of AD.

Neuroimaging measures contributing to the AD phenotype were processed using the same steps as used for the generation of the model. Each cortical volume and thickness measures was cleaned by removing outliers more extreme than eight times the median absolute deviation from the median (across both pre-pandemic and pandemic assessment visits). The measures were then de-confounded for age, estimated total intracranial volume, and sex, and rescaled using normalisation statistics from the ADNI training set. Software to generate these signatures was downloaded from https://github.com/tjiagoM/adni_phenotypes . PyTorch was used to apply the trained model (https://wandb.ai/tjiagom/adni_phenotypes/runs/2cxy59fk) to UK Biobank data.

Cognitive Scores

The UK Biobank includes a series of standard cognitive tests, but they can be individually
unreliable, and do not together correspond to established tests for general or dementia-
related cognitive scoring. A procedure to estimate a measure of general cognitive ability
from the UK Biobank data has previously been developed using an external validation
dataset [49]. We used the resulting score weights to calculate an overall score for cognitive
ability for participants in each assessment visit.

Statistical modelling

We assessed longitudinal change in proteomic biomarkers and the effect of SARS-CoV-2 on
this change using linear models which modelled protein levels in the pandemic assessment
samples in terms of the pre-pandemic protein concentrations, the time between assessment
visits, designation as a case or control and potential confounders including age and baseline
differences genetic backgrounds, sex, health status or social risk factors for COVID between
cases and controls[10]. We approached this is a staged fashion that allowed the impact of
confounder covariates (some which were available only in a proportion of subjects) to be
understood. Our basic model included covariates for Sex, Age, and the interval between
assessments:
741
742 Protein_post = Protein_pre + Interval_between_assessments +
743 Interval_between_assessments^2 + Case/control + Genetic_sex + Age_post
744
745 The extended models included further variables known or expected to change over the
746 assessment period and with COVID-19 (Table 4). Some variables presented considerable
747 missing data, such that the extended models had reduced power.
748
As the impact of SARS-CoV-2 infection is expected to vary according to certain
characteristics and vulnerabilities, we fit further models including interaction terms modelling
the interaction of SARS-CoV-2 status with characteristics including *APOE* status,
hospitalisation, and pre-pandemic neuroimaging AD score, to determine if these factors
modified the effect of SARS-CoV-2 on protein levels. Age is an important vulnerability
factor. Past modelling studies of brain structural changes associated with COVID-19 have
used a score of age-dependent vulnerability as an interaction term to account for the greater
vulnerability of the brain with age[10]. An exponential age-dependent vulnerability score term
($10^{Age-\times-0.0524--3.27}$) was derived from the modelling of the observed relationship between
age and COVID-19 neurological symptom severity[37]. All these interaction models included
both the original case-control variable and the original vulnerability terms in addition to the
interaction term, along with additional covariates used in the basic model.
761
In addition to protein levels, we modelled several additional outcome measures: disease risk
scores, a general cognitive ability score, a neuroimaging AD score, and reported level of
general health. These outcomes were modelled in the same manner as the protein levels
above. To assess whether changes in these outcomes were associated with changes in
protein scores, additional models were assessed which included as predictors the pre-
pandemic levels of proteins and their change across assessment visits. Linear models were
also used to assess relationships between protein levels and UK Biobank variables at

baseline. Age, sex, and interval between assessments were used as confound variables in
these models as above.

The models were fit using Ordinary Least Squares and tested model parameters using t-
tests. As we had strong hypotheses regarding direction of SARS-CoV-2-induced change in
AD protein levels based on prior literature, we used one-sided tests. Benjamini/Hochberg
False Discovery Rate with alpha=0.05 was used to control for statistical testing of multiple
hypothesis across proteins and/or covariates.

777   Data and code availability

778   All data is available upon application from the UK Biobank.

780   Code for analyses and figures will be shared from the UKDRI GitHub page.

782   Acknowledgements

This research has been conducted using the UK Biobank Resource under Application
Number 76059. We are grateful to the participants, researchers, and all UK Biobank staff for
their contributions to enabling this research. We particularly acknowledge the efforts of the
staff and the commitment of UK Biobank participants to allowing the acquisition of these data
during the COVID-19 pandemic. We also want to acknowledge support for the conception,
development, funding and interpretation of results of the study from Dr. Sally Johns of
Biogen. We thank the anonymous reviewers for their valuable suggestions. Figures were
created with BioRender.com.

EPD is supported jointly by funding to PMM from the UK Dementia Research Institute and
from the NIHR Imperial Biomedical Research Centre, in which he is a member of the
Multiple Long-Term Conditions theme. HZ is a Wallenberg Scholar supported by grants
from the Swedish Research Council (#2023-00356; #2022-01018 and #2019-02397), the
European Union's Horizon Europe research and innovation programme under grant
agreement No 101053962, Swedish State Support for Clinical Research (#ALFGBG-71320),
the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862), the AD
Strategic Fund and the Alzheimer's Association (#ADSF-21-831376-C, #ADSF-21-831381-
C, and #ADSF-21-831377-C), the Bluefield Project, Cure Alzheimer's Fund, the Olav Thon
Foundation, the Erling-Persson Family Foundation, Stiftelsen för Gamla Tjänarinnor,
Hjärnfonden, Sweden (#FO2022-0270), the European Union's Horizon 2020 research and
innovation programme under the Marie Skłodowska-Curie grant agreement No 860197
(MIRIADE), the European Union Joint Programme – Neurodegenerative Disease Research
(JPND2021-00694), the National Institute for Health and Care Research University College
London Hospitals Biomedical Research Centre, and the UK Dementia Research Institute at
UCL (UKDRI-1003). PMM acknowledges generous personal support from the Edmond J
Safra Foundation and Lily Safra and an NIHR Senior Investigator Award. His work
additionally is supported by the UK Dementia Research Institute, which receives its funding

from UK DRI Ltd., funded by the UK Medical Research Council, Alzheimer's Society and
Alzheimer's Research UK and, for this research specifically, Biogen.
812

Competing interests

HZ has served at scientific advisory boards and/or as a consultant for Abbvie, Acumen,
Alector, Alzinova, ALZPath, Annexon, Apellis, Artery Therapeutics, AZTherapies, Cognito
Therapeutics, CogRx, Denali, Eisai, Merry Life, Nervgen, Novo Nordisk, Optoceutics,
Passage Bio, Pinteon Therapeutics, Prothena, Red Abbey Labs, reMYND, Roche,
Samumed, Siemens Healthineers, Triplet Therapeutics, and Wave, has given lectures in
symposia sponsored by Alzecure, Biogen, Cellectricon, Fujirebio, Lilly, and Roche, and is a
co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU
Ventures Incubator Program (outside submitted work). PMM is a consultant for Biogen, Sudo
Therapeutics, Nimbus, Astex, GSK and Sangamo. He received research funding for aspects
of this work from Biogen and the UK DRI. He has research funding unrelated to this work
from Biogen and Bristol Meyers Squibb.
825

826   # References

827
1. Levine, K. S. *et al.* Virus exposure and neurodegenerative disease risk across national

biobanks. *Neuron* **111**, 1086-1093.e2 (2023).

2. Ou, Y.-N. *et al.* Associations of Infectious Agents with Alzheimer's Disease: A Systematic

Review and Meta-Analysis. *Journal of Alzheimer's Disease* **75**, 299–309 (2020).

3. Muzambi, R. *et al.* Assessment of common infections and incident dementia using UK

primary and secondary care data: a historical cohort study. *The Lancet Healthy*

Longevity **2**, e426–e435 (2021).

4. Muzambi, R. *et al.* Are infections associated with cognitive decline and neuroimaging

outcomes? A historical cohort study using data from the UK Biobank study linked to

electronic health records. *Transl Psychiatry* **12**, 1–11 (2022).

5. Debisarun, P. A. *et al.* Induction of trained immunity by influenza vaccination - impact on

COVID-19. *PLOS Pathogens* **17**, e1009928 (2021).

6. Franceschi, C. & Campisi, J. Chronic Inflammation (Inflammaging) and Its Potential

Contribution to Age-Associated Diseases. *The Journals of Gerontology: Series A* **69**,

S4–S9 (2014).

7. Tejera, D. *et al.* Systemic inflammation impairs microglial Aβ clearance through NLRP3

inflammasome. *The EMBO Journal* **38**, e101064 (2019).

8. Hosseini, S., Michaelsen-Preusse, K., Schughart, K. & Korte, M. Long-Term

Consequence of Non-neurotropic H3N2 Influenza A Virus Infection for the Progression

of Alzheimer's Disease Symptoms. *Frontiers in Cellular Neuroscience* **15**, (2021).

9. Xu, E., Xie, Y. & Al-Aly, Z. Long-term neurologic outcomes of COVID-19. *Nat Med* **28**,

2406–2415 (2022).

10. Douaud, G. *et al.* SARS-CoV-2 is associated with changes in brain structure in UK

Biobank. *Nature* **604**, 697–707 (2022).

11. Hampshire, A. *et al.* Cognitive deficits in people who have recovered from COVID-19.

eClinicalMedicine **39**, (2021).

12. Hampshire, A. *et al.* Cognition and Memory after Covid-19 in a Large Community

Sample. *N Engl J Med* **390**, 806–818 (2024).

13. Shan, D., Wang, C., Crawford, T. & Holland, C. Temporal Association between COVID-

19 Infection and Subsequent New-Onset Dementia in Older Adults: A Systematic

Review and Meta-Analysis. SSRN Scholarly Paper at

https://doi.org/10.2139/ssrn.4716751 (2024).

14. Xie, J. *et al.* Low-grade peripheral inflammation affects brain pathology in the AppNL-G-

Fmouse model of Alzheimer's disease. *acta neuropathol commun* **9**, 163 (2021).

15. Yang, A. C. *et al.* Dysregulation of brain and choroid plexus cell types in severe COVID-

19. *Nature* **595**, 565–571 (2021).

16. Taquet, M., Geddes, J. R., Husain, M., Luciano, S. & Harrison, P. J. 6-month

neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a

retrospective cohort study using electronic health records. *The Lancet Psychiatry* **8**,

416–427 (2021).

17. Zetterberg, H. & Blennow, K. Moving fluid biomarkers for Alzheimer's disease from

research tools to routine clinical diagnostics. *Molecular Neurodegeneration* **16**, 10

(2021).

18. Saunders, T. S. *et al.* Predictive blood biomarkers and brain changes associated with

age-related cognitive decline. *Brain Communications* **5**, fcad113 (2023).

19. Wang, L. *et al.* Plasma proteomics of SARS-CoV-2 infection and severity reveals impact

on Alzheimer's and coronary disease pathways. *iScience* **26**, (2023).

20. Gadd, D. A. *et al.* Blood protein levels predict leading incident diseases and mortality in

UK Biobank. 2023.05.01.23288879 Preprint at

https://doi.org/10.1101/2023.05.01.23288879 (2023).

21. Carrasco-Zanini, J. *et al.* Proteomic prediction of common and rare diseases.

2023.07.18.23292811 Preprint at https://doi.org/10.1101/2023.07.18.23292811 (2023).

22. You, J. *et al.* Plasma proteomic profiles predict individual future health risk. *Nat Commun*

14, 7817 (2023).

23. Oh, H. S.-H. *et al.* Organ aging signatures in the plasma proteome track health and

disease. *Nature* **624**, 164–172 (2023).

24. Jia, J. *et al.* Biomarker Changes during 20 Years Preceding Alzheimer's Disease. *New*

England Journal of Medicine **390**, 712–722 (2024).

25. Janelidze, S. *et al.* Plasma P-tau181 in Alzheimer's disease: relationship to other

biomarkers, differential diagnosis, neuropathology and longitudinal progression to

Alzheimer's dementia. *Nat Med* **26**, 379–386 (2020).

26. Baldeiras, I. *et al.* Addition of the Aβ42/40 ratio to the cerebrospinal fluid biomarker

profile increases the predictive value for underlying Alzheimer's disease dementia in mild

cognitive impairment. *Alzheimer's Research & Therapy* **10**, 33 (2018).

27. Fandos, N. *et al.* Plasma amyloid β 42/40 ratios as biomarkers for amyloid β cerebral

deposition in cognitively normal individuals. *Alzheimers Dement (Amst)* **8**, 179–187

(2017).

28. Palmqvist, S. *et al.* Cerebrospinal fluid and plasma biomarker trajectories with increasing

amyloid deposition in Alzheimer's disease. *EMBO Molecular Medicine* **11**, e11170

(2019).

29. Mattsson, N., Andreasson, U., Zetterberg, H., Blennow, K., & for the Alzheimer's

Disease Neuroimaging Initiative. Association of Plasma Neurofilament Light With

Neurodegeneration in Patients With Alzheimer Disease. *JAMA Neurology* **74**, 557–566

(2017).

30. Kim, K. Y., Shin, K. Y. & Chang, K.-A. GFAP as a Potential Biomarker for Alzheimer's

Disease: A Systematic Review and Meta-Analysis. *Cells* **12**, 1309 (2023).

31. Lai, Y.-J. *et al.* Biomarkers in long COVID-19: A systematic review. *Frontiers in Medicine*

10, (2023).

32. Frontera, J. A. *et al.* Comparison of serum neurodegenerative biomarkers among

hospitalized COVID-19 patients versus non-COVID subjects with normal cognition, mild

cognitive impairment, or Alzheimer's dementia. *Alzheimers Dement* **18**, 899–910 (2022).

33. Williams, S. A. *et al.* Plasma protein patterns as comprehensive indicators of health. *Nat*

Med **25**, 1851–1857 (2019).

34. Lindbohm, J. V. *et al.* Plasma proteins, cognitive decline, and 20-year risk of dementia in

the Whitehall II and Atherosclerosis Risk in Communities studies. *Alzheimer's &*

Dementia **18**, 612–624 (2022).

35. Sun, B. B. *et al.* Plasma proteomic associations with genetics and health in the UK

Biobank. *Nature* **622**, 329–338 (2023).

36. Schindler, S. E. *et al.* High-precision plasma β-amyloid 42/40 predicts current and future

brain amyloidosis. *Neurology* **93**, e1647–e1659 (2019).

37. Levin, A. T. *et al.* Assessing the age specificity of infection fatality rates for COVID-19:

systematic review, meta-analysis, and public policy implications. *European Journal of*

Epidemiology **35**, 1123 (2020).

38. Stocker, H. *et al.* Association of Kidney Function With Development of Alzheimer

Disease and Other Dementias and Dementia-Related Blood Biomarkers. *JAMA Network*

Open **6**, e2252387 (2023).

39. Gallo, G., Calvez, V. & Savoia, C. Hypertension and COVID-19: Current Evidence and

Perspectives. *High Blood Press Cardiovasc Prev* **29**, 115–123 (2022).

40. Biskaduros, A. *et al.* Longitudinal trajectories of Alzheimer's disease CSF biomarkers

and blood pressure in cognitively healthy subjects. *Alzheimer's & Dementia* **n/a**,.

41. Camerlingo, C. Vaccination to reduce severe COVID-19 and mortality in COVID-19

patients: a systematic review and meta-analysis. *European Review*

https://www.europeanreview.org/article/28248 (2022).

42. Safdari Lord, J., Soltani Rezaiezadeh, J., Yekaninejad, M. S. & Izadi, P. The association

of APOE genotype with COVID-19 disease severity. *Sci Rep* **12**, 13483 (2022).

43. Azevedo, T. *et al.* Identifying healthy individuals with Alzheimer neuroimaging

phenotypes in the UK Biobank. 2022.01.05.22268795 Preprint at

https://doi.org/10.1101/2022.01.05.22268795 (2022).

44. Klyucherev, T. O. *et al.* Advances in the development of new biomarkers for Alzheimer's

disease. *Transl Neurodegener* **11**, 25 (2022).

45. Vergallo, A. *et al.* Association of plasma YKL-40 with brain amyloid-β levels, memory

performance, and sex in subjective memory complainers. *Neurobiology of Aging* **96**, 22–

32 (2020).

46. Lennol, M. P. *et al.* Transient Changes in the Plasma of Astrocytic and Neuronal Injury

Biomarkers in COVID-19 Patients without Neurological Syndromes. *Int J Mol Sci* **24**,

2715 (2023).

47. Byeon, S. K. *et al.* Development of a multiomics model for identification of predictive

biomarkers for COVID-19 severity: a retrospective cohort study. *Lancet Digit Health* **4**,

e632–e645 (2022).

48. Gaudet, L. A. *et al.* Associations between SARS-CoV-2 infection and incidence of new

chronic condition diagnoses: a systematic review. *Emerg Microbes Infect* **12**, 2204166.

49. Fawns-Ritchie, C. & Deary, I. J. Reliability and validity of the UK Biobank cognitive tests.

PLOS ONE **15**, e0231627 (2020).

50. Atchison, C. J. *et al.* Long-term health impacts of COVID-19 among 242,712 adults in

England. *Nat Commun* **14**, 6588 (2023).

51. Mizrahi, B. *et al.* Long covid outcomes at one year after mild SARS-CoV-2 infection:

nationwide cohort study. *BMJ* **380**, e072529 (2023).

52. Boulo, S. *et al.* First amyloid β1-42 certified reference material for re-calibrating

commercial immunoassays. *Alzheimer's & Dementia* **16**, 1493–1503 (2020).

53. Virhammar, J. *et al.* Biomarkers for central nervous system injury in cerebrospinal fluid

are elevated in COVID-19 and associated with neurological symptoms and disease

severity. *European Journal of Neurology* **28**, 3324–3331 (2021).

54. Therriault, J. *et al.* Association of Phosphorylated Tau Biomarkers With Amyloid Positron

Emission Tomography vs Tau Positron Emission Tomography. *JAMA Neurol* **80**, 188–

199 (2023).

55. van Arendonk, J. *et al.* Plasma neurofilament light chain in relation to 10-year change in

cognition and neuroimaging markers: a population-based study. *GeroScience* (2023)

doi:10.1007/s11357-023-00876-5.

56. O'Connor, A. *et al.* Plasma GFAP in presymptomatic and symptomatic familial

Alzheimer's disease: a longitudinal cohort study. *J Neurol Neurosurg Psychiatry* **94**, 90–

92 (2023).

57. Diray-Arce, J. *et al.* Multi-omic longitudinal study reveals immune correlates of clinical

course among hospitalized COVID-19 patients. *CR Med* **4**, (2023).

58. Zhang, X., Fu, Z., Meng, L., He, M. & Zhang, Z. The Early Events That Initiate β-Amyloid

Aggregation in Alzheimer's Disease. *Front Aging Neurosci* **10**, 359 (2018).

59. Abbott, A. Are infections seeding some cases of Alzheimer's disease? *Nature* **587**, 22–

25 (2020).

60. Mawanda, F., Wallace, R. B., McCoy, K. & Abrams, T. E. Systemic and localized extra-

central nervous system bacterial infections and the risk of dementia among US veterans:

A retrospective cohort study. *Alzheimer's & Dementia: Diagnosis, Assessment &*

Disease Monitoring **4**, 109–117 (2016).

61. Sipilä, P. N. *et al.* Hospital-treated infectious diseases and the risk of dementia: a large,

multicohort, observational study with a replication cohort. *The Lancet Infectious*

Diseases **21**, 1557–1567 (2021).

62. Radke, J. *et al.* Proteomic and transcriptomic profiling of brainstem, cerebellum and

olfactory tissues in early- and late-phase COVID-19. *Nat Neurosci* **27**, 409–420 (2024).

63. Gordon, M. N. *et al.* Impact of COVID-19 on the Onset and Progression of Alzheimer's

Disease and Related Dementias: A Roadmap for Future Research. *Alzheimer's &*

Dementia **18**, 1038–1046 (2022).

64. Walker, K. A. *et al.* The role of peripheral inflammatory insults in Alzheimer's disease: a

review and research roadmap. *Molecular Neurodegeneration* **18**, 37 (2023).

65. Jorfi, M., Maaser-Hecker, A. & Tanzi, R. E. The neuroimmune axis of Alzheimer's

disease. *Genome Med* **15**, 6 (2023).

66. Holmes, C. *et al.* Systemic inflammation and disease progression in Alzheimer disease.

Neurology **73**, 768–774 (2009).

67. Morgan, D. G. & Mielke, M. M. Knowledge gaps in Alzheimer's disease immune

biomarker research. *Alzheimer's & Dementia* **17**, 2030–2042 (2021).

68. Kitazawa, M., Oddo, S., Yamasaki, T. R., Green, K. N. & LaFerla, F. M.

Lipopolysaccharide-Induced Inflammation Exacerbates Tau Pathology by a Cyclin-

Dependent Kinase 5-Mediated Pathway in a Transgenic Model of Alzheimer's Disease.

J. Neurosci. **25**, 8843–8853 (2005).

69. Walker, K. A. *et al.* Midlife systemic inflammatory markers are associated with late-life

brain volume: The ARIC study. *Neurology* **89**, 2262 (2017).

70. Jacob, F. *et al.* Human Pluripotent Stem Cell-Derived Neural Cells and Brain Organoids

Reveal SARS-CoV-2 Neurotropism Predominates in Choroid Plexus Epithelium. *Cell*

Stem Cell **27**, 937-950.e9 (2020).

71. Phetsouphanh, C. *et al.* Immunological dysfunction persists for 8 months following initial

mild-to-moderate SARS-CoV-2 infection. *Nat Immunol* **23**, 210–216 (2022).

72. Hur, J.-Y. *et al.* The innate immunity protein IFITM3 modulates γ-secretase in

Alzheimer's disease. *Nature* **586**, 735–740 (2020).

73. Gholami, M. *et al.* Interferon-Induced Transmembrane Protein 3 rs34481144 C/T

Genotype and Clinical Parameters Related to Progression of COVID-19. *J Immunol Res*

2023, 2345062 (2023).

74. Ashton, N. J. *et al.* Diagnostic Accuracy of a Plasma Phosphorylated Tau 217

Immunoassay for Alzheimer Disease Pathology. *JAMA Neurol* **81**, 255–263 (2024).

75. Janelidze, S. *et al.* Plasma β-amyloid in Alzheimer's disease and vascular disease. *Sci*

Rep **6**, 26801 (2016).

76. Xie, Y., Xu, E., Bowe, B. & Al-Aly, Z. Long-term cardiovascular outcomes of COVID-19.

Nat Med **28**, 583–590 (2022).

77. Wan, E. Y. F. *et al.* Association of COVID-19 with short- and long-term risk of

cardiovascular disease and mortality: a prospective cohort in UK Biobank.

Cardiovascular Research **119**, 1718–1727 (2023).

78. Marín, J. S. *et al.* Increased incidence of rheumatoid arthritis after COVID-19.

Autoimmunity Reviews **22**, 103409 (2023).

79. Rathmann, W., Kuss, O. & Kostev, K. Incidence of newly diagnosed diabetes after

Covid-19. *Diabetologia* **65**, 949–954 (2022).

80. Argentieri, M. A. *et al.* Proteomic aging clock predicts mortality and risk of common age-

related diseases in diverse populations. 2023.09.13.23295486 Preprint at

https://doi.org/10.1101/2023.09.13.23295486 (2023).

81. Livingston, G. *et al.* Dementia prevention, intervention, and care: 2020 report of the

Lancet Commission. *The Lancet* **396**, 413–446 (2020).

82. Wang, Y.-R. *et al.* Associations Between Hepatic Functions and Plasma Amyloid-Beta

Levels-Implications for the Capacity of Liver in Peripheral Amyloid-Beta Clearance. *Mol*

Neurobiol **54**, 2338–2344 (2017).

83. Lyu, H. *et al.* Plasma amyloid-beta levels correlated with impaired hepatic functions: An

adjuvant biomarker for the diagnosis of biliary atresia. *Front. Surg.* **9**, (2022).

84. Cousins, K. A. Q. *et al.* Elevated Plasma Phosphorylated Tau 181 in Amyotrophic Lateral

Sclerosis. *Annals of Neurology* **92**, 807–818 (2022).

85. Vacchiano, V. *et al.* Elevated plasma p-tau181 levels unrelated to Alzheimer's disease

pathology in amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* **94**, 428–435

(2023).

86. Schoeler, T. *et al.* Participation bias in the UK Biobank distorts genetic associations and

downstream analyses. *Nat Hum Behav* **7**, 1216–1227 (2023).

87. Janelidze, S. *et al.* Head-to-head comparison of 10 plasma phospho-tau assays in

prodromal Alzheimer's disease. *Brain* **146**, 1592–1601 (2022).

88. Ashton, N. J. *et al.* Plasma and CSF biomarkers in a memory clinic: Head-to-head

comparison of phosphorylated tau immunoassays. *Alzheimer's & Dementia* **19**, 1913–

1924 (2023).

89. Wang, L. *et al.* Association of COVID-19 with New-Onset Alzheimer's Disease. *J

Alzheimers Dis* **89**, 411–414 (2022).

90. Littlejohns, T. J. *et al.* The UK Biobank imaging enhancement of 100,000 participants:

rationale, data collection, management and future directions. *Nature Communications*

11, 2624 (2020).

91. Stevenson-Hoare, J. *et al.* Plasma biomarkers and genetics in the diagnosis and

prediction of Alzheimer's disease. *Brain* **146**, 690–699 (2023).

92. Inker Lesley A. *et al.* Estimating Glomerular Filtration Rate from Serum Creatinine and

Cystatin C. *New England Journal of Medicine* **367**, 20–29 (2012).

93. Bešević, J. *et al.* Persistence of SARS-CoV-2 antibodies over 18 months following

infection: UK Biobank COVID-19 Serology Study. *J Epidemiol Community Health* (2023)

doi:10.1136/jech-2023-220569.

94. Second-generation PLINK: rising to the challenge of larger and richer datasets |

GigaScience | Oxford Academic.

https://academic.oup.com/gigascience/article/4/1/s13742-015-0047-

8/2707533?login=false.

95. Miller, K. L. *et al.* Multimodal population brain imaging in the UK Biobank prospective

epidemiological study. *Nature Neuroscience* **19**, 1523–1536 (2016).

96. Alfaro-Almagro, F. *et al.* Image processing and Quality Control for the first 10,000 brain

imaging datasets from UK Biobank. *NeuroImage* **166**, 400–424 (2018).

97. Makaremi, S. *et al.* The role of IL-1 family of cytokines and receptors in pathogenesis of

COVID-19. *Inflamm. Res.* **71**, 923–947 (2022).

98. Dursun, E. *et al.* The interleukin 1 alpha, interleukin 1 beta, interleukin 6 and alpha-2-

macroglobulin serum levels in patients with early or late onset Alzheimer's disease, mild

cognitive impairment or Parkinson's disease. *Journal of Neuroimmunology* **283**, 50–57

(2015).

99. Yang, H.-S. *et al.* Plasma IL-12/IFN-γ axis predicts cognitive trajectories in cognitively

unimpaired older adults. *Alzheimers Dement* **18**, 645–653 (2022).

100.   Sutinen, E. M., Pirttilä, T., Anderson, G., Salminen, A. & Ojala, J. O. Pro-

inflammatory interleukin-18 increases Alzheimer's disease-associated amyloid-β

production in human neuron-like cells. *Journal of Neuroinflammation* **9**, 199 (2012).

101.   Brosseron, F. *et al.* Soluble TAM receptors sAXL and sTyro3 predict structural and

functional protection in Alzheimer's disease. *Neuron* **110**, 1009-1022.e4 (2022).

102.   Shin, J. J. *et al.* MIF is a common genetic determinant of COVID-19 symptomatic

infection and severity. *QJM* **116**, 205–212 (2023).

103.   Nasiri, E. *et al.* Key role of MIF-related neuroinflammation in neurodegeneration and

cognitive impairment in Alzheimer's disease. *Mol Med* **26**, 34 (2020).

1083
1084

1085

Supplementary Information

Tables not shown can be found in the accompanying Excel spreadsheet.

CLASS	COVARIATE	UKB DATA-FIELD ID(S)	CASE/CONTROL COMPARISON	BASIC MODEL	AD RISK FACTORS MODELS
DEMOGRAPHICS	*Age*	34,52,53	X	X	X
	Sex	31	X	X	X
	Height (cm)	50	X		
	Weight (kg)	21002	X		X
	Ethnicity (White)	21000	X		X
ASSESSMENT INFORMATION	*Date of assessment/ interval*	53	X	X	X
GENETICS	*APOE status*	Genotyping	X		X
HEALTH MEASURES	*Body Mass Index*	21001	X		X
	Hip/Waist Ratio	48,49	X		X
	Blood Pressure (Diastolic, Systolic)	4079,4080	X		X
	Hand grip strength	46,47	X		
	Chest Wheeze	2316	X		
	Est. Glomerular Filtration Rate	OLINK: P_CST3	X		X
	General Health	2178	X		X
LIFE STATUS	*Alcohol Intake Freq*	1558	X		X
	Smoking status	1239	X		X
	Years of education	845	X		X
	Deprivation	26410	X		
	Income	738	X		
	Household Size	709	X		X
	Employment status	6142	X		
	Key Worker Status	28063	X		
	Activity levels (min/day)	894,914	X		X
	Isolation (social visits)	1031,2110	X		X
COMORBIDITIES	*Type II Diabetes*	26206	X		X
	Heart Condition	20002	X		X
	Obesity	21001	X		X
	Hypertension	20002	X		X
	Depression	20002	X		X
	Irritable Bowel Syndrome	20002	X		X
	COPD	20002	X		X
	Emphysema	20002	X		X
	Renal conditions	20002	X		X
MEDICATION	*Blood pressure medication*	6153	X		
	Cholesterol medication	6153	X		
	Diabetes medication	6153	X		
	COVID vaccination	Health Records	X		
COGNITION	*General Cognitive Ability score*	Cognitive test fields	X		X
NEUROIMAGING	*AD imaging phenotype*	FreeSurfer IDPs	X		X

1093
1094
1095

Supplementary table 1. UK Biobank Covariates used in analyses. BMI: Body Mass Index; AD: Alzheimer's Disease, PRS: polygenic risk score. COPD - Chronic obstructive pulmonary disease

1096

Supplementary table 2 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session. Table shows model fits and p-values associated with case / control status (case=+1), age, sex
(male=+1), the time interval between sessions (days), and pre-pandemic protein level. Beta estimates have been
standardised. NfL – neurofilament light; GFAP (**) indicates significant, FDR corrected by column, alpha=0.05.

Supplementary table 3 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session with an extended set of confound variables. Beta estimates have been standardised. NfL – neurofilament
light; GFAP (**) indicates significant, FDR corrected by column, alpha=0.05.

Supplementary table 4 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session with additional terms modelling age-related vulnerability and its interaction with case/control status. Beta
estimates have been standardised. NfL – neurofilament light; GFAP (**) indicates significant, FDR corrected by
column, alpha=0.05.

Supplementary table 5 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session with additional terms modelling age-related vulnerability and its interaction with case/control status, and
an extended set of confound variables. Beta estimates have been standardised. NfL – neurofilament light; GFAP
(**) indicates significant, FDR corrected by column, alpha=0.05.

Supplementary table 6 Associations of protein biomarkers with UK Biobank variables at pre-pandemic
assessment visit. Associations were determined from linear models including age and sex covariates. (*) p<0.05
uncorrected (**) p<0.05 FDR correction (alpha = 0.05).

Supplementary table 7 Baseline protein level statistics for cases and controls. P-values reflect paired t-tests.

Supplementary table 8 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session with additional terms modelling covariates APOE variant status, smoking status, Hip/Waist ratio and
Diabetes. Beta estimates have been standardised. NfL – neurofilament light; GFAP (**) indicates significant, FDR
corrected by column, alpha=0.05.

Supplementary table 9 Associations of UK Biobank variables (measured at pre-pandemic assessment) with the
change in protein biomarkers across assessments (post-pandemic – pre-pandemic). Associations were
determined from linear models of protein level change including terms for age, sex, and interval between
assessments. (*) p<0.05 uncorrected (**) p<0.05 FDR correction (alpha = 0.05).

Supplementary table 10 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session with additional terms modelling covariates APOE variant status, smoking status, Hip/Waist ratio and
Diabetes. Beta estimates have been standardised. NfL – neurofilament light; GFAP (**) indicates significant, FDR
corrected by column, alpha=0.05.

Supplementary table 11 Parameter fits for linear model predicting proteomic levels in pandemic assessment
session with additional term modelling potential confound Glomerular Filtration Rate (GFR). GFR was available in
around half of participants. Beta estimates have been standardised. NfL – neurofilament light; GFAP (**)
indicates significant, FDR corrected by column, alpha=0.05.

Supplementary table 12 Parameter fits for interaction term with case-control status (case=+1) and associations
between UK Biobank variables at baseline with protein biomarker change. These terms identify UK Biobank
variables associated with the protein biomarkers in SARS-CoV-2 exposure specific manner. (*) p<0.05
uncorrected (**) p<0.05 FDR correction (alpha = 0.05).

Supplementary table 13 AD protein level change model fits for models including terms modelling AD
neuroimaging phenotype at baseline. Associations were determined from linear models including age and sex
covariates. (*) p<0.05 uncorrected (**) p<0.05 FDR correction (alpha = 0.05).

1139

1140

OLINK PROTEIN	COVID	AD
TUMOUR NECROSIS FACTOR ALPHA (TNF-α)	X[47]	X[64]
TNF SUPERFAMILY MEMBER 10 (TRAIL)	X[47]	X[42]
INTERLEUKIN(IL)-1α	X[97]	X[98]
IL-6	X[47,71]	X[98]
IL-12		X[99]

IL-18	X[97]	X[100]
IFN-B	X[71]	X[42,98]
INTERFERON GAMMA (IFNG)	X[71]	X[100]
COMPLEMENT C1QA		X[101]
COMPLEMENT C4BPB		X[101]
MACROPHAGE INHIBITORY FACTOR (MIF)	X[102]	X[101,103]
CHITINASE-3 LIKE-PROTEIN-1 (CHI3L1) - YKL-40		X[42]
PENTRAXIN 3 (PTX3)	X[71]	
CALCITONIN (CALCA)	X[47]	

1141

Supplementary table 14 Inflammatory proteins available from Olink panel identified as associated with COVID
and/or Alzheimer's Disease. X indicates reports of an association with the disease.

Supplementary table 15 Baseline Case-Control differences in Olink Inflammatory Proteins. P-values reflect paired
t-tests.

Supplementary table 16 Associations of Olink Inflammatory Protein levels (measured at baseline) with the
change in AD protein biomarkers across assessments (across cases and controls). Associations were
determined from linear models including terms for age, sex, and interval between assessments. (*) p<0.05
uncorrected (**) p<0.05 FDR correction (alpha = 0.05).

Supplementary table 17 Associations of an interaction term between Case/Control status and Olink Inflammatory
Protein levels (measured at baseline) with protein biomarker change. These identify Olink Inflammatory Proteins
associated with the AD protein biomarkers in a SARS-CoV-2 exposure specific manner. Associations were
determined from linear models including terms for age, sex, and interval between assessments. (*) p<0.05
uncorrected (**) p<0.05 FDR correction (alpha = 0.05).

Supplementary table 18 Parameter fits for linear model predicting Olink Inflammatory Protein levels in pandemic
assessment session. Beta estimates have been standardised. (**) indicates significant, FDR corrected by
column, alpha=0.05.

Supplementary table 19 SARS-CoV-2 related increases in Olink ProteinScore disease risk scores. Plots show
SARS-CoV-2 effect weighting parameter in model of change of scores between assessment visits. Due to the
ProteinScore disease risk training data, results are from participants <73 years. (*) reflects FDR corrected
significance, alpha=0.05.

Supplementary table 20 Correlations between Simoa Ultrasensitive neurology assays. Diagonal reflects
correlations of protein levels between pre-pandemic and pandemic assessment visits.

1164

1165