Detection of novel influenza viruses through community and healthcare testing: Implications
for surveillance efforts in the United States

Sinead E. Morris[1,2,*], Matthew Gilmer[1,2], Ryan Threlkel[1], Lynnette Brammer[1], Alicia P. Budd[1], A. Danielle
Iuliano[1], Carrie Reed[1], Matthew Biggerstaff[1]

[1]*Influenza Division, Centers for Disease Control and Prevention, Atlanta, GA*

[2]*Goldbelt Professional Services, Chesapeake, VA*

*Corresponding author: run7@cdc.gov

Short title: Healthcare testing of novel influenza viruses

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily
represent the views of the Centers for Disease Control and Prevention.

Data Availability Statement: All model inputs and code needed to perform the analysis are available at
https://github.com/CDCgov/novel-flu-detection. Data were used solely to inform model inputs and were
the result of secondary analyses; the original sources are cited in the text.

Conflict of Interest Statement: The authors have no conflicts of interest or funding sources to declare.

Abstract

Background Novel influenza viruses pose a potential pandemic risk and rapid detection of infections in
humans is critical to characterizing the virus and facilitating the implementation of public health
response measures.

Methods We use a probabilistic framework to estimate the likelihood that novel influenza virus cases
would be detected through testing in different community and healthcare settings (urgent care,
emergency department, hospital, and intensive care unit (ICU)) while at low frequencies in the United
States. Parameters were informed by data on seasonal influenza virus activity and existing testing
practices.

Results In a baseline scenario reflecting the presence of 100 novel virus infections with similar severity
to seasonal influenza viruses, the median probability of detecting at least one infection per month was
highest in urgent care settings (72%) and when community testing was conducted at random among the
general population (77%). However, urgent care testing was over 15 times more efficient (estimated as
the number of cases detected per 100,000 tests) due to the larger number of tests required for
community testing. In scenarios that assumed increased clinical severity of novel virus infection, median
detection probabilities increased across all healthcare settings, particularly in hospitals and ICUs (up to
100%) where testing also became more efficient.

Conclusions Our results suggest that novel influenza virus circulation is likely to be detected through
existing healthcare surveillance, with the most efficient testing setting impacted by the disease severity
profile. These analyses can help inform future testing strategies to maximize the likelihood of novel
influenza detection.

Keywords: influenza; H5N1; novel virus; detection; healthcare testing

Introduction

Novel influenza viruses are different from the seasonal influenza viruses currently circulating in humans
(A/H3N2, A/H1N1, and B/Victoria). Human infections with novel influenza viruses are generally rare and
isolated events that occur through exposure to infected animals (such as livestock) during recreational
or occupational activities. At the time of writing (10 May 2024), widespread avian influenza A(H5N1)
virus outbreaks occurring among wild and commercial birds since January 2022 have been associated
with just two detected human cases of H5N1 in the United States: one individual who was exposed to
infected poultry and one who was exposed to infected dairy cattle [1, 2]. The H5N1 viruses associated
with these outbreaks do not easily bind to receptors in the human upper respiratory tract and the risk to
the general public is currently low [1]. However, a novel influenza virus that transmits efficiently
between humans could pose a pandemic risk. Rapid detection of human infection with a novel influenza
virus is critical to characterizing the virus causing the infection and facilitating a rapid public health
response [3].

Testing is particularly important to distinguish novel influenza virus infection from seasonal influenza or
other respiratory virus infections with similar symptom profiles [4]. Although active monitoring and
testing of individuals with exposure to infected animals can identify new spillover infections [2], such

measures are not designed to detect cases in the wider community following sustained human-to-
human transmission. Public health surveillance systems must be equipped to detect novel influenza
cases through testing in the community or in healthcare settings where infected individuals might seek
care.

We use a probabilistic framework to estimate the likelihood of detection of novel influenza virus cases
once sustained human-to-human transmission is occurring at low frequencies within the United States
(i.e., 1,000 total cases or less). We consider testing of individuals presenting to different healthcare
settings with no known previous exposure to infected animals or humans and use information on testing
for seasonal influenza viruses to develop assumptions about plausible testing probabilities. Our findings
can help inform testing strategies to improve detection of novel influenza virus cases occurring at low
frequencies.

Methods

Model

We adapted an existing framework to estimate detection probabilities for a novel influenza virus in the
United States [5]. For a given case of novel influenza virus infection, the probability of detection in a
particular healthcare setting can be expressed as

$p_{detect} = P(test \,|case) \times p_d \times t_{sn} \times p_f,$

where $P(test \,|case)$ is the probability that someone is tested in that setting given that they are a case;
p_d is the probability that testing occurs while virus is still detectable; t_{sn} is the test sensitivity; and p_f is
the probability a positive test is forwarded to a public health laboratory for further testing. Most
commercial assays currently used for human influenza virus testing cannot distinguish novel influenza A
viruses from seasonal influenza A viruses. Thus, further testing at a public health laboratory is required
for a positive specimen to be identified as a novel virus (until tests specific for that virus become more
widely available). We initially assumed 50% of positive specimens are forwarded (i.e., p_f = 50%). This
was informed by the average percentage of influenza A hospitalizations that were subtyped between
2010–2019 [6]. However, we considered a range of forwarding levels (25, 50%, 75% and 100%) in
sensitivity analyses. All specimens forwarded for further testing were assumed to be correctly identified
as a novel influenza virus.

The per case probability of being tested is the combined probability that a case will develop symptoms
(p_{symp}), seek care for those symptoms in a particular healthcare setting (p_{seek}), and be tested in that
setting (p_{test}), i.e.,

$P(test \,|case) = p_{symp} \times p_{seek} \times p_{test}.$

For a certain incidence of novel cases each month, *I*, in a population of size *N* (where *I* is the fraction of
the population infected with the novel influenza virus), we estimate the probability of detecting at least
one novel case as 1 - the probability of detecting no cases among the entire population, or

$1 - (1 - I \times p_{detect})^N$

(see the supporting information for further details). The expected number of clinical tests used per
month, $E(T)$, is the combined number of tests conducted among cases and non-cases. Non-cases
represent individuals presenting at healthcare settings with respiratory illness symptoms that are not
due to novel influenza virus infection. The expected number of tests can be expressed as

$E(T) = I \times N \times P(test\,|case) + (1 - I) \times N \times P(test\,|not\,case),$

where $P(test\,|not\,case)$ is the probability that someone without novel influenza virus infection is
tested. The latter quantity is estimated as the background rate of presentation with respiratory illness
symptoms to a given healthcare setting among the general population (b_{seek}) multiplied by the
probability of being tested in that setting (p_{test}). To compare testing efficiency in different settings we
estimated the expected number of detected cases per 100,000 clinical tests conducted as
$(I \times N \times p_{detect}\,/\,E(T)) \times 100,000.$

Finally, we considered random testing in the general community as a supplemental strategy that could
be deployed in addition to healthcare testing. Given that community testing does not depend on
symptom presentation or care-seeking behavior, $P(test\,|case)$ was simply the frequency of community
tests conducted per month and the expected number of tests was $N \times P(test\,|case)$. Similarly, p_d was
the approximate time (in months) that virus would remain detectable and was parameterized to capture
individual variation in virus shedding dynamics. Since community testing would be initiated to seek out
novel influenza virus infection, we did not adjust for specimen forwarding (i.e., we assumed all
specimens would be tested to distinguish novel influenza virus from seasonal influenza viruses).

For each healthcare and community setting, we drew 10,000 parameter combinations from data-
informed distributions (outlined below) and calculated the quantities described above. All analyses and
visualizations were generated in R version 4.0.3 using the `data.table`, `truncnorm`, `here`,
`scales`, `patchwork`, `colorspace` and `tidyverse` packages [7-14].

Healthcare settings and model parameterization

We considered three distinct healthcare settings to reflect different care-seeking behaviors and testing
practices: (i) outpatient urgent care and emergency departments (UC/ED); (ii) inpatient hospital settings;
and (iii) intensive care units (ICU). Each setting was assumed independent such that a person presenting
to both (for example, a hospital admission followed by a subsequent ICU admission) could be tested in
both, according to the corresponding testing probabilities. Data were collated from various existing
influenza surveillance platforms to inform parameters for each setting (Table 1). We defined N = 330
million to approximate the U.S. population [15] and considered incidence values that corresponded to
100 and 1,000 total novel influenza cases.

Table 1. Baseline care-seeking and testing parameters. Surveillance platforms are Flu Near You (FNY), Outbreaks
Near Me (ONM), VISION Vaccine Effectiveness Network, FluSurv-Net, and IBM MarketScan® Commercial Claims
and Encounters Database (MarketScan) [16-19]. Further details of each platform are provided in the supporting
information.

Parameter	Assumed distribution	Source and available timeframe (if applicable)
Proportion of novel cases developing symptoms, p_{symp}	Uniform with range: 40–80%	[20]

Care-seeking and presentation of novel symptomatic cases at specific sites, p_{seek}: 　UC / ED 　Hospital 　ICU	Uniform with range: 10–20% of symptomatic cases 1–2% of symptomatic cases 15–20% of hospitalizations	 FNY, ONM: 2018–2023 CDC burden estimates: 2010-2021 [21] VISION: 2020-2021; FluSurv-Net: 2022–2023
Testing of individuals with ARI, p_{test}: 　UC / ED 　Hospital 　ICU	Truncated normal with mean / SD / range: 50% / 10% / 10–90% 53% / 10% / 20–95% 46% / 10% / 1–95%	VISION: December 2021–May 2022
Community testing as a proportion of the general population, regardless of symptoms	3–6% of general population per month	Assumption following [22]
Tests that occur while virus is detectable, p_d 　Healthcare settings 　Community settings	Uniform with range: 50–85% 25–50%	 Proportion seeking care ≤7 days after symptom onset [23] Proportion of month that virus is detectable [24]
Test sensitivity, t_{sn}	Uniform with range 80–100%	[25]
Proportion of positive specimens that are forwarded to a public health laboratory, p_f	50%	Assumption following [6]
Background occurrence in the general population, b_{seek}, of: 　ILI* 　Hospital ARI admissions 　ICU ARI admissions	Uniform with range: 0.6–6% 0.03–0.1% 0.02–0.03%	 FNY, ONM: 2019, 2022 MarketScan: 2015–2021 MarketScan: 2015–2021

Abbreviations: UC = urgent care; ED = emergency department; ICU = intensive care unit; ILI = influenza-like-illness; ARI = acute
respiratory illness; SD = standard deviation.
*Background ILI occurrence is multiplied by care-seeking probabilities in urgent care or emergency departments (p_{seek}) to
estimate the rate of presentation to urgent care or emergency departments with influenza symptoms in the general
population. We omitted data from 2020 and 2021 due to atypically low levels of respiratory virus circulation.

Our baseline scenario reflected a novel influenza virus with similar severity to seasonal influenza.
However, we also considered increased severity scenarios that ranged from severity that was similar to
COVID-19, to the severity of recent H5N1 virus infections in humans (Table 2). For these scenarios, we
assumed similar or increased probabilities of developing symptoms and seeking care in each healthcare
setting, while ensuring that the combined percentages did not exceed 100%. We initially assumed
testing probabilities were fixed (Table 1) but explored alternative scenarios with increased testing (p_{test}
mean = 90%) to compare the effect of enhanced surveillance across healthcare settings.

Table 2. Scenarios for increased symptom severity. All parameters are assumed to follow a Uniform distribution
with the reported range.

Scenario	Symptomatic	UC / ED[†]	Hospital	ICU[§]	Source(s)
Baseline	40–80%	10–20%	1–2% CHR[†]	15–20%	[20, 21]
COVID-like	40–80%	10–20%	1–2% IHR[‡]	20–30%	[22, 26]
Intermediate 1	25% > baseline*	25% > baseline*	4.5–5.5% IHR[‡]	30–40%	[22]

Intermediate 2	50% > baseline[*]	50% > baseline[*]	9.5–10.5% IHR[‡]	45–55%	[22]
Recent H5-like	50% > baseline[*]	50% > baseline[*]	60–70% IHR[‡]	75–85%	[1]

Abbreviations: UC = urgent care; ED = emergency department; ICU = intensive care unit; CHR = case-hospitalization ratio; IHR = infection-hospitalization ratio.
[*]Up to a maximum of 100%.
[†]Expressed as a percentage of symptomatic individuals.
[‡]Expressed as a percentage of infected individuals.
[§]Expressed as a percentage of hospitalizations.

We also considered scenarios in which testing practices changed according to seasonal influenza activity. For example, clinicians may be less likely to test for influenza viruses during summer months when background respiratory virus activity is low. To explore this, we first defined distinct probability distributions for the background rates of presentation to each healthcare setting, b_{seek}, during peak (November – February) and off-peak (May – August) time periods (Table 3). We then simulated the model for each time period and healthcare setting, assuming the care-seeking behavior of novel influenza cases did not change but that testing in off-peak periods was either equal to, or 50% of, testing in peak periods.

Table 3. Baseline occurrence of ILI or ARI symptoms partitioned by peak vs off-peak activity.

Parameter	Range of uniform distribution	Period	Source and available timeframe
Occurrence of: ILI[*]	1.0–6.0% 0.6–2.5%	Peak Off-peak	FNY, ONM: 2019, 2022
Hospital ARI admission	0.04–0.10% of general population 0.03–0.09% of general population	Peak Off-peak	MarketScan: 2015–2021
ICU ARI admission	0.014–0.035% of general population 0.010–0.030% of general population	Peak Off-peak	MarketScan: 2015–2021

Abbreviations: ILI = influenza-like-illness; ARI = acute respiratory illness; ICU = intensive care unit; FNY = Flu Near You; ONM = Outbreaks Near Me.
[*]Background ILI occurrence is multiplied by care-seeking probabilities in urgent care or emergency departments (p_{seek}) to estimate the rate of presentation to urgent care or emergency departments in the general population. We omitted data from 2020 and 2021 due to atypically low levels of respiratory virus circulation.

Results

We first simulated the model with baseline severity assumptions and no distinction between peak and off-peak time periods. At the lowest incidence (100 novel cases in the population), the median probability of detecting at least one case was highest in community and UC/ED settings, at 77% (95th percentile: 56–91%) and 72% (44–92%), respectively (Figure 1C; Baseline scenario). In comparison, median detection probabilities in hospital and ICU settings were less than 15%. The probability of detection increased across all settings when there were 1,000 assumed novel cases in the population, to 100% (100–100%) in UC/EDs and the community, 74% (47–94%) in hospitals, and 19% (9–35%) in ICUs. Testing in UC/ED settings was always most efficient and detected more cases per 100,000 tests than other settings (Figure 1E). Notably, community testing was least efficient due to the greater number of tests required (more than 10 million per month; Figure 1D), and no setting detected more than 3% of all

novel cases under our assumptions (Figure 1F). Increasing the percentage of influenza positive
specimens forwarded to public health laboratories to 75% or 100% increased detection probabilities and
test efficiency across all healthcare settings (Figure S1). For example, the median detection probability in
UC/EDs increased to 85% (58–98%) and 92% (69–99%) at the lowest incidence, respectively. Conversely,
a decrease in the percentage forwarded to 25% decreased detection probabilities and test efficiencies,
although the relative ordering of setting efficiency was preserved. Thus, for a novel influenza virus with
similar severity to seasonal influenza, UC/ED settings are likely to provide greatest opportunities for case
detection.

200
Figure 1. Probabilities of detection and test usage under different severity scenarios. (A) Assumed probabilities of
presentation to a particular setting, calculated as $p_{symp} \times p_{seek}$ for UC/ED, hospital, and ICU settings. All cases
are assumed to be in the community, resulting in a probability of one for that setting (not shown). (B) Assumed
proportion of individuals with ILI or ARI tested in UC/ED, hospital and ICU settings, or proportion of all individuals
tested in the community. (C) Estimated probability of detecting at least one novel case per month. Panels indicate

different assumed levels of incidence (100 and 1,000 novel cases). (D) Expected number of clinical tests used per
month. (E) Estimated test efficiency, calculated as the number of detected novel cases per 100,000 tests. (F)
Percent of all novel cases detected per month. In all panels, points represent median values across 10,000
simulations, inner shaded bands show 50th percentiles, and outer shaded bands show 95th percentiles.
Abbreviations: UC = urgent care; ED = emergency department; H = hospital; ICU = intensive care unit; Int 1 =
Intermediate 1; Int 2 = Intermediate 2; ILI = influenza-like illness; ARI = acute respiratory illness.

Given uncertainty in the potential severity of a novel influenza virus, we explored additional scenarios in
which cases were more likely to develop symptoms and/or present to a particular healthcare setting
than the baseline severity scenario (Table 2; Figure 1A). As severity increased, the probability of
detection also increased across all healthcare settings due to the greater probability of requiring medical
attention (Figure 1C). The difference between detection probabilities in UC/ED compared with hospital
and ICU settings also decreased as cases were more likely to be severe and require admission to the
latter. For example, median detection probabilities for ICU settings increased from 2% (1–4%) and 19%
(9–35%) at baseline with 100 and 1,000 novel cases, respectively, to 100% (98–100%) and 100% (100–
100%) in the "Recent-H5" scenario. There were also substantial increases in testing efficiency in hospital
and ICU settings (Figure 1E) and increases in the percent of novel cases detected (for example, from a
maximum of 0.3% in hospital settings at baseline to 16% in the Recent H5 scenario; Figure 1F). Test
usage is driven primarily by background seasonal influenza virus testing and thus did not change across
severity scenarios (Figure 1D). Simulating an increase in clinical testing probabilities (p_{test} mean = 90%)
substantially increased detection probabilities and test usage for all healthcare settings but did not
impact the relative performance among settings (Figure S2).

Finally, we assessed how seasonal changes in background activity could impact probabilities of case
detection and testing efficiency. Assuming testing practices did not change seasonally led to equal
probabilities of detection in peak and off-peak periods, although testing efficiencies were increased in
off-peak periods due to the lower number of background tests conducted (Figure S3). Conversely,
assuming a 50% reduction in testing across all healthcare settings in off-peak periods (Figure 2B)
reduced the corresponding probabilities of detection (Figure 2C). However, for the most severe
scenarios (Intermediate 1, Intermediate 2, and Recent H5) there was always at least one healthcare
setting with a median detection probability greater than 60% in off-peak periods at the lowest
incidence.

Discussion

We modeled the likelihood of detection of novel influenza virus cases occurring at low incidence in the
United States. We adapted a simple probabilistic framework that accounted for symptom severity, care-
seeking behavior, and testing practices in different healthcare settings, and used care-seeking and
testing information from recent influenza seasons to inform model parameters. We found that the most
efficient setting for detection depends on the severity profile of the novel influenza virus. Although the
percent of total novel influenza cases detected was relatively low, the probabilities of detecting at least
one case, and thus identifying novel influenza virus circulation, were high in at least one setting across a
range of different testing, severity, and specimen forwarding assumptions.

Figure 2. Probabilities of detection and test usage in healthcare settings assuming reduced testing probabilities
during periods of off-peak seasonal activity. Incidence is fixed at 100 novel cases in the population. (A) Assumed
probabilities of presentation to a particular setting, calculated as $p_{symp} \times p_{seek}$. Ranges are constant in peak and
off-peak periods. (B) Assumed proportion of individuals with ILI or ARI tested in peak and off-peak periods. Ranges
are constant across severity scenarios. (C) Estimated probability of detecting at least one novel case per month. (D)
Expected number of clinical tests used per month. (E) Estimated test efficiency, calculated as the number of
detected novel cases per 100,000 tests. (F) Percent of all novel cases detected per month. In all panels, points
represent median values across 10,000 simulations, inner shaded bands are the 50[th] percentiles, and outer shaded
bands are the 95th percentiles. Abbreviations: UC = urgent care; ED = emergency department; H = hospital; ICU =
intensive care unit; Int 1 = Intermediate 1; Int 2 = Intermediate 2; ILI = influenza-like illness; ARI = acute respiratory
illness.

The high probabilities of detecting at least one case that we have estimated here are relevant for public
health pandemic preparedness. The detection of one case would facilitate the implementation of public
health actions including increased testing strategies, further virus characterization, vaccine development

(if warranted), the implementation of appropriate public health control measures, and updated
recommendations for the use of influenza antiviral medications. One key parameter influencing the
detection probability was the probability of testing in each healthcare setting. We found that detection
probabilities could decrease if influenza testing is substantially reduced below in-season values (for
example, during off-peak months). However, it is also possible that clusters of cases and outbreaks could
be more likely to be detected and tested during off-peak months if clinicians remain vigilant for evidence
of atypical respiratory virus signs or symptoms. The detection probability was also influenced by
assumptions about the forwarding of clinical specimens. Our baseline value of 50% was informed by
subtyping information from hospitalized influenza infections between 2010–2019 [6]. However, we
included a lower bound of 25% to reflect recent post-COVID-19 pandemic trends and potentially
reduced forwarding in UC/ED outpatient settings [18]. We also included higher values up to 100% to
explore maximum attainable detection probabilities if all tests were forwarded and found a substantial
improvement in our estimates. Therefore, during the current H5N1 situation, it is critical that clinicians
maintain high testing frequencies and forward influenza A positive specimens to public health
laboratories for further testing when recommended. Finally, given the severity of prior H5N1 cases (for
example, there has been a 50% case-fatality proportion in cases identified since 1997 [1]), additional
strategies to increase testing in ICU settings may help increase the likelihood of detection and testing
efficiency, particularly during summer months when background acute respiratory illness rates are low.

Although the probability of detecting one case was generally high, the percent of total cases detected
was low, especially in the lower severity scenarios. This finding assumes there are no immediate
changes to testing or healthcare seeking behavior once the first case is detected, and arises because
detection of influenza through clinical settings requires someone to become symptomatic, seek care, be
tested in a timely manner, and have a positive specimen forwarded for further characterization.
Although community testing removes these barriers to identification, it is resource intensive and would
need to occur even in the absence of perceived novel influenza virus spread to be effective, potentially
requiring over 100 million tests per year at the level modeled in this analysis. Similarly, at-home or self-
administered tests could alleviate issues associated with care-seeking and clinical testing practices.
However, such tests would need to be specific to the novel influenza virus and undergo potentially
lengthy development and authorization procedures before being available for widespread use.
Pandemic planning efforts should therefore include strategies to rapidly increase testing of acute
respiratory illness cases in clinical settings once human-to-human spread of a novel influenza virus has
been identified or is likely. Such strategies should account for the possibility that many cases may not be
detected, even with increased testing.

There are several caveats to our modeling framework. First, as our primary aim was to estimate
detection capabilities once sustained human-to-human transmission is occurring within the United
States, we did not consider surveillance for earlier events that might spark such transmission, such as
spillover from infected animals or introductions from outside the United States. It is possible that these
events would be associated with a greater probability of testing due to relevant exposure histories, and
thus have a greater likelihood of being detected compared with our estimates. Second, we did not
stratify detection probabilities by age. The severity of seasonal influenza can vary substantially among
different age groups [27], and age patterns of severity may differ for a novel influenza virus compared to
seasonal influenza viruses due to immunological imprinting and age-related exposures to previous
circulating viruses [28, 29]. Age may also impact testing probabilities and healthcare seeking behavior

[30, 31], although mean testing probabilities for children <18 years were similar to those of adults ≥18
years in the VISION data used to parameterize p_{test} (for example, 55 vs. 50% in UC/ED for children and
adults, respectively). Including age in the current framework would require additional assumptions
regarding the cross-reactivity of the novel influenza virus with seasonal influenza viruses to infer age-
specific severity distributions, and thus reduce the generalizability of our results. Third, we did not
explicitly incorporate delays in case admission to hospital or ICU that could reduce the window for viable
virus detection relative to other settings. These delays are likely on the order of several days and are
captured within our conservative range for the proportion of care-seekers who are tested while virus is
still detectable [32]. Fourth, we modeled the United States as a single population and did not explicitly
consider spatial or other heterogeneities in care-seeking and testing practices. If such data were
available, our analysis could be replicated at finer resolution to assess local response and detection
capabilities. Fifth, data were not available to fully inform our test forwarding assumptions. Although we
considered a range in sensitivity analyses, further information would increase the accuracy of our
detection probability estimates. We also assumed perfect sensitivity and specificity for all forwarded
tests in line with evaluation of real-time RT-PCR tests for novel H1N1 variant influenza viruses [33].
Although minor reductions in sensitivity should not substantially impact our detection probability
estimates, reductions in specificity could lead to false positive results that we have not considered.
However, the number of false positive results is likely to be small unless testing reaches extremely high
levels, such as considered here in the community setting.

Finally, our assumed inputs for baseline testing and background activity were informed by previous
influenza seasons and may not reflect future changes to these values. Where possible, we developed
parameter distributions based on data from multiple influenza seasons, before and after the COVID-19
pandemic, to account for broad fluctuations in care-seeking behavior, testing practices, and seasonal
influenza dynamics. We also explored scenarios with increased testing to capture the potential impacts
of changes to healthcare surveillance following additional policy recommendations. More generally, our
estimates of detection probabilities and test efficiency reflect the combined uncertainty in each
underlying parameter value and should thus be robust to small changes in any single parameter.

Novel influenza viruses pose a potential pandemic risk, and prompt detection is critical to characterizing
the virus causing the infection and facilitating a rapid public health response. Here we demonstrate how
a simple probabilistic framework can be used to estimate novel influenza virus detection probabilities
through testing in different community and healthcare settings, and can help inform the targeting of
future testing efforts. Our work was motivated by the 2022–2024 H5N1 situation in the United States
but could be applied more broadly to other locations and/or other potential novel influenza viruses.

Author Contributions: Sinead E. Morris: Conceptualization; Formal analysis; Investigation;
Methodology; Visualization; Writing original draft; Writing - review and editing. **Matthew Gilmer**: Data
curation; Writing - review and editing. **Ryan Threlkel**: Data curation; Writing - review and editing.
Lynnette Brammer: Conceptualization; Writing - review and editing. **Alicia P. Budd**: Conceptualization;
Writing - review and editing. **A. Danielle Iuliano**: Data curation; Writing - review and editing. **Carrie
Reed**: Conceptualization; Investigation; Supervision; Writing - review and editing. **Matthew Biggerstaff**:
Conceptualization; Investigation; Methodology; Supervision; Writing - review and editing.

References

1. CDC. *H5N1 Bird Flu: Current Situation Summary*. 2024 [cited 2024 May 10]; Available from: https://www.cdc.gov/flu/avianflu/avian-flu-summary.htm.

2. Kniss, K., et al., *Risk for Infection in Humans after Exposure to Birds Infected with Highly Pathogenic Avian Influenza A(H5N1) Virus, USA, 2022.* Emerging Infectious Diseases, 2023. **29**(6).

3. CDC. *Pandemic Influenza*. 2023 [cited 2024 May 10]; Available from: https://www.cdc.gov/flu/pandemic-resources/index.htm.

4. NASPHV. *Zoonotic Influenza: Detection, Response, Prevention, and Control Reference Guide*. 2022 [cited 2024 May 10]; Available from: http://www.nasphv.org/documentsCompendiaZoonoticInfluenza.html.

5. Russell, S., et al., *Detecting Local Zika Virus Transmission in the Continental United States: A Comparison of Surveillance Strategies.* PLoS Curr, 2017. **9**.

6. Sumner, K., et al., *Severity of influenza-associated hospitalisations by influenza virus type and subtype in the USA, 2010-19: a repeated cross-sectional study*. Lancet Microbe, 2023.

7. Wickham, H., et al., *Welcome to the tidyverse.* Journal of Open Software, 2019. **4**(43): p. 1686.

8. Pedersen, T.L., *patchwork: The Composer of Plots.* R package version 1.1.1, 2020.

9. R Core Team, *R: A Language and Environment for Statistical Computing*. 2020: R Foundation for Statistical Computing.

10. Dowle, M. and A. Srinivasan *data.table: Extension of `data.frame`*. R package version 1.14.2, 2021.

11. A., Z., et al., *colorspace: A Toolbox for Manipulating and Assessing Colors and Palettes.* Journal of Statistical Software, 2020. **96**(1): p. 1-49.

12. Mersmann, O., et al. *truncnorm: Truncated Normal Distribution*. R package version 1.0.9, 2023.

13. Müller, K., *here: A Simpler Way to Find Your Files.* R package version 1.0.1, 2020.

14. Wickham, H. and D.P. Seidel *scales: Scale Functions for Visualization*. R package version 1.2.1, 2022.

15. United States Census Bureau. *Census Data*. 2023 [cited 2023 May 31]; Available from: https://data.census.gov/.

16. Outbreaks Near Me. *COVID-19 and Flu in your community.* 2023. [cited 2023 October 20]; Available from: https://outbreaksnearme.org/us/en-US.

17. CDC. *VISION Vaccine Effectiveness Network.* 2023 [cited 2023 April 17]; Available from: https://www.cdc.gov/flu/vaccines-work/vision-network.html.

18. CDC. *Influenza Hospitalization Surveillance Network (FluSurv-NET)*. 2023 [cited 2023 April 17]; Available from: https://www.cdc.gov/flu/weekly/influenza-hospitalization-surveillance.htm.

19. IBM MarketScan® Commercial Claims and Encounters Database [cited 2023 October 20]; Available from: https://www.ibm.com/downloads/cas/OWZWJ0QO.

20. Cohen, C., et al., *Asymptomatic transmission and high community burden of seasonal influenza in an urban and a rural community in South Africa, 2017–18 (PHIRST): a population cohort study.* The Lancet Global Health, 2021. **9**(6): p. e863-e874.

21. CDC. *Past Seasons Estimated Influenza Disease Burden*. 2023 [cited 2023 April 17]; Available from: https://www.cdc.gov/flu/about/burden/index.html.

22. UKHSA, *Investigation into the risk to human health of avian influenza (influenza A H5N1) in England: technical briefing 3*. 2023.

23. Biggerstaff, M., et al., *Influenza-like Illness, the Time to Seek Healthcare, and Influenza Antiviral Receipt During the 2010–2011 Influenza Season—United States.* The Journal of Infectious Diseases, 2014. **210**(4): p. 535-544.

24. Ip, D.K., et al., *Viral Shedding and Transmission Potential of Asymptomatic and*
Paucisymptomatic Influenza Virus Infections in the Community. Clin Infect Dis, 2017. **64**(6): p.
736-742.
25. Millman, A.J., et al., *Improving Accuracy of Influenza-Associated Hospitalization Rate Estimates.*
Emerg Infect Dis, 2015. **21**(9): p. 1595-601.
26. CDC. *COVID Data Tracker*. 2023 [cited 2023 April 17]; Available from:
https://covid.cdc.gov/covid-data-tracker/#datatracker-home.
27. Reed, C., et al., *Estimating influenza disease burden from population-based surveillance data in*
the United States. PLoS One, 2015. **10**(3): p. e0118369.
28. Shrestha, S.S., et al., *Estimating the burden of 2009 pandemic influenza A (H1N1) in the United*
States (April 2009-April 2010). Clin Infect Dis, 2011. **52 Suppl 1**: p. S75-82.
29. Gostic, K.M., et al., *Potent protection against H5N1 and H7N9 influenza via childhood*
hemagglutinin imprinting. Science, 2016. **354**(6313): p. 722-726.
30. Dalton, A.F., et al., *Patient and Epidemiological Factors Associated With Influenza Testing in*
Hospitalized Adults With Acute Respiratory Illnesses, 2016–2017 to 2019–2020. Open Forum
Infectious Diseases, 2023. **10**(4).
31. Baltrusaitis, K., et al., *Healthcare-Seeking Behavior for Respiratory Illness Among Flu Near You*
Participants in the United States During the 2015–2016 Through 2018–2019 Influenza Seasons.
The Journal of Infectious Diseases, 2020. **226**(2): p. 270-277.
32. Tenforde, M.W., et al., *Influenza Antiviral Use in Patients Hospitalized With Laboratory-*
Confirmed Influenza in the United States, FluSurv-NET, 2015-2019. Open Forum Infect Dis, 2023.
10(1): p. ofac681.
33. Ellis, J., et al., *Evaluation of four real-time PCR assays for detection of influenza A(H1N1)v viruses.*
Eurosurveillance, 2009. **14**(22): p. 19230.