1 **The effect of health insurance coverage on antenatal care utilization in Cambodia: A**
2 **secondary analysis of Cambodia Demographic and Health Survey 2021-2022**
3

4 Samnang Um[1,2*], Channnarong Phan[1], Khun Veha[3], Soklim Pay[4], Darapheak Chau[1]
5
6 1- National Institute of Public Health, Cambodia
7 2- Faculty of Social Science and Humanities, Royal University of Phnom Penh, Cambodia
8 3- Department of Payment Certification of National Payment Certification Agency, Cambodia
9 4- Technical Department of National Pediatric Hospital, Cambodia
10
11 * umsamnang56@gmail.com
12

Abstract

Health insurance is essential in reducing or eliminating the financial constraint to accessing

maternal health services caused by out-of-pocket payments. Also, it has a beneficial effect in

minimizing the number of maternal and child mortality. However, limited studies in Cambodia

examined the association between health insurance coverage on antenatal care (ANC) utilization.

Therefore, this study examined the effect of health insurance coverage on ANC utilization in

Cambodia. The study utilized data from the 2021-2022 Cambodia Demographic and Health

Surveys (CDHS). A total of 3,162 weighted women who gave birth within two years were

included in the study. Multiple logistic regression analysis was explored to assess the association

between health insurance coverage on women who attended four or more ANC visits. About 24.9%

of the women had health insurance coverage during 2021-2022. Most (86.1%) of women

attended four or more ANC visits. Women with health insurance coverage were statistically

significantly associated with attending four or more ANC visits with an adjusted odds ratio

(AOR = 1.6; 95% CI: 1.1–2.4). Other covariates significantly associated with women who

attended four or more ANC visits include women with higher education (AOR = 3.1; 95% CI:

1.2–7.7), secondary education (AOR = 2.3; 95% CI: 1.5–3.5), richest households (AOR = 3.2; 95%

CI: 1.5-6.8), and richer households (AOR = 1.9;95% CI: 1.2-2.8). Pregnant women with health

insurance coverage, who had completed at least secondary education and had a better wealth

index were more likely to attend at least four ANC visits. Thus, improving health insurance

coverage, and women's economic and education may be essential to improving women's access

to attended ANC utilization services in Cambodia.

Key work: Health insurance, Antenatal care utilization, Pregnant women, Cambodia

Introduction

Cambodia's maternal mortality rate has significantly declined in the past decade. Data from the 2021-2022 Cambodia Demographic and Health Surveys (CDHS) show that maternal mortality had declined dramatically, from 488 to 154 per 100,000 live births between 2000 and 2021-2022 [1,2]. By 2030, the global maternal death ratio is expected to drop to less than 70 per 100,000 live births, according to Sustainable Development Goal (SDG) 3.1[3]. This achievement can be attributed to the country's concerted effort to increase women's access to maternal health services, particularly the initiative to increase institutional births [4]. Institutional births dramatically increased, from 19.3% to 98%, while the proportion of pregnant women attending four or more antenatal care (ANC) appointments increased considerably, from 9% to 86.1%, between 2000 and 2021-2022 [5]. In several studies, women who had health insurance had higher rates of using maternal health treatments, such as timely ANC and attending four or more ANC visits [6-9].

In 2019, the total population of Cambodia was 15.55 million, with 17.8% living below the national poverty line [10]. Since 2016, Cambodia has been classified as a lower-middle-income country, with gross domestic product (GDP) per capita from 302 US dollars in 2000 to 1,625 US dollars in 2021 [11]. Also, current expenditures on health per capita significantly increased from 20 US dollars in 2000 to 116 US dollars in 2020 [12]. Globally, 50% of people cannot access essential health services, as the World Bank and World Health Organization (WHO) reported in 2017 [13].

Cambodian National Social Security Fund (NSSF) has provided health insurance coverage to formal sector workers [14]. And poor households are covered by the Health Equity Fund (HEF), the co-financing mechanism of the government and development partners [15]. By 2025, the government intends to expand the reach of the NSSF health insurance program to include the entire population [16]. Data from CDHS 2021-2022 indicated that 22% of women and 13% of men aged 15-49 years have any health insurance, respectively [5]. Health insurance coverage is expected to provide financial risk protection and reduce disparities in access by facilitating greater uptake of maternal health services [13]. To our knowledge, limited published peer-reviewed studies assess the association between health insurance coverage and access to

maternal health services among women of reproductive age in Cambodia using updated data.
One prior study on health insurance coverage and its impact on maternal healthcare utilization in
low- and middle-income countries utilized data from CDHS 2010 [8]. This study included all
women and men aged 15-49 and pooled Demographic and Health Survey (DHS) data in 30 low-
and middle-income countries (LMICs) [8]. An additional study aimed to assess levels of health
insurance coverage in 30 LMICs and examines the impact of health insurance status on the use
of maternal health care in eight countries spanning sub-Saharan Africa (Burundi et al., Namibia,
and Rwanda), West Asia (Albania), and South and Southeast Asia (Cambodia and Indonesia) [8].
Several pieces of evidence on the effect of health insurance coverage on ANC utilization have
been published [6-9,17-19]; results indicate that women with health insurance coverage had
higher odds attended four or more ANC visits than those without [6-9,17-19]. Moreover, those
women who reported exposure to media, married women, those with high education, those living
in wealthy economic families, those who are unemployed, and those living in urban areas were
also factors associated with attending four or more ANC visits [6-9,17-19]. Given the lack of
scholarship addressing this health concern among Cambodian women aged 15-49, we examined
the effects of health insurance coverage on antenatal care (ANC) utilization among women who
had a live birth in the past two years in Cambodia. The findings will provide a broader
perspective on levels of health insurance coverage and the impact of health insurance status on
the use of maternal health care in Cambodia. Additionally, the study will enable policymakers to
understand better health insurance coverage among the adult population in Cambodia and proffer
suggestions for improving universal health coverage in Cambodia.
88
89 **Material and Methods**
90
91 **Ethical statement**
92
The CDHS 2021–2022 is publicly available, with all personal identifiers of study participants
removed. Permission to analyze the data was granted by registering with the DHS program
website at (URL: https://dhsprogram.com/data/available-datasets.cfm). Written informed consent
was obtained from the parent/guardian of each participant under 18 before data collection. The
Cambodia National Ethics Committee for Human Health Research (NECHR) approved the data

collection tools and procedures for CHDS 2021-2022 for Health Research on 10 May 2021

(Reference number: 83 NECHR) and ICF's Institutional Review Board (IRB) in Rockville,

Maryland, USA.

Data source

We used data from the most recent CDHS (2021–2022), a household survey conducted every

five years nationally representative of the population [5]. The two-stage stratified cluster

sampling method collected the samples from all provinces. At the first stage, clusters, or

enumeration areas (EAs), that represent the entire country (urban and rural), are randomly

selected from the sampling frame using probability proportional (PPS) to cluster size. In the

second stage, a complete listing of households was selected from each cluster using an equal

probability systematic sampling, and then interviews were conducted with women aged 15–49

years who were born in the five years preceding the survey in the complete list selected

households [5]. In total, 19,496 women aged 15-49 who had given birth in the last five years

were interviewed face-to-face, using the survey standard questionnaire to collect information

from women on several health indicators such as maternal health care service utilization,

maternal and child health, nutrition, and reproductive health services [5]. Overall, 15,046 women

who had not given birth in the past two years were excluded. Data restriction resulted in women

who had a live birth in the past two years in a final analytic sample of 3,292 women (3,162

weighted women).

Measurements

Outcome variable

This study's outcome was the number of ANC visits during the last pregnancy among women

aged 15-49 years (coded as 0 = less than 4 ANC visits included women who reported no ANC

visits and 1 = four or more ANC visits) [6,18,20].

Independent variables

The primary independent variable is maternal health insurance coverage (coded as 0 = no

(reference and 1=yes), including public and private insurance. The confounding variables

included maternal factors: Women's age in years (coded as 1=15-30 (reference) and 2=31-49),

marital status (coded as 1=married (reference) and 2=not married), birth order (coded as 1=1

(reference), 2=2-3, and 3=4 or more), education (coded as 0 = no education (reference),

1=primary, and 2=secondary or higher), occupation (coded as 0=not working (reference),

1=professional, 2=sales or services, 3=agricultural, and 4=manual labor). Individual Household

factors included the household wealth index (coded as 1=poorest (reference), 2=poorest,

3=medium, 4=richer, and 5=richest) were calculated following the principal component analysis

(PCA) [5]. Cambodia's geographical regions were grouped into four categories (coded as

1=Plains (reference), 2=Tonle Sap, 3=Coastal/Sea, and 4=Mountains), and place of residences

(coded as 1=urban (reference) and 2=rural) was defined based on Cambodia's General

Population Census 2019 and adapted from the original CDHS 2021-2022 [5,10].

Statistical analysis

Statistical analysis was performed using STATA version 17 (StataCorp LLC). We applied for the

DHS standard sampling weight variable (**v005/1,000,000**). Then, we used the survey-specific

STATA command "**svy**" for descriptive and analytical analysis. Women's socio-economic and

demographic characteristics were described using weighted frequency and percentage

distributions.

Bivariate analysis using Chi-square tests assessed the association between the variables of

interest (maternal and individual household characteristics) and ANC visits. All independent

variables associated with ANC use at p-value ≤ 0.10, or that had a potential confounder variable

[6,18] were included in the multiple logistic regression analysis to determine the independent

factors related to ANC use [26]. Multicollinearity between original independent variables was

checked, including women's age, number of children ever born, education, wealth index,

occupation, marital status, coverage health insurance, and place of residence. The result of the

evaluating variance inflation factor (VIF) scores after fitting an Ordinary Least Squares

regression model with the mean value of VIF was 1.53, which is less than the cutoff point

indicating no collinearity correlation among the independent variables [27].

Results

Characteristics of the study population

Table 1 describes the socio-economic and demographic characteristics of the 3,162 women aged 15–49 born in the last two years preceding the survey. The mean age was 22.2 years old (SD = 4.2 years); the age group of 15–29 years old accounted for 94.3%. The majority (95%) were currently married. More than 33.4% of women had their first child. Half of the women completed at least secondary education, while 10.6% had no formal education. Only 6.5% of workers were professionals, and 31.2% were unemployed. Of the sample, 20.7% of women were from the poorest households, and 19.7% were from poorer households. Sixty-two percent of the women lived in rural areas. Only 786 (24.9%) women aged 15–49 had health insurance coverage. 86.1% of women attended at least four ANC visits during pregnancy.

Table 1. Socio-economic and demographic characteristics of women aged 15-49 years with a birth in the last two years preceding the survey (N = 3,162 weighted)

Variables	Freq.	%
Mean age at the time of birth (SD)	22.2(4.2)	
15-29	2,982	94.3
30-49	180	5.7
Marital status		
Married	3,004	95.0
Not married	158	5.0
Birth order		
1child	1055	33.4
2-3 children	1197	37.9
4 or more children	909	28.7
Educational		
No education	334	10.6
Primary	1253	39.6
Secondary	1361	43.0
Higher	214	6.8
Occupation		
Not working	986	31.2
Professional	205	6.5
Sales	579	18.3
Agricultural	445	14.1
Services	76	2.4
Manual labor	810	25.6
Wealth index		
Poorest	655	20.7

Poorer	623	19.7
Middle	626	19.8
Richer	683	21.6
Richest	574	18.2
Residence		
Urban	1202	38.0
Rural	1960	62.0
Region		
Plain	1532	48.5
Tonle Sap	996	31.5
Coastal	201	6.4
Plateau/Mountain	432	13.7
Covered by health insurance		
No	2376	75.1
Yes	786	24.9
Number of ANC visits		
< 4 ANC	440	13.9
$\geq$ 4 ANC	2722	**86.1**

Factors Associated with Four or more ANC visits in Chi-Square analysis

In bivariate analysis (**Table 3**), a higher proportion of women with health insurance coverage had a significant association with four or more ANC visits (91.6% vs. 84.2%, $p < 0.001$). Women aged 31–49 reported being more likely to attend four or more ANC visits (88.0% vs. 86.0%, $p < 0.001$). Also, married women reported four or more ANC visits than nonmarried women (86.6% vs. 76.2%, $p = 0.007$). Women with no education were less likely to attend four or more ANC visits than those with higher education (71.0% vs. 95.6%, $p < 0.001$). Four or more ANC visits were higher among women working in professional (95.3%) and service (96.4%), respectively, compared to unemployed women (84.3), with $p < 0.001$. Additionally, four or more ANC visits were higher among women from the richer and richest on the wealth index (94.8 and 89.4%, respectively), compared to the poorer and poorest (73.9 and 86.9%, respectively, with $p < 0.001$). Lastly, women living in urban areas reported higher four or more ANC visits than in rural areas (91.5 vs. 82.7%, $p < 0.001$).

Table 3. Maternal and household characteristics by women attending at least four antenatal care and delivery in a health facility (**N = 3,162**)

Variables	Number of ANC visits		p-value
	Four or more	**Less than four**	
	n=2,722	**n=440**	
	%	%	
Covered by health insurance			
No	84.2	15.8	<0.001
Yes	91.6	8.4	
Age at time of birth			
15-30	86.0	14.0	<0.001
31-49	88.0	12.0	
Marital status			
Married	86.6	13.4	0.007
Not married	76.2	23.8	
Birth order			
1 child	88.4	11.6	<0.001
2-3 children	88.6	11.4	
4 or more children	80.1	19.9	
Educational			
No education	71.0	29.0	<0.001
Primary	84.1	15.9	
Secondary	90.1	9.9	
Higher	95.6	4.4	
Occupation			
Not working	84.3	15.7	<0.001
Professional	95.3	4.7	
Sales	84.4	15.6	
Agricultural	79.0	21.0	
Services	96.4	3.6	
Manual labor	89.9	10.1	
Wealth index			
Poorest	73.9	26.1	<0.001
Poorer	86.9	13.1	
Middle	86.3	13.7	
Richer	89.4	10.6	
Richest	94.8	5.2	
Residence			
Urban	91.5	8.5	<0.001
Rural	82.7	17.3	
Region			
Plain	89.3	10.7	<0.001
Tonle Sap	86.6	13.4	
Coastal	89.0	11.0	
Plateau/Mountain	72.2	27.8	

207 *Notes: Survey weights are applied to obtain weighted percentages. **Plains:** Phnom Penh, Kampong Cham, Tbong Khmum, Kandal,*
208 *Prey Veng, Svay Rieng, and Takeo; **Tonle Sap:** Banteay Meanchey, Kampong Chhnang, Kampong Thom, Pursat, Siem Reap,*
209 *Battambang, Pailin, and Otdar Meanchey; **Coastal/sea:** Kampot, Kep, Preah Sihanouk, and Koh Kong; **Mountains:** Kampong Speu,*
210 *Kratie, Preah Vihear, Stung Treng, Mondul Kiri, and Ratanak Kiri.*
211
212
213
214

Association between health insurance and maternal Healthcare Services Utilization

Table 4 shows the results of the multiple logistic regression analysis of the association between health insurance coverage and maternal healthcare services utilization after controlling for the socio-demographic factors. Compared to women without health insurance, those with health insurance coverage were more likely to attend four or more ANC visits (AOR = 1.6, 95% CI: 1.1–2.4). Women with higher education (AOR = 3.1, 95% CI: 1.2–7.7), secondary education (AOR = 2.3, 95% CI: 1.5–3.5), and primary education (AOR = 1.7, 95% CI: 1.2–2.7) were more likely to have four or more ANC visits than women without any formal education. The odds of having four or more ANC visits were more significant for women from the wealthiest households than for those from the poorest households: richest households (AOR = 3.2; 95% CI: 1.5-6.8), richer households (AOR = 1.9; 95% CI: 1.2-2.8), and middle households (AOR = 1.5; 95% CI: 1.1-2.2). However, the odds of having four or more ANC visits were lower in unmarried women than in married women (AOR = 0.5; 95% CI: 0.3-0.8).

Table 4. Association between health insurance and four or more ANC visits in simple and multiple logistic regression model (**N = 3,162**)

| Variables | Four or more ANC visits | | | |
| | Unadjusted | | Adjusted | |
	OR	95% CI	AOR	95% CI
Covered by health insurance				
No	Ref.		Ref.	
Yes	2.0***	(1.4-2.9)	**1.6***	**(1.1-2.4)**
Age at time of birth				
15-30	Ref.		Ref.	
31-49	1.2	(0.7-2.1)	1.1	(0.6-2.0)
Marital status				
Married	Ref.		Ref.	
Not married	0.5**	(0.3-0.8)	**0.5****	**(0.3-0.8)**
Birth order				
1child	Ref.		Ref.	
2-3 children	1.0	(0.8-1.4)	1.1	(0.8-1.5)
4 or more children	0.5***	(0.4-0.7)	0.7*	(0.5-1.0)
Educational				
No education	Ref.		Ref.	
Primary	2.2***	(1.5-3.1)	**1.8****	**(1.2-2.7)**
Secondary	3.7***	(2.6-5.3)	**2.3****	**(1.5-3.5)**
Higher	8.9***	(4.3-18.3)	**3.1***	**(1.2-7.7)**
Occupation				
Not working	Ref.		Ref.	
Professional	3.8***	(2.0-7.0)	1.4	(0.7-2.8)

Sales	1.0	(0.7-1.4)	0.7	(0.5-1.0)
Agricultural	0.7[*]	(0.5-1.0)	1.0	(0.7-1.4)
Services	4.9[**]	(1.8-13.5)	2.6	(0.9-7.2)
Manual labor	1.7[**]	(1.2-2.3)	1.3	(0.9-1.9)
Wealth index				
Poorest	Ref.		Ref.	
Poorer	2.4[***]	(1.7-3.2)	**1.7[**]**	**(1.2-2.3)**
Middle	2.2[***]	(1.6-3.1)	**1.5[*]**	**(1.1-2.2)**
Richer	3.0[***]	(2.1-4.2)	**1.9[**]**	**(1.2-2.8)**
Richest	6.4[***]	(3.4-12.0)	**3.2[**]**	**(1.5-6.8)**
Residence				
Urban	Ref.		Ref.	
Rural	0.4[***]	(0.3-0.6)	0.8	(0.5-1.1)
Region				
Plain	Ref.		Ref.	
Tonle Sap	0.8	(0.6-1.1)	1.2	(0.9-1.7)
Coastal	1.0	(0.6-1.5)	1.3	(0.8-2.1)
Plateau/Mountain	0.3[***]	(0.2-0.4)	0.5[***]	(0.3-0.6)

Ref = reference value
[*] $p < 0.05$, [**] $p < 0.01$, [***] $p < 0.001$
*Notes: Survey weights are applied to obtain weighted percentages. ***Plains:** Phnom Penh, Kampong Cham, Tbong Khmum, Kandal,*
Prey Veng, Svay Rieng, and Takeo; **Tonle Sap:** *Banteay Meanchey, Kampong Chhnang, Kampong Thom, Pursat, Siem Reap,*
Battambang, Pailin, and Otdar Meanchey; **Coastal/sea:** *Kampot, Kep, Preah Sihanouk, and Koh Kong;* **Mountains:** *Kampong Speu,*
Kratie, Preah Vihear, Stung Treng, Mondul Kiri, and Ratanak Kiri.

Discussion

We analyzed the most recent 2021–2022 CDHS data to examine the relationship between health
insurance coverage and receiving four or more ANC visits during pregnancy. Overall, 24.9% of
women reported having health insurance coverage among women of reproductive age who gave
birth within two years of the survey. This finding is slightly similar to lower-middle-income
countries, where 27.3% of women had health insurance coverage [19]. This is higher than in
low-income countries, where 7.9% of women have health insurance coverage [19]. However,
lower than in upper-middle-income countries, 52.5% of women had health insurance coverage
[19]. Since the formal launch of the Cambodia National Social Security Fund (NSSF) with the
Health Insurance Scheme in 2008, the proportion of women with health insurance coverage has
increased from 16% in 2014 to 22% in 2021–2022 [5]. This proportion exponentially increased
due to the Royal Government of Cambodia's implementation of the NSSF for all workers in the
formal and informal sectors of the economy [10]. Moreover, it has plans to extend the healthcare
benefits under the NSSF to the family members of the employees as well [16].

This study found that women with health insurance coverage were 1.6 times more likely to
attend four or more ANC visits during pregnancy. Previous studies documented the positive
relationship between health insurance and the number of ANC visits among women of
reproductive age [8,9,19]. Health insurance eliminates the financial barrier to accessing maternal
health services caused by out-of-pocket payments. It has a beneficial effect in reducing the
number of low-birth-weight babies born and child mortality [17,21]. The result is more equitable
access to care, potentially improving maternal health outcomes [9,19]. The MoH has since raised
the minimum standard for ANC visits during pregnancy to at least four trips [20,22]. The
dramatically significant increase in the highest prevalence of four or more ANC visits was an
effort by the Royal Government of Cambodia, which has strengthened health facilities across the
country, particularly in rural areas, improved infrastructure, provided essential medical
equipment and supplies, increased the number of midwives, expanded antenatal care, and
provided more skilled medical practitioners at childbirth to ensure safe delivery practices.
Furthermore, to encourage early and routine ANC visits, the government is offering pregnant
women a monetary incentive of 20 US dollars for each visit during a maximum of four ANC
visits at any health facility with a contract with the National Social Security Fund (NSSF)
[14,23].

This study found that increased education and household wealth index increased the likelihood
of four or more ANC visits. Women's education and higher wealth index are more likely to attend
four or more ANC visits. This aligns with previous evidence around socio-economic inequalities
in maternal health service utilization in Cambodia and South Asia [24,25]. This may be because
the more educated women is the more aware of the importance of ANC for their health and baby
[26]. Additionally, education gives women the power to decide whether to seek medical attention
and enables them to recognize warning signs of pregnancy complications. Moreover, women
from higher-income households were more likely to be able to cover the costs of care-seeking,
including any related expenses and transportation [24,25]. Thus, in this study, women with
higher education levels and household wealth indexes had the highest proportion of health
insurance coverage.

This study has several strengths. First, it used the most recent women's data from the 2021–2022
CDHS, an extensive representative national population-based household survey with a high
response rate of 97%. Second, the recall bias has been minimized by limited analysis of women's
most recent deliveries within the last two years preceding the survey [5]. Third, the complex
survey design and sampling weights in the analysis accounted for both descriptive and analytics
methods that enabled us to generalize our findings to the population of WRA in Cambodia. In
addition, DHS data were collected using validated survey methods and highly trained data
collectors, contributing to improved data quality [27]. Last, to our knowledge, this is the first
study to report the association between health insurance coverage and ANC visits in Cambodia.
After controlling sociodemographic factors, we found significant associations between health
insurance coverage and several forums or more ANC visits. A key finding compares the
association with other studies in Southeast Asia and globally.

Despite this, there are several limitations. First, this study used a secondary analysis, so it did not
address health institution factors of antenatal care utilization and service availability; hence, this
study could not explore the quality of ANC services, though the quality of healthcare services
plays a vital role in patient satisfaction and use. Second, the study's cross-sectional nature could
not assist in the temporal relationship of variables including the number of years since women
joined health insurance on ANC utilization. Therefore, further study should be conducted to
identify factors related to health institutions. In addition, antenatal care utilization should be
performed based on the new WHO guidelines revised in 2016 at the national level [22].
Moreover, longitudinal studies that address comprehensive variables should be studied. Third,
we excluded other factors, such as maternal complications and women's empowerment indicators,
that could affect the use of maternal care. Lastly, CDHS did not assess a direct measure of
maternal health literacy.

Conclusion

This is the first study to report the association between health insurance coverage and ANC visits
in the healthcare setting in Cambodia. Cambodian pregnant women attend four or more antenatal
care visits, which is slightly high. However, it still needs to be satisfactory. Health among
coverage women of reproductive age in Cambodia is relatively low. Moreover, we found that

women with health insurance, women with education, and being rich in the wealth quintile were strong predictors of women attending four or more ANC visits. There is a need to pay close attention to improving the uptake of health insurance among women of reproductive age, especially targeting women with no education, from low-income families, and women who reside in rural areas. Policymakers may need to prioritize women of reproductive age in designing and implementing health insurance programs to increase their uptake. This would provide financial risk protection, facilitate access to maternal health services, and possible attainment of Cambodia's SDG 3 targets.

Acknowledgments

The authors would like to thank DHS-ICF, who approved the data used for this paper.

References

1.   National Institute of Statistics, Directorate General for Health, and ICF International. Cambodia Demographic and Health Survey 2014. Phnom Penh, Cambodia, and Rockville, Maryland, USA: National Institute of Statistics, Directorate General for Health, and ICF International. Available online: https://dhsprogram.com/pubs/pdf/fr312/fr312.pdf. 2015.

2.   National Institute of Statistics PP, Cambodia, Ministry of Health Phnom Penh, Cambodia, The DHS Program, ICF, Rockville, Maryland, USA. Cambodia Demographic and Health Survey 2021–22. Phnom Penh, Cambodia, and Rockville, Maryland, USA: NIS, MoH, and ICF. 2022. Available from: https://dhsprogram.com/pubs/pdf/FR377/FR377.pdf.

3.   The United Nations Development Programme. The Sustainable Development Goals by 2016-2030. Available from: https://www.undp.org/content/undp/en/home/sustainable-development-goals.html.

4.   Acuin CS, Khor GL, Liabsuetrakul T, Achadi EL, Htay TT, Firestone R, et al. Maternal, neonatal, and child health in southeast Asia: towards greater regional collaboration. Lancet. 2011;377(9764):516-25. Epub 20110125. doi: 10.1016/s0140-6736(10)62049-1. PubMed PMID: 21269675; PubMed Central PMCID: PMCPMC7159081.

5.   National Institute of Statistics (NIS) MoHM, and ICF. . Cambodia Demographic and Health Survey 2021–22 Final Report. Phnom Penh, Cambodia, and Rockville, Maryland, USA: NIS, MoH, and ICF. 2023. Available from: https://dhsprogram.com/pubs/pdf/PR136/PR136.pdf.

6.   Merga BT, Raru TB, Deressa A, Regassa LD, Gamachu M, Negash B, et al. The effect of health insurance coverage on antenatal care utilizations in Ethiopia: evidence from national survey. Front Health Serv. 2023;3:1101164. Epub 20231006. doi: 10.3389/frhs.2023.1101164. PubMed PMID: 37869683; PubMed Central PMCID: PMCPMC10587574.

7.   Aboagye RG, Okyere J, Ahinkorah BO, Seidu AA, Zegeye B, Amu H, et al. Health insurance coverage and timely antenatal care attendance in sub-Saharan Africa. BMC Health Serv Res. 2022;22(1):181. Epub 20220211. doi: 10.1186/s12913-022-07601-6. PubMed PMID: 35148769; PubMed Central PMCID: PMCPMC8840787.

8.   Wang W, Gheda Temsah, and Lindsay Mallick. . Health Insurance Coverage and Its Impact on Maternal Health Care Utilization in Low- and Middle-Income Countries. DHS Analytical Studies No. 45. Rockville, Maryland, USA: ICF International. . 2014.

9.   Ahuru RR, Omon IJ, Nzoputam CI, Ekomoezor E. Health insurance ownership and maternal health service uptake among Nigerian women. HEALTH. 2021;12.

10.   National Institute of Statistics. Ministry of Planning. General population census of the Kingdom of   Cambodia   2019.   Available   from: https://www.nis.gov.kh/nis/Census2019/Final%20General%20Population%20Census%202019-English.pdf.

11.   The   World   Bank   In   Cambodia.   Overview   2019.   Available   from: https://www.worldbank.org/en/country/cambodia/overview.

12.   World Health Organization (WHO). Global Health Expenditure Database 2019. Available from: https://apps.who.int/nha/database/PHC_Country_profile/Index/en.

13.   World Health Organization (WHO); World Bank and WHO. Half the world lacks access to essential health services, 100 million still pushed into extreme poverty because of health expenses 2017. 2017. Available from: http://www.who.int/mediacentre/news/releases/2017/half-lacks-access/en/.

14.   National Social Security Fund (NSSF). Health Insurance Scheme 2016. Available from: https://www.nssf.gov.kh/health-care-scheme-2/.

15.   Nakamura H, Amimo F, Yi S, Tuot S, Yoshida T, Tobe M, et al. Implementing a sustainable health insurance system in Cambodia: a study protocol for developing and validating an efficient household income-level assessment model for equitable premium collection. Int J Equity Health. 2020;19(1):17. Epub 20200131. doi: 10.1186/s12939-020-1126-8. PubMed PMID: 32005237; PubMed Central PMCID: PMCPMC6995079.

16.   The Royal Government of Cambodia. National Social Protection Policy Framework 2016–2025. Phnom Penh: The Council of Ministers; 2017.

17.   Likka MH, Handalo DM, Weldsilase YA, Sinkie SO. The effect of community-based health insurance schemes on utilization of healthcare services in low- and middle-income countries: a systematic review protocol of quantitative evidence. JBI Database System Rev Implement Rep. 2018;16(3):653-61. doi: 10.11124/jbisrir-2017-003381. PubMed PMID: 29521866.

18.   Dadjo J, Ahinkorah BO, Yaya S. Health insurance coverage and antenatal care services utilization in West Africa. BMC Health Serv Res. 2022;22(1):311. Epub 20220307. doi: 10.1186/s12913-022-07698-9. PubMed PMID: 35255895; PubMed Central PMCID: PMCPMC8899447.

19.   Hooley B, Afriyie DO, Fink G, Tediosi F. Health insurance coverage in low-income and middle-income countries: progress made to date and related changes in private and public health expenditure. BMJ Glob Health. 2022;7(5). doi: 10.1136/bmjgh-2022-008722. PubMed PMID: 35537761; PubMed Central PMCID: PMCPMC9092126.

20.   Ministry of Health. Guideline for implementation the service package of Antenatal Care Delivery and   Post   natal   Care   2019   [updated   2019].   3:[Available   from: https://niph.org.kh/niph/uploads/library/pdf/GL220_NMCHC_anc_pnc_package_eng.pdf.

21.   Imo CK, De Wet-Billings N, Isiugo-Abanihe UC. The impact of maternal health insurance coverage and adequate healthcare services utilisation on the risk of under-five mortality in Nigeria: a cross-sectional study. Arch Public Health. 2022;80(1):206. Epub 20220913. doi: 10.1186/s13690-022-00968-2. PubMed PMID: 36100949; PubMed Central PMCID: PMCPMC9472384.

22.   WHO Guidelines Approved by the Guidelines Review Committee. WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience. Geneva: World Health Organization

Copyright © World Health Organization 2016.; 2016.

23.   Ministry of Information. Government Raises Allowances For Mothers And Children. Available from: https://www.information.gov.kh/articles/108509.

24.   Chham S, Radovich E, Buffel V, Ir P, Wouters E. Determinants of the continuum of maternal health care in Cambodia: an analysis of the Cambodia demographic health survey 2014. BMC Pregnancy Childbirth. 2021;21(1):410. Epub 20210602. doi: 10.1186/s12884-021-03890-7. PubMed PMID: 34078318; PubMed Central PMCID: PMCPMC8170811.

25.   Lee HY, Oh J, Kim R, Subramanian SV. Long-term trend in socioeconomic inequalities and
geographic variation in the utilization of antenatal care service in India between 1998 and 2015. Health
Serv Res. 2020;55(3):419-31. Epub 20200304. doi: 10.1111/1475-6773.13277. PubMed PMID: 32133652;
PubMed Central PMCID: PMCPMC7240766.
26.   Thakkar N, Alam P, Saxena D. Factors associated with underutilization of antenatal care in India:
Results from 2019-2021 National Family Health Survey. PLoS One. 2023;18(5):e0285454. Epub
20230508. doi: 10.1371/journal.pone.0285454. PubMed PMID: 37155665; PubMed Central PMCID:
PMCPMC10166529.
27.   Allen C, Fleuret J, Ahmed J, editors. Data quality in demographic and health surveys that used
long and short questionnaires2020: ICF.

418