Causal relationship between mental disorders and abdominal aortic aneurysm: insights

from the genetic perspective

Short title: Mental disorders and abdominal aortic aneurysm

Ming-Gang Deng[1,2], Chen Chai[3], Kai Wang[4], Zhi-Hui Zhao[5], Jia-Qi Nie[6], Fang Liu[7], Yuehui

Liang[7], Jiewei Liu[1,2]

[1] Department of Psychiatry, Wuhan Mental Health Center, Wuhan 430012, Hubei, China.

[2] Department of Psychiatry, Wuhan Hospital for Psychotherapy, Wuhan 430012, Hubei,

China.

[3] Emergency Center, Hubei Clinical Research Center for Emergency and Resuscitation,

Zhongnan Hospital of Wuhan University, Wuhan 430071, Hubei, China.

[4] Department of Public Health, Wuhan Fourth Hospital, Wuhan 430000, Hubei, China.

[5] School of Nursing, Shanghai Jiao Tong University, Shanghai 200025, China.

[6] Xiaogan Municipal Center for Disease Control and Prevention, Xiaogan 432000, Hubei,

China.

[7] School of Public Health, Wuhan University, Wuhan 430071, Hubei, China.

Correspondence:

Ming-Gang Deng

ORCID: 0000-0002-3154-6909

Email: deng.minggang@outlook.com

Address: Department of Psychiatry, Wuhan Mental Health Center, Wuhan 430012, Hubei,

China.

Jiewei Liu

ORCID: 0009-0009-8265-8984

Email: liujiewei@hust.edu.cn

Address: Department of Psychiatry, Wuhan Mental Health Center, Wuhan 430012, Hubei,

China.

Word count: 5950

Abstract

Objective: This study investigated the genetic link between mental disorders—depression,

schizophrenia (SCZ), and bipolar disorder (BIP)—and abdominal aortic aneurysm (AAA).

Methods: We used global/local genetic correlation analyses and identified shared genomic

loci. Bidirectional univariable Mendelian Randomization (UMR) assessed causal directions,

with multivariable MR (MVMR) refining the analysis. Mediation analyses examined if

known AAA risk factors influenced this relationship.

Results: Global correlations showed positive links between depression, SCZ, and AAA, but

not BIP. Local analyses identified specific genomic regions of correlation. We found 26, 141,

and 10 shared loci for AAA with depression, SCZ, and BIP, respectively. UMR indicated

significant associations between genetically predicted depression (OR 1.270; 95% CI

1.071-1.504; $p = 0.006$) and SCZ (OR 1.047; 95% CI 1.010-1.084; $p = 0.011$) with AAA, but

not BIP. These results were confirmed by MVMR analyses. Mediation analyses showed that

smoking, hypertension, hyperlipidemia, and coronary atherosclerosis mediated the impact of

depression on AAA while smoking mediated SCZ's impact.

Conclusion: This study provides evidence that genetically predicted depression and SCZ are

linked to an increased risk of AAA, mediated by traditional AAA risk factors.

Keywords: Mental disorders; Abdominal aortic aneurysm; Mendelian Randomization;

Genetic association.

1. Introduction

Abdominal aortic aneurysm (AAA) is a prevalent vascular disease marked by localized dilation of the abdominal aorta and is linked to an approximately 90% mortality risk due to rupture[1]. While recent open surgical repair and endovascular stent graft therapies have notably enhanced the treatment of AAA, there is currently no approved and effective pharmacotherapy to restrain and prevent AAA progression and rupture[2,3]. Consequently, primary preventive measures targeting the risk factors of AAA are necessary to reduce its incidence and adverse effects on health.

Mental disorders, such as depression, schizophrenia (SCZ), and bipolar disorder (BIP), have been found to lead to various adverse outcomes. For example, our previous studies revealed that individuals with depression or SCZ were more likely to be frail[4,5]. In addition, current evidence suggests that mental disorders are linked to higher risks of cardiovascular disease, which include but are not limited to heart failure, coronary artery disease, and hypertension[6-8]. Therefore, investigating the association between mental disorders and AAA is of vital importance for public health.

To our knowledge, the associations of AAA with SCZ and BIP are currently unclear, as no studies have reported them. In the case of depression and AAA, a prospective study based on the Korean National Health Insurance System database revealed that patients with AAA were more likely to be depressed[9]. Conversely, depressed individuals were found to have a higher risk of incident AAA in the Norwegian Nord-Trøndelag Health Study[10]. Despite this, the causative association between depression and AAA needs further investigation, as observational studies were susceptible to confounding bias, reverse causality, etc.

Recent large-scale genome-wide association studies (GWASs) have identified numerous genetic loci associated with depression[11], SCZ[12], BIP[13], and AAA[14]. These findings enhance the statistical power and offer the opportunity to explore their genetic associations. In addition, Mendelian Randomization (MR), which uses genetic variants (single nucleotide polymorphisms, SNPs) as instruments, is a powerful method for obtaining robust causal inference[15] and has been widely utilized in recent epidemiological research[16-18]. As a result, this study aims to investigate the genetic correlation between AAA and mental disorders and to utilize the MR design to evaluate their causality.

2. Methods and materials

2.1. Study design

An overview of this research is presented in **Fig 1**. Initially, we explored the genetic

association between AAA and mental disorders from three perspectives: global genetic

correlation, local genetic correlation, and identification of shared genomic loci. Following that,

we utilized a bidirectional MR design to evaluate the causal association between AAA and

mental disorders. Furthermore, mediation MR analyses were performed to ascertain whether

the causality was mediated by known risk factors.

2.2. Data sources and ethical approval

The data sets utilized in this research were all publicly available from the previous studies,

and the details were described in the **Supplementary Methods**. The ethical approvals were

all obtained by the original papers, and thus the ethical approval for our study could be

waived.

2.3. Genetic association

2.3.1. Global genetic correlation

We conducted a global genetic correlation analysis to estimate the correlation between the

GWAS summary statistics of AAA and three mental disorders. Two different software tools

were utilized to perform this analysis. The first one is Linkage disequilibrium score regression

(LDSC, https://github.com/bulik/ldsc)[19]. We used the cross-trait LDSC which only requires

the GWAS summary statistics that account for sample overlap. Cross-trait LDSC analysis

requires the GWAS summary statistics to be from the same ethnic group. Since the GWAS

summary statistics used in this study are largely from Europeans, we used the 1000 Genomes

Project European samples as the reference sample for LD score calculation.

Another complementary method for estimating global genetic correlation is the Genetic

Covariance   Analyzer   (GNOVA)[20].   GNOVA   (available   at

https://github.com/xtonyjiang/GNOVA) also requires only the GWAS summary statistics as

input to perform global genetic correlation analysis. The major advantage of GNOVA is its

ability to include omic datasets (e.g., transcriptomic and epigenomic) in the GNOVA model.

The LD score was also calculated using the 1000 Genomes European samples.

2.3.2. Local genetic correlation

In addition to estimating the global genetic correlation between the GWAS summary statistics

of AAA and mental disorders, we also attempted to estimate the local genetic correlation

between them in specific genomic regions. To perform this analysis, we used SUPERGNOVA

(available at https://github.com/qlu-lab/SUPERGNOVA). SUPERGNOVA was developed

within the previous GNOVA statistical framework, which includes functional annotation when

estimating genetic correlation[21]. Similar to GNOVA, only GWAS summary statistics and an

LD calculation reference panel were needed as input to run SUPERGNOVA.

2.3.3. Shared genomic loci

To enhance the statistical power of detecting loci associated with both AAA and mental

disorders, we utilized the conjunctional false discovery rate (conjFDR) method for this

analysis. This method is based on standard FDR analysis and aims to identify SNPs that are

associated with both target traits (e.g., SNPs that are jointly associated with AAA and SCZ).

The conjFDR analysis is implemented in the pleioFDR software (available at

https://github.com/precimed/pleiofdr), and it runs in the MATLAB environment[22]. The

analysis followed the instructions of the pleioFDR tool with default parameters.

2.4. Causal inference

2.4.1. Univariate Mendelian Randomization

We initially performed the univariate Mendelian Randomization (UMR) to assess the causal

association between mental disorders and AAA. The statistical analyses were conducted in

two directions: 1) mental disorders as the exposures to check the effect of mental disorders on

AAA, and 2) AAA as exposures to evaluate whether people with AAA had a likelihood to

develop mental disorders.

Independent SNPs (R^2 < 0.001, window size = 10MB) were firstly screened at the

genome-wide significance threshold ($p < 5\times10^{-8}$) to ensure their robust association with

exposure and freedom from linkage disequilibrium. Subsequently, the F statistic for each SNP

was calculated using the formula $beta^2/se^2$, and those with an F < 10 were considered weak

instruments and were excluded. The remaining SNPs were harmonized with the outcome data,

and the palindromic genetic variants were eliminated at this stage. If any genetic variants were

unavailable in the outcome data, they will be replaced by the proxied SNPs if possible. After

the aforementioned procedures, the remaining SNPs were used for causal inference.

The inverse variance weighted (IVW) method was used as the primary approach to obtain the

causal estimate, assuming the absence of average pleiotropic effects, and in this scenario, it is

the most efficient method. Cochran's Q statistic was computed to assess the heterogeneity

among the SNPs. If significant heterogeneity was detected ($p < 0.05$), the IVW method would

be applied under a random-effects model; otherwise, it would be applied under a fixed-effects

model. Alongside the IVW method, several other approaches, such as MR-Egger, weighted

median, MR-cML (constrained maximum likelihood), and MR-RAPS (robust adjusted profile

score), were employed to evaluate the robustness of the findings. In the meantime, the

MR-Egger regression intercept term was utilized to evaluate the presence of horizontal

pleiotropy among the instruments.

2.4.2. Multivariate Mendelian Randomization

The multivariate Mendelian Randomization (MVMR), which employs SNPs associated with

various and potentially related exposures, is an extension of UMR that can detect the causal

effects of many risk factors simultaneously[23]. Considering that the instrumental variables in

the UMR may meanwhile be linked to secondary phenotypes and even to the risk factor of

outcomes, which may violate the critical assumption of MR, we further carried out the

MVMR to ascertain if the UMR estimates were affected by any confounders. The

PhenoScanner V2, an expanded tool for searching human genotype-phenotype associations,

was employed in advance to screen the secondary phenotypes with which the instrumental

variables were associated[24]. The significant confounding phenotype ($p < 1 \times 10^{-5}$) would then be

included as a covariate in the MVMR model. Similar to the UMR analyses, the

multivariable-adjusted effect sizes were primarily estimated by the IVW method, and

supplemented by the MR-Egger, weighted median, and MR-cML methods.

2.4.3. Mediation effect of known risk factors

After establishing the causal relationship between mental disorders and AAA, our objective is

to delve deeper into whether this causality is mediated by known risk factors. For example,

smoking, hypertension, hyperlipidemia, and coronary atherosclerosis (CA) have been

identified as risk factors for AAA[3]. To assess their mediation role, a two-step UMR analysis

was performed. The first step involved estimating the causal effects of mental disorders on

smoking, hypertension, hyperlipidemia, and coronary atherosclerosis, and the second step

involved estimating the causal effects of smoking, hypertension, hyperlipidemia, and CA on

AAA. The mediation proportion was calculated based on the causal effect estimated by the

IVW method, while other methods, including MR-Egger, weighted median, MR-cML, and

MR-RAPS, were employed as sensitivity analyses.

3. Results

3.1. Genetic association

3.1.1. Global genetic correlation

The estimates of the global genetic correlation between mental disorders and AAA are

presented in **Fig 2**. The LDSC method indicated that genetic variants related to depression

[$r_g = 0.157$, se = 0.023, $p = 3.52 \times 10^{-12}$] and SCZ ($r_g = 0.042$, se = 0.020, $p = 0.042$) were

positively correlated with AAA, while BIP was not ($r_g = 0.013$, se = 0.023, $p = 0.591$).

Additionally, the GNOVA method estimates further supported these findings, indicating a

positive association of AAA with depression ($r_g = 0.152$, se = 0.019, $p = 8.41 \times 10^{-16}$) and SCZ

(r_g= 0.053, se = 0.017, p = 0.002). Conversely, BIP did not demonstrate a significant

association with AAA (r_g= 0.021, se = 0.019, p = 0.272).

3.1.2. Local genetic correlation

The local genetic correlation analyses revealed that mental disorders exhibited local

correlations with AAA in various genomic regions, as detailed in **Table 1**. Following

correction for multiple tests, depression was identified to have a positive association with

AAA in four genomic regions (chr9: 126927204-128926989, chr10: 106139294-107874146,

chr11: 42244050-42854352, and chr18: 77151867-78017074). Additionally, SCZ

demonstrated a positive correlation with AAA in 9 out of 10 significant regions. While the

global correlation did not indicate an association between BIP and AAA, the local analyses

revealed correlations in 7 genomic regions, with 6 of them also showing a positive

association.

3.1.3. Shred genetic variants

In line with the preceding analyses, a multitude of shared genetic variants have been discerned

between mental disorders and AAA. The findings are meticulously outlined in **Fig. 3** and

Table S1. Specifically, there are 26 independent SNPs shared between depression and AAA,

with 21 of them demonstrating a consistent directional effect on both conditions. Similarly,

SCZ and AAA exhibit the sharing of 141 independent loci, among which 21 loci manifest an

opposing effect. Notably, only 10 genetic variants are concurrently associated with BIP and

AAA, with 4 of them displaying the same directional effect.

3.2. Causal inference

3.2.1. Univariate Mendelian Randomization

We initially examined the bidirectional causal relationship between mental disorders and AAA

using the UMR, and the results are illustrated in **Fig. 4**. In the forward analyses, the

MR-Egger regression intercept suggested the absence of horizontal pleiotropy among the

selected SNPs (p values all > 0.05). The Cochran's Q test revealed significant heterogeneity

among the instruments concerning depression (Cochran's Q = 163.448, p < 0.001), SCZ

(Cochran's Q = 333.578, p < 0.001), and BIP (Cochran's Q =32.604, p = 0.002). Consequently,

the primary causal inference utilized the random-effect IVW method. The IVW method

revealed that people with depression (OR: 1.270, 95% CI: 1.071-1.504, p = 0.006) and SCZ

(OR: 1.047, 95% CI: 1.010-1.084, p = 0.011) have a heightened susceptibility to AAA, while

no causal effect of BIP on AAA was found (OR: 0.980, 95% CI: 0.896-1.121, p = 0.653).

The findings mentioned above are supported by the weighted median, MR-CML, and

MR-RAPS methods.

For the reverse analyses, the MR-Egger regression intercept also demonstrated there was no

horizontal pleiotropy among the AAA-associated instrumental variables, and obvious

heterogeneity was identified. The IVW method revealed that genetically predicted AAA

showed no association with the risks of depression (OR: 1.010, 95% CI: 0.995-1.025, p =

0.190), SCZ (OR: 1.007, 95% CI: 0.968-1.048, p = 0.740), as well as BIP (OR: 0.989, 95% CI:

0.944-1.037, p = 0.646), and these null associations were consistently supported by estimates

from the MR-Egger, weighted median, MR-cML, and MR-RAPS methods.

3.2.2. Multivariate Mendelian Randomization

After performing the UMR, we proceeded with the MVMR to evaluate potential bias in the

causal estimates due to confounding effects. The PhenoScanner revealed that instruments

associated with mental disorders were predominantly linked to body mass index (BMI) **(Table**

S2). Additionally, SNPs associated with AAA were found to be correlated with BMI, coronary

heart disease (CAD), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein

cholesterol (LDL-C), and total cholesterol (TC) **(Table S3)**. Consequently, we conducted

bidirectional MVMR to evaluate the multivariate-adjusted causal estimates between mental

disorders and AAA. The forward analyses aimed to assess the effect of mental disorders on

AAA with adjustments for BMI, while the reverse analyses aimed to evaluate the effect of

AAA on mental disorders with adjustments for BMI, CAD, HDL-C, LDL-C, and TC.

The results of MVMR are presented in **Table S4**. The IVW method revealed that individuals

with depression (OR: 1.315, 95% CI: 1.095-1.578, p = 0.003) and SCZ (OR: 1.042, 95%

CI:1.0004-1.084, p = 0.048) were correlated with a higher risk of AAA even after controlling

the effect of BMI, while people with BIP were not (OR: 1.014, 95% CI: 0.955- 1.076, p =

0.652). Consistent with the results of UMR, the IVW methods also presented that genetically

predicted AAA still showed no association with the risks of depression, SCZ, and BIP after

adjustments for BMI, CAD, HDL-C, LDL-C, and TC. These bidirectional relationships were

supported by the sensitivity analysis methods.

3.2.3. Mediation Mendelian Randomization

Since the UMR and MVMR only supported the causal effects of depression and SCZ on AAA,

we would only investigate whether these two causalities were mediated by the known risk

factors of AAA. As described in the methods section, we selected smoking, hypertension,

hyperlipidemia, and CA as the four candidate mediating factors.

The results of mediation MR analyses are presented in **Table S5**. The first step MR analyses

presented that genetically predicted depression was associated with increased odds ratio of

smoking (OR: 1.236, 95% CI: 1.140-1.340, p <0.001), hypertension (OR: 1.206, 95% CI:

1.069-1.361, p = 0.002), hyperlipidemia (OR: 1.200, 95% CI: 1.037-1.388, p = 0.015), and

CA (OR: 1.210, 95% CI: 1.078-1.359, p = 0.001). In contrast, genetically predicted SCZ was

only associated with an increased likelihood of smoking (OR: 1.047, 95% CI: 1.023-1.071, p

<0.001). The second step MR analyses demonstrated that genetically predicted smoking (OR:

1.610, 95% CI: 1.449-1.788, p <0.001), hypertension (OR: 1.218, 95% CI: 1.142-1.300, p

<0.001), hyperlipidemia (OR: 1.547, 1.288-1.857, p <0.001), and CA (OR: 1.385, 95% CI:

1.204-1.594, p <0.001) were all associated with elevated risks of AAA. Based on the results

of mediation MR findings, the causal effect of depression on AAA was mediated by smoking,

hypertension, hyperlipidemia, and CA, while the causal effect of SCZ on AAA was mediated

by smoking only (**Fig 5**).

4. Discussion

We conducted an investigation into the relationship between mental disorders and AAA from

the genetic perspective. Our findings provided solid evidence of the genetic correlation

between AAA and depression as well as SCZ. Besides, the causal effects of depression and

SCZ on AAA were identified, with known risk factors for AAA playing a mediating role.

Few studies have investigated the associations between mental disorders and AAA, with the

evidence supporting the correlations between depression and AAA being limited. Furthermore,

a study involving 574 participants from four centers in Australia suggested an insignificant

association between depression and AAA growth[25], contradicting conclusions drawn in

previous studies from another perspective. Given the scarcity of research and the

inconsistency of results, we conducted comprehensive genetic association, and univariate,

multivariate as well as mediating MR to evaluate the correlation and causal links between

mental disorders and AAA.

The global genetic correlation analyses indicated positive associations of AAA with

depression and SCZ, and meanwhile, numerous locally correlated genomic regions and shared

loci were identified. The genetic association analyses presented that AAA had plenty of

genetic basis with mental disorders. The results of univariate and multivariate MR analyses

demonstrated robust and statistically significant associations of depression, SCZ, and AAA,

and we further extended our investigation through mediation analysis to delineate potential

pathways underlying these associations. By incorporating well-established risk factors of

AAA as potential mediators, significant mediation effects of smoking, hyperlipidemia, CA,

and hypertension were identified in the causal pathways between depression and AAA, and

the proportions of mediation effects were calculated as 42.22%, 32.20%, 26.05%, and 15.41%,

respectively. In addition, smoking exhibited a substantial mediation effect proportion of 47.60%

in the association between SCZ and AAA. These findings underscore the pivotal role of

smoking in the nexus between mental disorders and AAA, offering crucial insights for further

etiological explorations.

While the pathological mechanism of AAA remains elusive, existing literature has elucidated

a series of contributing processes, notably the infiltration and subsequent destruction of the

aortic wall mediated by immune and inflammatory mechanisms[26]. Both innate and adaptive

immune cells, including macrophages and T cells, have undergone extensive scrutiny in the

pathogenesis of AAA, with several inflammasomes suggested to be involved in both the

inhibition and amelioration of AAA[27,28]. NF-κB, the primary regulator of inflammation and

immune response, is activated by multiple factors including angiotensin II, triggering

macrophage infiltration, inflammation, and subsequent aortic deterioration[29]. Numerous

studies have explored the impact of pharmacological inhibition of nucleotide-binding

oligomerization domain-like receptor family protein 3 (NLRP3) inflammasome on

experimental AAAs, revealing preventive effects characterized by reduced AAA diameter and

incidence[30-32], etc. Furthermore, the remarkable parallels between the immune system and the

brain, both intricately integrated and complex systems associated with memory, imply

intimate connections between immune responses and mental disorders. For instance, some

researchers consider depression as an inflammatory disease[33,34], with inflammation likely

playing a critical role in heightening individual susceptibility to depression[35]. Similarly,

extensive reviews have highlighted the central role of inflammation and immunity in SCZ,

findings supported by both human and animal studies[36,37]. Several two-sample MR studies

indicate the causal associations of inflammatory biomarkers such as C-reactive protein and

soluble interleukin-6 receptor with SCZ, with anti-inflammation strategies emerging as

therapeutic considerations[38,39]. Therefore, inflammation and immune responses may function

as a conduit linking mental disorders and AAA.

Similar to other cardiovascular diseases (CVD), the relationship between mental disorders and

AAA may also be explained by the shared etiological factors, encompassing genetic,

biological, and behavioral mechanisms. Early in 2002, Bondy et al. found that the same allelic

combination of two genes could increase the vulnerability for depressive disorder while also

being associated with a higher risk of myocardial infarction[40], suggesting the genetic

associations between depression and CVD. Molecular genetic studies have further

corroborated modest genetic correlations between severe mental illness and CVD[41].

Furthermore, biological constituents such as neutrophil gelatinase associated lipocalin[42] and

physiological processes like platelet receptors[43] and function have been implicated as

mediating factors linking CVD and depression. Moreover, the overlap of behavioral

determinants including smoking, physical inactivity, obesity, and poor diet was observed

among participants affected by mental disorders and CVD. Similarly, GWAS studies have

unveiled extensive genetic convergence between SCZ and CVD risk factors, with notable

associations observed with smoking initiation and body mass index[44].

Our study had several strengths worth pointing out. Firstly, we utilized the genetic data from

the large-scale GWASs, which could significantly enhance the statistical power and avoid the

confounding bias and reverse causality of observational studies. Additionally, we conducted

two parts of analyses, including genetic correlation and causality inference, and the results

provided consistent evidence on the causal effects of depression and SCZ on AAA.

Furthermore, we identified the mediation effect, which highlighted the interactive impacts of

mental disorders and other risk factors on the risk of AAA.

Despite these advantages, our findings should be interpreted in the context of limitations.

Firstly, the most recent GWAS of BIP have identified various significant SNPs[45], but the

samples overlapped with the AAA. To avoid the causal inference bias due to sample overlap,

we selected the genetic variants associated with BIP from the GWAS with relatively lower

sample sizes[13]. Further studies were warranted to check the relationship between BIP and

AAA. In addition, only the summary-level data could be obtained thus making us unable to

perform the stratified analyses. At the same time, current knowledge indicates that the

prevalence of AAA in males was higher than in females. Further research was needed to

confirm our findings in specific sub-populations. Lastly, the samples were all of European

ancestry, making our findings may not apply to other populations.

5. Conclusion

345 Genetically predicted depression and schizophrenia were associated with elevated risk of

346 abdominal aortic aneurysm, and the known risk factors of abdominal aortic aneurysm played

347 mediation roles.

6. Declarations

6.1. Acknowledgement

350 None.

6.2. Ethical approval

The datasets used in this study were all publicly available from previous studies, and the

ethical approvals were obtained by the original researchers. Therefore, the ethical approval for
this study could be waived.

6.3. Consent for publication

Not applicable

6.4. Data availability

The datasets utilized in this study are publicly available; the details can be found in the
supplementary method.

6.5. Code availability

ldscr (https://github.com/bulik/ldsc), GNOVA (https://github.com/xtonyjiang/GNOVA), SU
PERGNOVA (https://github.com/qlu-lab/SUPERGNOVA), pleioFDR (https://github.com/pr
ecimed/pleiofdr), TwoSampleMR (https://mrcieu.github.io/TwoSampleMR/), MendelianRa
ndomization (https://github.com/cran/MendelianRandomization), and mr.raps (https://githu
b.com/qingyuanzhao/mr.raps).

6.6. Competing interests

The authors declare no conflict of interest.

6.7. Funding

This work receives no external funding.

6.8. Author contributions

Ming-Gang Deng: Conceptualization, Methodology, Data Curation, Formal Analysis, and
Writing - Original Draft; **Chen Chai:** Writing - Original Draft; Kai Wang: Writing - Original
Draft; **Zhi-Hui Zhao:** Writing – review & editing; **Jia-Qi Nie**: Writing – review & editing;
Fang Liu: Writing – review & editing; **Yuehui Liang:** Writing – review & editing; **Jiewei**
Liu: Conceptualization, and Writing - Original Draft.

7. Reference

1.   Summers KL, Kerut EK, Sheahan CM, Sheahan MG, 3rd. Evaluating the prevalence of
abdominal aortic aneurysms in the United States through a national screening database.
Journal of vascular surgery. 2021;73(1):61-68.
2.   Davis FM, Rateri DL, Daugherty A. Abdominal aortic aneurysm: novel mechanisms and
therapies. *Current opinion in cardiology*. 2015;30(6):566-573.
3.   Zhang F, Li K, Zhang W, et al. Ganglioside GM3 Protects Against Abdominal Aortic
Aneurysm by Suppressing Ferroptosis in Vascular Smooth Muscle Cells. *Circulation*.
2023.

4.   Deng MG, Liu F, Liang Y, Wang K, Nie JQ, Liu J. Association between frailty and
depression: A bidirectional Mendelian randomization study. *Science advances.*
2023;9(38):eadi3902.

5.   Deng M-G, Wang K, Liu F, Zhou X, Liu J. The Relationship Between Frailty and
Schizophrenia: A Genetic Association and Mendelian Randomization Study. 2023.

6.   Li GH, Cheung CL, Chung AK, et al. Evaluation of bi-directional causal association
between depression and cardiovascular diseases: a Mendelian randomization study.
Psychological medicine. 2022;52(9):1765-1776.

7.   Veeneman RR, Vermeulen JM, Abdellaoui A, et al. Exploring the Relationship Between
Schizophrenia and Cardiovascular Disease: A Genetic Correlation and Multivariable
Mendelian Randomization Study. *Schizophrenia bulletin.* 2022;48(2):463-473.

8.   So HC, Chau KL, Ao FK, Mo CH, Sham PC. Exploring shared genetic bases and causal
relationships of schizophrenia and bipolar disorder with 28 cardiovascular and
metabolic traits. *Psychological medicine.* 2019;49(8):1286-1298.

9.   Kim MH, Yoo JH, Cho HJ, et al. Increased depression risk in patients with abdominal
aortic aneurysm: a nationwide cohort study. *Annals of surgical treatment and research.*
2021;101(5):291-298.

10.   Nyrønning L, Stenman M, Hultgren R, Albrektsen G, Videm V, Mattsson E. Symptoms
of Depression and Risk of Abdominal Aortic Aneurysm: A HUNT Study. *Journal of the*
American Heart Association. 2019;8(21):e012535.

11.   Howard DM, Adams MJ, Clarke TK, et al. Genome-wide meta-analysis of depression
identifies 102 independent variants and highlights the importance of the prefrontal brain
regions. *Nature neuroscience.* 2019;22(3):343-352.

12.   Trubetskoy V, Pardiñas AF, Qi T, et al. Mapping genomic loci implicates genes and
synaptic biology in schizophrenia. *Nature.* 2022;604(7906):502-508.

13.   Stahl EA, Breen G, Forstner AJ, et al. Genome-wide association study identifies 30 loci
associated with bipolar disorder. *Nature genetics.* 2019;51(5):793-803.

14.   Roychowdhury T, Klarin D, Levin MG, et al. Genome-wide association meta-analysis
identifies risk loci for abdominal aortic aneurysm and highlights PCSK9 as a therapeutic
target. *Nature genetics.* 2023;55(11):1831-1842.

15.   Emdin CA, Khera AV, Kathiresan S. Mendelian Randomization. *Jama.*
2017;318(19):1925-1926.

16.   Deng MG, Cui HT, Nie JQ, Liang Y, Chai C. Genetic association between circulating
418   selenium level and the risk of schizophrenia in the European population: A two-sample
419   Mendelian randomization study. *Frontiers in nutrition.* 2022;9:969887.

420   17.   Deng MG, Liu F, Wang K, Liang Y, Nie JQ, Chai C. Genetic association between
421   coffee/caffeine consumption and the risk of obstructive sleep apnea in the European
422   population: a two-sample Mendelian randomization study. *European journal of nutrition.*
423   2023;62(8):3423-3431.

424   18.   Yu X, Deng MG, Tang ZY, Zhang ZJ. Urticaria and increased risk of rheumatoid
425   arthritis: a two-sample Mendelian randomisation study in European population. *Modern*
426   *rheumatology.* 2022;32(4):736-740.

427   19.   Bulik-Sullivan B, Finucane HK, Anttila V, et al. An atlas of genetic correlations across
428   human diseases and traits. *Nature genetics.* 2015;47(11):1236-1241.

429   20.   Lu Q, Li B, Ou D, et al. A Powerful Approach to Estimating Annotation-Stratified
430   Genetic Covariance via GWAS Summary Statistics. *American journal of human*
431   *genetics.* 2017;101(6):939-964.

432   21.   Zhang Y, Lu Q, Ye Y, et al. SUPERGNOVA: local genetic correlation analysis reveals
433   heterogeneous etiologic sharing of complex traits. *Genome biology.* 2021;22(1):262.

434   22.   Andreassen OA, Thompson WK, Schork AJ, et al. Improved detection of common
435   variants associated with schizophrenia and bipolar disorder using pleiotropy-informed
436   conditional false discovery rate. *PLoS genetics.* 2013;9(4):e1003455.

437   23.   Sanderson E. Multivariable Mendelian Randomization and Mediation. *Cold Spring*
438   *Harbor perspectives in medicine.* 2021;11(2).

439   24.   Kamat MA, Blackshaw JA, Young R, et al. PhenoScanner V2: an expanded tool for
440   searching human genotype-phenotype associations. *Bioinformatics (Oxford, England).*
441   2019;35(22):4851-4853.

442   25.   Thanigaimani S, Phie J, Quigley F, et al. Association of Diagnosis of Depression and
443   Small Abdominal Aortic Aneurysm Growth. *Annals of vascular surgery.*
444   2022;79:256-263.

445   26.   Márquez-Sánchez AC, Koltsova EK. Immune and inflammatory mechanisms of
446   abdominal aortic aneurysm. *Frontiers in immunology.* 2022;13:989933.

447   27.   Yuan Z, Lu Y, Wei J, Wu J, Yang J, Cai Z. Abdominal Aortic Aneurysm: Roles of
448   Inflammatory Cells. *Frontiers in immunology.* 2020;11:609161.

28.   Raffort J, Lareyre F, Clément M, Hassen-Khodja R, Chinetti G, Mallat Z. Monocytes
and macrophages in abdominal aortic aneurysm. *Nature reviews Cardiology.*
2017;14(8):457-471.

29.   Li Z, Kong W. Cellular signaling in Abdominal Aortic Aneurysm. *Cellular signalling.*
2020;70:109575.

30.   Tsuruda T, Kato J, Hatakeyama K, et al. Adventitial mast cells contribute to
pathogenesis in the progression of abdominal aortic aneurysm. *Circulation research.*
2008;102(11):1368-1377.

31.   Yamanouchi D, Morgan S, Kato K, Lengfeld J, Zhang F, Liu B. Effects of caspase
inhibitor on angiotensin II-induced abdominal aortic aneurysm in apolipoprotein
E-deficient mice. *Arteriosclerosis, thrombosis, and vascular biology.*
2010;30(4):702-707.

32.   Kaneko H, Anzai T, Morisawa M, et al. Resveratrol prevents the development of
abdominal aortic aneurysm through attenuation of inflammation, oxidative stress, and
neovascularization. *Atherosclerosis.* 2011;217(2):350-357.

33.   Berk M, Williams LJ, Jacka FN, et al. So depression is an inflammatory disease, but
where does the inflammation come from? *BMC medicine.* 2013;11:200.

34.   Halaris A. Inflammation and depression but where does the inflammation come from?
Current opinion in psychiatry. 2019;32(5):422-428.

35.   Beurel E, Toups M, Nemeroff CB. The Bidirectional Relationship of Depression and
Inflammation: Double Trouble. *Neuron.* 2020;107(2):234-256.

36.   Müller N. Inflammation in Schizophrenia: Pathogenetic Aspects and Therapeutic
Considerations. *Schizophrenia bulletin.* 2018;44(5):973-982.

37.   Khandaker GM, Cousins L, Deakin J, Lennox BR, Yolken R, Jones PB. Inflammation
and immunity in schizophrenia: implications for pathophysiology and treatment. *The*
lancet Psychiatry. 2015;2(3):258-270.

38.   Hartwig FP, Borges MC, Horta BL, Bowden J, Davey Smith G. Inflammatory
Biomarkers and Risk of Schizophrenia: A 2-Sample Mendelian Randomization Study.
JAMA psychiatry. 2017;74(12):1226-1233.

39.   Williams JA, Burgess S, Suckling J, et al. Inflammation and Brain Structure in
Schizophrenia and Other Neuropsychiatric Disorders: A Mendelian Randomization
Study. *JAMA psychiatry.* 2022;79(5):498-507.

40.   Bondy B, Baghai TC, Zill P, et al. Combined action of the ACE D- and the G-protein
beta3 T-allele in major depression: a possible link to cardiovascular disease? *Molecular*
psychiatry. 2002;7(10):1120-1126.
41.   Amare AT, Schubert KO, Klingler-Hoffmann M, Cohen-Woods S, Baune BT. The
genetic overlap between mood disorders and cardiometabolic diseases: a systematic
review of genome wide and candidate gene studies. *Translational psychiatry.*
2017;7(1):e1007.
42.   Gouweleeuw L, Naudé PJ, Rots M, DeJongste MJ, Eisel UL, Schoemaker RG. The role
of neutrophil gelatinase associated lipocalin (NGAL) as biological constituent linking
depression and cardiovascular disease. *Brain, behavior, and immunity.* 2015;46:23-32.
43.   Amadio P, Zarà M, Sandrini L, Ieraci A, Barbieri SS. Depression and Cardiovascular
Disease: The Viewpoint of Platelets. *International journal of molecular sciences.*
2020;21(20).
44.   Rødevand L, Rahman Z, Hindley GFL, et al. Characterizing the Shared Genetic
Underpinnings of Schizophrenia and Cardiovascular Disease Risk Factors. *The*
American journal of psychiatry. 2023;180(11):815-826.
45.   Mullins N, Forstner AJ, O'Connell KS, et al. Genome-wide association study of more
than 40,000 bipolar disorder cases provides new insights into the underlying biology.
Nature genetics. 2021;53(6):817-829.
500

Table 1. Locally correlated genomic regions between mental disorders and abdominal aortic aneurysm.

Trait1&Trait2	chr	start	end	rho	corr	h^2_1	h^2_2	var	m	*p*-value	*p*.adjust
Depression & AAA	9	126927204	128926989	7.98E-05	0.945	7.26E-05	9.81E-05	2.47E-10	644	3.78E-07	8.57E-04
Depression & AAA	10	106139294	107874146	6.17E-05	1.112	1.40E-05	2.19E-04	1.72E-10	758	2.58E-06	2.93E-03
Depression & AAA	11	42244050	42854352	4.31E-05	1.130	2.05E-05	7.07E-05	9.28E-11	277	7.83E-06	5.92E-03
Depression & AAA	18	77151867	78017074	4.62E-05	1.159	2.74E-05	5.79E-05	1.12E-10	323	1.30E-05	7.37E-03
SCZ & AAA	2	73174848	74121000	7.41E-05	0.907	1.61E-05	4.15E-04	3.73E-10	277	1.24E-04	2.84E-02
SCZ & AAA	2	232791948	233964632	-1.38E-04	-1.021	2.56E-05	7.19E-04	6.93E-10	514	1.44E-07	1.64E-04
SCZ & AAA	5	73508509	75240469	1.21E-04	0.792	9.10E-05	2.57E-04	9.36E-10	766	7.55E-05	2.46E-02
SCZ & AAA	10	106139294	107874146	1.16E-04	1.355	1.40E-05	5.21E-04	7.46E-10	759	2.28E-05	1.30E-02
SCZ & AAA	10	130282817	131192968	1.08E-04	1.746	3.02E-05	1.27E-04	7.75E-10	576	1.02E-04	2.59E-02
SCZ & AAA	11	16790799	18788126	1.76E-04	0.722	6.85E-05	8.64E-04	1.90E-09	974	5.51E-05	2.19E-02
SCZ & AAA	12	2176265	2886299	9.31E-05	0.815	6.39E-06	2.04E-03	5.36E-10	367	5.76E-05	2.19E-02
SCZ & AAA	14	102341650	104759919	2.23E-04	0.671	8.11E-05	1.37E-03	2.12E-09	801	1.28E-06	9.75E-04
SCZ & AAA	15	78640760	79259009	2.80E-04	0.777	1.88E-04	6.91E-04	1.27E-09	202	3.75E-15	8.56E-12
SCZ & AAA	17	820511	1480393	1.05E-04	1.043	2.83E-05	3.56E-04	7.27E-10	264	1.01E-04	2.59E-02
BIP & AAA	2	73174848	74121000	8.47E-05	1.036	1.61E-05	4.15E-04	4.64E-10	277	8.37E-05	3.01E-02
BIP & AAA	8	24457122	25461315	-2.03E-04	-0.944	7.38E-05	6.29E-04	1.71E-09	466	9.08E-07	1.04E-03
BIP & AAA	12	47505424	48267360	1.90E-04	1.252	5.57E-05	4.15E-04	1.27E-09	379	9.37E-08	2.14E-04
BIP & AAA	15	66948951	68640917	2.27E-04	1.046	9.96E-05	4.72E-04	2.86E-09	789	2.18E-05	1.10E-02
BIP & AAA	15	78640760	79259009	1.71E-04	0.873	1.88E-04	2.03E-04	1.63E-09	202	2.41E-05	1.10E-02
BIP & AAA	19	10030690	11279257	2.63E-04	0.585	1.35E-04	1.49E-03	4.52E-09	482	9.23E-05	3.01E-02
BIP & AAA	22	43187900	43645335	1.21E-04	1.231	1.47E-05	6.53E-04	8.14E-10	257	2.31E-05	1.10E-02

Abbreviations: SCZ, schizophrenia; AAA, abdominal aortic aneurysm; BIP, bipolar disorder; chr, chromosome; start/end, the start or end position of the genomic region from the input genome partition file; rho, the estimation of local genetic covariance; corr, the estimation of local genetic correlation; h^2_1/h^2_2, The estimation of local heritability of trait 1/2; var, the variance of the estimation of local genetic covariance; m, the number of SNPs involved in the estimation of local genetic covariance in the genomic region.

Figure Legend

Fig 1. An overview of study design

Fig 2. Global genetic correlation between mental disorders and abdominal aortic aneurysm.
LDSC, linkage disequilibrium score regression; GNOVA, GeNetic cOVariance Analyzer; SCZ,
schizophrenia; BIP, bipolar disorder; *** indicates $p < 0.001$; ** indicates $p < 0.01$; * indicates $p <$
0.05; [ns] indicates no significance.

Fig 3. ConjFDR Manhattan plots. The red dashed line represents the threshold of p-value for
conjFDR < 0.05, and the black points are the independent SNPs; SCZ, schizophrenia; BIP,
bipolar disorder; AAA, abdominal aortic aneurysm.

Fig 4. The univariate bidirectional causal inference between mental disorders and abdominal
aortic aneurysm. MDD, major depression disorder; SCZ, schizophrenia; BIP, bipolar disorder;
AAA, abdominal aortic aneurysm.

Fig 5. Mediation effects of risk factors in the causal associations of abdominal aortic
aneurysm with depression and schizophrenia. SNPs, single nucleotide polymorphisms; AAA,
abdominal aortic aneurysm; CA, coronary atherosclerosis.

Exposure	Outcome	Methods	OR (95% CI)	p-value		Cochran's Q	p value for heterogeneity* or pleiotropy†
Forward analyses							
Depression	AAA	IVW	1.270 (1.071, 1.504)	0.006		163.448	<0.001
		MR-Egger	0.865 (0.348, 2.149)	0.754			0.404
		Weighted median	1.229 (1.063, 1.420)	0.005			
		MR-cML	1.184 (1.040, 1.349)	0.011			
		MR-RAPS	1.302 (1.190, 1.424)	<0.001			
SCZ	AAA	IVW	1.047 (1.010, 1.084)	0.011		333.578	<0.001
		MR-Egger	1.018 (0.882, 1.174)	0.809			0.692
		Weighted median	1.041 (1.002, 1.082)	0.041			
		MR-cML	1.041 (1.004, 1.079)	0.029			
		MR-RAPS	1.049 (1.025, 1.074)	<0.001			
BIP	AAA	IVW	0.980 (0.896, 1.071)	0.653		32.604	0.002
		MR-Egger	1.165 (0.656, 2.069)	0.601			0.559
		Weighted median	1.024 (0.936, 1.121)	0.607			
		MR-cML	0.987 (0.915, 1.065)	0.737			
		MR-RAPS	0.978 (0.924, 1.035)	0.448			
Reverse analyses							
AAA	Depression	IVW	1.010 (0.995, 1.025)	0.190		183.629	<0.001
		MR-Egger	1.006 (0.969, 1.046)	0.743			0.844
		Weighted median	1.001 (0.982, 1.020)	0.897			
		MR-cML	1.010 (0.995, 1.024)	0.187			
		MR-RAPS	1.010 (0.998, 1.022)	0.097			
AAA	SCZ	IVW	1.007 (0.968, 1.048)	0.740		328.312	<0.001
		MR-Egger	0.989 (0.891, 1.096)	0.832			0.711
		Weighted median	1.004 (0.962, 1.048)	0.843			
		MR-cML	0.985 (0.951, 1.020)	0.390			
		MR-RAPS	1.007 (0.983, 1.032)	0.562			
AAA	BIP	IVW	0.989 (0.944, 1.037)	0.646		183.276	<0.001
		MR-Egger	1.005 (0.889, 1.137)	0.940			0.786
		Weighted median	1.011 (0.950, 1.075)	0.735			
		MR-cML	0.991 (0.946, 1.039)	0.704			
		MR-RAPS	0.989 (0.952, 1.027)	0.560			

*p-value for heterogeneity based on Cochran's Q statistic.

†p-value or pleiotropy based on MR-Egger regression intercept.

0.8 1 1.2 1.4 1.6