Spatial Multi-Omics Connect SHROOM3 and COL18A1 in Chronic Kidney Disease

Pukhraj S. Gaheer^{1,2}, Nikhil Uppal^{1,2}, Amy Paul³, Pedrum Mohammadi-Shemirani¹, Nicolas Perrot¹, Guillaume Paré^{1,2,3}, Darren Bridgewater³, Matthew B. Lanktree^{1,2,4,5}

- 1. Population Health Research Institute, Hamilton, Ontario, Canada
- Department of Health Research Methodology, Evidence & Impact, McMaster University, Hamilton, Ontario, Canada
- Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario, Canada
- 4. Department of Medicine, McMaster University, Hamilton, Ontario, Canada
- 5. Division of Nephrology, St. Joseph's Healthcare Hamilton, Hamilton, Ontario, Canada

Abstract word count: 232

Manuscript word count: 4237

Figures: 6

Tables: 2

Supplementary Tables: 15

Running Head: Kidney compartment-specific multi-omics of SHROOM3

Address for Correspondence: Matthew B. Lanktree St. Joseph's Healthcare Hamilton M446, 50 Charlton Ave. E. Hamilton, ON, Canada Email: lanktree@mcmaster.ca

Abstract

Shroom Family Member 3 (SHROOM3) encodes an actin-binding protein that impacts kidney development. Genome-wide association studies (GWAS) identified CKDassociated common variants around SHROOM3 and Shroom3 knock-out mice develop glomerular abnormalities. We sought to evaluate the impact of genetically predicted SHROOM3 expression on kidney traits and the circulating proteome, and validate findings in a mouse model. Genetic instruments for SHROOM3 expression in distinct kidney compartments (glomerular n=240, tubulointerstitial n=311) were constructed using single cell sequencing data from NephQTL2. Using two-sample Mendelian randomization, we evaluated the effects of glomerular and tubulointerstitial SHROOM3 expression on kidney traits and the concentration of 1,463 plasma proteins in the UK Biobank and CKDGen Consortium. Genetically predicted tubulointerstitial SHROOM3 expression colocalized with the genetic signals for eGFR and albuminuria. A 34% reduction in genetically predicted tubulointerstitial SHROOM3 expression was associated with a 0.3% increase in cross-sectional eGFR ($P = 6.8 \times 10^{-4}$), a 1.5% increase in albuminuria (P = 0.01), and a 2.2% reduction in plasma COL18A1 concentration ($P = 1.2 \times 10^{-5}$). In contrast, genetically predicted glomerular SHROOM3 expression showed neither colocalization nor significant Mendelian randomization results. Using immunofluorescence, heterozygous Shroom3 knockout mice had a concordant reduction of Col18a1 in their kidneys, primarily around the tubules. Thus, reduced tubulointerstitial SHROOM3 expression, but not glomerular, is associated with increased cross-sectional eGFR, increased uACR, and reduced plasma COL18A1 and Shroom3 knockout leads to reduced kidney Col18a1, agnostically linking SHROOM3 and COL18A1 in CKD pathogenesis.

Keywords: Mendelian randomization, Genomics, Tissue-specific, CKD

Lay Summary

The *SHROOM3* gene is consistently linked to kidney disease, but we have yet to fully understand why. We looked at how genetic changes affecting *SHROOM3* impact different regions of the kidney. We found that less *SHROOM3* in kidney tubules led to increased kidney filtration, but also increased leakage of protein into the urine. After looking at over 1000 proteins, we identified a new link between *SHROOM3* and a collagen protein called COL18A1. We then confirmed the link between SHROOM3 and COL18A1 by imaging the kidneys of mice designed to have less SHROOM3. Our results suggest an interaction of SHROOM3 and COL18A1 leads to increased pressure on the kidney filtration system.

Introduction

Chronic kidney disease (CKD) impacts 10% of the global population, culminating in heightened risk of kidney failure, cardiovascular disease, and death.¹ Populationscale genome-wide associations studies (GWAS) have identified genetic variants within and around Shroom Family Member 3 (SHROOM3) to be associated with CKD, crosssectional estimated glomerular filtration rate (eGFR), urinary albumin-to-creatinine ratio (uACR), and blood urea nitrogen (BUN).²⁻⁵ The associated variants are located in a linkage disequilibrium block around the transcription start site of SHROOM3, suggesting altered quantity of SHROOM3 gene expression could play a causal pathophysiological role.⁶ Transcriptome-wide association analyses, exploring links between gene expression and kidney phenotypes, have demonstrated an association between kidney SHROOM3 expression and cross-sectional eGFR.⁷ Analysis of RNA-sequencing from 50 human kidney transplant biopsies revealed a weak association between SHROOM3 expression in the glomerular and non-glomerular regions, but exhibiting opposing correlations with eGFR in each region (increased SHROOM3 expression in glomerular tissue and increased eGFR, P = 0.03; decreased SHROOM3 expression in nonglomerular tissue and increased eGFR, P = 0.05).⁸ During embryonic development, both *Shroom3* homozygous and heterozygous knockout mice exhibit abnormal nephron development and a Shroom3 dose-dependent reduction in glomerular number.⁹ Following acute kidney injury, Shroom3 heterozygous knockout mice displayed increased mortality, worsened kidney function, and heightened fibrosis.¹⁰ Studies in different animal models show similar results: post-natal Shroom3 knockdown induces albuminuria in mice, reintroducing the normal Shroom3 gene into hypertensive rats

3

alleviates albuminuria and glomerulosclerosis, and introducing wild-type *Shroom3* into *Shroom3*-deficient zebrafish remedies glomerular abnormalities.^{11,12} Despite these insights, the comprehensive impact of the *SHROOM3* gene on kidney disease, relevant biomarkers, and underlying mechanism remains incompletely understood.

Mendelian randomization (MR) is a powerful tool for evaluating causal relationships by utilizing the genetic determinants of a risk factor.^{13,14} If a genetic variant is associated with a lifelong increase in an exposure, and greater exposure is causative for a phenotype, then the variant associated with greater exposure should also be associated with the phenotype. MR has been commonly used to assess traditional risk factors such as lipids, uric acid, hypertension, and body mass index,¹⁵⁻¹⁸ but can also be used to assess molecular phenotypes such as gene expression as in transcriptomewide association studies (TWAS), and protein biomarkers as in proteome-wide MR studies (also referred to as proteo-genomic or proteome-wide association studies (PWAS)).^{19,20} Phenome-wide MR further explores the effects of genetically influenced changes in a risk factor on a wide range of phenotypes. By simultaneously investigating a broad range of phenotypes, pleiotropic effects (when a gene or genetic variant influences multiple seemingly unrelated traits) can be investigated to identify shared biological pathways. Colocalization provides evidence for the presence of a shared causal variant between traits, suggesting a shared biological mechanism underlying the association.²¹ MR examines the impact of an expected change in the *quantity* of the risk factor on a specific phenotype. Conversely, rare exonic variant analyses can explore the impact of changes in protein function.

We sought to further evaluate the impact of changes in *SHROOM3* quantity and function using population-scale biobank genotyping, sequencing, phenome-wide, and proteome-wide data, coupled with kidney compartment-specific gene expression data, MR, colocalization, and rare variant analyses, with validation of findings in a *Shroom3* knockout model (Figure 1).

Methods

Phenotype Definition in the UK Biobank

All UK biobank outcomes were defined using either: 1) diagnosis of acute, chronic, unspecified, or end-stage kidney disease using the International Classification of Diseases, Tenth Revision (ICD-10) (UK Biobank data field 41270) or algorithmicallydefined end-stage kidney disease (UK Biobank data field 42046), or 2) eGFR_{Crea} or eGFR_{Cys} less than 60 ml/min/1.73m² at baseline. eGFR_{Crea}, eGFR_{Cys}, and eGFR_{CreaCys} (eGFR calculated from serum cystatin C and creatinine) using the 2021 CKD-Epi equations,²² or 3) uACR. Urinary albumin was determined in a spot urine sample. uACR and eGFR were log-transformed and standardized. Individual-level analysis was restricted to white British ancestry participants in the UK Biobank. Plasma protein concentrations for 1463 proteins in 36,458 UK Biobank participants were standardized. Data analyses in UK Biobank were performed under application number 15255 and has approval from the Northwest Multi-Center Research Ethics Committee.

SHROOM3 eQTL Instrument Selection

As *SHROOM3* isn't expressed in lymphocytes, results from the expression quantitative trait loci (eQTL) genetics consortium (eQTLGen) arising from whole blood were not applicable.²³ Thus, our first genetic instrument representing multi-tissue *SHROOM3* expression was generated through an inverse variance weighted metaanalysis of eQTLs across 48 tissues in the Genotype-Tissue Expression (GTEx) Project v8. Tissues from GTEx with a sample size of \geq 70 were included, excluding whole blood due to the low *SHROOM3* expression. This instrument represented genetically predicted *SHROOM3* expression in every tissue.

We analyzed eQTLs from the distinct kidney tubulointerstitial and glomerular compartments using single cell sequencing data from NephQTL2. The Nephrotic Syndrome Study Network (NEPTUNE) cohort is an observational study that enrolls individuals with non-inflammatory glomerular diseases.²⁴ Using glomerular and tubulointerstitial data of 332 participants of the NEPTUNE cohort, the NephQTL group identified eQTL genetic variants associated with the expression of 24,474 genes from 240 glomerulus and 311 tubulointerstitial tissue samples.²⁴ 55% of participants in NEPTUNE self-reported race as White and 61% of participants were males. We excluded genetic variants more than 500 kilobases away from *SHROOM3*, focusing on cis-eQTLs.²⁵ Palindromic and ambiguous variants, and non-synonymous and splicing variants were removed using ANNOVAR and RefSeq annotation.^{26,27} Pruning and thresholding was done within 250 kilobase regions using a linkage disequilibrium threshold of $r^2 < 0.01$ to ensure variants were independent. r^2 was calculated using the 1000 genomes phase 3 European ancestry reference panel.²⁸ F-statistic was used to

evaluate the strength of each genetic instrument.²⁹ Each genetic instrument was passed through Phenoscanner to assess MR assumptions.³⁰

Two-Sample MR

Three assumptions must be met for a MR analysis to be valid: (1) genetic variants should explain a significant proportion of variation in the exposure (relevance); (2) genetic variants in the instrument should be independent of confounders (independence); and (3) genetic variants must not have an alternative direct association with the outcome (exclusion, or absence of pleiotropy).¹³ Careful selection of variants to be included in a genetic instrument and detailed sensitivity analyses are required to evaluate these assumptions. Independent genetic variants not in linkage disequilibrium are combined into scores at the individual-level (individual-level allele score MR), or the effects of numerous independent variants are jointly assessed using regression (summary-level MR).

After generating genetic instruments, outlier variants were removed using MR Pleiotropy RESidual Sum and Outlier (MR-PRESSO) analysis.³¹ Inverse variance weighted (IVW) MR was the primary analysis to infer the relationship between *SHROOM3* expression and kidney traits, with weighted median, MR-Egger, and MR-PRESSO serving as sensitivity analyses.^{32,33} A *P* value < 0.05 for the MR-Egger yintercept indicates directional pleiotropy. Statistical analyses were performed using R software version 4.3.0 (https://www.r-project.org) with the *TwoSampleMR* package (version 0.5.6).³⁴ All analyses were conducted in accordance with STROBE-MR: guidelines for strengthening the reporting of MR studies (Supplementary Table S1).

Summary-level MR Data

To evaluate eGFR and BUN, we utilized summary-level data from the 2021 Stanzick et al. GWAS of cross-sectional eGFR_{Crea}, eGFR_{Cys}, and BUN (n = 1,201,930) and summary-level GWAS data from 2022 Liu et al. trans-ethnic GWAS of eGFR_{Crea} (n = 1,508,659).^{3,4} Diabetes-stratified kidney function was also investigated (n with diabetes mellitus = 178,691).³⁵ To evaluate longitudinal eGFR decline we analyzed summary-level data from the 2022 Gorski et al. GWAS of longitudinal eGFR decline (n = 343,339).³⁶ The eGFR decline GWAS was unadjusted for baseline eGFR, avoiding spurious associations due to collider bias.³⁷ Summary statistics from the 2019 Teumer et al. European ancestry GWAS of uACR (n = 348,954) and 2021 Liu et al., European ancestry GWAS of kidney volume (n = 32,860) were also analyzed.^{5,38} CKD summarylevel data was obtained from the 2019 Wuttke et al. CKD Genetics (CKDGen) Consortium GWAS of stage 3 CKD or higher (eGFR_{Crea} < 60 ml/min/1.73m²) (41,395 cases and 439,303 controls) which does not include UK Biobank samples.² Thus, we also conducted an inverse variance-weighted meta-analysis of our individual-level MR results with Wuttke et al. results.

Individual-level Allele Score MR

The cumulative effect of these genetic variants was evaluated on 333,735 participants of the UK Biobank by performing linear and logistic regression between individual-level *SHROOM3* genetic risk score and kidney phenotypes adjusted for age, sex, assessment center, and the first 20 genetic principal components of ancestry. Individual-level genotype data was used to create an effect-weighted genetic risk score

representing genetically predicted *SHROOM3* expression. For each variant in the genetic instrument, expression-increasing alleles were coded as 0, 1, and 2 and multiplied by the normalized effect on *SHROOM3* expression.

1,463 protein biomarker concentrations were measured in plasma using proximity extension assays developed by Olink (https://www.olink.com; Uppsala, Sweden) in 36,458 European ancestry participants of the UK Biobank. We tested if a genetic risk score representing genetically altered *SHROOM3* expression was associated with altered biomarker concentration, adjusted for age, sex and genetic principal components of ancestry, using two-sample MR analysis. A Bonferroni-adjusted significance threshold for the proteome analysis was $3x10^{-5}$ (P < 0.05/1463).

Phenome-Wide MR

To evaluate the breadth of phenotypes impacted by modified *SHROOM3* expression, we performed an MR of 228 binary phenotypes in the UK Biobank. Data was obtained from the 2018 Zhou et al. GWAS catalog.³⁹ Analysis included European ancestry participants, phenotypes with a minimal case-to-control ratio of 0.01 and used a Bonferroni-corrected p-value threshold of $2x10^{-4}$ (P < 0.05/228).

Colocalization Analysis

Colocalization analysis to examine the overlap of genetic signals for *SHROOM3* expression with $eGFR_{Crea}$, $eGFR_{Cys}$, CKD, uACR, and BUN was performed using the Bayesian colocalization *coloc* R package (version 5.1).⁴⁰ Five hypotheses are tested: null hypothesis (H₀), no association with either outcome or exposure; H₁, association

with outcome, but not exposure; H₂, association with exposure but not outcome; H₃, association with exposure and outcome with distinct causal variants; H₄, shared causal variants associated with outcome and exposure.⁴⁰ A posterior probability for H₄ > 0.8 is strong evidence of colocalization. The prior probabilities $p_1 = p_2 = 10^{-4}$ and $p_{12} = 10^{-5}$ were set at a conservative threshold.⁴¹

Evaluation of Rare Exonic SHROOM3 Genetic Variants

We evaluated the prevalence of rare (minor allele frequency below 1%) predicted loss-of-function (nonsense, frameshift, and splice donor or acceptor) variants by performing a weight-based meta-analysis of allele counts across gnomAD v2.1.1 (non-TOPMED) (122,431 genomes; 13,300 exomes) and BRAVO (132,345 genomes) (bravo.sph.umich.edu/freeze8/hg38/).⁴² An ancestry-specific prevalence estimate of loss-of-function variants was also calculated in gnomAD. Next, rare pathogenic SHROOM3 variants were tested for association with kidney phenotypes in 157,528 UK Biobank participants with available phenotype and exome sequencing data. Pathogenic SHROOM3 variants included predicted loss-of-function variants and non-synonymous variants predicted deleterious by the Mendelian Clinically Applicable Pathogenicity algorithm (M-CAP score > 0.025).⁴³ The cohort allelic sum test is a rare variant association burden test that aggregates rare genetic variants in an individual into a dichotomous score.⁴⁴ A dichotomous score was incorporated in a linear regression model to test the association between SHROOM3 rare variant carrier status, and several kidney phenotypes adjusted for age, sex, assessment center, chip type, and the first 20 genetic principal components of ancestry.

10

Immunofluorescence Staining of Col18a1

We performed immunofluorescence staining of Col18a1 in Shroom3 heterozygous knockout mice (Shroom3^{Gt/+}) previously generated using a gene trap cassette.^{45,46} Whole kidneys from postnatal 3-month-old Shroom3^{Gt/+} and CD1 wild type mice were resected, fixed in 4% paraformaldehyde at 4°C for 48 hours, and embedded in paraffin. Kidneys were sectioned to 5 µm on a Leica microtome, mounted on Superfrost plus microscope slides (VWR, Mississauga, Ontario) and incubated at 37°C overnight. For immunofluorescence, samples were blocked with 7.5% normal goat serum and 4.5% bovine serum albumin at room temperature for 1 hour, then incubated with primary antibody Col18a1 (1:500), overnight at 4°C.^{47,48} An incubation buffer containing 1% PBS, 3% bovine serum albumin, 5% normal goat serum, and 0.3% Tween20 was used to dilute the primary antibody. Tissue samples were washed 3 times in PBS, followed by incubation with secondary antibody Alexa Fluor 488 anti-rabbit (Life Technologies, 1:300, Milton, Ontario) in incubation buffer at room temperature for 1 hour, washed in PBS, counterstained with DAPI (Sigma Aldrich, Oakville, Ontario), washed in PBS again, then fixed using Fluoromount mounting medium (Sigma Aldrich, Oakville, Ontario). Images were captured using the Olympus BX80 light and fluorescence microscope, and CellSens image acquisition software.

Results

SHROOM3 eQTL Genetic Instruments

Our multi-tissue *SHROOM3* genetic instrument did not colocalize with any of the assessed kidney phenotypes, displaying significant heterogeneity. Thus, we redirected

our focus to kidney compartment-specific genetic instruments. After clumping, removing palindromic variants, and removing the variant rs4859682 for violating the exclusion assumption, 12 independent variants were associated with tubulointerstitial *SHROOM3* expression and 10 independent variants were associated with glomerular *SHROOM3* expression (Supplementary Table S2 & S3). For the only overlapping variant, the minor allele (T) of variant rs28584831 reduced *SHROOM3* expression in both kidney compartments and was present in both genetic instruments. Using MR, we observed no association (P > 0.7) between either genetic instrument with *SHROOM3* expression in the opposite kidney compartment (Supplementary Table S4 and S8). Both genetic instruments were strong: the tubulointerstitial genetic instrument explained 27% of the variation in *SHROOM3* expression in the kidney tubulointerstitium, and the kidney glomeruli genetic instrument explained 30% of the variation in *SHROOM3* expression in the kidney Table S4 & S8).

Tubulointerstitial SHROOM3 MR Analysis

We found the tubulointerstitial *SHROOM3* expression genetic instrument was associated with cross-sectional eGFR_{Crea} ($\beta = 0.3\%$ increase per 34% relative decrease in *SHROOM3* genetic instrument, $P = 6.8\times10^{-4}$) and eGFR_{Cys} ($\beta = 0.4\%$ increase per 34% relative decrease in *SHROOM3* genetic instrument, $P = 1.4\times10^{-4}$) (Figure 2 & Supplementary Table 4). We obtained similar results in our trans-ancestry eGFR_{Crea} analysis ($\beta = 1.4\%$ increase per 34% relative decrease in *SHROOM3* genetic instrument, $P = 2.7\times10^{-3}$). Summary-level MR for eGFR were concordant with the individual-level allele score MR results in UK biobank (Supplementary Table S4 & S5).

The impact of genetically predicted tubulointerstitial *SHROOM3* expression on eGFR was comparable between individuals with or without diabetes (Supplementary Table S4). Genetically predicted tubulointerstitial *SHROOM3* expression was also associated with uACR (β = 1.5% increase per 34% relative decrease in *SHROOM3* genetic instrument, *P* = 0.01) (Figure 2). We did not observe an association between genetically predicted tubulointerstitial *SHROOM3* expression and longitudinal eGFR decline (*P* = 0.47), risk of CKD (52,204 CKD cases and 884,322 controls; *P* = 0.4), or kidney volume (*P* = 0.19) (Supplementary Table S4 & S6). Phenome-wide MR demonstrated no association between genetically predicted tubulointerstitial *SHROOM3* expression and any of the 228 phenotypes following Bonferroni correction (Supplementary Table S7).

Glomerular SHROOM3 MR Analysis

MR analysis indicated a nominal association between genetically predicted glomerular *SHROOM3* expression and cross-sectional eGFR_{Crea} in the opposite direction as the tubulointerstitial instrument (β = 0.13% decrease per 34% relative decrease in *SHROOM3* genetic instrument, *P* = 0.03; *P*_{heterogeneity} with tubulointerstitial instrument < 0.001) (Figure 3 & Supplementary Table S8). In contrast, genetically predicted glomerular *SHROOM3* expression and uACR were nominally associated in the same direction as the tubulointerstitial instrument (β = 1.03% increase per 34% relative decrease in *SHROOM3* genetic instrument, *P* = 0.03, *P*_{heterogeneity} with tubulointerstitial instrument = 0.5) (Supplementary Table S8). Individual-level allele score MR from the UK biobank did not show an association between genetically predicted glomerular *SHROOM3* expression and kidney phenotypes (Supplementary Table S9). We did not observe any association between genetically predicted glomerular *SHROOM3* expression and eGFR decline, BUN, CKD, or kidney volume (all P > 0.05). Moreover, our phenome-wide MR (Supplementary Table S10) did not reveal any phenotypes associated with genetically predicted glomerular *SHROOM3* expression.

Colocalization of SHROOM3 Genetic Variants and Kidney Outcomes

We observed strong evidence for colocalization of genetic signals for tubulointerstitial *SHROOM3* expression and eGFR_{Crea} (H₄ = 96%), eGFR_{Cys} (H₄ = 95%), uACR (H₄ = 83%), and kidney volume (H₄ = 92%; Figure 4). This colocalization suggests the presence of a shared causal variant influencing both tubulointerstitial *SHROOM3* expression and eGFR, uACR, and kidney volume. In contrast, there was no colocalization between genetically predicted glomerular *SHROOM3* expression and eGFR_{Crea} or uACR (H₄ < 80%) (Supplementary Table S11), nor tubulointerstitial *SHROOM3* expression with glomerular *SHROOM3* expression or multi-tissue *SHROOM3* expression.

SHROOM3 Rare Genetic Variant Analysis

No monogenic phenotype has been reported from rare pathogenic variants in *SHROOM3*. Homozygous *SHROOM3* loss-of-function variants were absent in the population sequencing data from gnomAD and BRAVO, including whole genome and exome data. Meta-analysis of 270,000 population sequencing participants revealed a prevalence rate of heterozygous loss-of-function *SHROOM3* variants at 0.07% (95% CI:

0.05% – 0.08%) in the general population (Supplementary Table S12), consistently across all eight available ancestries (Supplementary Table S13).

In UK Biobank exome sequencing data, we identified 3435 individuals carrying a rare loss-of-function or predicted pathogenic missense variant in *SHROOM3*. These variants were nominally associated with eGFR_{Cys} (β = 3.44% increase in trait in *SHROOM3* rare variant carriers, *P* = 0.03) but not with uACR (*P* = 0.26) (Table 1).

Tubulointerstitial SHROOM3 Impacts on the Circulating Proteome

In a novel analysis looking at the impact of genetically predicted *SHROOM3* expression on the circulating proteome, the three most significant associations with genetically predicted tubulointerstitial *SHROOM3* expression were with proteins encoded by genes located on chromosome 4 and within 500 kilobases of *SHROOM3* (*NAAA, SCARB2,* and *ART3*) (Table 2 & Supplementary Table S14). Reduced collagen type XVIII alpha one (COL18A1), a protein biomarker located on chromosome 21, was also associated with lower genetically predicted tubulointerstitial *SHROOM3* expression and surpassed Bonferroni-adjusted significance ($\beta = -2.15\%$ per 34% relative decrease in *SHROOM3* genetic instrument, $P = 1.2 \times 10^{-5}$). Genetically predicted glomerular *SHROOM3* expression was not associated with changes in a circulating biomarker (Supplementary Table S15).

Immunofluorescence Staining Validation of COL18A1 Association

The association of *SHROOM3* expression with COL18A1 was reinforced experimentally by visualizing the expression pattern of Col18a1 in the kidneys of 3-

month-old *Shroom3^{Gt/+}* knockout mice. Wild type mice kidneys demonstrated robust Col18a1 protein expression surrounding the parietal epithelium of the renal corpuscle (glomerulus and Bowman's capsule) and around the tubular epithelium. *Shroom3^{Gt/+}* knockout kidneys exhibited a similar, but reduced, pattern of Col18a1 protein expression around the renal corpuscle. Notably, Col18a1 expression was even more reduced around the cortical and medullary tubular epithelium (Figure 5). The staining patterns indicate a reduction in Col18a1 protein levels in the kidneys resulting from *Shroom3* heterozygous knockout.

Discussion

Variants that impact *SHROOM3* expression in the kidney tubulointerstitial and glomerular regions do not colocalize, suggesting different genetic regulatory mechanisms. Leveraging this distinction, we were able to evaluate genetic expression specific to these compartments by creating separate genetic instruments for targeted spatial evaluation in multi-omic MR analyses. We observed a 34% relative reduction in genetically predicted tubulointerstitial *SHROOM3* expression led to a 0.28% increase in cross-sectional eGFR, 1.52% increase in uACR, and no detectable association with eGFR decline nor prevalent CKD. Colocalization analysis supported genetically predicted tubulointerstitial expression, rather than glomerular, drove the regional GWAS associations. Evaluating 1,463 biomarkers in the human circulating proteome, lower genetically predicted *SHROOM3* expression led to less COL18A1 in the blood. We then validated this causal relationship by immunofluorescence staining the kidneys of

Shroom3 knockout mice, confirming the link and temporal sequence of reduced Col18a1 from reduced *Shroom3* expression.

Cross-sectional eGFR is determined by a combination of two factors: maximally attained eGFR, determined by nephron endowment, and the rate of eGFR decline due to age or disease.⁴⁹ Gorski *et al.*'s 2022 study categorized variants associated with eGFR into two groups: 1) those associated with both maximally attained eGFR and its subsequent decline, and 2) those associated with maximally attained eGFR but not with eGFR decline.³⁶ Three genes, *GATM*, *CPS1*, and *SHROOM3* fell into the second category, highlighting a potential role in affecting nephron endowment. Building on these findings, we found reduced genetically predicted *SHROOM3* expression in the tubulointerstitial is associated with increased uACR and increased cross-sectional eGFR rather than eGFR decline. The question arises: does *SHROOM3* influence nephron endowment and, if so, what is the result?

Prior research identified *SHROOM3* as a key factor in kidney development. Human embryonic kidneys show the highest *SHROOM3* expression in the nephron progenitor cells, which give rise to all segments of the nephron.^{50,51} Both *Shroom3* heterozygous and homozygous knockout mice demonstrate significant dose-dependent reductions in glomerular number compared to wild type mice at embryonic day 18.5, with many glomeruli becoming cystic or collapsing.⁹ Although the exact mechanism is unknown, it is possible that the reduction of *SHROOM3* in the nephron progenitor cells results in the improper development of nephrons. We also demonstrated an association between *SHROOM3* and COL18A1, an extracellular collagen protein with a known role in kidney development. In embryonic mice, the absence of Col18a1 leads to altered

17

kidney development, including a shorter ureteric tree, smaller kidney volume, reduced nephron count, and fewer terminal branch points and ureteric tips.⁴⁸ Although MR analysis did not reveal a significant association between tubulointerstitial *SHROOM3* expression and total kidney volume, genetic colocalization of the phenotypes suggests the possibility of a mediating factor, possibly COL18A1. The connection between Shroom3 and Col18a1 in the tubulointerstitial area of postnatal mice suggests changes in Shroom3 expression could impact nephron formation via Col18a1. One possible hypothesis is disruptions in the ability of developing and mature nephrons to connect to the extracellular matrix. The observed interactions between *SHROOM3* and COL18A1 warrant further investigation to understand the pathways that influence kidney morphology.

COL18A1 and its cleavage product, endostatin, an antiangiogenic protein derived from the C-terminal of COL18A1, have significant roles in kidney disease pathology and function.⁵² In young mice with induced glomerulonephritis, *Col18a1* knockout led to heightened inflammatory responses and a decrease in glomerular and interstitial vascular density.⁵³ A study of 7-month-old *Col18a1* knockout mice demonstrated podocyte foot process effacement and loosening of the proximal tubular basement membrane.⁵⁴ In adult diabetic mice, administering endostatin peptide resulted in significant reductions in glomerular hypertrophy, hyperfiltration, and albuminuria.⁵⁵ Although *SHROOM3* and COL18A1 have imperfectly understood individual roles in CKD, the interplay between both in CKD pathogenesis, embryonically and post-natally, warrants further exploration.

Altered nephron endowment exerts a distinctive impact on kidney phenotypes. According to the Brenner hypothesis, a diminished nephron count leads to a reduction in the total filtration surface area, prompting an adaptive increase in single nephron glomerular filtration rate (snGFR).^{56,57} This adaptive response leads to 'relative' hyperfiltration to normalize the overall eGFR despite a reduced number of functioning nephrons.⁵⁸ The increase in snGFR is relative to the amount of renal mass that has been lost. Over years or decades, an increased snGFR causes sclerosis in the glomeruli of the remaining nephrons, proteinuria, hypertension, and CKD. In contrast to CKD, where low eGFR is associated with higher ACR, hyperfiltration manifests initially with increased eGFR and elevated uACR.^{58,59} This pattern of increased eGFR and uACR is also observed in early stages of diabetic nephropathy and focal segmental glomerulosclerosis.^{59,60} Our findings support hyperfiltration, with reduced tubulointerstitial SHROOM3 expression associated with both increased uACR and cross-sectional eGFR, likely stemming from a reduction in nephron mass during kidney development (Figure 6).

The absence of homozygous loss-of-function *SHROOM3* variants in population sequencing of 270,000 human participants is consistent with neonatal lethality observed in *Shroom3* homozygous knockout mice.⁴⁵ Due to the lethality of homozygous knockout, our rare variant analysis was limited to individuals with heterozygous variants. Rather than examining the quantity of *SHROOM3*, rare exonic missense variants in *SHROOM3* would be expected to impact protein function and were nominally associated with higher cross-sectional eGFR, but were not found to be associated with prevalent CKD in the healthy UK Biobank population.

19

Despite widespread expression of *SHROOM3* RNA and protein across human tissues, our phenome-wide MR did not uncover significant associations between altered genetically predicted *SHROOM3* expression and any of the 228 examined dichotomous outcomes. The lack of findings may be attributable to the tissue specificity of the *SHROOM3* genetic instruments. Complex diseases are more susceptible to the influence of tissue-specific effects, particularly those tissues relevant to the disease.⁶¹ Our kidney compartment-specific genetic instruments are more potent in assessing relationships with kidney diseases compared to other traits. While a multi-tissue genetic instrument has more power and might theoretically represent gene expression in all tissues, in reality, serious heterogeneity arises from varying eQTL effects across tissues.^{62,63} Furthermore, recent expansions in human kidney single cell RNA sequencing databases have allowed for the development of sufficiently powered kidney compartment-specific genetic instruments.^{4,24}

Strengths of this work include large non-overlapping population-based cohorts, utilization of tissue compartment-specific eQTL genetic instruments for spatial multi-omics, integration of colocalization, phenome-wide, proteome-wide and rare variant analyses of *SHROOM3* and validation of findings in a *Shroom3* knockout mouse. There is an increasing recognition of the pivotal role of tissue-specific and cell-type-specific eQTLs in gene regulation, which are more likely to colocalize with and explain GWAS signals.⁶⁴⁻⁶⁶ Recently, tissue-specific analyses identified kidney tubulointerstitial *DAB2* expression as a causal risk factor for CKD, a connection not previously recognized when analyzing whole kidney expression.⁶⁶ Utilizing MR offers an advantage to GWAS, as it employs multiple variants to examine the correlation between the effects of these

20

variants on gene expression and their influence on various outcomes rather than evaluating each variant independently as in GWAS yielding a theoretical improvement in power. Furthermore, MR is inherently robust against reverse causation and confounding factors, a notable strength compared to conventional epidemiological methods.¹³ Limitations of this study include reliance on largely European ancestry GWAS data, though we did validate our findings with eGFR_{Crea} in a trans-ancestry GWAS, and the constrained sample size for the kidney compartment eQTL data.

In conclusion, human population and knockout mice studies support a novel interaction between SHROOM3 and COL18A1 leading to impaired kidney development and hyperfiltration.

Disclosures: MBL has received speaker and advisory fees from Otsuka, Reata, Bayer, and Sanofi Genzyme. GP has received consulting fees from Bayer, Sanofi, Bristol-Myers Squibb, Lexicomp, and Amgen and support for research through his institution from Sanofi and Bayer.

Funding: MBL is supported by a McMaster University Early Career Research Award and a CIHR project grant (201909-PJT). This work was supported by a Kidney Foundation of Canada grant to DB and MBL.

Role of the Funder: Funders had no role in concept and design, acquisition, analysis, or interpretation of data, drafting, review, or approval of the manuscript for publication.

Data Availability: CKD Genetics Consortium summary-level data is publicly available (https://ckdgen.imbi.uni-freiburg.de/) and UK Biobank individual-level data is available with application (https://www.ukbiobank.ac.uk/enable-your-research/apply-for-access). UK Biobank phenome-wide summary statistics are available (https://www.leelabsg.org/resources). eQTL summary-level data obtained from NephQTL2 and GTEx is also publicly available (http://nephqtl2.org) (https://gtexportal.org/home/).

Acknowledgements: We would like to thank Dr. Thomas Drysdale at the University of Western Ontario for gifting the heterozygous *Shroom3* knockout mice and Dr. Satu Kuure at the University of Helsinki and Dr. Takako Sasaki at Oita University for gifting the Col18a1 antibody. The authors thank the participants and investigators of the contributing consortiums, including NEPTUNE, UK Biobank, the CKD Genetics Consortium, and GTEx. The GTEx Project was supported by the Common Fund of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained from: the GTEx Portal on 11/11/2022.

Author Contributions: MBL, DB, and PSG designed the project. AP and DB performed the mice experiments and imaging. PSG performed all statistical analyses and drafted the manuscript. MBL revised the manuscript. All authors provided intellectual content and approved the final version of the manuscript.

List of Supplementary Materials

Supplementary Table S1. STrengthening the Reporting of OBservational studies in Epidemiology using Mendelian Randomization (STROBE-MR) Checklist.

Supplementary Table S2. Genetic instrument for tubulointerstitial SHROOM3 expression and glomerular SHROOM3 expression.

Supplementary Table S3. Phenoscanner V2 search of tubulointerstitial and glomerular genetic instrument variants.

Supplementary Table S4. Summary-level Mendelian randomization results using tubulointerstitial SHROOM3 genetic instrument.

Supplementary Table S5. Individual-level allele score Mendelian randomization using UK Biobank phenotype data and tubulointerstitial genetic instrument.

Supplementary Table S6. Inverse variance weighted meta-analysis of CKD Mendelian randomization results across CKD Genetics Consortium and UK Biobank.

Supplementary Table S7. Phenome-wide Mendelian randomization of 228 binary phenotypes in the UK Biobank using the tubulointerstitial genetic instrument.

Supplementary Table S8. Summary-level Mendelian randomization results using glomerular SHROOM3 genetic instrument.

Supplementary Table S9. Individual-level allele score Mendelian randomization using UK Biobank phenotype data and glomerular genetic instrument.

Supplementary Table S10. Phenome-wide Mendelian randomization of 228 binary phenotypes in the UK Biobank using the glomerular genetic instrument.

Supplementary Table S11. Colocalization of tubulointerstitial and glomerular genetic instruments with kidney phenotypes.

Supplementary Table S12. Prevalence of predicted loss-of-function SHROOM3 variants in the general population.

Supplementary Table S13. Prevalence of predicted loss-of-function SHROOM3 variants across eight ancestries present in gnomAD v2.1.

Supplementary Table S14. Proteome-wide individual-level allele score Mendelian randomization using UK Biobank biomarker data and tubulointerstitial genetic instrument.

Supplementary Table S15. Proteome-wide individual-level allele score Mendelian randomization using UK Biobank biomarker data and glomerular genetic instrument.

References

- 1. Kovesdy CP. Epidemiology of chronic kidney disease: an update 2022. Kidney Int Suppl (2011). 2022;12(1):7-11. doi:10.1016/j.kisu.2021.11.003
- Wuttke M, Li Y, Li M, et al. A catalog of genetic loci associated with kidney function from analyses of a million individuals. *Nat Genet.* 2019;51(6):957-972. doi:10.1038/s41588-019-0407-x
- Stanzick KJ, Li Y, Schlosser P, et al. Discovery and prioritization of variants and genes for kidney function in >1.2 million individuals. *Nat Commun.* 2021;12(1):4350. Published 2021 Jul 16. doi:10.1038/s41467-021-24491-0
- Liu H, Doke T, Guo D, et al. Epigenomic and transcriptomic analyses define core cell types, genes and targetable mechanisms for kidney disease. *Nat Genet*. 2022;54(7):950-962. doi:10.1038/s41588-022-01097-w
- 5. Teumer A, Li Y, Ghasemi S, et al. Genome-wide association meta-analyses and fine-mapping elucidate pathways influencing albuminuria. *Nat Commun*. 2019;10(1):4130. Published 2019 Sep 11. doi:10.1038/s41467-019-11576-0
- Paul A, Lawlor A, Cunanan K, et al. The Good and the Bad of SHROOM3 in Kidney Development and Disease: A Narrative Review. Canadian Journal of Kidney Health and Disease. 2023;10. doi:10.1177/20543581231212038
- Hellwege JN, Velez Edwards DR, Giri A, et al. Mapping eGFR loci to the renal transcriptome and phenome in the VA Million Veteran Program. *Nat Commun.* 2019;10(1):3842. Published 2019 Aug 26. doi:10.1038/s41467-019-11704-w
- Wei C, Banu K, Garzon F, et al. SHROOM3-FYN Interaction Regulates Nephrin Phosphorylation and Affects Albuminuria in Allografts. *J Am Soc Nephrol.* 2018;29(11):2641-2657. doi:10.1681/ASN.2018060573
- Khalili H, Sull A, Sarin S, et al. Developmental Origins for Kidney Disease Due to Shroom3 Deficiency. J Am Soc Nephrol. 2016;27(10):2965-2973. doi:10.1681/ASN.2015060621
- 10. Li A, Cunanan J, Khalili H, et al. Shroom3, a Gene Associated with CKD, Modulates Epithelial Recovery after AKI. *Kidney360*. 2021;3(1):51-62. Published 2021 Nov 3. doi:10.34067/KID.0003802021
- Banu K, Lin Q, Basgen JM, et al. AMPK mediates regulation of glomerular volume and podocyte survival. *JCI Insight*. 2021;6(19):e150004. Published 2021 Oct 8. doi:10.1172/jci.insight.150004
- 12. Yeo NC, O'Meara CC, Bonomo JA, et al. Shroom3 contributes to the maintenance of the glomerular filtration barrier integrity. *Genome Res.* 2015;25(1):57-65. doi:10.1101/gr.182881.114
- Burgess S, Davey Smith G, Davies NM, et al. Guidelines for performing Mendelian randomization investigations. *Wellcome Open Res.* 2020;4:186. Published 2020 Apr 28. doi:10.12688/wellcomeopenres.15555.2
- Dobrijevic E, van Zwieten A, Kiryluk K, Grant AJ, Wong G, Teixeira-Pinto A. Mendelian randomization for nephrologists. *Kidney Int*. 2023;104(6):1113-1123. doi:10.1016/j.kint.2023.09.016
- 15. Lanktree MB, Thériault S, Walsh M, Paré G. HDL Cholesterol, LDL Cholesterol, and Triglycerides as Risk Factors for CKD: A Mendelian Randomization Study. *Am J Kidney Dis*. 2018;71(2):166-172. doi:10.1053/j.ajkd.2017.06.011

- Park S, Lee S, Kim Y, et al. Mendelian randomization reveals causal effects of kidney function on various biochemical parameters. *Commun Biol.* 2022;5(1):713. Published 2022 Jul 18. doi:10.1038/s42003-022-03659-4
- 17. Yu Z, Coresh J, Qi G, et al. A bidirectional Mendelian randomization study supports causal effects of kidney function on blood pressure. *Kidney Int.* 2020;98(3):708-716. doi:10.1016/j.kint.2020.04.044
- 18. Kjaergaard AD, Teumer A, Witte DR, et al. Obesity and Kidney Function: A Two-Sample Mendelian Randomization Study. *Clin Chem.* 2022;68(3):461-472. doi:10.1093/clinchem/hvab249
- 19. Pietzner M, Wheeler E, Carrasco-Zanini J, et al. Mapping the proteo-genomic convergence of human diseases. *Science*. 2021;374(6569):eabj1541. doi:10.1126/science.abj1541
- Schlosser P, Zhang J, Liu H, et al. Transcriptome- and proteome-wide association studies nominate determinants of kidney function and damage. *Genome Biol.* 2023;24(1):150. Published 2023 Jun 26. doi:10.1186/s13059-023-02993-y
- 21. Zuber V, Grinberg NF, Gill D, et al. Combining evidence from Mendelian randomization and colocalization: Review and comparison of approaches. *Am J Hum Genet*. 2022;109(5):767-782. doi:10.1016/j.ajhg.2022.04.001
- 22. Inker LA, Eneanya ND, Coresh J, et al. New Creatinine- and Cystatin C-Based Equations to Estimate GFR without Race. *The New England Journal of Medicine*. 2021;385(19):1737-1749. doi:https://doi.org/10.1056/NEJMoa2102953
- 23. Võsa U, Claringbould A, Westra HJ, et al. Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and polygenic scores that regulate blood gene expression. *Nat Genet*. 2021;53(9):1300-1310. doi:10.1038/s41588-021-00913-z
- 24. Han SK, McNulty MT, Benway CJ, et al. Mapping genomic regulation of kidney disease and traits through high-resolution and interpretable eQTLs. *Nat Commun*. 2023;14(1):2229. Published 2023 Apr 19. doi:10.1038/s41467-023-37691-7
- 25. Gkatzionis A, Burgess S, Newcombe PJ. Statistical methods for cis-Mendelian randomization with two-sample summary-level data. *Genet Epidemiol.* 2023;47(1):3-25. doi:10.1002/gepi.22506
- 26. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* 2010;38(16):e164. doi:10.1093/nar/gkq603
- 27. O'Leary NA, Wright MW, Brister JR, et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016;44(D1):D733-D745. doi:10.1093/nar/gkv1189
- 28.1000 Genomes Project Consortium, Auton A, Brooks LD, et al. A global reference for human genetic variation. *Nature*. 2015;526(7571):68-74. doi:10.1038/nature15393
- 29. Pierce BL, Ahsan H, Vanderweele TJ. Power and instrument strength requirements for Mendelian randomization studies using multiple genetic variants. *Int J Epidemiol*. 2011;40(3):740-752. doi:10.1093/ije/dyq151

- 30. Kamat MA, Blackshaw JA, Young R, et al. PhenoScanner V2: an expanded tool for searching human genotype-phenotype associations. *Bioinformatics*. 2019;35(22):4851-4853. doi:10.1093/bioinformatics/btz469
- Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases [published correction appears in Nat Genet. 2018 Aug;50(8):1196]. *Nat Genet.* 2018;50(5):693-698. doi:10.1038/s41588-018-0099-7
- 32. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol.* 2016;40(4):304-314. doi:10.1002/gepi.21965
- Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. Int J Epidemiol. 2015;44(2):512-525. doi:10.1093/ije/dyv080
- 34. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic causal inference across the human phenome. *Elife*. 2018;7:e34408. Published 2018 May 30. doi:10.7554/eLife.34408
- 35. Winkler TW, Rasheed H, Teumer A, et al. Differential and shared genetic effects on kidney function between diabetic and non-diabetic individuals. *Commun Biol.* 2022;5(1):580. Published 2022 Jun 13. doi:10.1038/s42003-022-03448-z
- 36. Gorski M, Rasheed H, Teumer A, et al. Genetic loci and prioritization of genes for kidney function decline derived from a meta-analysis of 62 longitudinal genomewide association studies. *Kidney Int.* 2022;102(3):624-639. doi:10.1016/j.kint.2022.05.021
- 37. Khan A, Kiryluk K. Kidney disease progression and collider bias in GWAS. *Kidney Int*. 2022;102(3):476-478. doi:10.1016/j.kint.2022.06.018
- 38. Liu Y, Basty N, Whitcher B, et al. Genetic architecture of 11 organ traits derived from abdominal MRI using deep learning. *Elife*. 2021;10:e65554. Published 2021 Jun 15. doi:10.7554/eLife.65554
- 39. Zhou W, Nielsen JB, Fritsche LG, et al. Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nat Genet*. 2018;50(9):1335-1341. doi:10.1038/s41588-018-0184-y
- 40. Giambartolomei C, Vukcevic D, Schadt EE, et al. Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet*. 2014;10(5):e1004383. Published 2014 May 15. doi:10.1371/journal.pgen.1004383
- 41. Wallace C. Eliciting priors and relaxing the single causal variant assumption in colocalisation analyses. *PLoS Genet*. 2020;16(4):e1008720. Published 2020 Apr 20. doi:10.1371/journal.pgen.1008720
- Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans [published correction appears in Nature. 2021 Feb;590(7846):E53] [published correction appears in Nature. 2021 Sep;597(7874):E3-E4]. *Nature*. 2020;581(7809):434-443. doi:10.1038/s41586-020-2308-7

- 43. Jagadeesh KA, Wenger AM, Berger MJ, et al. M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. *Nat Genet*. 2016;48(12):1581-1586. doi:10.1038/ng.3703
- 44. Morgenthaler S, Thilly WG. A strategy to discover genes that carry multi-allelic or mono-allelic risk for common diseases: a cohort allelic sums test (CAST). *Mutat Res.* 2007;615(1-2):28-56. doi:10.1016/j.mrfmmm.2006.09.003
- 45. Hildebrand JD, Soriano P. Shroom, a PDZ domain-containing actin-binding protein, is required for neural tube morphogenesis in mice. *Cell*. 1999;99(5):485-497. doi:10.1016/s0092-8674(00)81537-8
- 46. Lawlor A, Cunanan K, Cunanan J, et al. Minimal Kidney Disease Phenotype in Shroom3 Heterozygous Null Mice. Can J Kidney Health Dis. 2023;10:20543581231165716. Published 2023 Jun 7. doi:10.1177/20543581231165716
- 47. Lian X, Bond JS, Bharathy N, et al. Defining the Extracellular Matrix of Rhabdomyosarcoma. *Front Oncol.* 2021;11:601957. Published 2021 Feb 23. doi:10.3389/fonc.2021.601957
- 48. Rinta-Jaskari MM, Naillat F, Ruotsalainen HJ, et al. Temporally and spatially regulated collagen XVIII isoforms are involved in ureteric tree development via the TSP1-like domain. *Matrix Biol.* 2023;115:139-159. doi:10.1016/j.matbio.2023.01.001
- 49. Lanktree MB. Incorporating Linear Mixed Models into GWAS of Kidney Function Decline. *J Am Soc Nephrol*. 2023;34(9):1473-1475. doi:10.1681/ASN.00000000000175
- 50. Wu H, Malone AF, Donnelly EL, et al. Single-Cell Transcriptomics of a Human Kidney Allograft Biopsy Specimen Defines a Diverse Inflammatory Response. J Am Soc Nephrol. 2018;29(8):2069-2080. doi:10.1681/ASN.2018020125
- 51. Lindström NO, De Sena Brandine G, Tran T, et al. Progressive Recruitment of Mesenchymal Progenitors Reveals a Time-Dependent Process of Cell Fate Acquisition in Mouse and Human Nephrogenesis. *Dev Cell*. 2018;45(5):651-660.e4. doi:10.1016/j.devcel.2018.05.010
- 52. Lanktree MB, Perrot N, Smyth A, et al. A novel multi-ancestry proteome-wide Mendelian randomization study implicates extracellular proteins, tubular cells, and fibroblasts in estimated glomerular filtration rate regulation. *Kidney Int.* 2023;104(6):1170-1184. doi:10.1016/j.kint.2023.08.025
- 53. Hamano Y, Okude T, Shirai R, et al. Lack of collagen XVIII/endostatin exacerbates immune-mediated glomerulonephritis. *J Am Soc Nephrol*. 2010;21(9):1445-1455. doi:10.1681/ASN.2009050492
- 54. Kinnunen AI, Sormunen R, Elamaa H, et al. Lack of collagen XVIII long isoforms affects kidney podocytes, whereas the short form is needed in the proximal tubular basement membrane. *J Biol Chem.* 2011;286(10):7755-7764. doi:10.1074/jbc.M110.166132
- 55. Ichinose K, Maeshima Y, Yamamoto Y, et al. Antiangiogenic endostatin peptide ameliorates renal alterations in the early stage of a type 1 diabetic nephropathy model [published correction appears in Diabetes. 2006 Sep;55(9):2665]. *Diabetes*. 2005;54(10):2891-2903. doi:10.2337/diabetes.54.10.2891

- 56. Brenner BM, Lawler EV, Mackenzie HS. The hyperfiltration theory: a paradigm shift in nephrology. *Kidney Int*. 1996;49(6):1774-1777. doi:10.1038/ki.1996.265
- 57. Gurusinghe S, Tambay A, Sethna CB. Developmental Origins and Nephron Endowment in Hypertension. *Front Pediatr*. 2017;5:151. Published 2017 Jun 29. doi:10.3389/fped.2017.00151
- 58. Helal I, Fick-Brosnahan GM, Reed-Gitomer B, Schrier RW. Glomerular hyperfiltration: definitions, mechanisms and clinical implications. *Nat Rev Nephrol.* 2012;8(5):293-300. Published 2012 Feb 21. doi:10.1038/nrneph.2012.19
- 59. Ellis JW, Chen MH, Foster MC, et al. Validated SNPs for eGFR and their associations with albuminuria. *Hum Mol Genet*. 2012;21(14):3293-3298. doi:10.1093/hmg/dds138
- 60. Rosenberg AZ, Kopp JB. Focal Segmental Glomerulosclerosis [published correction appears in Clin J Am Soc Nephrol. 2018 Dec 7;13(12):1889]. *Clin J Am Soc Nephrol.* 2017;12(3):502-517. doi:10.2215/CJN.05960616
- 61. Fu J, Wolfs MG, Deelen P, et al. Unraveling the regulatory mechanisms underlying tissue-dependent genetic variation of gene expression. *PLoS Genet*. 2012;8(1):e1002431. doi:10.1371/journal.pgen.1002431
- 62. Nica AC, Parts L, Glass D, et al. The architecture of gene regulatory variation across multiple human tissues: the MuTHER study. *PLoS Genet*. 2011;7(2):e1002003. Published 2011 Feb 3. doi:10.1371/journal.pgen.1002003
- 63. GTEx Consortium; Laboratory, Data Analysis & Coordinating Center (LDACC)— Analysis Working Group; Statistical Methods groups—Analysis Working Group; Genetic effects on gene expression across human tissues [published correction appears in Nature. 2017 Dec 20;:]. *Nature*. 2017;550(7675):204-213. doi:10.1038/nature24277
- 64. Dimas AS, Deutsch S, Stranger BE, et al. Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science*. 2009;325(5945):1246-1250. doi:10.1126/science.1174148
- Donovan MKR, D'Antonio-Chronowska A, D'Antonio M, Frazer KA. Cellular deconvolution of GTEx tissues powers discovery of disease and cell-type associated regulatory variants [published correction appears in Nat Commun. 2020 Sep 1;11(1):4426]. *Nat Commun.* 2020;11(1):955. Published 2020 Feb 19. doi:10.1038/s41467-020-14561-0
- 66. Qiu C, Huang S, Park J, et al. Renal compartment-specific genetic variation analyses identify new pathways in chronic kidney disease. *Nat Med.* 2018;24(11):1721-1731. doi:10.1038/s41591-018-0194-4
- 67. Prokop JW, Yeo NC, Ottmann C, et al. Characterization of Coding/Noncoding Variants for *SHROOM3* in Patients with CKD. *J Am Soc Nephrol*. 2018;29(5):1525-1535. doi:10.1681/ASN.2017080856

Tables

Table 1: Rare variant association analysis of *SHROOM3* pathogenic variants shows weak association with estimated glomerular filtration rate using serum cystatin C (eGFR_{cys}). Rare variants were classified as predicted loss-of-function (nonsense, frameshift, and splice donor or acceptor) variants or missense variants with an MCAP score of > 0.025 indicating pathogenicity. Beta (β) represents % change in trait if carrying a *SHROOM3* rare variant. N represents the number of samples included in the analysis. 3,435 rare pathogenic variants were present in the UK Biobank exome sequencing database. N, sample size; CKD, chronic kidney disease; uACR, urinary albumin-to-creatinine ratio eGFR_{Crea}, estimated glomerular filtration rate from serum creatinine; eGFR_{CreaCys}, estimated glomerular filtration rate from serum cystatin C and creatinine.

Outcome	OR/ß	SE	Р	Cases	Controls	Ν
СКД	0.90	0.11	0.36	4,854	149,643	154,497
uACR	3.50%	0.03	0.26			48,516
eGFR _{Cys}	3.44%	0.02	0.03			154,433
eGFR _{Crea}	0.80%	0.02	0.63			154,354
eGFR _{CreaCys}	2.76%	0.02	0.09			154,238

Table 2: The impact of genetically predicted tubulointerstitial *SHROOM3* expression on proteome-wide serum biomarkers in European ancestry UK Biobank participants. Beta (β) represents percent change in biomarker per 34% relative decrease in *SHROOM3* genetic instrument score. N represents sample size. Chr represents chromosome of gene encoding biomarker. *NAAA*, *SCARB2*, and *ART3* are genes located adjacent to the *SHROOM3* gene (chr4:77,356,253-77,704,406). Only biomarkers surpassing the Bonferroni-corrected threshold of *P* < 3x10⁻⁵ are shown. Proteome-wide results can be found in Supplementary Table S14.

Biomarker		Transcription Start Site (b37)	ß	95% CI	Р	Ν
NAAAN-Acylethanolamine Acid Amidase (NAAA)	4	76,831,809	-7.64%	[-8.65%, -6.63%]	6.6x10 ⁻⁵⁰	36,458
Scavenger Receptor Class B Member 2 (SCARB2)	4	77,079,890	2.39%	[1.44%, 3.35%]	9.5x10 ⁻⁷	36,458
ADP-Ribosyltransferase 3 (ART3)	4	76,932,337	-2.17%	[-3.12%, -1.25%]	4.3x10 ⁻⁶	36,458
Collagen Type XVIII Alpha 1 Chain (COL18A1)	21	46,825,052	-2.15%	[-3.11%, -1.19%]	1.2x10 ⁻⁵	36,458

Figure Captions

Figure 1: Overall study design of a spatial multi-omic investigation of SHROOM3 quantity and function. Genetic variants affecting glomerular and tubulointerstitial gene expression were downloaded from GTEx and NephQTL2 database. Kidney phenotype summary statistics retrieved from the CKD Genetics Consortium or calculated in UK Biobank. n, sample size; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; uACR, urinary albumin-to-creatinine ratio; BUN, blood urea nitrogen. Created with BioRender.com.

Figure 2: Two-sample Mendelian randomization plots of tubulointerstitial *SHROOM3* expression on (A) estimated glomerular filtration rate (eGFR_{Crea})³ and (B) urinary albumin-to-creatinine ratio (uACR)⁵. Each dot represents a genetic variant within the genetic instrument. X axis represents standard deviation (SD) change in *SHROOM3* expression. Y axis represents the SD change in kidney trait. Beta represents the percent change in trait per 34% relative decrease in *SHROOM3* genetic instrument score. Full results can be found in Supplementary Table S4.

Figure 3: Forest plot comparing the impact of *SHROOM3* expression on kidney traits in the glomerulus versus the tubulointerstitial region reveals compartmentspecific effects. Beta represents the percent change in each trait per one standard deviation decrease in the *SHROOM3* genetic instrument score. Results depicted are from the inverse variance weighted analysis. Heterogeneity P-Value determined using Cochrane's Q statistic. eGFR_{Cys}, estimated glomerular filtration rate from serum cystatin C; eGFR_{Crea}, estimated glomerular filtration rate from serum creatinine; uACR, urinary albumin-to-creatinine ratio; BUN, blood urea nitrogen.

Figure 4: Colocalization of genetically predicted (A) tubulointerstitial *SHROOM3* expression with (B) estimated glomerular filtration rate (eGFRCrea),³ (C) kidney volume,³⁸ (D) urinary albumin-to-creatinine ratio (uACR),⁵ (E) glomerular *SHROOM3* expression, and (F) multi-tissue *SHROOM3* expression reveals compartment-specific colocalization. Degree of redness represents the posterior probability of colocalization of the genetic variant. Purple dots represent genetic variants included in *SHROOM3* genetic instruments. The cyan shaded region represents the location of the *SHROOM3* gene on chromosome 4, while the dashed cyan lines represent the three transcription start sites within *SHROOM3* corresponding to three unique *SHROOM3* isoforms.⁶⁷ Colocalization results can be found in Supplementary Table S11.

Figure 5: Immunofluorescence staining of Col18a1 protein in *Shroom3* **heterozygous knockout mice shows reduced Col18a1 protein expression around the kidney glomeruli and tubules.** Col18a1 immunofluorescence staining was performed in adult 3-month-old wild type and *Shroom3* heterozygous knockout (*Shroom3*^{Gt/+}) kidneys (n=2). Fluorescent green staining represents Col18a1 protein expression. **(A-C)** In wild type kidneys, Col18a1 protein expression prominently surrounds the renal corpuscle (glomerulus and Bowman's capsule) (arrow) and tubules

(arrowhead) in the cortical and medullary kidney regions. **(D-F)** *Shroom3*^{*Gt*/+} kidneys show substantially reduced expression of Col18a1 around the glomeruli and even less expression around the tubules. Scale bars = 100 µm (A, D). Scale bars = 50 µm (B, E). Scale bars = 20 µm (C, F).

Figure 6: Proposed pathway depicting the impact of *SHROOM3* gene expression on estimated glomerular filtration rate (eGFR) and urinary albumin-to-creatinine ratio (uACR) through kidney development and hyperfiltration. The association of *Shroom3* expression with altered kidney development was previously demonstrated in Khalili et al., 2016,⁹ while the association of Col18a1 expression with altered kidney development was shown in Rinta-Jaskari et al., 2023.⁴⁸ Created with BioRender.com.





Phenotype	ß [95% CI]					
eGFRCrea	0.28% [0.44; 0.12] -0.13% [-0.01; -0.25]	<				
eGFRCys	0.39% [0.59; 0.19] -0.02% [0.23; -0.27]	<				
eGFR Decline	0.74% [2.77; -1.29] 1.03% [2.78; -0.72]					
eGFR (w/ Diabetes)	0.49% [0.98; 0] -0.15% [0.34; -0.64]					
eGFR (w/o Diabetes)	0.25% [0.48; 0.02] -0.16% [-0.03; -0.29]					
UACR	1.52% [2.69; 0.35] 1.03% [1.93; 0.13]					
BUN	-0.24% [0.06; -0.54] 0.1% [0.31; -0.11]					
Kidney Volume	2.51% [6.43; -1.41] -1.74% [0.73; -4.21]					





Variant Position

Variant Position



























Brenner Hypothesis

Altered Kidney Development ↓ Nephron or Glomerular Count ↓ Kidney Volume

