Bacterial Bloodstream Infections in Cameroon: A Systematic Review and Meta-Analysis of
Prevalence, and Antibiotic Resistance.

Moise Matakone[1*‡], Patrice Landry Koudoum[2‡], Ravalona Jessica Zemtsa[3], Sen Claudine Henriette
Ngomtcho[2,4], Isaac Dah[1,5] and Michel Noubom[2,7]

[1]National Veterinary Laboratory (LANAVET), Yaoundé, Cameroon;
[2]Faculty of Medicine and Pharmaceutical Sciences, University of Dschang, Dschang P.O. Box 96,
Cameroon;
[3]Department of Biochemistry, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I,
Yaoundé, Cameroon;
[4]National Public Health Laboratory, Yaoundé, Cameroon;
[5]School of Veterinary Medicine and Science, University of Ngaoundéré, Ngaoundéré, Cameroon;
[6]Annex Regional Hospital of Dschang (ARHD), Dschang, Cameroon

*Corresponding author: moisematakone19@gmail.com;
‡ These authors contributed equally.

Abstract

Background: The paucity of data on the epidemiology of bloodstream infection (BSI) in Low-and middle-income countries (LMICs) countries hampers its effective prevention and management. This review sought to provide the current prevalence, bacteriological and antimicrobial resistance profiles of bacteria implicated in BSI in Cameroon.

Methods: PubMed and Google Scholar databases were searched to identify relevant articles and screened according to the PRISMA guidelines. Data was analysed using comprehensive meta-analysis software. The I^2 was used to evaluate heterogeneity between studies, funnel plot and Egger's test to evaluate publication bias and random-effects analysis models to calculate pooled prevalence.

Results: The 10 eligible studies were carried out in only two of the 10 regions of Cameroon and investigated 4,223 participants, of whom 920 were children under two years old. The overall pooled prevalence of bacterial BSI was 26.36% (95%CI= 17.51%−37.64%). *Klebsiella* spp. (28.20%; 95%CI= 2.80%−63.60%), *Escherichia coli* (18.94%; 95%CI= 13.02%−26.72%) and *Staphylococcus aureus* (14.40%; 95%CI= 8.31%−23.81%) were the most incriminated bacteria species. *E. coli* and *Klebsiella* spp. displayed the highest resistance to amoxicillin (82.6%; 95%CI= 63.3%−92.9 vs 86.4%; 95%CI= 55.9%−97.0%), amoxicillin + clavulanic acid (71.7%; 95%CI= 0.44%−89.2% vs 73.1%; 95%CI= 38.7%−92.1%) and trimethoprim/sulfamethoxazole (76.2%; 95%CI= 51.3%−90.7% vs 69.5%; 95%CI= 52.3−82.6%) respectively. However, meropenem (26.7%; 95%CI= 20.8%−33.7%) and fosfomycin (14.9%; 95%CI= 9.1%−23.4%) were the least resistant in *E. coli* and *Klebsiella* spp. respectively. *Staphylococcus aureus* showed high resistance to penicillin (84.4%; 95%CI= 68.1%−93.2%), erythromycin (41.5%; 95%CI= 28.0%−56.5%) and oxacillin (37.4%; 95%CI= 8.8%−78.7%) and low resistance to rifampicin (2.9%; 95%CI= 0.6%−13.4%), fusidic acid (8.1%; 95%CI= 3.4%−18.0%) and vancomycin (13.6%; 95%CI= 2.2%−52.9%).

Conclusion: This study reports a high prevalence of bacterial BSIs in Cameroon and the high resistance of these bacteria to common antibiotics. There is a pressing need to conduct BSI surveillance studies in all regions of Cameroon to generate data for evidence-based measures regarding BSI prevention and management.

Keywords: Bloodstream infection, epidemiology, *Enterobacterales,* antibiotic resistance, Cameroon.

Introduction

Bloodstream infection (BSI) is characterised by positive blood cultures in patients with systemic symptoms and signs of infection [1]. BSI can eventually lead to sepsis which is a complication characterised by a life-threatening organ dysfunction caused by a dysregulated host response to infection [1]. Sepsis and BSI are significantly associated with morbidity and mortality worldwide and represent a huge public health concern [2]. According to the World Health Organization (WHO), sepsis accounted for around 20% of all global deaths in 2017, affected 49 million people, and resulted in 11 million fatalities [2, 3]. Vulnerable populations such as pregnant women, elderly persons, patients with severe comorbidities, newborns, and those in low-resource settings are disproportionately affected [4, 5]. Cases of sepsis and sepsis-related deaths in low-and middle-income countries (LMICs) account for 85% of the global sepsis incidence [3] and hospital mortality due to sepsis is estimated at 19% for moderate sepsis and 39% for severe sepsis in sub-Saharan Africa (SSA) [6]. Bacteria, especially *Enterobacterales* including *Escherichia coli*, *Klebsiella* spp., *Salmonella enterica* and Gram-positive cocci such as *Staphylococcus aureus* and coagulase-negative *Staphylococcus* are all major species implicated in BSI and sepsis [7, 8].

Inappropriate use of antibiotics is known to exacerbate the burden of Antimicrobial Resistance (AMR) in bacterial pathogens [9]. Globally, there is an increase in drug resistance to front-line antibiotics including β-lactams, fluoroquinolones, aminoglycosides, and macrolides [10, 11]. Therefore, AMR is a major public health concern and is expected to be the leading cause of mortality by 2050 with 10 million deaths per year if no immediate measures are taken [12]. AMR is a serious challenge in hospital settings as it worsens the effects of bacterial infections in general especially sepsis, from extended hospital stays to increased mortality [13, 14].

In Cameroon, the bacterial BSI and sepsis and the AMR rate of implicated bacteria are still poorly documented. Notwithstanding, some studies have been published, it would be important to summarise these already available data to inform on context-specific epidemiology of BSI in terms of aetiology, AMR rate and to direct future BSI and sepsis prevention and management efforts in Cameroon. Thus, in this review, we estimate the pooled proportion of culture-confirmed bacterial BSIs and their AMR rates in Cameroon.

Methods

We conducted this systematic review and meta-analysis following the Preferred Reporting Items for Systematic Review and Meta-Analysis statement (PRISMA) 2020 guidelines [15] and, the protocol was registered in the Prospero database (Ref: CRD42023482760, available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023482760).

Search strategy

The National Library of Medicine database (MEDLINE) was searched through PubMed, along with Google Scholar and hand search. These databases were searched to find all articles reporting the prevalence of culture-confirmed bacteria-driven bloodstream infections. Library research was

performed using Medical Subject Headings (MeSH) and keywords paired with Boolean operators (**Fig. 1)**.

Eligibility criteria

Three authors independently searched the literature and evaluated the titles and abstracts of papers. Articles included were studies which (1) reported culture-confirmed bacteria-driven bloodstream infections in Cameroon, (2) published in peer-reviewed journals and (3) published in English or French from 2010 up to December 2023. Articles were excluded if (1) the full texts were not assessable and (2) they were case reports or case series.

Study selection

After the literature search, we uploaded all articles to an EndNote (version 20.6) database and duplicates were removed. Two reviewers did the screening of titles and abstracts to assess whether specific inclusion criteria were met and non-relevant citations and those that did not meet inclusion criteria were excluded. Disagreements on whether or not an article should be included or not were resolved by a third reviewer.

Data extraction

Two authors used a Microsoft Excel spreadsheet (2016 version) to extract data from included articles independently. The first author, publication year, region of study, study design, sample size, age of study population, the prevalence of BSIs, the proportion of each bacteria causing BSIs and AMR rates of bacteria groups were extracted. Inconsistencies were resolved through a group consensus.

Quality assessment

Quality of included studies was done using the Joanna Briggs Institute revised Critical Appraisal Checklist [16]. The checklist was applied to each study to assess the eight items with each item awarded a point for "yes" and the final score for each article added and fell between zero and eight. Any disagreement among authors was resolved by consensus. A study was considered low-risk when it scored ≥ 4 points.

Data analysis

We exported the extracted data to Comprehensive Meta-Analysis Software (version 3.0) for analysis. Begg and Egger's regression tests were used to objectively assess publication bias while the funnel plot was used to visualise this bias [17]. A p-value < 0.05 was considered statistically significant [18]. Cochrane Q statistics assessed the presence and magnitude of heterogeneity between studies. Quantification of heterogeneity was done using I^2 expressed in percentages. Values of 75, 50 and 25% were considered as high, medium and low heterogeneity [19] respectively. Subgroup meta-analysis was conducted according to participant's ages into all ages and paediatric (children under 2 years).

Results

Results of the literature search

The online database search from PubMed and Google Scholar yielded 329 articles and two articles through the hand search. The duplicates were removed and 314 articles were enrolled for the title and abstract screening. Finally, 25 articles were screened for full-text review and 10 which responded to the study objectives were included (**Fig. 2**).

Publication bias

When BSI among study participants was analysed, a visual inspection of the funnel plot showed that there was no evidence of publication bias among the included studies. Similarly, the coefficient (4.89, 95% CI= -5.23, 15.01) for Egger's regression test was not statistically significant (*p-value* = 0.298).

Characteristics of included studies

The 10 included studies described a total of 4223 blood cultures, of which 920 (21.79%) were performed exclusively in the paediatric population (children under 2 years). The studies were conducted in two of the 10 regions of Cameroon and included patients from only two cities Douala (n=2547; 60.31%) [20-22] and Yaoundé (n=1676; 39.69%) [23-29]. Most of the studies were cross-sectional (n=08) [20, 22-26, 28, 29], one longitudinal [27] and one retrospective [21] study but all were of low-risk quality. Only the cross-sectional studies reported the age of the patients which ranged from 0 to 92 years **(Table 1).**

Pooled prevalence and bacterial aetiology of bloodstream infections

The heterogeneity among the included studies was statistically significant (I^2= 97,63%; p=0.000). Thus, we used Random Effect Models to calculate the pooled prevalence of culture-confirmed bacteria-driven bloodstream infections in Cameroon and their 95% confidence intervals were represented on forest plots. Blood culture was performed in 4223 patients and 888 were positive for a bacterial BSI. The bacterial BSI prevalence in this study ranged from 11.93% up to 67.89% and the overall pooled prevalence was 26.36% (95%CI= 17.51%−37.64%). Additionally, the prevalence in children and all age subgroups was similar (26.54% and 26.22% respectively) (**Fig. 3**).

The main bacterial groups reported in this study were *Enterobacterales, Staphylococci* and *Streptococci*. Among the *Enterobacterales* group, *Klebsiella* spp. (28.20%; 95%CI= 2.80%−63.60%) was the most implicated followed by *Escherichia coli* (18.94%; 95%CI= 13.02%−26.72%), *Enterobacter* spp. (7.66%; 95%CI= 4.46%−12.83%) and *Salmonella enterica* (7.01%; 95%CI= 3.62%−13.15%) (**Fig.4**). Seven studies reported gram-positive cocci, of which *Staphylococcus aureus* was the most implicated (14.40%; 95%CI= 8.31%−23.81%), followed by *Coagulase-negative Staphylococcus* (8.23%; 95%CI= 2.68%−22.65%) (**Fig. 5**) and *Streptococcus* spp. (4.07%; 95%CI= 2.64%−6.21%) as presented in **Fig. 6**. Non-fermenting gram-negative bacilli were also recorded in both age groups and *Acinetobacter* spp. (3.93%; 95%CI=2.13%−7.29%) was the commonly reported followed by *Pseudomonas* spp. (3.17%; 95%CI= 1.96%−5.07%) (**Fig. 7**).

Antimicrobial resistance rates of bacterial BSI

This study reported only the pooled AMR rate of the most represented bacteria in each group. Most of the included studies failed to report the AST of several antibiotics. Overall, both *E. coli* and *Klebsiella* spp. were highly resistant to amoxicillin 82.6% (95%CI= 63.3%–92.9%) vs 86.4% (95%CI= 55.9%–97.0%), amoxicillin + clavulanic acid 71.7% (95%CI= 0.44%–89.2%) vs 73.1% (95%CI= 38.7%–92.1%) and trimethoprim/sulfamethoxazole 76.2% (95%CI= 51.3%–90.7%) vs 69.5% (95%CI= 52.3%–82.6%) respectively. Additionally, resistance to third and fourth-generation cephalosporins and monobactam was high among *E. coli*: cefotaxime 46.5% (95%CI= 26.8%–67.3%), ceftazidime 49.9% (95%CI= 29.5%–70.3%), cefepime 51.5% (95%CI= 10.9%–62.0%), aztreonam 49.9% (95%CI= 40.6%-59.3%) and *Klebsiella* spp.: cefotaxime 30.4% (95%CI= 9.8%–63.6%), ceftazidime 45.4% (95%CI= 16.8%–77.1%) and aztreonam 54.2% (95%CI= 34.2%–73.0%). However, meropenem 26.7% (95%CI= 20.8–33.7%) and fosfomycin 14.9% (95%CI= 9.1%–23.4%) were the most effective in *E. coli* and *Klebsiella* spp. respectively (**Table 2)**. *Staphylococcus aureus* showed high resistance to penicillin 84.4% (95%CI= 68.1%–93.2%), erythromycin 41.5% (95%CI= 28.0%–56.5%) and oxacillin 37.4% (95%CI= 8.8%–78.7%). However, a low resistance rate of *Staphylococcus aureus* was recorded in non-β-lactams antibiotics including rifampicin 2.9% (95%CI= 0.6%–13.4%), fusidic acid 8.1% (95%CI= 3.4%–18.0%) and vancomycin 13.6% (95%CI= 2.2%–52.9%) and cefoxitin 18.9% (95%CI= 12.0%–82.1%) was the most effective β-lactams. In the *CONS* group, a high resistance rate was registered against penicillin 89.9% (95%CI= 66.0%–97.6%), trimethoprim/sulfamethoxazole 52.9% (95%CI= 42.1%–63.5%) and oxacillin 50.7% (95%CI= 25.0%–76.1%) while cefoxitin 14.9% (95%CI= 11%–73.6%) and fusidic acid 25.5% (95%CI= 17.4%–35.6%) were the most effective (**Table 3)**.

Discussion

This review aimed to determine the pooled prevalence of bacterial BSIs and the AMR rates of implicated bacteria in Cameroon. The 10 studies selected for this review described 4223 blood cultures and the overall pooled prevalence of BSI in was 26.36%. This prevalence is higher than those reported in Africa by Reddy *et al* (2010) and in Africa and Asia by Marchello *et al* (2019) which were 7.4% and 10.8% respectively [30, 31]. However, studies conducted in Ethiopia by Bitew *et al* (2023), Legese *et al* (2022) and Abebe *et al* (2021) documented a similar prevalence of BSI [32-34]. This review also recorded a high prevalence of 26.54% of BSI in children under 2 years old. A similar pooled prevalence of neonatal sepsis in East Africa was documented by Abate *et al* (2020) [35] while Belachew and Tewabe (2020) recorded a pooled prevalence of neonatal sepsis in Ethiopia of 49.98% [36]. The observed difference and the high prevalence either in children and all age groups in this review may be due to the small sample size of many included studies [20, 22, 27, 28], the study design as well as the study periods. The study revealed that *Enterobacterales* including *Klebsiella* spp. (28.20%) and *Escherichia coli* (18.94%) were the leading bacteria implicated in the BSI. Studies conducted in several low and middle-income countries have demonstrated the high implication of these bacteria in bloodstream infections and sepsis especially in hospital settings [10, 34, 37]. However, these results were contradictory to those recorded by Wattal *et al* (2020) and Reddy *et al* (2010) where *Salmonella* spp. was the leading

cause of BSI and sepsis [11, 30]. This contradiction might be due to the bacterial identification methods used poorly described in some studies which can considerably vary depending on the laboratory [20, 21, 27].

The most frequent *Enterobacterales* reported in this review showed a high resistance to β-lactam antibiotics especially third and fourth-generation cephalosporins and monobactam. This might indicate the production of β-lactamases particularly the extended-spectrum β-lactamase enzymes by most of these bacteria even though most of the included studies failed to report the resistance phenotypes.

Limitations

The statements and conclusions in this review cannot be generalised to Cameroon as the included studies investigated only two regions and were conducted in a small sample size. Also, the study was unable to precise whether the BSI was community or hospital-acquired. Finally, the study failed to report AMR among several bacteria and the resistance phenotypes of implicated bacteria.

Conclusion

The prevalence of bacterial bloodstream infection in this study is elevated in both paediatric and adult populations. *Enterobacterales* including *Klebsiella* spp. and *Escherichia coli* and Gram-positive cocci particularly *Staphylococcus aureus* remain the leading causes of bloodstream infection. Antibiotic resistance was higher to β-lactam in both *Enterobacterales* and Gram-positive cocci. This study may help clinicians optimise their antibiotic prescription protocol in the early management of bacterial bloodstream infection in the absence of antibiotic susceptibility testing results. The review also calls for researchers to make more effort to produce AMR data to permit policymakers and stakeholders a better bloodstream infection management and prevention in Cameroon.

Conflict of interest

The authors declare no conflict of interest

Authors contributions

MM conceived and designed the study. MM and PLK coordinated the literature search. PKL, RJZ and MM undertook data extraction, quality assessment and statistical analysis. MM, PLK, and RJZ prepared the first draft of the manuscript. ID, SCHN and MN critically reviewed the manuscript. All authors read and approved the final version of the manuscript.

Funding

The study was not funded.

References

[1]. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M*, et al.* The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). Jama. 2016;315(8):801-10.

[2].   WHO. Global report on the epidemiology and burden of sepsis: Current evidence, identifying

gaps and future directions. 2020.

[3].   Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, *et al.* Global, regional,

and national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease

Study. Lancet. 2020;395(10219):200-11.

[4].   Cheng AC, West TE, Limmathurotsakul D, Peacock SJ. Strategies to reduce mortality from

bacterial sepsis in adults in developing countries. PLoS Med. 2008;5(8):e175.

[5].   Rhee C, Jones TM, Hamad Y, Pande A, Varon J, O'Brien C, *et al.* Prevalence, Underlying

Causes, and Preventability of Sepsis-Associated Mortality in US Acute Care Hospitals. JAMA Netw

Open. 2019;2(2):e187571.

[6].   Lewis JM, Feasey NA, Rylance J. Aetiology and outcomes of sepsis in adults in sub-Saharan

Africa: a systematic review and meta-analysis. Critical Care. 2019;23(1):212.

[7].   Timsit JF, Ruppé E, Barbier F, Tabah A, Bassetti M. Bloodstream infections in critically ill

patients: an expert statement. Intensive Care Med. 2020;46(2):266-84.

[8].   Mun SJ, Kim S-H, Kim H-T, Moon C, Wi YM. The epidemiology of bloodstream infection

contributing to mortality: the difference between community-acquired, healthcare-associated, and

hospital-acquired infections. BMC Infectious Diseases. 2022;22(1).

[9].   Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of antibiotic

resistance. Nat Rev Microbiol. 2015;13(1):42-51.

[10].   Ombelet S, Kpossou G, Kotchare C, Agbobli E, Sogbo F, Massou F, *et al.* Blood culture

surveillance in a secondary care hospital in Benin: epidemiology of bloodstream infection pathogens

and antimicrobial resistance. BMC Infect Dis. 2022;22(1):119.

[11].   Wattal C, Goel N. Pediatric Blood Cultures and Antibiotic Resistance: An Overview. Indian J

Pediatr. 2020;87(2):125-31.

[12].   O'Neill J. Tackling drug-resistance infections globally: final report and recommendations, 2016.

Available: https://amr-review.org/sites/default/files/160518_Final%20paper_with%20cover.pdf

[Accessed on 1 Jul 2023] 2016 [

[13].   Chaurasia S, Sivanandan S, Agarwal R, Ellis S, Sharland M, Sankar MJ. Neonatal sepsis in

South Asia: huge burden and spiralling antimicrobial resistance. Bmj. 2019;364:k5314.

[14].   Murray CJL, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, *et al.* Global

burden of bacterial antimicrobial resistance in 2019: a systematic analysis. The Lancet.

2022;399(10325):629-55.

[15].   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, *et al.* The PRISMA

2020 statement: An updated guideline for reporting systematic reviews. PLoS Med.

2021;18(3):e1003583.

[16].   Joanna Briggs Institute. Critical-Appraisal-Tools—Critical Appraisal Tools | JBI. 2022. Available

online: https://jbi.global/critical-appraisal-tools (accessed on 07 November 2023).

[17].   Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple,

graphical test. Bmj. 1997;315(7109):629-34.

[18].   Fletcher J. What is heterogeneity and is it important? Bmj. 2007;334(7584):94-6.

[19].   Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses.
Bmj. 2003;327(7414):557-60.

[20].   Kemeze S, Moudze B, Chiabi A, Eposse C, Kaya A, Mbangue M*, et al.* Profil clinique et
bactériologique des infections néonatales bactériennes à l'Hôpital Laquintinie de Douala, Cameroun.
The Pan African Medical Journal. 2016;23.

[21].   Okalla Ebongue C, Mefo'o JP, Te N, Moukoko E, Adiogo D, Beyiha G. Profil Bactériologique et
sensibilité aux antibiotiques des isolats d'hémoculture (2006 – 2011) à Douala, Cameroun. Revue
Malienne d'Infectiologie et de Microbiologie. 2014;2.

[22].   Teghonong J, Okalla Ebongue C, Yadufashije C, Kojom LP, Adiogo D. Phenotypic
Characteristics of Klebsiella pneumoniae Extended Spectrum β-Lactamases Producers Isolated in
Hospitals in the Littoral Region, Cameroon. European Journal of Clinical Microbiology. 2020;6:9-13.

[23].   Carole Suzie Lacmago Titsamp HKG, Simon Ngamli Fewou. Bloodstream infections initiated
by ESBL-producing Escherichia coli in neonates and infants at two hospitals in Yaoundé, Cameroon.
International Journal of Biological and Chemical Sciences. 2021;15(3).

[24].   Chiabi A, Djoupomb M, Mah E, Nguefack S, Mbuagbaw L, Zafack J*, et al.* The clinical and
bacteriogical spectrum of neonatal sepsis in a tertiary hospital in yaounde, cameroon. Iran J Pediatr.
2011;21(4):441-8.

[25].   Djuikoue CI, Djouela Djoulako PD, Wouambo RK, Lacmago ST, Dayomo A, Kamga HG*, et al.*
Prevalence of Escherichia coli Producing Extended Spectrum Beta-Lactamase (ESBL) Driven
Septicaemia in Children Aged 0–2 Years in Two Districts Hospitals in Yaounde, Cameroon. Bacteria.
2022;1(4):294-301.

[26].   Kamga H, Njunda A, Nde P, Assob J, Nsagha D, Weledji P. Prevalence of septicaemia and
antibiotic sensitivity pattern of bacterial isolates at the University Teaching Hospital, Yaoundé,
Cameroon. African Journal of Clinical and Experimental Microbiology. 2011;12(1).

[27].   Nouetchognou JS, Ateudjieu J, Jemea B, Mesumbe EN, Mbanya D. Surveillance of nosocomial
infections in the Yaounde University Teaching Hospital, Cameroon. BMC Res Notes. 2016;9(1):505.

[28].   Yimtchi LCN, Pokam BDT, Baiye AW, Betbeui AC, Gonsu HK, Djakissam W. Antibiotic
Susceptibility Patterns of Bacteria Associated with Sepsis among Hospitalized Patients in the Yaoundé
University Teaching Hospital—Cameroon. Open Journal of Medical Microbiology. 2023;13(1):101-15.

[29].   Zefack JT, Ambe NF, Bobga TP, Ketum AS, Awambeng DN, Kelly CN*, et al.* Bacteriological
Profile and Antimicrobial Sensitivity Pattern of Blood Culture Isolates among Septicemia Suspected
Patients in the University Teaching Hospital (UTH) Yaoundé, Cameroon. International Journal of Trend
in Scientific Research and Development.2020;4(5).

[30].   Reddy EA, Shaw AV, Crump JA. Community-acquired bloodstream infections in Africa: a
systematic review and meta-analysis. The Lancet infectious diseases. 2010;10(6):417-32.

[31].   Marchello CS, Dale AP, Pisharody S, Rubach MP, Crump JA. A Systematic Review and Meta-
analysis of the Prevalence of Community-Onset Bloodstream Infections among Hospitalized Patients
in Africa and Asia. Antimicrob Agents Chemother. 2019;64(1).

[32]. Bitew A, Adane A, Abdeta A. Bacteriological spectrum, extended-spectrum β-lactamase
production and antimicrobial resistance pattern among patients with bloodstream infection in Addis
Ababa. Scientific Reports. 2023;13(1).

[33]. Abebe W, Tegene B, Feleke T, Sharew B. Bacterial Bloodstream Infections and their
Antimicrobial Susceptibility Patterns in Children and Adults in Ethiopia: a 6-Year Retrospective Study.
Clin Lab. 2021;67(11).

[34]. Legese MH, Asrat D, Swedberg G, Hasan B, Mekasha A, Getahun T, *et al.* Sepsis: emerging
pathogens and antimicrobial resistance in Ethiopian referral hospitals. Antimicrobial Resistance &
Infection Control. 2022;11(1):83.

[35]. Abate BB, Kasie AM, Reta MA, Kassaw MW. Neonatal sepsis and its associated factors in East
Africa: a systematic review and meta-analysis. International Journal of Public Health. 2020;65(9):1623-
33.

[36]. Belachew A, Tewabe T. Neonatal sepsis and its association with birth weight and gestational
age among admitted neonates in Ethiopia: systematic review and meta-analysis. BMC Pediatrics.
2020;20(1).

[37]. Solomon S, Akeju O, Odumade OA, Ambachew R, Gebreyohannes Z, Van Wickle K, *et al.*
Prevalence and risk factors for antimicrobial resistance among newborns with gram-negative sepsis.
PLOS ONE. 2021;16(8):e0255410.

(antimicrobial OR antibacterial OR antibiotics OR antibiotiques OR antimicrobial resistance OR antibiotic resistance OR multiresistance OR multidrug resistance) OR (bacteria OR bactéries OR *Enterobacterales* OR *Enterobacteriaceae* OR entérobactéries OR *Escherichia coli* OR *E. coli* OR *Klebsiella* OR *Staphylococcus* OR *Salmonella* OR Typhoid OR *Enterobacter* OR *Pseudomonas* OR *Acinetobacter* OR *Streptococcus* OR coagulase negative OR gram negative OR gram positive) AND (bloodstream infections OR hémoculture OR hemoculture OR sepsis OR septicemia OR septicaemia OR septicémie OR neonatal sepsis) AND (Cameroon OR Cameroun).

Fig. 1: Search terms used to identify relevant literature from PubMed and Google Scholar

Fig. 2: PRISMA flow chart of the study selection process

Study		Pooled prevalence (%)	95% CI (%)		Weight (%)
All age					
Ebongue et al, 2014		12,81	11,51	14,23	17,43
Kamga et al, 2011		28,28	24,06	32,92	17,23
Nouetchognou et al, 2016		20,34	11,93	32,49	15,33
Teghonong et al, 2020		67,89	58,58	75,97	16,57
Zefack et al, 2020		25,49	20,52	31,20	17,04
Yimtchi et al, 2023		16,00	10,96	22,76	16,41
Subtotal (I^2 = 97.51%, p = 0.000)		26,22	14,97	41,78	
Children (≤ 2 years) only					
Chiabi et al, 2011		11,93	8,25	16,94	24,64
Kemeze et al, 2016		16,35	10,41	24,73	23,67
Titsamp et al, 2021		43,29	37,77	48,98	25,84
Djuikoue et al, 2022		43,33	37,83	49,00	25,85
Subtotal (I^2 = 96.06%, p = 0.000)		26,54	14,19	44,11	
Overall (I^2 = 97.63%, p = 0.000)		26,36	17,51	37,64	100

0.0   40.0   80.0

Note: Weights are from random effect analysis

Fig. 3: Forest plot of pooled prevalence of culture-confirmed bacterial bloodstream infections in Cameroon

A: Klebsiella spp.

Study	Pooled prevalence (%)	95% CI (%)		Weight (%)
All age				
Ebongue et al, 2014	27,76	22,98	33,11	18,16
Kamga et al, 2011	7,27	3,68	13,87	16,74
Nouetchognou et al, 2016	66,67	37,59	86,91	14,44
Zefack et al, 2020	67,89	58,58	75,97	17,83
Yimtchi et al, 2023	12,31	6,28	22,73	16,65
Subtotal (I² = 94.76%, p = 0.000)	33,33	17,63	53,88	16,18
Children (≤ 2 years) only				
Chiabi et al, 2011	34,62	14,29	54,51	
Kemeze et al, 2016	5,88	19,09	54,29	59,78
Subtotal (I² = 73.00%, p = 0.054)	18,31	0,82	32,03	40,22
Overall (I² = 92.97%, p = 0.000)	28,20	2,80	63,60	

0.0   40.0   80.0

Note: Weights are from random effect analysis

B: Escherichia coli

Study	Pooled prevalence (%)	95% CI (%)		Weight (%)
All age				
Ebongue et al, 2014	12,04	8,81	16,24	35,58
Kamga et al, 2011	5,45	2,47	11,61	20,86
Nouetchognou et al, 2016	3,85	0,24	40,32	3,24
Zefack et al, 2020	10,77	5,22	20,91	21,88
Yimtchi et al, 2023	25,00	11,69	45,64	18,44
Subtotal (I² = 53.05%, p = 0.074)	11,18	6,90	17,62	
Children (≤ 2 years) only				
Chiabi et al, 2011	19,23	8,24	38,70	21,38
Kemeze et al, 2016	35,29	16,79	59,59	21,13
Titsamp et al, 2021	69,77	61,31	77,07	28,73
Djuikoue et al, 2022	69,23	60,79	76,56	28,76
Subtotal (I² = 88.06%, p = 0.000)	50,96	31,34	70,30	
Overall (I² = 96.40%, p = 0.000)	18,94	13,02	26,72	100

0.0   40.0   80.0

Note: Weights are from random effect analysis

C: Enterobacter spp.

Study	Pooled prevalence (%)	95% CI (%)		Weight (%)
All age				
Ebongue et al, 2014	11,71	8,52	15,87	41,58
Kamga et al, 2011	6,36	3,06	12,75	28,36
Nouetchognou et al, 2016	16,67	4,20	47,72	12,98
Zefack et al, 2020	1,54	0,22	10,12	8,64
Yimtchi et al, 2023	4,17	0,58	24,35	8,45
Subtotal (I² = 48.41%, p = 0.101)	8,05	4,41	14,26	
Children (≤ 2 years) only				
Chiabi et al, 2011	7,69	1,93	26,07	79,16
Kemeze et al, 2016	2,78	0,17	32,21	20,84
Subtotal (I² = 0.00%, p = 0.507)	6,25	1,81	19,39	
Overall (I² = 31.75%, p = 0.186)	7,66	4,46	12,83	

0.0   40.0   80.0

Note: Weights are from random effect analysis

D: Salmonella enterica

Study	Pooled prevalence (%)	95% CI (%)		Weight (%)
All age				
Ebongue et al, 2014	8,36	5,71	12,08	42,25
Kamga et al, 2011	15,45	9,83	23,46	39,27
Nouetchognou et al, 2016	3,85	0,24	40,32	6,07
Zefack et al, 2020	0,76	0,05	10,98	6,24
Yimtchi et al, 2023	2,00	0,12	25,13	6,17
Subtotal (I² = 48.41%, p = 0.101)	8,18	4,06	15,79	
Children (≤ 2 years) only				
Chiabi et al, 2011	1,85	0,11	23,64	50,24
Kemeze et al, 2016	2,78	0,17	32,21	49,76
Subtotal (I² = 0.00%, p = 0.507)	2,27	0,32	14,42	
Overall (I² = 31.75%, p = 0.186)	7,01	3,62	13,15	

0.0   40.0   80.0

Note: Weights are from random effect analysis

Fig. 4: Forest plot of pooled prevalence of *Enterobacterales*-driven BSI in Cameroon.

A: *Klebsiella* spp. ; **B:** *Escherichia coli;* **C:** *Enterobacter* spp. and **D:** *Salmonella enterica*

Fig. 5: Forest plot of pooled prevalence of *Staphylococci*-driven BSI in Cameroon.

E: *Staphylococcus aureus* and **F:** Coagulase negative *Staphylococci*

Fig. 6: Forest plot of pooled prevalence of *Streptococcus* spp. driven bloodstream infections in Cameroon

Fig. 7: Forest plot of pooled prevalence of non-fermenting Gram-negative bacteria-driven bloodstream infections in Cameroon

Table 1: Summary characteristics of studies included in the meta-analysis

Study period	Study type	City	Population size	Age range	Positive cultures (%)	Quality	Authors Study period
January-April 2018	Cross-sectional	Yaoundé	255	0-82 years	65 (25.50)	Low-risk	Zefack et *al*, 2020
August-October 2017	Cross-sectional	Yaoundé	298	0-48 weeks	129 (43.30)	Low-risk	Titsamp et *al*, 2021
August 2019-March 2020	Cross-sectional	Yaoundé	300	0-2 years	130 (43.33)	Low-risk	Djuikoue *et al*, 2022
March-June 2015	Cross-sectional	Douala	104	0-28 days	17 (16.34)	Low-risk	Kemeze et *al*, 2016
September 2013 - March 2014	Longitudinal	Yaoundé	59	NA	12 (20.34)	Low-risk	Nouetchognou et *al*, 2016
2006-2011	Retrospective	Douala	2334	NA	299 (12.81)	Low-risk	Ebongue et *al*, 2014
January-June 2010	Cross-sectional	Yaoundé	396	0->76 years	112 (28.30)	Low-risk	Kamga et *al*, 2011
November 2008-May 2009	Cross-sectional	Yaoundé	218	0-28 days	26 (11.92)	Low-risk	Chiabi *et al*, 2011
November 2021-March 2022	Cross-sectional	Yaoundé	150	10-92 years	24 (16)	Low-risk	Yimtchi *et al*, 2023
January 2016-February 2017	Cross-sectional	Douala	109	>1-71 years	74 (67.89)	Low-risk	Teghonong et al, 2020

Table 2: Pooled prevalence of antibiotic resistance of *E. coli* and *Klebsiella* spp. isolated from bloodstream infections in Cameroon

Bacteria	Antibiotics	Number of studies	Number of isolates	Pooled prevalence of AMR (95% CI)
E. coli N = 281	Amoxicillin	5	275	82.6 (63.3-92.9)
	Amoxicillin + Clavulanic acid	5	276	71.7 (0.44-89.2)
	Cefotaxime	3	263	46.5 (26.8-67.3)
	Ceftazidime	5	274	49.9 (29.5-70.3)
	Cefepime	3	187	51.5 (10.9-62.0)
	Cefoxitin	6	281	33.2 (18.5-52.0)
	Aztreonam	4	108	49.9 (40.6-59.3)
	Meropenem	3	180	26.7 (20.8-33.7)
	Amikacin	5	274	37.5 (18.7-61.1)
	Gentamicin	6	281	49.5 (37.3-61.7)
	Ciprofloxacin	5	191	41.7 (32.6-51.4)
	Nalidixic acid	3	185	34.6 (28.1-41.7)
	Trimethoprim/Sulfamethoxazole	5	275	76.2 (51.3-90.7)
Klebsiella spp. N = 115	Amoxicillin	4	107	86.4 (55.9-97.0)
	Amoxicillin + Clavulanic acid	4	99	73.1 (38.7-92.1)
	Cefotaxime	3	99	30.4 (9.8-63.6)
	Ceftazidime	3	99	45.4 (16.8-77.1)
	Cefoxitin	4	107	35.8 (19.5-56.3)
	Aztreonam	3	24	54.2 (34.2-73.0)
	Fosfomycin	3	99	14.9 (9.1-23.4)
	Amikacin	4	107	26.8 (6.1-67.2)
	Gentamicin	5	115	44.8 (35.6-54.3)
	Ciprofloxacin	5	115	56.5 (29.9-79.8)
	Trimethoprim/Sulfamethoxazole	4	107	69.5 (52.3-82.6)

Table 3: Pooled prevalence of antibiotic resistance of *S. aureus* and Coagulase Negative *Staphylococcus* (CONS) isolated from bloodstream infections in Cameroon

Bacteria	Antibiotics	Number of studies	Total number of isolates	Pooled prevalence of AMR (95% CI)
S. aureus N = 68	Penicillin	2	61	84.4 (68.1-93.2)
	Oxacillin	3	62	37.4 (8.8-78.7)
	Cefoxitin	3	29	18.9 (12.0-82.1)
	Gentamicin	4	68	30.2 (20.3-42.3)
	Erythromycin	4	68	41.5 (28.0-56.5)
	Lincosamin	2	62	27.4 (3.6-79.1)
	Tetracycline	4	68	29.3 (5.6-74.2)
	Rifampicin	3	61	2.9 (0.6-13.4)
	Trimethoprim/Sulfamethoxazole	3	62	29.8 (19.6-42.5)
	Fusidic acid	3	62	8.1 (3.4-18.0)
	Vancomycin	3	68	13.6 (2.2-52.9)
CONS N = 93	Penicillin	3	93	89.9 (66.0-97.6)
	Oxacillin	3	93	50.7 (25.0-76.1)
	Cefoxitin	2	71	14.9 (11-73.6)
	Gentamicin	3	93	36.7 (18.7-59.4)
	Erythromycin	3	93	46.8 (27.3-67.4)
	Lincosamin	3	93	42.0 (32.2-52.5)
	Tetracycline	3	93	41.7 (27.0-58.1)
	Trimethoprim/Sulfamethoxazole	3	93	52.9 (42.1-63.5)
	Fusidic acid	3	93	25.5 (17.4-35.6)
	Vancomycin	3	93	30 (14.4-53.2)