Characterization of the common genetic variation in the Spanish population of Navarre

Authors:

Alberto Maillo, alberto.ruizdeinfante@kaust.edu.sa, Spain[1,2]

Estefania Huergo, ehuergoi@navarra.es, Spain[1,#]

María Apellániz-Ruiz, mv.apellaniz.ruiz@navarra.es, Spain[3,#]

Edurne Urrutia, eurrutil@navarra.es, Spain[1,3,#]

María Miranda, mmirandp@navarra.es, Spain[3]

Josefa Salgado, jsalgadg@navarra.es, Spain[3,4,5]

Sara Pasalodos-Sánchez, spasalos@navarra.es, Spain[3]

Luna Delgado-Mora, lunadelde@gmail.com, Spain[3,6]

Óscar Teijido, oteijidh@navarra.es, Spain[3]

Ibai Goicoechea, igoicoeo@nasertic.es, Spain[7]

Rosario Carmona, rosariom.carmona@juntadeandalucia.es, Spain[8,9,10,11]

Javier Perez-Florido, javier.perez.florido.sspa@juntadeandalucia.es, Spain[8,9,10,11]

Virginia Aquino, virginia.aquino@juntadeandalucia.es, Spain[8]

Daniel Lopez-Lopez, daniel.lopez.lopez@juntadeandalucia.es, Spain[8,9,11]

María Peña-Chilet, maria_pena@iislafe.es, Spain[8,11]

Sergi Beltrán, sergi.beltran@cnag.eu, Spain[12,13,14]

Joaquín Dopazo, joaquin.dopazo@juntadeandalucia.es, Spain[8,9,10,11]

Iñigo Lasa, ilasa@unavarra.es, Spain[15]

Juan José Beloqui, jj.beloqui.lizaso@navarra.es, Spain[3]

NAGEN-scheme, (*see list at the end*),

Ángel Alonso, aalonsos@navarra.es, Spain[3*]

David Gomez-Cabrero, david.gomez.cabrero@navarra.es, Spain[1,2*]

Affiliations:

1: Translational Bioinformatics Unit, Navarrabiomed, Hospital Universitario de Navarra (HUN), Universidad Pública de Navarra (UPNA), IdiSNA, Pamplona, Spain

2: Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia.

3: Genomics Medicine Unit, Navarrabiomed, Hospital Universitario de Navarra (HUN), Universidad Pública de Navarra (UPNA), IdiSNA, Pamplona, Spain

4: Servicio de Genética Médica-Hospital Universitario de Navarra (HUN), Pamplona, Spain.

5: Dp. Bioquímica y Biología Molecular-Universidad Pública de Navarra (UPNA), Pamplona, Spain.

6: Instituto de Genética Médica y Molecular (INGEMM), Hospital Universitario La Paz, Madrid, Spain.

7: Department of Personalized Medicine, NASERTIC, Government of Navarra, Pamplona, Spain.

8: Computational Medicine Platform, Andalusian Public Foundation Progress and Health-FPS, Sevilla, Spain.

9: Institute of Biomedicine of Seville, IBiS, University Hospital Virgen del Rocio/CSIC/University of Sevilla, Sevilla, Spain.

10: FPS/ELIXIR-ES, Fundación Progreso y Salud (FPS), CDCA, Hospital Virgen del Rocio, Sevilla, Spain.

11: Biomedical Research Networking Center in Rare Diseases (CIBERER), Health Institute Carlos III, 28029 Madrid, Spain.

12: CNAG-CRG, Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Barcelona, Spain.

13: Universitat Pompeu Fabra (UPF), Barcelona, Spain.

14: Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona (UB), Barcelona, Spain.

15: Laboratory of Microbial Pathogenesis, Navarrabiomed, Pamplona, Spain.

*corresponding. #contributed equally

ABSTRACT

Large-scale genomic studies have significantly increased our knowledge of genetic variability

across populations. Regional genetic profiling is essential for distinguishing common benign

variants from disease-causing ones. To this end, we conducted a comprehensive characterization

of exonic variants in the population of Navarre (Spain), utilizing whole genome sequencing data

from 358 unrelated individuals of Spanish origin. Our analysis revealed 61,410 biallelic single

nucleotide variants (SNV) within the Navarrese cohort, with 35% classified as common (MAF >

1%). By comparing allele frequency data from 1000 Genome Project (excluding the Iberian cohort

of Spain, IBS), Genome Aggregation Database, and a Spanish cohort including IBS individuals as

well as data from Medical Genome Project, we identified 1,069 SNVs common in Navarre but rare

(MAF ≤ 1%) in all other populations. This observation was further corroborated by a second

regional cohort of 239 unrelated exomes, which confirmed 676 of the 1,069 SNVs as common in

Navarre. In conclusion, this study highlights the importance of population-specific

characterization of genetic variation to improve allele frequency filtering in sequencing data

analysis to identify disease-causing variants.

Keywords: personalized medicine, whole genome sequencing WGS, whole exome sequencing

WES, single nucleotide variant SNV, population frequencies, genetic variability.

 INTRODUCTION

In recent years, the use of NGS in patient healthcare has increased due to technological advances,

cost reduction, and enhanced efficiency (1). The advancement of NGS spans a spectrum of

applications, encompassing whole exome/genome sequencing (WES/WGS). These technologies

revealed a wealth of genetic variants, necessitating the implementation of filters to narrow down

the list of candidate variants. In this regard, the availability of population-specific catalogues of

common variants enables the identification of rare variants (2), such as the international

initiatives 1000 Genome Project (1KGP) (3) and Genome Aggregation Database (gnomAD) (4).

Moreover, various countries like the UK (5), USA (6), and Japan (7) have established their

databases. In Spain, for instance, the Medical Genome Project (MGP) compiles data from

unrelated healthy individuals (8)(9).

In Navarre, a 650,000 population region of north-eastern Spain, the local Government supported

the "*NAGEN scheme*" to integrate genomic data analysis into the regional public healthcare

system. Nowadays, NAGEN has generated numerous WES/WGS and associated phenoclinical

profiles in seven projects, including *NAGEN1000*, focused on rare diseases, and *pharmaNAGEN*

on pharmacogenomics in patients with inflammatory bowel diseases (10). The NAGEN strategy's

success hinges on identifying population-specific common variants to establish a comprehensive

Navarrese population frequency catalogue.

In this study (Fig. 1), we aimed to identify and characterize common exonic variants specific to

the Navarrese population. Firstly, we identified common single nucleotide variants (SNVs) in

Navarre and rare in other populations. Secondly, we validated the allele frequency of these

variants in another Navarrese cohort with only exome data. Finally, the resulting variants were

annotated using genomic databases, and their clinical and pharmacological effects and

pathogenicity were assessed. Additionally, we conducted functional enrichment analyses to

provide further insights. The results will significantly contribute to advancing personalized

medicine in Navarre.

MATERIAL AND METHODS

Please refer to the Supplementary information for detailed material and methods.

Referenced population projects

Population frequencies were obtained from 1) gnomAD: genomes v2.1.1; 2) *1KGP_noIBS*: mean

1KGP (phase3) population's frequencies, excluding the IBS cohort; and 3) *spain*: combining the

MGP and IBS frequencies.

Variant annotation

Variants were annotated using ANNOVAR (11). Clinical and pharmacological relevance was

assessed using ClinVar (12), Online Mendelian Inheritance in Man (OMIM) (13) and PharmGKB

(14). Variant classification was conducted following the American College of Medical Genetics

(ACMG) guidelines (15).

RESULTS

Navarrese discovery cohort

The *NAGEN1000* WGS Navarrese project comprised 688 individuals from 294 families (mainly trios) with a rare disease. The WGS was conducted with a mean coverage of 30X, providing comprehensive genomic data across the entire genome. For our study, we kept a subset of this cohort satisfying two criteria: unrelatedness and Spanish ancestry. This result yielded 358 individuals, referred to as NAVARREsel.

Then, biallelic SNVs on chromosomes 1 to 22, from the exonic region covered by Nextera Exome Enrichment kit, were extracted. Subsequently, we filtered out variants with read depth < 10, genotype quality < 50, or missing genotype in at least one sample. Additionally, sites significantly deviated from Hardy-Weinberg equilibrium (HWE, p-value < 10^{-5}) were removed (16). Finally, 61,410 SNVs remained; 21,174 were identified as common variants (MAF > 1%). We observed that including additional individuals did not reveal new common variants, and 21,174 were achieved when considering over 100 individuals (Fig. S1a).

Genetic variation between Navarre, Spanish, and global populations

We performed a principal component analysis on the shared variants between NAVARREsel, 1KGP, and MGP to depict its relationship. We observed a clear distinction between Navarre and Asian/African populations, reflecting established genetic differences (Fig. 2a). Conversely, an overlap is observed between Navarre and European populations, emphasizing their genetic affinity. Thus, focusing on European populations (Fig. 2b), we observed that Navarrese individuals are close to the Spanish populations (IBS and MGP) and exhibit proximity to Italian individuals.

To further analyse the genetic differentiation, we calculated the mean pairwise F_{ST} values. The

lower F_{ST} value indicates greater similarity between populations. This occurred when comparing

Navarre with the Spanish ($F_{ST(Navarre-IBS)}$ = 0.0001 and $F_{ST(Navarre-MGP)}$ = 0.0007) and Italian ($F_{ST(Navarre-}$

$_{TSI)}$ = 0.0014) populations. In contrast, the highest differentiation was observed against East-Asian

and African populations ($F_{ST(Navarre-EAS)}$ = 0.0328, $F_{ST(Navarre-AFR)}$ = 0.0434).

These findings, aligning with biological expectations, underscore the regional and continental

genetic affinities, providing insights into historical populations and evolutionary dynamics.

Exclusive common variants in Navarre

To identify exclusive Navarrese common variants, we examined allele frequency among Navarre

population and the referenced population: *1KGP_noIBS*, gnomAD, and *spain*. A comparison of the

MAF revealed that most variants (17,532 SNVs) were classified as common (MAF > 1%) across the

four populations. However, 835 variants exhibited higher prevalence solely in Spanish cohorts

(Navarre and *spain*). Specifically, 1,069 SNVs were identified as common in Navarre, and rare

(MAF ≤ 1%) in the rest (Fig. 3a).

To validate these 1,069 variants, we used the NAVARREval cohort, a subset of 239 WES unrelated

individuals of Spanish descent from the current Navarrese population and diagnosed with Crohn's

disease (159/239) or ulcerative colitis (86/239) (*pharmaNAGEN* project). Before validation, we

assessed the association of these SNVs with these conditions by cross-referencing them with

reported variants in the Inflammatory Bowel Disease database, which catalogues variants highly

linked to the mentioned diseases (17). The absence of the 1,069 SNVs in this database ensured

an unbiased and robust validation process.

Among the 1,069 variants initially identified, 998 were detected in NAVARREval with a call rate

greater than 80% and demonstrated conformity to HWE. Notably, 676/998 of these SNVs (68%;

p-value = $2.2e^{-16}$) were consistently classified as common in NAVARREval, confirming their

prevalence within the Navarrese population (variants' information in Table S2). The validation

cohort was sufficient to validate the Navarrese common variants, reaching a plateau in Fig. S1b.

On the contrary, within the non-validated subset (322/998, 32%), 134 SNVs exhibited MAFs in

NAVARREsel that did not exceed a 2-fold difference in NAVARREval, indicating close MAF between

both datasets (Fig. S2). This exploration of MAF patterns ensures a comprehensive understanding

of the genetic landscape within the Navarre population and its stability across different datasets.

Characterization of common Navarrese variants

The annotation of the 676 common Navarrese SNVs revealed 227 synonymous, 371 missense,

and five loss-of-function (LoF) variants. These LoF variants were not reported in ClinVar database

(12) and were located in five distinct genes without an associated phenotype, according to OMIM

(13). Following the ACMG guidelines for variant classification, four were classified as variants of

uncertain significance (VUS) and one as benign (15).

Clinically, 264/676 reported in ClinVar: 1/264 as a risk factor, 181/264 as benign/likely-benign,

32/264 as VUS, 48/264 as having conflicting interpretations, and 2/264 as likely-pathogenic.

These likely-pathogenic missense variants were in *SCNN1B* [c.1688G>A p.Arg563Gln;

$MAF_{NAVARREsel}=0.013$, $MAF_{NAVARREval}=0.016$] and in *PTGIS* [c.824G>A p.Arg275Gln;

$MAF_{NAVARREsel}=0.013$, $MAF_{NAVARREval}=0.021$], associated with "low renin hypertension" and

"childhood-onset schizophrenia" respectively, according to ClinVar. The evidence supporting

these associations is limited, with a score of 1 out of 4 and reviewed by a single submitter record.

Additionally, given its notable prevalence in the Navarrese population, observed in healthy and

affected (not related to this phenotype) individuals, these variants might be reconsidered and

reclassified as VUS under the ACMG guidelines.

Moreover, common Navarrese variants showed no impact on drug metabolism/efficacy,

according to PharmGKB (14). Further in silico and enrichment analysis are detailed in

Supplementary information.

Refining disease-causing variant identification in the Navarrese population

We identified common variants in the Navarrese population, highlighting the importance of

population-specific in advancing personalized medicine. The aim was to improve the

identification of disease-causing variants during genetic diagnosis using NGS. Therefore, we

selected 127 WGS patients from Navarre diagnosed with rare disorders and extracted exonic SNVs

on chromosomes 1 to 22, averaging 8,871 variants per patient.

We refined the variant list by excluding common variants from *1KGP_noIBS*, gnomAD, *spain*, and

Navarre. The Navarrese filtering emerged as the most stringent, resulting in 2.1% of the initial set,

compared to 2.7% with gnomAD, 2.9% with *spain* frequencies, and the least restrictive, 4.9% with

1KGP_noIBS (Fig. 3). This underscores the effectiveness of Navarrese-specific filter in prioritizing

and streamlining genetic investigations.

DISCUSSION

In this study, we aimed to enhance diagnostic precision in the Navarrese population by exploring common population-specific variants. Utilizing WGS data from 358 individuals of Navarre, we identified 61,410 SNVs, with 21,174 common. Genetic analysis shows affinity with European populations and low differentiation with Spanish populations.

Focusing on exclusively common variants in Navarre compared with referenced populations, we obtained 1,069 SNVs, of which 676 were validated in another Navarrese cohort. None showed clinical or pharmacological relevance beyond what was observed in the Spanish population (18). This aligns with the expectation that common population variants are less likely to be associated with disease ethology.

Our findings underscore the relevance of considering population-specific factors in genomic diagnostics, which provides complementary insights alongside pangenome references (19). By identifying and excluding common variants within the Navarrese population, we have successfully refined the identification of potential disease-causing variants, contributing to the advancement of personalized medicine for individuals from Navarre. Further research will enhance these insights for broader applications.

214 DATA AVAILABILITY

215 Data is available from the corresponding author upon reasonable request.

REFERENCES

1.   Satam H, Joshi K, Mangrolia U, Waghoo S, Zaidi G, Rawool S, et al. Next-Generation Sequencing Technology: Current Trends and Advancements. Biology (Basel). 2023 Jul 13;12(7):997.

2.   Fattahi Z, Beheshtian M, Mohseni M, Poustchi H, Sellars E, Nezhadi SH, et al. Iranome: A catalog of genomic variations in the Iranian population. Hum Mutat. 2019 Nov 17;40(11):1968–84.

3.   1000 Genomes Project Consortium, Abecasis GR, Altshuler D, Auton A, Brooks LD, Durbin RM, et al. A map of human genome variation from population-scale sequencing. Nature. 2010 Oct 28;467(7319):1061–73.

4.   Gudmundsson S, Singer-Berk M, Watts NA, Phu W, Goodrich JK, Solomonson M, et al. Variant interpretation using population databases: Lessons from gnomAD. Hum Mutat. 2022 Aug 16;43(8):1012–30.

5.   Smetana J, Brož P. National Genome Initiatives in Europe and the United Kingdom in the Era of Whole-Genome Sequencing: A Comprehensive Review. Genes (Basel). 2022 Mar 21;13(3):556.

6.   Ramirez AH, Sulieman L, Schlueter DJ, Halvorson A, Qian J, Ratsimbazafy F, et al. The All of Us Research Program: Data quality, utility, and diversity. Patterns. 2022 Aug;3(8):100570.

7.   Mitsuhashi N, Toyo-oka L, Katayama T, Kawashima M, Kawashima S, Miyazaki K, et al. TogoVar: A comprehensive Japanese genetic variation database. Hum Genome Var. 2022 Dec 12;9(1):44.

8. Dopazo J, Amadoz A, Bleda M, Garcia-Alonso L, Alemán A, García-García F, et al. 267

Spanish Exomes Reveal Population-Specific Differences in Disease-Related Genetic

Variation. Mol Biol Evol. 2016;33(5):1205–18.

9. Peña-Chilet M, Roldán G, Perez-Florido J, Ortuño FM, Carmona R, Aquino V, et al. CSVS, a

crowdsourcing database of the Spanish population genetic variability. Nucleic Acids Res.

2021 Jan 8;49(D1):D1130–7.

10. NAGEN | Navarrabiomed [Internet]. [cited 2023 Nov 28]. Available from:

https://www.navarrabiomed.es/en/nagen

11. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from

high-throughput sequencing data. Nucleic Acids Res. 2010 Sep 1;38(16):e164–e164.

12. Landrum MJ, Lee JM, Riley GR, Jang W, Rubinstein WS, Church DM, et al. ClinVar: public

archive of relationships among sequence variation and human phenotype. Nucleic Acids

Res. 2014 Jan;42(Database issue):D980-5.

13. Amberger J, Bocchini CA, Scott AF, Hamosh A. McKusick's Online Mendelian Inheritance in

Man (OMIM(R)). Nucleic Acids Res. 2009 Jan 1;37(Database):D793–6.

14. Thorn CF, Klein TE, Altman RB. PharmGKB: The Pharmacogenomics Knowledge Base. In

2013. p. 311–20.

15. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines

for the interpretation of sequence variants: a joint consensus recommendation of the

American College of Medical Genetics and Genomics and the Association for Molecular

Pathology. Genetics in Medicine. 2015 May;17(5):405–24.

16. Anderson CA, Pettersson FH, Clarke GM, Cardon LR, Morris AP, Zondervan KT. Data quality

control in genetic case-control association studies. Nat Protoc. 2010 Sep 26;5(9):1564–73.

17. Khan F, Radovanovic A, Gojobori T, Kaur M. IBDDB: a manually curated and text-mining-

enhanced database of genes involved in inflammatory bowel disease. Database. 2021 Apr

30;2021.

18. Nunez-Torres R, Pita G, Peña-Chilet M, López-López D, Zamora J, Roldán G, et al. A

Comprehensive Analysis of 21 Actionable Pharmacogenes in the Spanish Population:

From Genetic Characterisation to Clinical Impact. Pharmaceutics. 2023 Apr

19;15(4):1286.

19. Liao WW, Asri M, Ebler J, Doerr D, Haukness M, Hickey G, et al. A draft human

pangenome reference. Nature. 2023 May 11;617(7960):312–24.

FUNDING
272
273 NAGEN1000 and PharmaNAGEN were supported by Navarra Gov (Dirección General de Industria,
274 Energia y Proyectos Estrategicos S3). GRANTS_NUMBERS: 0011-1411-2017-000032, 0011-1411-
275 2018-000047.
276

AUTHOR INFORMATION

Authors and Affiliations

Translational Bioinformatics Unit, Navarrabiomed, Hospital Universitario de Navarra (HUN), Universidad Pública de Navarra (UPNA), IdiSNA, Pamplona, Spain

Alberto Maillo, Estefania Huergo, Edurne Urrutia & David Gomez-Cabrero

Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia

Alberto Maillo & David Gomez-Cabrero

Genomics Medicine Unit, Navarrabiomed, Hospital Universitario de Navarra (HUN), Universidad Pública de Navarra (UPNA), IdiSNA, Pamplona, Spain

María Apellániz-Ruiz, Edurne Urrutia, María Miranda, Josefa Salgado, Sara Pasalodos-Sánchez, Luna Delgado-Mora, Óscar Teijido, Juan José Beloqui & Ángel Alonso

Servicio de Genética Médica-Hospital Universitario de Navarra (HUN), Pamplona, Spain

Josefa Salgado

Dp. Bioquímica y Biología Molecular-Universidad Pública de Navarra (UPNA), Pamplona, Spain

Josefa Salgado

Instituto de Genética Médica y Molecular (INGEMM), Hospital Universitario La Paz, Madrid, Spain

Luna Delgado-Mora

Department of Personalized Medicine, NASERTIC, Government of Navarra, Pamplona, Spain.

Ibai Goicoechea

Computational Medicine Platform, Andalusian Public Foundation Progress and Health-FPS, Sevilla, Spain.

Rosario Carmona, Javier Perez-Florido, Virginia Aquino, Daniel Lopez-Lopez, María Peña-Chilet & Joaquín Dopazo

Institute of Biomedicine of Seville, IBiS, University Hospital Virgen del Rocio/CSIC/University of Sevilla, Sevilla, Spain.

Rosario Carmona, Javier Perez-Florido, Daniel Lopez-Lopez & Joaquín Dopazo

FPS/ELIXIR-ES, Fundación Progreso y Salud (FPS), CDCA, Hospital Virgen del Rocio, Sevilla, Spain.

Rosario Carmona, Javier Perez-Florido & Joaquín Dopazo

Biomedical Research Networking Center in Rare Diseases (CIBERER), Health Institute Carlos III, 28029 Madrid, Spain

Rosario Carmona, Javier Perez-Florido, Daniel Lopez-Lopez, María Peña-Chilet & Joaquín Dopazo

CNAG-CRG, Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and
Technology, Barcelona, Spain
Sergi Beltrán

Universitat Pompeu Fabra (UPF), Barcelona, Spain
Sergi Beltrán

Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de
Barcelona (UB), Barcelona, Spain
Sergi Beltrán

Laboratory of Microbial Pathogenesis, Navarrabiomed, Pamplona, Spain
Iñigo Lasa
321

Contributions

Conceptualisation (AM, AA, DGC); Clinical and sample collection: (MM, LDM, OT, JS, MAR, SPS);

Formal analysis (AM, RC, JPF, VA, DLL, MPC); Data curation (AM); Investigation (AM, EH, MAR,

EU, DGC); Funding acquisition (JB, AA); Visualisation (AM, EH, MAR, EU); Writing-original draft

(AM, EH, MAR, EU, DGC) Writing-review & editing (AM, MAR, EH, EU, SB, SPS, IG, JD, IL, JB, AA,

DGC).

Corresponding authors

Correspondence to Ángel Alonso & David Gomez-Cabrero.

NAGEN scheme

Name	Department	NAGEN project
Anda Apiñaniz, Emma	Servicio de Endocrinología y Nutrición, HUN	NAGEN 1,000

Artigas López, Mercedes	Servicio de Genética Médica, HUN	NAGEN 1,000
Bandrés Elizalde, Eva	Servicio de Hematología, HUN	NAGEN 1,000
Basurte Elorz, Mª Teresa	Servicio de Cardiología, CHN	NAGEN 1,000
Brennan, Paul	NENC NHS Genomic Medicine Centre. Newcastle upon Tyne, UK.	NAGEN 1,000
Celaya Lecea, Concepción	Subdirección de Farmacia, SNS-O	PharmaNAGEN
Cuesta Zorita, Manuel Jesús	Servicio de Psiquiatría, Salud mental	NAGEN 1,000
Curi Chercoles, Sergio Miguel	Servicio de Neumología, HUN	NAGEN 1,000
De la Cruz Sánchez, Susana	Servicio de Oncología Médica, HUN	NAGEN 1,000
Erviti López, Juan	Subdirección de Farmacia, SNS-O	PharmaNAGEN
Fanlo Mateo, Patricia	Servicio de Medicina Interna, HUN	NAGEN 1,000
González, Luis Angel	AVANTIA 400+.	NAGEN 1,000
Gonzalo Etayo	Navarra de Servicios y Tecnología NASERTIC. Spain	NAGEN 1,000
Gorricho Mendívil, Javier	Subdirección de Farmacia, SNS-O	PharmaNAGEN

Guerra Lacunza, Ana	Servicio de Aparato Digestivo, HUN	NAGEN 1,000
Gut, Ivo	Centro Nacional de Análisis Genómicos CNAG. Spain	NAGEN 1,000
Ibáñez Bosch, Rosario	Servicio de Endocrinología y Nutrición, HUN	NAGEN 1,000
Jiménez, Jorge	Navarra de Servicios y Tecnología NASERTIC. Spain	NAGEN 1,000
Lasheras, Gorka	Navarra de Servicios y Tecnología NASERTIC. Spain	NAGEN 1,000
Lorea Bueno	Pharmamodelling	PharmaNAGEN
Maite Sarobe Carricas	Servicio de Farmacia Hospitalaria, HUN	PharmaNAGEN
Mendioroz Iriarte, Maite	Servicio de Neurología, HUN	NAGEN 1,000
Molinuevo Ruiz de Zarate, José Ignacio	Servicio de Oftalmología, HUN	NAGEN 1,000
Montes Díaz, Marta	Servicio de Anatomía Patológica, HUN	NAGEN 1,000
Navarro, Adela	Servicio de Cardiología, HUN	PharmaNAGEN
Onintza Sayar	Pharmamodelling	PharmaNAGEN
Pinillos, Iñaki	Navarra de Servicios y Tecnología NASERTIC. Spain	NAGEN 1,000

Purroy Irurzon, Carolina Eugenia	Servicio de Nefrología, HUN	NAGEN 1,000
Sagaseta de Ilurdoz Uranga, Mª Josefa	Servicio de Pediatría, HUN	NAGEN 1,000
Santesteban Muruzabal, Raquel	Servicio de Dermatología, HUN	NAGEN 1,000
Vicuña, Miren	Servicio de Digestivo, HUN	PharmaNAGEN
Viguria, Mª Cruz	Servicio de Hematología, HUN	PharmaNAGEN
Yoldi Petri, Mª Eugenia	Servicio de Pediatría, HUN	NAGEN 1,000
Zubicaray Ugarteche, José Jacinto	Servicio de Otorrinolaringología, HUN	NAGEN 1,000
Zudaire, Maite	Servicio de Hematología, HUN	PharmaNAGEN

ETHICS DECLARATIONS

Competing interests:

The authors declare no competing interests.

Ethics approval:

NAGEN1000 and PharmaNAGEN were approved by Navarra Ethics Committee for Clinical

Research (CEIC Navarra).

FIGURES LEGENDS

Figure 1: Workflow of this study. Abbreviations: *MGP*, Medical Genome Project; *1KGP*, 1000 Genomes Project; *1KGP_noIBS*, 1000 Genomes Project without Iberian population; *gnomAD,* Genome Aggregation Database; *MAF*, minor allele frequency; *SNV*, single nucleotide variant; *SNP*, single nucleotide polymorphism; *WGS*, whole genome sequencing; *WES*, whole exome sequencing; *LoF*, Loss-of-function; HPO, Human Phenotype Ontology; BP, biological process.

Figure 2: a) Principal component analysis of overlapped variants between NAVARREsel, MGP, and 1KGP (including all populations), and coloured by superpopulations. **b)** Principal component analysis of overlapped variants between NAVARREsel, MGP, and 1KGP (including exclusively European populations). Abbreviations: *PCA*, principal component analysis; AFR, African populations; AMR, American populations; EAS, east-Asian populations; SAS, south-Asian populations; EUR, European populations; IBS, Iberian populations in Spain; MGP, Medical Genome Project; TSI, Toscani in Italy; CEU, Utah residents with Northern and Western European ancestry; GBR, British in England and Scotland; FIN, Finnish in Finland.

Figure 3: a) Upset plot of common variants (MAF > 1%) of each population: NAVARREsel, *spain*, *1KGP_noIBS*, and gnomAD **b)** Resulting percentage of variants per patient (n=127) after removing common variants from Navarre, *spain*, gnomAD, or *1KGP_noIBS* populations. In the box plots, the median, upper, and lower quartiles are represented by the centre line and box bounds, respectively. Whiskers display the largest and smallest values within 1.5 times the interquartile range from the quartiles. Abbreviations: *1KGP_noIBS*, 1000 Genomes Project without Iberian population; gnomAD*,* Genome Aggregation Database; *spain*, integration of IBS and MGP populations.