Brain eQTLs of European, African American, and Asian ancestry improve interpretation of

schizophrenia GWAS

Yu Chen[1,2,16], Sihan Liu[1,3,16], Zongyao Ren[1], Feiran Wang[1], Yi Jiang[1], Rujia Dai[4], Fangyuan

Duan[1], Yan Xia[2], Cong Han[1], Zhilin Ning[5], Miao Li[1], Kai Yuan[2], Wenying Qiu [7], Xiao-Xin Yan

[8], Jiapei Dai[9], Richard F. Kopp[4], Jufang Huang[8], Shuhua Xu[10], Beisha Tang[14], Eric R.

Gamazon[11], Tim Bigdeli[12], Gershon Elliot[13], Hailiang Huang[2], Chao Ma[7*], Chunyu Liu[1,4*], and

Chao Chen [1,14,15*]

[1] MOE Key Laboratory of Rare Pediatric Diseases & Hunan Key Laboratory of Medical Genetics,

School of Life Sciences, and Department of Psychiatry, The Second Xiangya Hospital, Central

South University, Changsha, Hunan 410000, China.

[2] Broad Institute of MIT and Harvard, Cambridge, MA, USA.

[3] Institute of Rare Diseases, West China Hospital, Sichuan University, Chengdu, China.

[4] Department of Psychiatry, SUNY Upstate Medical University, Syracuse, NY, USA.

[5] Key Laboratory of Computational Biology, Shanghai Institute of Nutrition and Health,

University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai, China.

[6] Hunan Key Laboratory of Molecular Precision Medicine, Department of Neurosurgery, Xiangya

Hospital, and Hunan Key Laboratory of Medical Genetics, School of Life Sciences, Central South

University, Changsha, Hunan, China.

[7] Institute of Basic Medical Sciences, Neuroscience Center, National Human Brain Bank for

Development and Function, Chinese Academy of Medical Sciences; Department of Human

Anatomy, Histology and Embryology, School of Basic Medicine, Peking Union Medical College,

Beijing, China.

[8] Department of Human Anatomy and Neurobiology, Xiangya School of Medicine, Central South

University, Changsha, China.

[9] Wuhan Institute for Neuroscience and Engineering, South-Central University for Nationalities,

Wuhan, China.

[10] State Key Laboratory of Genetic Engineering, Center for Evolutionary Biology, Collaborative

Innovation Center of Genetics and Development, School of Life Sciences, Fudan University,

Shanghai, China.

[11] Division of Genetic Medicine Vanderbilt University School of Medicine Nashville, USA

[12] Institute for Genomics in Health, SUNY Downstate Health Sciences University, Brooklyn, New

York, USA

[13] Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL,

USA

[14] National Clinical Research Center for Geriatric Disorders, Xiangya Hospital, Central South

University, Changsha, China.

[15] Hunan Key Laboratory of Animal Models for Human Diseases, Central South University,

Changsha, China.

[16] These authors contributed equally: Yu Chen, Sihan Liu.

*Corresponding author. Email: machao@ibms.cams.cn (C.M.); LiuCh@upstate.edu (C.L.);

chenchao@sklmg.edu.cn (C.C.)

Abstract

Research on brain expression quantitative trait loci (eQTLs) has illuminated the genetic underpinnings of schizophrenia (SCZ). Yet, the majority of these studies have been centered on European populations, leading to a constrained understanding of population diversities and disease risks. To address this gap, we examined genotype and RNA-seq data from African Americans (AA, n=158), Europeans (EUR, n=408), and East Asians (EAS, n=217). When comparing eQTLs between EUR and non-EUR populations, we observed concordant patterns of genetic regulatory effect, particularly in terms of the effect sizes of the eQTLs. However, 343,737 cis-eQTLs (representing ~17% of all eQTLs pairs) linked to 1,276 genes (about 10% of all eGenes) and 198,769 SNPs (approximately 16% of all eSNPs) were identified only in the non-EUR populations. Over 90% of observed population differences in eQTLs could be traced back to differences in allele frequency. Furthermore, 35% of these eQTLs were notably rare (MAF < 0.05) in the EUR population. Integrating brain eQTLs with SCZ signals from diverse populations, we observed a higher disease heritability enrichment of brain eQTLs in matched populations compared to mismatched ones. Prioritization analysis identified seven new risk genes (*SFXN2*, *RP11-282018.3*, *CYP17A1*, *VPS37B*, *DENR*, *FTCDNL1*, and *NT5DC2*), and three potential novel regulatory variants in known risk genes (*CNNM2*, *C12orf65*, and *MPHOSPH9*) that were missed in the EUR dataset. Our findings underscore that increasing genetic ancestral diversity is more efficient for power improvement than merely increasing the sample size within single-ancestry eQTLs datasets. Such a strategy will not only improve our understanding of the biological underpinnings of population structures but also pave the way for the identification of novel risk genes in SCZ.

Keywords: eQTLs; Schizophrenia; GWAS signals; Population diversity

Introduction

Genome-wide association studies (GWAS) have identified 287 risk loci associated with schizophrenia (SCZ)[1]. Yet, the underlying mechanisms of these loci in disease development and progression remain poorly understood. Primarily, over 80% of GWAS risk loci reside in non-coding regions, devoid of protein-coding sequences, making it challenging to attribute them to specific genes. Moreover, predicting the regulatory effect of these loci proves challenging due to their tendency for gene-specific and tissue-specific effects. One effective strategy for gaining insights into their functions involves the integration of SCZ GWAS signals with expression quantitative trait loci (eQTLs), utilizing genotype and expression data from post-mortem brains. These brain eQTLs establish crucial links between risk genomic regions and gene expression levels, prioritizing potential disease risk genes through methods such as colocalization and transcriptome-wide association study (TWAS).

Past brain eQTL studies primarily focused on European (EUR) ancestry[2–6]. Global population diversity has not been adequately represented. Cross-population studies have shown that these European ancestry based models do not effectively predict gene expression in other ancestral groups[7]. This limitation weakens the power to detect TWAS associations in genetically diverse samples. While multi-ancestry eQTL meta-analyses in the human brain improve statistical power in uncovering risk loci shared across populations, key genetic variants regulating expression in specific underrepresented populations remain largely uncharted. The benefits of having brain eQTLs in diverse populations have not been thoroughly documented. Identifying eQTLs specific to biomedically underrepresented groups like African Americans (AA) and East Asians (EAS) can better understand the genetic contributions to disease susceptibilities and outcomes in these populations[7]. These populations have unique genetic variants and linkage disequilibrium (LD)

patterns. Additionally, previous studies have shown that combining eQTLs from different

ancestries can enable fine-mapping of causal variants and uncover potential novel mechanisms of

brain disorders[8,9]. Thus, the question of how to effectively leverage difference to uncover potential

novel mechanisms of brain disorders is a significant topic in the field.

To enhance the diversity in brain eQTL mapping and improve the interpretation of SCZ GWAS

across populations, we performed brain eQTL mapping in three major ancestries. Our data pool

comprised genotype and RNA-seq data of AA (n= 158) and EUR (n= 408) from the

PsychENCODE Consortium and EAS (n= 217) from the Chinese Human Brain Bank

(Supplementary Table 1). We juxtaposed non-EUR results against EUR to systematically examine

differences and similarities in the brain eQTLs. Further, we investigated the contributing factors

for eQTL differences across populations. By applying diverse population brain eQTLs to TWAS

and colocalization analysis of SCZ GWAS, we identified new risk genes and pathways. Lastly, we

identified likely causal variants by multi-ancestry fine-mapping. The two key questions we sought

to answer are: 1) what drives the brain eQTL differences across populations? 2) what do we gain

by studying brain eQTLs in diverse populations?

Results

To capture brain eQTLs across diverse populations, we utilized high-density genotype data

alongside high-throughput RNA-sequencing from prefrontal cortices. We derived AA (n = 158)

and EUR (n = 408) data from the BrainGVEx project of the PsychENCODE Consortium. We

generated EAS (n = 217) data from the Chinese Human Brain Bank (Fig. 1). Following rigorous

quality checks and preprocessing (Extended Data Fig. 1 and Extended Data Fig. 2), we compiled

expression data for 18,939 genes and genotype data at 6.4 million autosomal single nucleotide

polymorphisms (SNPs) across the three groups. Aligning the samples with the 1000 Genomes

Project reference populations, principal component analysis (PCA) confirmed the ancestry origins

of donors (Fig. 2a). We ensured sample identity consistency by comparing the genotypes from the

DNA and RNA samples. See methods for additional details.

Characterizing the cis-acting eQTLs in European, East Asian, and African American

population

We separately conducted cis-eQTLs mapping in the EUR, EAS, and AA samples using a 5%

empirical gene-level false discovery rate (FDR) threshold. This yielded 1,966,209 significant

eQTL signals covering 11,622 genes (eGenes) and 1,226,769 SNPs (eSNPs) across the populations

(see Fig. 2b and Supplementary Table 2). Specifically, we identified 1,616,818 eQTL signals

spanning 10,236 eGenes and 1,025,004 eSNPs for EUR; 562,058 eQTLs incorporating 5,000

eGenes and 416,025 eSNPs for EAS; and 143,736 eQTLs covering 3,039 eGenes and 121,079

eSNPs for AA. To identify credible SNP sets harboring plausible causal variants in cis-eQTLs, we

applied a fine-mapping method named SuSiE[10] to each population's eQTL results. The results

showed 966 credible SNP sets for 757 eGenes in the EUR cohort, 826 sets for 726 eGenes in EAS,

and 847 sets for 746 eGenes in AA (Supplementary Tables 3-5).

To investigate the genomic features of these cis-eQTLs, we evaluated the SNP distributions and

locations relative to various functional regions. 20% of cis-eQTLs in both EUR and non-EUR

populations were located within 10kb of transcription start site (TSS) regions (Fig. 2c). According

to the chromatin states predicted by Genomic Regulatory Elements and Gwas Overlap algoRithm

(GREGOR)[11] for prefrontal cortical tissue, eSNPs from the non-EUR populations were

significantly enriched in TSSs, promoters, and transcribed regulatory promoters or enhancers

($P_{Bonferroni} < 0.05$), identical to the observation in the EUR data. Moreover, using transcription

factor binding site (TFBS) annotation for 51 TFs, 46 and 49 TFs were significantly enriched with

cis-eQTLs in the AA and EAS populations respectively ($P_{Bonferroni} < 0.05$). All these TFBS were

also significantly enriched with cis-eQTLs in the EUR population.

To maximize the power of our population-based datasets, we employed METAL to amalgamate

the cis-eQTLs data from all three populations. This meta-analysis generated 598,193 eQTLs,

correlating 436,456 eSNPs with 5,209 eGenes (with a stringent P<2.5e-6) as depicted in Fig.

2b. However, meta-analysis does not consider the LD differences among the populations. We thus

used SuSiEx[16] to identify likely causal variants regulating expression by incorporating the LD

reference data from different populations. In total, SuSiEx identified 2,121 credible SNP sets for

1,801 eGenes in the 3-population combined data. Further details from the meta-analysis and fine-

mapping results can be found in Supplementary Table 6.

Population shared eQTLs showed similar regulatory effect across population

To evaluate the effect sizes across populations, we conducted a correlation test of effect size values

between the EUR and non-EUR populations. The effect sizes of the shared eQTLs between the

non-EUR and EUR were highly concordant (0.910 for eQTLs comparing EAS to EUR and 0.944

for AA to EUR; Fig 3a). To assess the robustness of the concordant effect size, we examined eQTL

slopes in the different populations. We looked at eQTLs obtained from the nominal, permutation,

and conditional tests, separately. eQTLs with smaller p-values or larger effect sizes showed greater

consistency across populations (Extended Data Fig. 3). Considering that sample size and

heterogeneity may influence the results, we randomly down-sampled the EUR data to match the

size of the non-EUR data. The results were similar to the results comparing all samples, showing

highly concordant effect sizes across populations (see Fig. 3c, Extended Data Fig. 3). In addition,

we compared the slope of the down-sampled EUR data with GTEx data (also of EUR). We

randomly sub-sampled 100 times and obtained a distribution of correlation values (R^2). The mean

R^2 was 0.94, which was not significantly different from the correlation between the EUR and non-

EUR population. We therefore concluded that the effect sizes of eQTLs in diverse populations

were mostly stable across human populations.

We evaluated the replicated rate (π_1), which gauges the true positive rate for the eQTLs identified

in the non-EUR populations that were also associated in the EUR population. The replicated rate

was π_1(EAS-EUR) = 0.86 and π_1(AA-EUR) = 0.91 (see Fig.3b). The π_1 for the non-EUR

populations in EUR was slightly but significantly lower than the π_1 between two EUR cohorts, as

represented by the GTEx cis-eQTL data (prefrontal cortex) in our EUR eQTL data (π_1(GTEx-

EUR) = 0.86, p-value = 0.023). To ensure a fair comparison of the replication rate of detected cis-

eQTLs in non- EUR data, we adjusted the EUR data to reflect the smaller sample size of the non-

EUR data. This adjustment enabled us to determine how many non-EUR cis-eQTLs were

confirmed in the adjusted EUR dataset. The adjusted results revealed a concordant trend: the

replicated rate between different populations was still slightly lower than that within the same

population assuming the same sample size (EUR-nonEUR average π_1 = 0.68, $EUR_{adjusted}$-EUR

average π_1 = 0.72, p-value = 0.037) (see Fig. 3d).

Population differences in brain cis-eQTLs are mainly caused by differences in SNP allele frequency while differences in effect size are small and uncertain

Here we defined those eQTLs that were exclusively observed in a single population as population-specific eQTLs. Upon analyzing the cis-eQTLs overlapping between populations, we identified 343,737 cis-eQTLs that were exclusively observed in the non-EUR populations, as detailed in Supplementary Table 2. This number represents approximately 17% of all eQTL pairs. These eQTLs involved 1,276 genes (about 10% of all eGenes) and 198,769 SNPs (around 16% of all eSNPs, Fig. 2d-f.). Specifically, there were 292,254 cis-eQTLs involving 165,300 eSNPs and 937 eGenes that were observed only in the EAS population and 51,483 cis-eQTLs involving 33,469 eSNPs and 339 eGenes that were observed only in the AA population.

To further characterize these non-EUR-specific eQTLs, we analyzed the variance, taking into account both the eQTL slope (effect size) and differences in allele frequency between populations. We found that more than 90% of the population differences in variance were attributable to differences in allele frequency. Moreover, to delve deeper into the distinctive characteristics of the eQTLs exclusive to the non-EUR groups, we leveraged two statistics, the fixation index (FST) and the minor allele frequency (MAF), retrieved from the 1000 Genomes Selection Browser[12]. A high FST value indicates that the measured locus has diverged over time in the populations. As expected, eSNPs detected only in the EAS or AA population displayed a significantly elevated FST when juxtaposed against eSNPs shared across populations (Wilcoxon test P < 2.2e-16). Meanwhile, the non-EUR-specific eSNPs showed higher MAF values in their respective source populations (Wilcoxon test P < 2.2e-16;) than in EUR. Of the 343,737 eQTLs absent in the EUR data, 309,363

were likely due to inadequate statistical power because they have smaller MAF in the EUR than

non-EUR population. Furthermore, 70,980 were rare in the EUR population ($MAF_{1KG} < 0.05$).

For the rest the eQTLs for which population differences could not be explained by differences in

MAF, a test for differences in eQTL slopes (effect sizes) was also conducted between the EUR

and non-EUR populations. The z-score of each independent eQTLs from conditional analysis was

calculated based on effect size and its standard deviation. Here the null hypothesis was that the

difference in eQTL effect size between the populations equals zero. No eQTL pairs detected by

conditional analysis could reject the null hypothesis. We then investigated if any eQTLs exhibited

opposite effect directions across populations. None of the independent eQTLs from the conditional

analysis displayed such effects. We relaxed our eQTL threshold using a nominal p-value < 0.05.

Only 534 eQTLs involving eighteen genes exhibited opposing eQTL effects between the EUR and

non-EUR populations. For example, the bitter taste receptor gene *TAS2R31* showed opposite

directions of eQTLs in EAS and EUR (Fig. 3e), which could be replicated using the blood eQTLs

from a previous study[7,13].

In conclusion, the variance in population differences can be largely attributed to differences in

allele frequency. The influence of effect size differences, on the other hand, appears to be minimal

and inconclusive.

Brain eQTLs from matched population can improve interpretation of SCZ GWAS

To determine if eQTLs detected from a specific population could explain the disease GWAS

signals and SNP-based disease heritability better than eQTLs from non-matching populations, we

undertook a two-step analysis. Firstly, we gathered SCZ GWAS summary statistics for the EUR,

EAS, and AA populations from previously published studies[14–16]. We employed the partitioned

LD-score regression (LDSR)[17] approach to assess the GWAS signal enrichment of these eQTLs.

eQTLs identified in the EAS population demonstrated a markedly higher enrichment in EAS-based

GWAS signals than eQTLs identified in the EUR population. Conversely, eQTLs identified in the

EUR population showed a greater enrichment for EUR-based GWAS signals than the eQTLs from

the EAS cohort. Both of these enrichments were statistically significant (Welch Modified Two-

Sample t-Test P-value < 0.001, Fig. 4a and 4b).

Besides the SNP heritability enrichment of all eQTLs, we also compared the significance of the

GWAS signals for population-unique eSNPs. We found that population-unique eSNPs tended to

have smaller p-values of disease association (i.e., stronger associations) in the corresponding

population than in the mismatched population (Welch Modified Two-Sample t-Test $P < 0.001$),

indicating the ability of population-unique eSNPs to explain the disease association and propose

the relevant gene, which will be missed when only one population is studied.

SCZ risk genes identified using eQTLs and GWAS from non-EUR population

To uncover risk genes and pathways for SCZ in non-EUR populations, we used TWAS, regulatory

Trait Concordance (RTC)[18], and summary data-based mendelian randomization (SMR)[19] to

prioritize SCZ candidate risk genes in non-EUR populations and compared them with risk genes

identified in the EUR (see Methods). In total, we prioritized ten risk genes in the EAS

(Supplementary Tables 7-9). It is worth noting that our TWAS analysis of AA data did not reveal

any significant associations. This lack of association in AA data might be attributed to the

relatively small sample size available from the AA SCZ GWAS.

Seven SCZ candidate risk genes (*SFXN2*, *RP11-282018.3*, *CYP17A1*, *VPS37B*, *DENR*, *FTCDNL1*,

and *NT5DC2*) uniquely discovered in the EAS population were assessed for allele frequency. From

these seven genes, two of them were located in chromosome 10 and five of them were located in

the chromosome 12. The eSNPs for these genes showed lower allele frequency in the EUR

population than in EAS. For instance, the GWAS signal chr12:123286491:A:G in the gene

VPS37B was found to be significant in the EAS population with a high AF of 0.48. In contrast,

this association was not significant in the EUR population with a markedly lower allele frequency

of 0.04. A parallel pattern emerged with the eSNP for *VPS37B*, with a markedly higher frequency

(chr12:123306558:G:A, MAF=0.24) in EAS than in the EUR (MAF = 0.04). These results further

confirm that allele frequency differences between populations can explain most of the

discrepancies between the EUR and EAS GWAS and the eQTL results (Supplementary Table 10).

Novel potential SCZ regulatory variations were refined utilizing brain eQTLs from non-

EUR population

Three of the ten risk genes identified in the EAS population were shared with the EUR population

(*CNNM2*, *C12orf65*, and *MPHOSPH9*), but differences in genetic architecture between

populations were still apparent. For example, two distinct significant SNPs in EAS and in EUR

were associated with SCZ on chromosome 10 (GWAS$_{EUR}$: chr10:104850632:G:A with GWAS p-

value = 6.4E-13; GWAS$_{EAS}$: chr10:104657300:T:C with GWAS p-value = 5.4E-12).   Using

eQTLs with GWAS signals in EAS and EUR separately, colocalization and SMR analysis

prioritized these two distinct GWAS SNPs to the same risk gene *CNNM2* in the two populations,

respectively.

To further investigate if these signals are located within any regulatory elements, we utilized the

non-coding variant annotation database (NCAD)[20] to annotate their regulatory information. Our

findings revealed that all the EAS GWAS risk SNPs are situated in enhancer regions

(Supplementary Table 11). The EAS GWAS risk SNPs near the *CNNM2* showed the strongest

evidence. Furthermore, we used the Lineage-specific Brain Open Chromatin Atlas[21] to investigate

whether this enhancer region shows different effects in major brain cell types. The results did not

show any cell type differences, which indicates the enhancer effect exists universally in major

brain cell types (Extended Data Fig. 4a-c). Integrating these insights, we discovered strong

evidence for multiple regulatory regions among the EAS eSNPs-chr10:104654577:T:C, which

have a high LD with *CNNM2* GWAS SNPs (LD R^2 = 1, p-value < 0.00001). Additionally, our

dual luciferase reporter assay results confirmed that the EAS eSNPs C-allele at

chr10:104654577:T:C significantly enhances luciferase activity compared to the reference vector,

as detailed in the Extended Data Fig. 4d-e.

High-confidence putative causal variants of SCZ using multi-ancestry brain eQTLs

To identify high-confidence putative causal variants from multiple populations, we applied

colocalization to our fine-mapped eQTLs and SCZ GWAS signals. In total, we identified four

SNP-gene-disease triplets in which the SNP colocalized with both gene expression and SCZ

GWAS (Supplementary Table 12, PiPcoloc = $PiP_{GWAS} \times PiP_{cis\text{-}eQTLs}$ > 0.1). The top genes with

PIP_{coloc} > 0.1 include *FURIN*, *ZNF823*, *RHOA*, and *VWA5A* (Fig. 4c). As an example, we identified

the strongest putative SCZ causal SNP for *FURIN*-chr15:91426560:G:A. This SNP is located in

the 3' untranslated regions (UTRs) of *FURIN*. Notably, this variant did not reach genome-wide

significance in the EAS population (P = 1.06E-3) likely due to limited statistical power. Our result

strongly supported that this causal variant is shared across populations, with causal probabilities

of 1. Previous study has also implicated the variant in both the EUR and EAS populations[13].

Discussion

In this study, we have created a brain transcriptome resource and identified eQTLs in the prefrontal cortex, specifically focusing on non-European populations. Our findings address the initial inquiries raised in the introduction. Firstly, we investigated the driver behind the variation in brain eQTLs across different populations. We found that differences in allele frequency and LD are instrumental in connecting disease susceptibility to gene expression regulation. This finding greatly augments our comprehension of genetic influences on gene expression in the human brain. Secondly, when examining brain eQTLs from diverse populations, we gained power to explain the GWAS heritability, uncover novel risk genes, and fine-map risk variants. We observed a pronounced enrichment of disease heritability among eQTLs in matched populations. In the non-EUR cohort, the allele frequencies and LD configurations facilitated the identification of seven novel SCZ risk genes. Additionally, we identified four high-confidence putative causal SCZ variants. These results highlight the utility of studying non-European cohorts.

Population differences appear to be more pronounced at the allele frequency level but are less so at the effect size level. In general, the estimated π_1 of eQTLs from non-EUR populations in EUR is lower compared to the rate observed between down-sampling-EUR and the EUR population cohort. Despite the relatively small sample size and statistical power, we still detected a significant number of 343,737 cis-eQTLs including 232,254 EAS eQTLs and 51,483 AA eQTLs that were significant only in the non-EUR populations. While over half of eSNPs in our non-EUR dataset were population-unique, 80% of eGenes identified in the non-EUR were also eGenes in the EUR data, but associated with different SNPs. The consistency of our observations with prior research involving diverse populations, including studies on gene expression[22,23], methylation[24], and chromatin accessibility[25], confirms the shared regulatory patterns across different populations.

Interestingly, some eQTLs showed contrasting effects across populations. ~0.1% of the non-EUR-

specific eQTLs displayed opposing directions in effect size. A notable example of this is the eQTL

rs2599400-*TAS2R31*, which showed opposite effects in different populations. Blood eQTLs from

EAS[26], EUR, and AFR[7] also support this observation. Prior studies have underscored the

population-specific variations in *TAS2R31*, linking these variations to differing sensitivity to the

bitter taste[27]. It is important to further investigate these discrepancies in gene expression regulation

among populations as they influence phenotypes. However, it should be noted that lack of

replication across all these studies limited our power to detect opposite-effect eQTLs.

Our findings underscore that enhancing genetic ancestral diversity is more efficient for power gain

than increasing the sample size within large-scale eQTLs datasets. Through our benchmarking of

eQTLs across three populations, we have established robust capabilities for identifying eQTLs

with a MAF greater than 0.2 and an effect size of 0.6 (Extended Data Fig. 5). Our power analysis

indicates that more than 30,000 individuals of European ancestry is needed to uncover all eQTLs

with MAF of 0.01 in this population, based on the estimated effect size of eQTLs exclusively

observed in non-EUR populations (Extended Data Fig. 5). For example, one the eQTL pair

(chr12:123306558:G:A-VPS37B) would require 3,215 EUR samples based on the power estimate

because of the low frequency in the EUR population (MAF = 0.04). However, the MAF of this

eSNP is 0.22 in EAS, which reduces the required sample size from 3,215 to 246. Thus,

incorporating a more diverse population would not only reveal numerous regulatory variants that

are rarer in EUR but more prevalent in non-EUR groups. Advancing towards a broader, more

diverse human reference dataset will facilitate more comprehensive investigations into the impact

of human demography on eQTL detection, thereby deepening our understanding of the distribution

and influence of genetic regulation in the human brain.

Differences in the genetic architecture underlying gene expression can help us to prioritize new

risk genes. Notably, prior research has reported that disease-associated loci tend to be skewed

towards variants with higher allele frequency in the discovery population, indicating that limited

statistical power may result in "missing" disease-association signals. Incorporating diverse

samples can enhance our ability to uncover the etiology of the disease. In our study, we identified

seven new SCZ risk genes using the non-EUR population, including *VPS37B* in the EAS

population. *VPS37B* is associated with calcium-dependent protein binding, providing new

evidence to support the involvement of the calcium-related pathway in SCZ risk in the EAS

population[28]. Another interesting candidate highlighted in our study was *CYP17A1* (RTC = 0.99).

The corresponding GWAS signal was significant in the EAS and EUR populations (P_{EAS} = 4.5E-

8; MAF_{EAS} = 0.48; P_{EUR} = 2.6E-13; MAF_{EUR} =0.30), while the corresponding eSNP in EAS

population (MAF=0.48) showed extremely low frequency in EUR population (MAF<0.001).

CYP17A1 notably serves as an enzyme important for the production of glucocorticoids and sex

hormones, such as estrogen, which have been linked to schizophrenia[29–31] .

Besides enhancing the power for detecting risk genes, the inclusion of brain eQTLs from diverse

populations improves the ability to fine-map SCZ GWAS loci, identifying new regulatory variants

which have the potential to regulate downstream gene expression. This approach aids in

interpretation, thereby facilitating subsequent computational and experimental functional

investigations. Our result revealed a potential novel regulatory region near the population-shared

risk gene *CNNM2*. This discovery showcases the power of leveraging diverse populations.

By leveraging the multi-ancestry information, trans-ancestry fine-mapping also helped us identify

high-confidence putative causal variants. In addition to previously validated genes, our study

uncovered another significant finding at chr3:50297330:A:G-*RHOA*-SCZ through trans-ancestry

colocalization. *RHOA* encodes a member of the Rho family of small GTPases, pivotal in signal

transduction cascades by toggling between inactive GDP-bound and active GTP-bound states.

Some limitations of our study merit attention. Our sample size is still relatively small. Our analysis

suggests that expanding the sample size would capture a larger set of eQTLs. The small sample

size could have significantly impacted the comprehensiveness of our findings. The modest sample

size of the AA eQTL dataset and the SCZ GWAS cohort likely led to the failure of TWAS in the

AA population.

In conclusion, we presented a novel genome-wide map of human brain gene expression regulation.

Importantly, this resource bridges the gap between neuropsychiatric GWAS and brain gene

expression profiling in non-European populations. Our study emphasizes the significance of this

new atlas of brain gene expression regulation in non-European populations for advancing our

understanding of human diversity, addressing health disparities, and developing precision

medicine.

Materials and methods

Sample collection and sequencing

We collected 217 prefrontal cortical samples of Han Chinese ancestry from the National Human

Brain Bank for Development and Function[32,33]; the samples were handled according to the

standardized operational protocol of the China Human Brain Banking Consortium, under the

approval of the Institutional Review Board of the Institute of Basic Medical Sciences, Chinese

Academy of Medical Sciences, Beijing, China (Approval Number: 009-2014).

We sequenced 217 samples following the BGISEQ-500 protocol outsourced to BGI. 1μg genomic

DNA was randomly fragmented by Covaris, the fragmented DNA was selected by Agencourt

AMPure XP-Medium kit to an average size of 200-400bp, followed by adapter ligation, PCR

amplification, and the products were recovered by the AxyPrep Mag PCR clean up kit. The double-

stranded PCR products were heat-denatured and circularized by the splint oligo sequence. The

single-strand circle DNA (ssCir DNA) was formatted as the final library and qualified by QC.

Sequencing was performed on BGISEQ-500 platform with an average depth of 10X.

Total RNA was extracted from the brain tissue using Trizol (Invitrogen, Carlsbad, CA, USA)

according to manufacturers' instructions. Then, total RNA was qualified and quantified using a

Nano Drop and Agilent 2100 bioanalyzer (Thermo Fisher Scientific, MA, USA). Ribo-zero

method was used to remove the rRNA. Purified mRNA was fragmented into small pieces with

fragment buffer at an appropriate temperature. The cDNAs were purified by magnetic beads. After

purification, A-Tailing Mix and RNA Index Adapters were added by incubating to carry out end

repair. The cDNA fragments with adapters were amplified by PCR, and the products were purified

by Ampure XP Beads. The library was validated on the Agilent Technologies 2100 bioanalyzer

for quality control. The final library was amplified with phi29 (Thermo Fisher Scientific, MA,

USA) to make DNA nanoball (DNB), DNBs were loaded into the patterned nanoarray and single

end 50 base reads were generated on BGISEQ500 platform.

Data quality control

Raw sequencing reads were filtered to get clean reads by using SOAPnuke (v1.5.6)[34], and

FastQC[35] was used to evaluate the quality of sequencing data via several metrics, including

sequence quality per base, sequence duplication levels, and quality score distribution for each

sample. The average quality score for overall DNA and RNA sequences was above 30, indicating

that a high percentage of the sequences had high quality.

Variant identification

Clean DNA sequencing reads were mapped to the human reference genome hg19 (GRCh37) using

BWA-MEM algorithm (BWA v. 0.7.128)[36]. Ambiguously mapped reads (MAPQ <10) and

duplicated reads were removed using SAMtools v. 1.29[37] and PicardTools v. 1.1 respectively.

Genomic variants were called following the Genome Analysis Toolkit software (GATK v. 3.4.4.6)

best practices.

Population validation, imputation, and filtering

We used PLINK to infer the genomic ancestry of each sample in this study by combining our

genotype data and the genotype data from the 1000 Genomes Project[38]; no sample was excluded.

Using Michigan Imputation Server[41], EAS genotypes were imputed into the 1000 Genomes Project

phase 3 EAS reference panel by chromosome and subsequently merged. Imputed genotypes were

filtered for LD $R^2 < 0.3$, Hardy-Weinberg equilibrium p-value $< 10e-6$ and MAF < 0.05, resulting

in ~ 6 million autosomal SNPs.

For AA population, genotypes were imputed into the 1000 Genomes Project phase 3 AA reference

panel by chromosome and subsequently merged. To further confirm the ancestry of the African

American samples, all AA samples were evaluated for their ancestry with three broad population

groups with PC1 ≥ 25% AFR and < 25% AMR, < 25% EAS, < 25% SAS; clustering of individuals

in each broad population group with the 1000 Genomes Project reference populations are shown

in Fig.2a.

Sex check and sample swap identification

The sex of each sample was inferred with SNPs using PLINK. In the EAS cohort, two samples

were identified as sex-mismatched and were subsequently removed in downstream analysis.

Quality control was performed on genotypes using sample Binary Alignment Map (BAM) files to

detect any sample identity swaps between the RNA and DNA experiments. The QTLtools match

function[39] confirmed that all samples were appropriately matched.

Gene expression quantification and quality control

The RNA-sequencing reads were mapped using STAR (2.4.2a)[40] and the genes and transcripts

quantification was performed using RSEM (1.3.0)[41]. Raw read counts were log-transformed using

R package VOOM[42], thereafter filtering those with log2(CPM) < 0 in more than 75% of the

samples. Mitochondrial DNA and X and Y chromosome-derived transcripts were excluded.

Samples with a Z-score (measured for inter-sample connectivity) less than -3 were also discarded.

Finally, quantile normalization was utilized to equalize distributions across samples.

Covariate selection

To measure technical covariates, quality control metrics were collected using STAR, PicardTools

v1.139 and RNASeQC. Principal components of the metrics data were calculated and included as

SeqPCs for covariate selection. Hidden covariates were measured using probabilistic estimation

of expression residuals (PEER)[43] and found to be significantly correlated with technical and

biological covariates such as experimental batch, RNA Integrity Number (RIN), sex, and age of

death. Based on the Bayesian information criterion (BIC) score, redundant covariates were

removed to avoid overfitting. A forward and backward selection procedure was followed, and

PEER hidden factors and known covariates were added. The covariate with the higher BIC score

was selected for subsequent QTL mapping. PEER was run with varying numbers of inferred

hidden factors: 5, 10, 15, 20, 25, 30, 35, 40, 45 and 50, independently.

cis-eQTL mapping

Cis-eQTL mapping was performed using QTLtools, accounting for PEER factors, with a defined cis window spanning one megabase upstream and downstream of the gene/intron cluster body. Using the PEER from gene expression matrix, twenty hidden covariates were identified and adjusted. To detect all available QTLs, QTLtools was conducted in nominal pass mode. To identify the best nominal associated SNP per phenotype, QTLtools was executed in the permutation pass mode. Additionally, to identify SNPs with independent effects on regulating gene expression, QTLtools was run in the conditional pass mode. SNPs with P-values below the threshold of the permutation threshold is classified as significant QTLs.

To address potential sample size disparities that could impact the results, the EUR data were randomly sampled with various sample size (150, 200, 250, 300, 350, and 400) and applied the same analytical pipeline while exploring the relationship between sample size and number of QTLs.

eQTL meta-analysis

The SNP-gene pairs with a genome-wide P value reaching the threshold of 10e-6 for each population were collected and performed the meta-analysis. Standard fixed-effects meta-analysis were used to combine all data into a single regression model by METAL[44]. The meta-analysis assumes a fixed-effects size, as well as constant error variance, across all data.

eQTL fine-mapping

The initial step of fine-mapping involved using the in-sample LD of the three populations. We extracted common variants with MAF > 5% from each group and used PLINK to determine the LD regions of these common variants for each population. To eliminate strand flipping and alignment issues, multi-allelic variations and indels were removed. Next, SuSiEx was applied to

merge the eQTLs summary statistics from the three groups. Credible set is defined as a set of

putative causal variants. A credible set was discarded if it lacked genetic variants reaching genome-

wide significance ($p < 1e-6$) in either the population-unique eQTLs or cross-population meta-

eQTLs. By considering prior knowledge and the observed data, this method provides a posterior

probability (PIP) for each variant being the causal one in the associated region. Variants with high

PIPs are then considered strong candidates for functional follow-up studies.

Functional enrichment

To determine functional enrichment, GREGOR[48] was performed to test the eQTL enrichment.

GREGOR calculated the enrichment value based on the observed and expected overlap within

each annotation. To conduct our analysis, the 15-state ChromHMM model BED (Browser

Extensible Data) files from the Roadmap Epigenetics Project[45], and 78 consensus transcription

factor and DNA-protein binding site BED files existing in multiple cells were downloaded. Fifty

binding proteins showed cortical brain expression in EAS and AA populations data[46].

The fraction of shared eQTLs between non-EUR and EUR populations

Sharing rate was assessed based on significant eQTLs in the discovery dataset by estimating the

proportion of true associations (π_1) on the distribution of corresponding p-values of the overlapping

eQTLs in the replication dataset[47].

F_{ST} and MAF analysis

Fixation index (F_{ST}) was estimated using vcftools following the Weir and Cockerham approach

for each eSNP[48]. The population-divergent SNPs were defined as those with $F_{ST} >= 0.05$ and

population-shared SNPs as those with $F_{ST} < 0.05$. To generate the list of population-unique QTLs

and population-shared QTLs, we collected the overlap of eQTLs from the pairwise comparisons

of the list of AA eQTLs, EAS eQTLs, and EUR eQTLs. Finally, Fisher's exact test was performed

between population-unique QTLs and population-shared QTLs to test the contribution of MAF in

the QTL comparison.

Variance explained

Variance explained, which combines the effect size (beta) and frequency of the allele (f), can be

considered an approximate measure of a causal variant's importance within a population. Variance

is approximated using the formula $2f(1 - f)\log(beta)^2/(\pi^2/3)$[49]. Although these variants often exhibit

similar odds ratios across populations, their allele frequencies may differ. By considering both the

effect size (OR) and the frequency of the risk allele (f), the variance explained offers a valuable

approximation of a causal variant's significance within a given population.

Power estimation

We used R to calculate the sample size needed to achieve a given power level in a chi-square test,

based on an assumed effect size and a significance threshold. This analysis starts by setting initial

values for power, effect size, and p-value threshold. Then, the critical chi-square statistic required

to meet the power level is computed. A function, *calculate_ncp*, is defined to calculate the non-

centrality parameter from the p-value and degrees of freedom, adjusting for the critical chi-square

statistic. Subsequently, the non-centrality parameter is computed for the given power and p-value

threshold. Another function, *af_n_relation*, is created to determine the relationship between allele

frequency and sample size, incorporating the effect size and the non-centrality parameter. Finally,

the code iteratively solves for the sample size corresponding to a range of allele frequencies, thus

enabling the determination of the necessary sample size for different allele frequencies to maintain

the specified power level in the chi-square test.

Partitioned LDSR

Partitioned LD score regression v1.0.1[50] was used to measure the enrichment of GWAS summary

statistics in each functional category by accounting for LD. Brain QTL annotations were created

by eSNP, mapped to the corresponding 1000 Genome reference panel. LD scores were calculated

for each SNP in the QTL annotation using an LD window of 1cM in 1000 Genomes European

Phase 3 and 1000 Genomes Asian Phase 3 separately. Enrichment for each annotation was

calculated by the proportion of heritability explained by each annotation divided by the proportion

of SNPs in the genome falling in that annotation category. We then applied Welch Modified Two-

Sample t-Test on enrichment values generated from QTLs in the two populations.

Colocalization

Conditional association was used to test for evidence of colocalization. This method compares the

p-value of association for the lead SNP of an eQTL before and after conditioning on the GWAS

hit. The equation for the regulatory trait concordance (RTC) Score is as follows: RTC= (N_{SNPs} in

an LD block/$Rank_{GWAS\_SNP}$)/ N_{SNPs} in an LD block. The rank denoted the number of SNPs, which

when used to correct the expression data, has a higher impact on the QTL than the GWAS SNPs.

RTC values close to 1.0 indicated causal regulatory effects. A threshold of 0.9 was used to select

causal regulatory elements.

Colocalization of fine-mapped variations from complex traits and cis-eQTL correlations were

performed. Based on complex trait and cis-eQTL fine-mapping data, a posterior inclusion

probability of colocalization for a variant was calculated as a product of PIP for GWAS and PIP

for the cis-eQTLs (PIPcoloc = PIP_{GWAS} * $\text{PIP}_{\text{cis-eQTLs}}$).

Summary-data-based Mendelian randomization

SMR[19] was applied on SCZ GWAS summary data to prioritize candidate genes. Significant QTLs

identified in the previous analysis (FDR < 0.05), were combined with filtered GWAS summary

data (p < 5e-8) to perform the SMR test. In general, we used the default parameters suggested by

the developers of the SMR software. These included the application of heterogeneity independent

instruments (HEIDI) testing, filtering out hits that arose from significant linkage with

pleiotropically associated variants (LD cutoff of P = 0.05 in the HEIDI test, as suggested by SMR).

Genes with an empirical P that passed Bonferroni correction in the SMR test and a P > 0.05 in the

HEIDI test were considered as risk genes.

Prioritizing genes underlying GWAS hits using TWAS

In this research, we initially developed gene expression prediction models for distinct populations

using MetaXcan software[51]. Following this, we integrated these models with GWAS (Genome-

Wide Association Studies) summary statistics specifically focused on schizophrenia. This

integration aimed to generate gene-level z-scores representing the association of the genetically-

determined expression for a gene from its prediction model with the phenotype. TWAS enabled

us to compute P-values and subsequently prioritize genes in relation to their association with

schizophrenia risk.

Plasmid construction

We obtained the 55 bp SNP-centered DNA sequence from UCSC Genome Browser

(GRCh38/hg38), then added the sticky end of restriction enzymes KpnI and NheI at both ends of

the 55bp sequence to synthesis primers. Primer annealing to obtained double-strand sequence and

then inserted into pGL3-Promoter Vector (Promega) using FastDigest enzymes (ThermoFisher)

and T4 DNA Ligase (Invitrogen). We valid the vector sequence using sanger sequence by primer

GLP2 and RVP3.

Dual Luciferase Reporter Assay

For transfection, we used SH-SY5Y and HS-683 cells to perform the experiments. Transfecting

cells at 50-60% confluency, cells were co-transfected with 500ng reconstruction vector and 10ng

pRL-TK using Lipofectamine 3000 Transfection Reagent (ThermoFisher) in 24 well plates. After

48h transfection, using Dual Luciferase Reporter Assay Kit (Promega) to measure the firefly

luciferase activity and renilla luciferase activity, the luminescence was detected using Tube

Luminometer (Berthold Sirius).

Fig. 1. Study design. We examined genotype and RNA-seq data from African Americans (AA),

Europeans (EUR), and East Asians (EAS) to identify expression quantitative trait loci (eQTLs)

unique to non-European populations and their role in schizophrenia risk.

Fig. 2: Identification and characterization of eQTLs. a) PCA plot showing the population structure

of individuals in our study as well as the 1000 Genomes Project. AFR: African; AMR: American;

EAS: East Asian; EUR: European; b) Circos manhattan plot of significant eQTL genes among the

three populations with highlighted top 50 fine-mapped eGenes. d-f) Upset plot showed overlap

among the significantly associated eQTL pairs (d) eGenes (e) as well as eSNPs (f) between

populations.

580 Fig. 3: Analysis of the regulatory patterns. a) relationship between sample size and the # of

581 detected eQTLs. b) Effect sizes for common (MAF> 1%) sentinel cis-eQTLs across EA and AA

582 populations; c) Comparison of MAF between population-shared and non-EUR unique eSNPs; d)

583 downsampling to estimate π_1 between non-European and European eQTLs . e) example of opposite

584 effect eQTL *TAS2R31*- rs2599400.

Fig. 4: Explanation of SCZ GWAS signals and prioritization of candidate genes. a) integration strategies; b) GWAS enrichment results from LDSR. ***: Welch Modified Two-Sample t-Test P < 0.001; c) Fine-mapped colocalization results. Each point represents an eQTL pais, the x-axis represents the GWAS pip for that eSNP, and the y-axis represents the eQTL pip for that eSNP. Red points represent $pip_{GWAS}*pip_{eQTL}>0.1$ and are labeled with the eGene

Data availability

The raw sequence data reported in this paper have been deposited in the Genome Sequence Archive in BIG Data Center, Beijing Institute of Genomics (BIG), Chinese Academy of Sciences, under accession numbers HRA000108, HRA000108 that can be accessed at https://bigd.big.ac.cn/gsa-human. eQTLs summary results can be downloaded while requesting to the corresponding author.

Code availability

Codes are available at https://github.com/liusihan/population-compare-pipeline.

Acknowledgments

We thank all donors and their families. Tissue was provided by the Human Brain Bank, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China, the Chinese Brain Bank Center, and the Xiangya School of Medicine Brain Bank. We acknowledge Stanley Center for Psychiatric Research for supporting Hailiang Huang and Yu Chen working on this project.

Funding

This study was supported by National Natural Science Foundation of China (Grants Nos. 82022024, 31571312, and 91632116), the National Key R&D Project of China (Grants No. 2016YFC1306000), The Science and Technology Innovation Program of Hunan Province (2021RC4018, 2021RC5027), and NIH grant 1R01MH126459-01A1.

Author contributions

Y.C. and S.L. wrote the manuscript, analyzed the data and performed all computations. F.W. and

Y. J. contributed to eQTLs analysis of EUR population. Z.R., F.D., C.H and M.L. extracted DNA

and RNA, as well as collected sample information. R.D., Y.X., and R.K. substantively revised the

manuscript. Z.N., S.X. and H.H. participated in the design of comparing the brain regulatory

architecture. K.Y., W.Q., C.M., X.Y., A.B., J.D., J.H., B.T. and C.L. provided primary sequenced

samples and data including their clinical information. C.M., C.L. and C.C. conceived, designed

and supervised the study and modified the manuscript.

Conflict of Interest

All the authors declare no competing financial interests.
632

633 **Ethnics**
634 Ethics Committee of Central South University gave ethical approval for this work

References

1. Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* **604**, 502–508 (2022).

2. Peterson, R. E. *et al.* Genome-wide Association Studies in Ancestrally Diverse Populations: Opportunities, Methods, Pitfalls, and Recommendations. *Cell* **179**, 589–603 (2019).

3. Hindorff, L. A. *et al.* Prioritizing diversity in human genomics research. *Nat. Rev. Genet.* **19**, 175–185 (2018).

4. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**, 580–585 (2013).

5. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* **12**, e1001779 (2015).

6. Wang, L. *et al.* Brain Banks Spur New Frontiers in Neuropsychiatric Research and Strategies for Analysis and Validation. *Genomics Proteomics Bioinformatics* **17**, 402–414 (2019).

7. Mogil, L. S. *et al.* Genetic architecture of gene expression traits across diverse populations. *PLoS Genet.* **14**, e1007586 (2018).

8. de Klein, N. *et al.* Brain expression quantitative trait locus and network analyses reveal downstream effects and putative drivers for brain-related diseases. *Nat. Genet.* **55**, 377–388 (2023).

9. Zeng, B. *et al.* Multi-ancestry eQTL meta-analysis of human brain identifies candidate causal variants for brain-related traits. *Nat. Genet.* **54**, 161–169 (2022).

10. Zou, Y., Carbonetto, P., Wang, G. & Stephens, M. Fine-mapping from summary data with the 'Sum of Single Effects' model. *PLoS Genet.* **18**, e1010299 (2022).

11. Schmidt, E. M. *et al.* GREGOR: evaluating global enrichment of trait-associated variants in

epigenomic features using a systematic, data-driven approach. *Bioinforma. Oxf. Engl.* **31**,

2601–2606 (2015).

12. Pybus, M. *et al.* 1000 Genomes Selection Browser 1.0: a genome browser dedicated to

signatures of natural selection in modern humans. *Nucleic Acids Res.* **42**, D903-909 (2014).

13. Ning, Z. *et al.* Expression profiles of east-west highly differentiated genes in Uyghur

genomes. *Natl. Sci. Rev.* **10**, nwad077 (2023).

14. Lam, M. *et al.* Comparative genetic architectures of schizophrenia in East Asian and

European populations. *Nat. Genet.* **51**, 1670–1678 (2019).

15. Bipolar Disorder and Schizophrenia Working Group of the Psychiatric Genomics

Consortium. Electronic address: douglas.ruderfer@vanderbilt.edu & Bipolar Disorder and

Schizophrenia Working Group of the Psychiatric Genomics Consortium. Genomic

Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell* **173**,

1705-1715.e16 (2018).

16. Bigdeli, T. B. *et al.* Contributions of common genetic variants to risk of schizophrenia

among individuals of African and Latino ancestry. *Mol. Psychiatry* **25**, 2455–2467 (2020).

17. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide

association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).

18. Nica, A. C. *et al.* Candidate causal regulatory effects by integration of expression QTLs with

complex trait genetic associations. *PLoS Genet.* **6**, e1000895 (2010).

19. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex

trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).

20. Feng, X., Liu, S., Li, K., Bu, F. & Yuan, H. NCAD v1.0: A database for non-coding variant

annotation and interpretation. *J. Genet. Genomics Yi Chuan Xue Bao* S1673-8527(23)00257–

6 (2023) doi:10.1016/j.jgg.2023.12.005.

21. Fullard, J. F. *et al.* An atlas of chromatin accessibility in the adult human brain. *Genome Res.*

28, 1243–1252 (2018).

22. Kachuri, L. *et al.* Gene expression in African Americans, Puerto Ricans and Mexican

Americans reveals ancestry-specific patterns of genetic architecture. *Nat. Genet.* **55**, 952–963

(2023).

23. Lappalainen, T. *et al.* Transcriptome and genome sequencing uncovers functional variation

in humans. *Nature* **501**, 506–511 (2013).

24. Fraser, H. B., Lam, L. L., Neumann, S. M. & Kobor, M. S. Population-specificity of human

DNA methylation. *Genome Biol.* **13**, R8 (2012).

25. Tehranchi, A. *et al.* Fine-mapping cis-regulatory variants in diverse human populations. *eLife*

8, e39595 (2019).

26. Ning, Z. *et al.* Expression profiles of east-west highly differentiated genes in Uyghur

genomes. *Natl. Sci. Rev.* **10**, nwad077 (2023).

27. Wooding, S. P. & Ramirez, V. A. Global population genetics and diversity in the TAS2R

bitter taste receptor family. *Front. Genet.* **13**, 952299 (2022).

28. Andrade, A. *et al.* Genetic Associations between Voltage-Gated Calcium Channels and

Psychiatric Disorders. *Int. J. Mol. Sci.* **20**, 3537 (2019).

29. Heringa, S. M., Begemann, M. J. H., Goverde, A. J. & Sommer, I. E. C. Sex hormones and

oxytocin augmentation strategies in schizophrenia: A quantitative review. *Schizophr. Res.*

168, 603–613 (2015).

30. de Boer, J., Prikken, M., Lei, W. U., Begemann, M. & Sommer, I. The effect of raloxifene

augmentation in men and women with a schizophrenia spectrum disorder: a systematic

review and meta-analysis. *NPJ Schizophr.* **4**, 1 (2018).

31. Chiappelli, J. *et al.* Disrupted glucocorticoid--Immune interactions during stress response in

schizophrenia. *Psychoneuroendocrinology* **63**, 86–93 (2016).

32. Yan, X.-X., Ma, C., Bao, A.-M., Wang, X.-M. & Gai, W.-P. Brain banking as a cornerstone

of neuroscience in China. *Lancet Neurol.* **14**, 136 (2015).

33. Qiu, W. *et al.* Standardized Operational Protocol for Human Brain Banking in China.

Neurosci. Bull. **35**, 270–276 (2019).

34. Chen, Y. *et al.* SOAPnuke: a MapReduce acceleration-supported software for integrated

quality control and preprocessing of high-throughput sequencing data. *GigaScience* **7**, 1–6

(2018).

35. Trivedi, U. H. *et al.* Quality control of next-generation sequencing data without a reference.

Front. Genet. **5**, 111 (2014).

36. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinforma. Oxf. Engl. **25**, 1754–1760 (2009).

37. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**,

2078–2079 (2009).

38. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based

linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

39. Delaneau, O. *et al.* A complete tool set for molecular QTL discovery and analysis. *Nat.*

Commun. **8**, 15452 (2017).

40. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinforma. Oxf. Engl.* **29**, 15–

21 (2013).

41. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

42. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model

analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29 (2014).

43. Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of

expression residuals (PEER) to obtain increased power and interpretability of gene

expression analyses. *Nat. Protoc.* **7**, 500–507 (2012).

44. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of

genomewide association scans. *Bioinforma. Oxf. Engl.* **26**, 2190–2191 (2010).

45. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human

epigenomes. *Nature* **518**, 317–330 (2015).

46. Arbiza, L. *et al.* Genome-wide inference of natural selection on human transcription factor

binding sites. *Nat. Genet.* **45**, 723–729 (2013).

47. Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proc. Natl.*

Acad. Sci. U. S. A. **100**, 9440–9445 (2003).

48. Bhatia, G., Patterson, N., Sankararaman, S. & Price, A. L. Estimating and interpreting FST:

the impact of rare variants. *Genome Res.* **23**, 1514–1521 (2013).

49. Pawitan, Y., Seng, K. C. & Magnusson, P. K. E. How many genetic variants remain to be

discovered? *PloS One* **4**, e7969 (2009).

50. Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies

disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).

51. Barbeira, A. N. *et al.* Exploring the phenotypic consequences of tissue specific gene

expression variation inferred from GWAS summary statistics. *Nat. Commun.* **9**, 1825 (2018).

a

b

c