Mitral annular plane systolic excursion to left atrial volume ratio

- a strainless relation with left ventricular filling pressures

Thomas Lindow MD PhD[1,2,3], Hande Oktay Tureli MD PhD[4], Charlotte Eklund Gustafsson, Bsc, PhD,[2] Daniel Manna MD,[2] Björn Wieslander MD PhD[2,5], Per Lindqvist BSc PhD[4], Ashwin Venkateshvaran, MSc, PhD[1]

1. Clinical Physiology, Clinical Sciences, Lund University, Lund, Sweden
2. Department of Clinical Physiology, Department of Research and Development, Region Kronoberg, Växjö Central Hospital, Växjö, Sweden
3. Kolling Institute, Royal North Shore Hospital, and University of Sydney, Sydney, Australia
4. Department of Clinical Physiology, Umeå University Hospital, Umeå, Sweden
5. Department of Clinical Physiology, Karolinska University Hospital, and Karolinska Institutet, Stockholm, Sweden

Funding: This work has been funded in part by grants to TL from Region Kronoberg and the Swedish Heart Lung Association, to PL from The Swedish Heart and Lung foundation (20160787, 20200160) and the Swedish Research Council (2019-01338).

Disclosures: TL, HT, CEG, DM, BW, AV: None. PL: Consulting fees from Pfizer.

Data availability: The data underlying this article can be shared upon reasonable request to the corresponding author.

Word count: Total: 4472. Main text: 1966; abstract: 249; figure legends: 205; references: 942; tables: 1110.

Corresponding author: Thomas Lindow, MD, PhD, Thomas.akesson_lindow@med.lu.se, Department of Clinical Physiology, Växjö Central Hospital, Region Kronoberg, 351 88 Växjö, Sweden, +46470587574

Background: Left atrial reservoir strain (LASr) offers diagnostic and prognostic value in patients with heart failure. However, LASr may be technically challenging and is not routinely evaluated in clinical practice. Since LASr is a consequence of left atrial (LA) stretch during apical descent of the mitral annulus, we hypothesized that a ratio between mitral annular plane systolic excursion (MAPSE) and LA volume (LAV) may offer similar diagnostic value as LASr.

Aim: To investigate the relationship between MAPSE/LAV and LASr and evaluate the diagnostic performance of MAPSE/LAV to identify patients with elevated LV filling pressure.

Methods: MAPSE/LAV and LA strain measures were obtained in patients referred for echocardiography due to aortic stenosis, and in patients who had undergone clinically indicated right heart catheterization (RHC) with simultaneous echocardiography.

Results: In 93 patients with moderate aortic stenosis, MAPSE/LAV was moderately correlated with LASr (r=0.57) but was lower in patients with elevated compared to normal LV filling pressure by echocardiography (0.11 vs. 0.16 mm/mL, p<0.001). In 72 patients who had undergone RHC and simultaneous echocardiography, MAPSE/LAV and LASr correlated weakly with pulmonary artery wedge pressure (PAWP) (r=-0.44 and r=0.37). MAPSE/LAV was lower in patients with elevated (>15 mmHg) vs. normal PAWP (0.14 mm/mL vs. 0.27 mm/mL). Accuracy for detection of elevated PAWP was similar for MAPSE/LAV (area under the curve MAPSE/LAV: 0.75 [0.58–0.92] and LASr: 0.75 [0.57–0.90]).

Conclusions: Despite a moderate correlation with LASr, MAPSE/LAV provided similar diagnostic value as LASr regarding LV filling pressures as determined by echocardiography and RHC.

Keywords: left ventricular filling pressures; heart failure; echocardiography; atrioventricular displacement; left atrial strain; left atrial reservoir strain

Background

In recent years, peak left atrial longitudinal strain (LASr) has emerged as an accurate

diagnostic and prognostic tool in the assessment of patients with increased left ventricular

filling pressures (1-8). However, strain measurements require advanced software and the

technique is afflicted by vendor variability (9). Since longitudinal strain measurements are

determined by basic volumetric changes, the information obtained through strain techniques

are often available from other measurements, often included in standard examinations. The

left atrial (LA) reservoir function is determined by the LA size and its increase during

ventricular systole (10, 11), i.e. the mitral annular plane systolic excursion (MAPSE), Fig. 1.

MAPSE is an easily acquired, reliable measure, which is often feasible to acquire when image

quality is sub-optimal, and is an independent prognostic marker in heart failure (12). LA size

is included as a standard measurement in echocardiographic examinations (13).

We hypothesized that a ratio between mitral annular plane systolic excursion (MAPSE) and

LA volume (LAV) could offer similar diagnostic value as LASr. Therefore, we aimed to

describe the correlation between MAPSE/LAV and LASr, and to describe interobserver

variability for both measures. Further, we aimed to determine the diagnostic value of

MAPSE/LAV, in comparison to LASr, regarding detection of elevated left ventricular filling

pressure.

Methods

Two datasets were used for this study and ethical approvals were obtained from the Human

Research Ethics Committees for each cohort, respectively. The correlation between

MAPSE/LAV and LASr, as well as interobserver variability, were initially studied in an

existing datset including 100 patients clinical referral for echocardiography due to moderate

aortic stenosis (aortic valve peak velocity >3.0 to <4 m/s) (14). In the original study, these

patients were either retrospectively identified (n=70) from a set starting a date and consecutively included dating backward, or prospectively (n=30) included based on the presence of a moderate aortic stenosis with the aim to evaluate interobserver variability in aortic stenosis measurements. Patients with non-sinus rhythms were excluded. A comprehensive transthoracic echocardiographic exam was performed in all patients using a Vivid E9 system (GE Medical Systems, Horten, Norway). Off-line analyses were performed using commercially available image analysis software (EchoPAC, General Electric, Waukesha, Wisconsin, USA), and standard measurements were obtained (15). For the purpose of this study, MAPSE and LA strain measurements were obtained by two observers. LA strain was measured on a single-beat in both 4- and 2-chamber images using the semi-automatic Automated Functional Imaging Left Atrium (AFI) tool in EchoPAC. LA contours were extrapolated across pulmonary veins and the LA appendage orifice. Zero reference strain was set at end-diastole (16). MAPSE measurements were performed on 4-chamber images on which M-mode was applied to the septal and lateral points of the mitral ring.

LA volumes were determined through delineation of the LA in 4- and 2-chamber views at end-systole along the inner contours of the LA, excluding pulmonary veins and LA appendage, with a straight line along the mitral annulus, and calculated based on the modified Simpson's biplane method (15). MAPSE/LAV was then calculated for all patients.

Patients with poor image quality, defined as at least two out six non-visualized LA segments from either the 4- or 2-chamber views or obvious foreshortening of the LA in either view, were excluded (n=7).

For the purpose of mirroring clinical reality when determining interobserver variability, no explicit measurement instructions were provided to the two observers, apart from baseline knowledge of echocardiographic guidelines for standard acquisitions, and no mutual agreement between observers were made prior to the commencement of the study. Also, both

observers could freely select from the available images which images to include in their measurements. Of note, Observer 1 is an experienced reader of echocardiogram (TL; consultant, > 15 years of experience of clinical echocardiography) and Observer 2 a less experienced reader (CEG; biomedical scientist).

Echocardiographic assessment of left ventricular filling pressures was determined according to the American Society of Echocardiography(ASE)/European Association Cardiovascular Imaging (EACVI) guidelines based on the presence/absence of LA enlargement, increased tricuspid regurgitation velocity, reduced septal or lateral tissue doppler myocardial velocities (e'), and mitral peak early velocity (E)/e' ratio (13). In addition, quantitative estimation of left ventricular filling pressures was performed according to a recently described non-invasive method (ePAWP), which is based on LAV indexed to body surface area (LAVI), E and pulmonary vein systolic velocities (ePAWP = $0.179 \times$ LAVI + $2.672 \times$ mitral E/PVs + 2.7, in which ePAWP is given in mmHg, LAVI in mL/m^2, and mitral E and PVs in the same units of velocity (e.g. both in m/s)) (17).

Patients in sinus rhythm, with less than moderate mitral regurgitation/stenosis were included from a database with patients who had undergone clinically indicated right heart catherization (RHC) at Umeå University Hospital, Sweden, between 2010 and 2015 ($n = 154$) (18). RHC was performed using a Swan–Ganz thermodilution catheter inserted through the right internal jugular vein, a medial cubital vein, or the right femoral vein. Pulmonary artery wedge pressure (PAWP) was defined as the mean PAWP and recorded at end-expirium during spontaneous breathing. Increased PAWP was defined as mean PAWP > 15 mmHg (19). Main and/or contributing diagnoses for this cohort are presented in Supplements (Table S1).

Statistical analysis

Data are presented as mean±standard deviation (SD) or median (interquartile range) based on normal distribution. Normality was assessed using the Shapiro Wilk test. Correlations between LASr and MAPSE/LAV, LASr and PAWP, MAPSE/LAV and PAWP, respectively, were described using the Pearson correlation coefficient. Differences in means between multiple groups were performed using the analysis of variance (ANOVA) test. Differences in median between groups were assessed using the Wilcoxon Rank Sum Test or the Kruskal Wallis test when comparisons of more than two groups were performed. Interobserver variability was described a the mean±SD difference, and coefficients of variation (SD of the differences between observer measurements/mean of observer measurements $\times$ 100 with 95% confidence intervals).

Diagnostic performance in detection of elevated PAWP (>15 mmHg) was evaluated using receiver operating characteristics (ROC) analysis, and presented as sensitivity, specificity, accuracy, positive likelihood ratio (LR+) and negative likelihood ratio (LR-).

Statistical significance was accepted at the level of $P < 0.05$ (two-sided). Statistical analysis was performed using R version 4.2.1 (R Core Team, Vienna, Austria).

Results

After excluding 7 patients due to poor image quality, MAPSE/LAV and LASr were obtained in 93 patients with moderate aortic stenos. Baseline characteristics are described in Table 1 for the patients with moderate aortic stenosis and for patients who underwent simultaneous echocardiography in Table 2.

MAPSE/LAV was moderately correlated with LASr (r=0.57, p<0.001]), Fig. 2. MAPSE/LAV was lower for patients with increased left ventricular filling pressure, both according to the ASE/EACVI algorithm and ePAWP, Table 3 and Fig. 3.

Mean difference between observers was 0.02±0.04 mm/mL for MAPSE/LAV with a coefficient of variation of 24 [19–29]%. For MAPSE and LAV alone, coefficients of variation were 14 [11–17]%, and 18 [14 – 22]%, respectively. For LASr mean difference between observers was 4.4±3.8%, and the coefficient of variation was 18 [14–22]%.

After excluding patients with non-sinus rhythm (n = 34), at least moderate mitral valve lesions (n = 21), missing LA strain measurements (n = 16), 72 patients who had undergone RHC and simultaneous echocardiography were finally included. Among these, MAPSE/LAV and LASr were weakly correlated with PAWP (r=-0.44, p<0.001 and r=-0.38, p=0.001), Table 4. MAPSE/LAV was lower in patients with elevated (>15 mmHg) vs. normal PAWP, Table 3 and Fig. 4. Accuracy for detection of elevated PAWP was similar for MAPSE/LAV (area under the curve MAPSE/LAV: 0.75 [0.57–0.92] and LASr: 0.75 [0.57–0.90]), Fig. 5 and Table 5.

Discussion

A surrogate measure of LASr obtained through measurements of mitral annular displacement and LAV (MAPSE/LAV) provided similar diagnostic information as its strain-based counterpart. Both MAPSE/LAV and LASr were lower in patients with elevated left ventricular filling pressures as determined both by the ASE/EACVI algorithms, a quantitative echocardiographic PAWP estimation, and invasively measured PAWP. LASr has previously been shown to provide diagnostic information regarding left ventricular filling pressures (4, 8, 20, 21). For example, when LASr was incorporated into the ASE/EACVI algorithm, both feasibility and accuracy in detection of elevated filling pressures were improved, and LASr could effectively be used as replacement when either E/e' or tricuspid regurgitation velocity were unavailable (8). Although, LAV constitutes an important part of LASr, LASr improves

prediction of elevated PAWP beyond that obtained from LAV indexed-to-body surface area (22).

Interobserver variability was substantial for both MAPSE/LAV and LASr, although higher for MAPSE/LAV. Possibly, this difference is caused by the semi-automatic procedure used for LA strain, in contrast to MAPSE/LAV, which likely lowers variability. However, coefficients of variation were similar for LAV and LASr, indicating differences in LA delineation between the two observers. The addition of another measure (MAPSE) increased variability further. The high variability observed in this study contrasts previous reports of repeatability of LA strain measures, which have suggested low interobserver variability (23). This can be explained by the use of single-beat measurements used in this study, suggesting that averaging beats over multiple cycles are needed to obtain adequate reproducibility. Also, we actively refrained from synchronizing measurement strategies between observers, to mirror clinical reality. For both MAPSE/LAV and LASr there was a systematic bias between observers, possibly caused by the different contour delineation of the LA or differences in placement of the annular detachments for both measures.

LASr is determined, to a large extent, by atrioventricular plane displacement and LA dimensions. This has been observed previously (10, 11). Mălăescu showed excellent correlations between LV and LA strain curves across different cardiovascular pathologies. Similarly, Inoue, et al, showed LV global strain and LAV to be important determinants of LASr (11). LV global strain describes the systolic longitudinal shortening of the LV in relation to the length of LV cavity. Thus, the main difference between MAPSE and LV global strain in relation to their association with LA strain is that LV global strain also incorporates information of LV dimensions, which is likely to be redundant. Despite this, we did not find a stronger relation between MAPSE and LASr compared to between LV global strain and LASr. Possibly because of a closer relationship of measurement techniques between the two

strain modalities compared to the M-mode based measurement, or due to measurement variability.

Both MAPSE/LAV and LASr were lower in the cohort of patients with moderate aortic stenosis, compared to the cohort of patients who underwent clinical RHC. This could be explained by older patients in the aortic stenosis cohort (24). Also, reduced longitudinal function commonly occurs in patients with significant aortic stenosis (25).

Limitations

The cohort of patients with aortic stenosis did not include individuals with suspected heart failure, representing a limitation. However, the primary objective of analysing this cohort was to describe the correlation between MAPSE/LAV and LASr, and to determine reproducibility. Nonetheless, the potential clinical utility of MAPSE/LAV requires evaluation in settings that encompass patients with suspected or confirmed heart failure. Although such patients were part of the invasively examined study cohort, those patients constitute a mixed population with advanced disease unlikely to be representative of the typical population in which non-invasive determination of left ventricular filling pressures is sought. Nevertheless, our findings lend support to the hypothesis that comparable diagnostic information, as derived from strain-based LA measures, can be achieved without the use of advanced software.

An additional limitation to this study is the lack of core lab measurements. Interestingly, MAPSE/LAV were lower among patients in the non-invasive patient cohort than in the invasive cohort. This can either be caused by differences in measurements of acquisition or based on differences in patient material, and further highlights the need for future studies.

Conclusion

MAPSE/LAV was lower in patients with elevated LV filling pressures as

determined both by echocardiography and RHC. MAPSE/LAV provided similar diagnostic information as LASr in relation to LV filling pressures, and may serve as a substitute for LASr when such measures are unavailable, but was afflicted by high interobserver variability.

Table 1. Baseline characteristics among patients who underwent clinical echocardiography due to moderate aortic stenosis

	All	Normal	Indeterminate	Increased	p
		Left ventricular filling pressure according to the ASE/EACVI guidelines (13)			
N (%)	93 [100]	30 [32.0]	17 [18.3]	41 [44.1]	
Age, years	77 [72–83]	71.71[57.84–77.14]	74.06 [72.14–82.72]	82.63 [76.29–84.18]	0.01
Male sex, n (%)	54 [62.8]	24 (77.4)	9 (56.2)	21 (53.8)	0.11
BMI, kg/m^2	27 [26–30]	27 [26–30]	28 [26–32]	28 [25–30]	0.69
BSA, m2	1.9±0.2	2.0±0.2	2.0±0.3	1.9±0.2	0.03
Echocardiographic measures					
LVEF, %	60 [60–63]	60 [60–63]	60 [60–60]	60 [60–65]	0.91
MAPSE, mm	11±3	12±3	12±2	11±3	0.14
LA volume, mL	83±21	72±18	80±22	92±19	<0.001
LAVI, mL/m^2	43±11	36±8	41±10	49±10	<0.001
LV diameter, mm	49±6	51±5	50±5	48±5	0.02
LV mass, g/m^2	87±21	90±16	85±19	86±26	0.67
E/A, unitless	0.9 [0.7–1.1]	0.9 [0.7–1.0]	0.7 [0.6–0.9]	1.0 [0.8–1.4]	0.01
S/D, unitless	1.3 [1.0–1.6]	1.3 [1.1–1.5]	1.5 [1.2–2.0]	1.2 [0.7–1.6]	0.16
E/e', unitless	13 [10–17]	10 [8–13]	13 [9–14]	17 [15–21]	<0.001
TR velocity, m/s	2.6 [2.5–2.8]	2.5 [2.4–2.6]	2.7 [2.4–2.8]	2.8 [2.5–3.1]	0.001
AV$_{peak}$, m/s	3.4 [3.2–3.7]	3.3 [3.2–3.5]	3.3 [3.3–3.6]	3.6 [3.3–3.8]	0.02
AV$_{mean,}$ mmHg	28 [24–33]	26 [23–30]	28 [25–33]	32 [26–35]	0.01

Data are presented as mean±standard deviation or median [interquartile range] based on normal distribution–or as n [%].
Abbreviations: AV$_{peak}$: aortic valve peak velocity; AV$_{mean}$: aortic valve mean gradient; BMI: body mass index; BSA: body surface area; LA: left atrial; LAVI: left atrial volume indexed to BSA; LVEF: left ventricular ejection fraction; LVP: left ventricular filling pressures; MAPSE: mitral annular plane systolic excursion; TR: tricuspid regurgitation

Table 2. Baseline characteristics among patients who underwent echocardiography simultaneously with clinically indicated right heart catherization (n=70)

	All	PAWP ≤15 mmHg	PAWP >15 mmHg	p
N [%]	73 [100]	58 [79.5]	15 [20.5]	
Age, years	64 [55–74]	64 [55–74]	64 [58–73]	1.0
Male sex, n [%]	23 [31.5]	16 (27.6)	7 (46.7)	0.27
Systolic blood pressure, mmHg	132 [120–145]	132 [122–144]	135 [120–155]	0.71
Diastolic blood pressure, mmHg	77 [70–82]	77 [70–83]	76 [73–80]	0.99
NT-ProBNP, ng/L	518 [217–1787]	375 [178–1260]	1623 [864–2013]	0.02
BMI, kg/m^2	27±6	26±5	30±6	0.23
BSA, m^2	1.8±0.2	1.8±0.2	1.9±0.3	0.12
Hypertension, n [%]	21 [28.8]	16 [27.6]	5 [33.3]	0.91
Diabetes mellitus, n [%]	6 [8.2]	2 [3.4]	4 [26.7]	0.02
Ischemic heart disease, n [%]	11 [15.1]	10 [17.2]	1 [6.7]	0.54
Betablockers, n [%]	22 [30.1]	15 [25.9]	7 [46.7]	0.21
ACE/ARB, n [%]	30 [41.1]	21 [36.2]	9 [60.0]	0.17
Calcium antagonist, n [%]	3 [4.1]	3 [5.2]	0 [0.0]	0.87
Diuretics, n [%]	33 [45.2]	22 [37.9]	11 [73.3]	0.03
Right heart catherization measures				
PVR, Wood units	3.0 [1.9–5.0]	3.0 [1.8–5.8]	2.5 [2.0–3.2]	0.48
mPAP, mmHg	33±16	32±17	38±9	0.20
PAWP, mmHg	12±6	10±4	22±4	<0.001
Echocardiographic measures				
LVEF, %	56 [44–65]	61 [47–66]	47 [33–55]	0.02
GLS, %	16±6	17±6	12±5	0.004
MAPSE, mm	12 [10–13]	12 [11–13]	10 [8–12]	0.009
LA volume, mL	46 [32–70]	44 [31–58]	70 [46–97]	0.02
LAVI, mL/m^2	26 [19–33]	23 [17–31]	32 [27–51]	0.05
LV diameter, mm	47 [42–51]	47 [42–50]	47 [44–59]	0.38
LV mass, g/m^2	80 [66–88]	80 [66–88]	77 [71–86]	0.91

E/A, unitless	1.1 [0.8–1.4]	1.0 [0.8–1.3]	1.6 [1.4–2.6]	0.004
S/D, unitless	1.1 [0.8–1.5]	1.1 [0.9–1.6]	0.8 [0.6–0.9]	0.01
E/e', unitless	9 [7–12]	9 [7–12]	11 [9–13]	0.18
SPAP, mmHg	49 [35–67]	53 [33–68]	47 [39–59]	0.49
ePAWP, mmHg	11 [9–14]	11 [9–12]	15 [14–18]	0.004

Data are presented as mean±standard deviation or median [interquartile range] based on normal distribution–or as n [%] for categorical variables.

Abbreviations: BMI: body mass index; BSA: body surface area; GLS: global longitudinal strain; LA: left atrial; LASr: left atrial reservoir strain; LAVI: left atrial volume indexed to BSA; LVEF: left ventricular ejection fraction; MAPSE: mitral annular plane systolic excursion; MAPSE/LAV: mitral annular plane systolic excursion; PAWP: pulmonary artery wedge pressure; SPAP: systolic pulmonary arterial pressure (by echocardiography)

Table 3. Mitral annular plane systolic excursion to left atrial volume ratio (MAPSE/LAV) and left atrial reservoir strain (LASr) stratified by left ventricular filling pressure according to echocardiography or right heart catherization

Non-invasively assessed LV filling pressure among patients with a clinically indicated echocardiography due to aortic stenosis (n=93)

	LV filling pressure according to the ASE/EACVI algorithm (13)			
	Normal	Indeterminate	Increased	p
MAPSE/LAV, mm/mL	0.16 [0.13–0.20]	0.13 [0.13–0.17]	0.11 [0.09–0.15]	<0.001
LASr, %	20 [17–23]	22 [16–24]	16 [14–18]	0.007

	LV filling pressure according to ePAWP (17)		
	Normal	Increased	
MAPSE/LAV, mm/mL	0.15 [0.13–0.18]	0.10 [0.08–0.14]	<0.001
LASr, %	20 [16–23]	17 [14–21]	0.08

	LV filling pressure according to invasively measured PAWP (19)		
	Normal	Increased	
MAPSE/LAV, mm/mL	0.27 [0.19–0.37]	0.14 [0.09–0.23]	0.003
LASr, %	19 [12–24]	8 [6–18]	0.006

Data are presented as median [interquartile range]
Abbreviations: LASr: left atrial reservoir strain; LV: left ventricular; MAPSE/LAV: mitral annular plane systolic excursion to left atrial volume ratio

Table 4. Correlation between MAPSE/LAV and LASr with pulmonary artery wedge pressure, left ventricular function, E/e ratio and NT-proBNP

	MAPSE/LAV		LASr	
	r	p	r	p
PAWP	-0.44	<0.001	-0.37	0.002
LVEF	0.24	0.045	0.27	0.02
GLS	0.19	0.11	0.49	<0.001
LA volume	-0.78	<0.001	-0.21	0.08
MAPSE	0.38	0.001	0.50	<0.001
E/e'	-0.23	0.08	-0.29	0.03
NT-proBNP (log)	-0.33	0.006	-0.29	0.02

Abbreviations: GLS: global longitudinal strain; LA: left atrial; LVEF: left ventricular ejection fraction; MAPSE: mitral annular plane systolic excursion; NT-proBNP: N-terminal pro–B-type natriuretic peptide; PAWP: pulmonary artery wedge pressure

MAPSE/LAV	Sensitivity (%)	Specificity (%)	Accuracy (%)	LR+	LR-
0.42	93	17	33	1.1	0.39
0.25	80	59	64	1.9	0.35
0.21	73	72	74	2.8	0.36
0.16	67	83	81	4.2	0.39
0.13	47	90	82	5.3	0.58
0.10	33	98	85	16.5	0.68
LASr	**Sensitivity (%)**	**Specificity (%)**	**Accuracy (%)**	**LR+**	**LR-**
24	92	23	36	1.2	0.35
20	85	50	56	1.7	0.30
18	77	59	62	1.9	0.39
13	69	73	72	2.6	0.42
9	54	86	80	3.9	0.53
6	38	95	84	7.6	0.65

Table 5. Sensitivity, specificity, accuracy, and likelihood ratios for different cutoffs for MAPSE/LAV and LASr in detection of invasively measured elevated pulmonary arterial wedge pressure (n=70).

Figure legends

Figure 1. Mitral annular plane systolic excursion (MAPSE, upper panel, orange) and left atrial reservoir strain (LASr, lower panel, orange) measurements from echocardiography. Notice the close relation between MAPSE curves and LASr. The amplitude of the LASr depends on the left atrial size.

Figure 2. Scatterplot describing the relation between left atrial (LA) reservoir function (LASr) and mitral annular plane systolic excursion (MAPSE)/LA volume (LAV).

Figure 3. Box plots of mitral annular plane systolic excursion (MAPSE)/left atrial (LA) volume (left panels) and LA reservoir strain (LASr) (right panels) among patients with normal (green), indeterminate (orange) or increased (red) left ventricular pressure according to the ASE/EACVI guidelines for assessment of diastolic function (upper panels) and estimated pulmonary artery wedge pressure using echocardiography (17) (ePAWP; lower panels).

Figure 4. Box plots of mitral annular plane systolic excursion (MAPSE)/left atrial (LA) volume (upper left panel), LA reservoir strain (LASr) (upper right panel), MAPSE (lower left panel) and LA volume (lower right panel) among patients with normal (green) or high (red) pulmonary artery wedge pressure (PAWP) in patients who had undergone simultaneous echocardiography and right heart catheterization.

Figure 5. Receiver operating curve for the detection of elevated pulmonary artery wedge pressure (> 15 mmHg) according to right heart catherization.

References

1.	Lacalzada-Almeida J, Izquierdo-Gómez MM, García-Niebla J, Elosua R, Jiménez-Sosa A, Baranchuk A, et al. Advanced interatrial block is a surrogate for left atrial strain reduction which predicts atrial fibrillation and stroke. *Ann Noninvasive Electrocardiol*. 2019 Jul;**24**(4):e12632.

2.	Cau R, Bassareo P, Suri JS, Pontone G, Saba L. The emerging role of atrial strain assessed by cardiac MRI in different cardiovascular settings: an up-to-date review. *Eur Radiol*. 2022 Jul;**32**(7):4384-94. E

3.	Mandoli GE, Pastore MC, Benfari G, Bisleri G, Maccherini M, Lisi G, et al. Left atrial strain as a pre-operative prognostic marker for patients with severe mitral regurgitation. *Int J Cardiol*. 2021 Feb 1;**324**:139-45.

4.	Cameli M, Mandoli GE, Loiacono F, Dini FL, Henein M, Mondillo S. Left atrial strain: a new parameter for assessment of left ventricular filling pressure. *Heart Fail Rev*. 2016 Jan;**21**(1):65-76.

5.	Morris DA, Belyavskiy E, Aravind-Kumar R, Kropf M, Frydas A, Braunauer K, et al. Potential Usefulness and Clinical Relevance of Adding Left Atrial Strain to Left Atrial Volume Index in the Detection of Left Ventricular Diastolic Dysfunction. *JACC Cardiovasc imaging*. 2018 Oct;**11**(10):1405-15.

6.	Park JH, Hwang IC, Park JJ, Park JB, Cho GY. Prognostic power of left atrial strain in patients with acute heart failure. *Eur Heart J Cardiovasc Imaging*. 2021 Jan 22;**22**(2):210-9.

7.	Rausch K, Shiino K, Putrino A, Lam AK, Scalia GM, Chan J. Reproducibility of global left atrial strain and strain rate between novice and expert using multi-vendor analysis software. *Int J Cardiovasc Imaging*. 2019 Mar;**35**(3):419-26.

8.   Venkateshvaran A, Tureli HO, Faxén UL, Lund LH, Tossavainen E, Lindqvist P. Left atrial reservoir strain improves diagnostic accuracy of the 2016 ASE/EACVI diastolic algorithm in patients with preserved left ventricular ejection fraction: insights from the KARUM haemodynamic database. *Eur Heart J Cardiovasc Imaging*. 2022 Aug 22;**23**(9):1157-68.

9.   Mouselimis D, Tsarouchas AS, Pagourelias ED, Bakogiannis C, Theofilogiannakos EK, Loutradis C, et al. Left atrial strain, intervendor variability, and atrial fibrillation recurrence after catheter ablation: A systematic review and meta-analysis. *Hellenic J Cardiol*. 2020 May-Jun;**61**(3):154-64.

10.   Mălăescu GG, Mirea O, Capotă R, Petrescu AM, Duchenne J, Voigt JU. Left Atrial Strain Determinants During the Cardiac Phases. *JACC Cardiovasc imaging*. 2022 Mar;**15**(3):381-91.

11.   Inoue K, Khan FH, Remme EW, Ohte N, García-Izquierdo E, Chetrit M, et al. Determinants of left atrial reservoir and pump strain and use of atrial strain for evaluation of left ventricular filling pressure. *Eur Heart J Cardiovasc Imaging*. 2021 Dec 18;**23**(1):61-70.

12.   Xue H, Artico J, Davies RH, Adam R, Shetye A, Augusto JB, et al. Automated In-Line Artificial Intelligence Measured Global Longitudinal Shortening and Mitral Annular Plane Systolic Excursion: Reproducibility and Prognostic Significance. *J Am Heart Ass*. 2022 Feb 15;**11**(4):e023849.

13.   Nagueh SF, Smiseth OA, Appleton CP, Byrd BF, 3rd, Dokainish H, Edvardsen T, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Eur Heart J Cardiovasc Imaging*. 2016 Dec;**17**(12):1321-60.

14.   Manna D, Eliasson M, Bech-Hanssen O, Lindow T. Reproducibility of Echocardiographic Measures of Aortic Stenosis Severity and Its Impact on Grading of Severity. *J Am Soc Echocardiogr*. 2023 Nov 14. Epub 2023/11/17.

15.   Mitchell C, Rahko PS, Blauwet LA, Canaday B, Finstuen JA, Foster MC, et al. Guidelines for Performing a Comprehensive Transthoracic Echocardiographic Examination in Adults: Recommendations from the American Society of Echocardiography. *J Am Soc Echocardiogr*. 2019 Jan;**32**(1):1-64.

16.   Badano LP, Kolias TJ, Muraru D, Abraham TP, Aurigemma G, Edvardsen T, et al. Standardization of left atrial, right ventricular, and right atrial deformation imaging using two-dimensional speckle tracking echocardiography: a consensus document of the EACVI/ASE/Industry Task Force to standardize deformation imaging. *Eur Heart J Cardiovasc Imaging*. 2018 Jun 1;**19**(6):591-600.

17.   Lindow T, Manouras A, Lindqvist P, Manna D, Wieslander B, Kozor R, et al. Echocardiographic estimation of pulmonary artery wedge pressure - invasive derivation, validation, and prognostic association beyond diastolic dysfunction grading. *Eur Heart J Cardiovasc Imaging*. 2023 Nov 9. Epub 2023/11/11.

18.   Tossavainen E, Henein MY, Grönlund C, Lindqvist P. Left Atrial Intrinsic Strain Rate Correcting for Pulmonary Wedge Pressure Is Accurate in Estimating Pulmonary Vascular Resistance in Breathless Patients. *Echocardiogr*. 2016 Aug;**33**(8):1156-65.

19.   Humbert M, Kovacs G, Hoeper MM, Badagliacca R, Berger RMF, Brida M, et al. 2022 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Resp J*. 2023 Jan;**61**(1).

20.   Kurt M, Tanboga IH, Aksakal E, Kaya A, Isik T, Ekinci M, et al. Relation of left ventricular end-diastolic pressure and N-terminal pro-brain natriuretic peptide level with left atrial deformation parameters. *Eur Heart J Cardiovasc Imaging*. 2012 Jun;**13**(6):524-30.

21.   Cameli M, Sparla S, Losito M, Righini FM, Menci D, Lisi M, et al. Correlation of Left Atrial Strain and Doppler Measurements with Invasive Measurement of Left Ventricular End-Diastolic Pressure in Patients Stratified for Different Values of Ejection Fraction. *Echocardiogr*. 2016 Mar;**33**(3):398-405.

22.   Bytyçi I, Bajraktari G, Lindqvist P, Henein MY. Compromised left atrial function and increased size predict raised cavity pressure: a systematic review and meta-analysis. *Clin Physiol Funct Imaging*. 2019 Sep;**39**(5):297-307.

23.   Cameli M, Caputo M, Mondillo S, Ballo P, Palmerini E, Lisi M, et al. Feasibility and reference values of left atrial longitudinal strain imaging by two-dimensional speckle tracking. *Cardiovasc Ultrasound*. 2009 Feb 8;**7**:6.

24.   Støylen A, Mølmen HE, Dalen H. Relation between Mitral Annular Plane Systolic Excursion and Global longitudinal strain in normal subjects: The HUNT study. *Echocardiogr*. 2018 May;**35**(5):603-10.

25.   Vollema EM, Amanullah MR, Prihadi EA, Ng ACT, van der Bijl P, Sin YK, et al. Incremental value of left ventricular global longitudinal strain in a newly proposed staging classification based on cardiac damage in patients with severe aortic stenosis. *Eur Heart J Cardiovasc Imaging*. 2020 Oct 20;**21**(11):1248-58.