Differences in HIV-1 reservoir size, landscape characteristics and decay dynamics in acute and chronic treated HIV-1 Clade C infection.

Kavidha Reddy[1], Guinevere Q. Lee[2], Nicole Reddy[1,3], Tatenda J.B. Chikowore[1,4], Kathy Baisley[1,5], Krista L. Dong[6,7,8], Bruce D. Walker[6,7,8], Xu G. Yu[6,8], Mathias Lichterfeld[6,8,9], Thumbi Ndung'u[1,3,4,6,7,*]

[1]Africa Health Research Institute, Durban, South Africa, [2]Weill Cornell Medical College, New York, USA, [3]University of KwaZulu-Natal, Durban, South Africa, [4]University College of London, London, UK, [5]London School of Hygiene and Tropical Medicine, London, UK, [6]Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, USA, [7]HIV Pathogenesis Programme (HPP), The Doris Duke Medical Research Institute, University of KwaZulu-Natal, Durban, South Africa, [8]Harvard Medical School, Boston, Massachusetts, USA, [9]Brigham and Women's Hospital, Boston, MA, USA.

*Correspondence: Thumbi Ndung'u (thumbi.ndungu@ahri.org)
Africa Health Research Institute, K-RITH Tower Building, Level 3, Nelson R. Mandela School of Medicine, 719 Umbilo Road, Congella, Durban, 4001, South Africa.

Abstract

Background
Persisting HIV reservoir viruses in resting CD4 T cells and other cellular subsets are the main barrier to cure efforts. Antiretroviral therapy (ART) intensification by early initiation has been shown to enable post-treatment viral control in some cases but the underlying mechanisms are not fully understood. We hypothesized that ART initiated during the hyperacute phase of infection before peak will affect the size, decay dynamics and landscape characteristics of HIV-1 subtype C viral reservoirs.

Methods
We studied 35 women at high risk of infection from Durban, South Africa identified with hyperacute HIV infection by twice weekly testing for plasma HIV-1 RNA. Study participants included 11 who started ART at a median of 456 (297-1203) days post onset of viremia (DPOV), and 24 who started ART at a median of 1 (1-3) DPOV. We used peripheral blood mononuclear cells (PBMC) to measure total HIV-1 DNA by ddPCR and to sequence reservoir viral genomes by full length individual proviral sequencing (FLIP-seq) from onset of detection of HIV up to 1 year post treatment initiation.

Results
Whereas ART in hyperacute infection blunted peak viremia compared to untreated individuals (p<0.0001), there was no difference in total HIV-1 DNA measured contemporaneously (p=0.104). There was a steady decline of total HIV DNA in early treated persons over 1 year of ART (p=0.0004), with no significant change observed in the late treated group. Total HIV-1 DNA after one year of treatment was lower in the early treated compared to the late treated group (p=0.02). Generation of 697 single viral genome sequences revealed a difference in the longitudinal proviral genetic landscape over one year between untreated, late treated and early treated infection: the relative contribution of intact genomes to the total pool of HIV-1 DNA after 1 year was higher in untreated infection (31%) compared to late treated (14%) and early treated infection (0%). Treatment initiated in both late and early infection resulted in a more rapid decay of intact (13% and 51% per month) versus defective (2% and 35% per month) viral genomes. However, intact genomes were still observed one year post chronic treatment initiation in contrast to early treatment where intact genomes were no longer detectable. Moreover,

48   early ART reduced phylogenetic diversity of intact genomes and limited the seeding and persistence of
49   cytotoxic T lymphocyte immune escape variants in the reservoir.
50

51 **Conclusions**
52 Overall, our results show that whereas ART initiated in hyperacute HIV-1 subtype C infection did not
53 impact reservoir seeding, it was nevertheless associated with more rapid decay of intact viral genomes,
54 decreased genetic complexity and immune escape in reservoirs, which could accelerate reservoir
55 clearance when combined with other interventional strategies.
56

57 **Keywords**
58 HIV-1, Clade C, reservoir, viral dynamics
59

Introduction
Although antiretroviral therapy (ART) effectively controls HIV, it does not eradicate persisting virus
from resting CD4 T cells and other cellular subsets. ART is therefore lifelong, with many side effects
and may be financially unsustainable particularly in low- to middle- income countries that are most
affected by the HIV epidemic. A cure is therefore necessary but difficult to achieve as the persistent
HIV reservoir, which is established soon after infection [1-3] hinders these efforts through its complex
nature [4].
67

The HIV-1 reservoir is heterogenous and dynamic in genotypic, phenotypic and cellular composition
and also differs between individuals and populations, making the source of persistent infection difficult
to target. The lability of the viral reservoir has been demonstrated by evidence of clonal expansion that
occurs by homeostatic proliferation [5, 6], antigenic stimulation [7] or integration into genic regions
controlling cell growth [8-10]. Additionally, its size and genetic composition changes over time with
intact and defective proviral genomes displaying variability in their rate of decay during suppressive
ART [11-13]. The heterogeneity of the HIV-1 reservoir landscape is further obscured by the integration
patterns of virus-intact genomes that determine their ability to generate replication competent viral
particles when stimulated [8]. Moreover, recent studies show that the viral reservoir is not restricted to
CD4 T cells but persists in other cell types such as monocytes, macrophages [14-19] and follicular
dendritic cells [20-23] with each having distinct biological characteristics.
79

Among the many studies directed towards HIV-1 cure, some have demonstrated that early ART
initiation may be a crucial step in limiting the viral reservoir and achieving post-treatment viral control
[24, 25]. Even though early treatment does not prevent reservoir establishment, it does result in a rapid
decline of viremia and substantially reduces HIV reservoir size [11, 26-28] which delays viral load
rebound and results in sustained virological remission following discontinuation of ART [25, 29, 30].
Other described benefits of starting therapy soon after primary infection include preservation of immune
function and restriction of viral diversification [26, 28, 31-41]). However, there is a paucity of data that
describes the impact of the timing of treatment initiation on the characteristics of the reservoir.
88

HIV-1 reservoir studies in African populations are limited, particularly in subtype C HIV-1 infection,
which is the most prevalent form of HIV-1 globally and predominates in southern Africa. The design
of a globally applicable HIV cure strategies and interventions to target the viral reservoir depend on a
deeper understanding of the variability in the size, composition and characteristics of the genetic
landscapes of persisting reservoir genomes in non-subtype B HIV infections. Moreover, data are lacking
on reservoir characteristics in women, despite known sex differences in immune responses and in viral
load during primary infection that could potentially impact the reservoir [42-44].

In this study we performed an extensive longitudinal analysis of HIV-1 subtype C reservoir characteristics in a unique hyperacute infection cohort in Durban, South Africa [45, 46]. The cohort was designed to identify acute infection before peak viremia (Fiebig stages I to III) providing HIV-1 testing twice a week to young women at high risk for HIV-1 infection in a region with high population prevalence [45]. Following changes in treatment guidelines in South Africa that allowed for ART initiation regardless of CD4 counts, all study participants were offered ART, including those who were newly detected with acute infection who received ART on average a day after first detection of plasma viremia. Study participants underwent frequent clinical follow-up and sampling following infection and initiation of ART, allowing us to study HIV-reservoir establishment and evolution from the earliest possible stages of infection. We hypothesized that the timing of ART will impact HIV-1 reservoir characteristics in terms of the size, genetic composition, and decay dynamics. The findings of this study provide insights into HIV-1 reservoirs characteristics that could inform viral targeting strategies for reservoir control in African populations.

Results

Total proviral DNA load kinetics following early and late treatment

In this analysis we included 35 participants (Supplementary Table 1), of whom 11 first initiated treatment during chronic infection at a median of 456 days (297-1203) post detection of viremia and 24 who were treated during acute infection at a median of 1 day (1-3) post detection of viremia. All participants were female and 31 (89%) were identified with acute infection at Fiebig stage I. Additional participant characteristics are shown in Table 1.

We first quantified total HIV-1 DNA which incorporates all forms of intracellular HIV-1 DNA, both intact and defective, including integrated and unintegrated forms, as well as linear and circularized 2-LTR and 1-LTR forms. Total HIV-1 DNA measurements were performed longitudinally, from baseline (1-3 days following detection of HIV), at the time of peak viral load and was also assessed at 6- and 12-months post-infection for untreated participants and 6- and 12-months post-treatment initiation for late and early treated participants. As expected, treatment during acute infection resulted in a significantly reduced peak plasma viral load (median = 4.18 log copies/ml, IQR, 3.40-4.87) compared to untreated acute infection (median = 7.06 log copies/ml, IQR, 6.83-7.54) (p<0.0001, Figure 1A top panel). However, at time of peak viremia, the untreated and the early treated groups did not differ in total proviral DNA load (Figure 1A, bottom panel). Longitudinal measurements showed that treatment initiated during chronic infection resulted in a significant decline in plasma viral load to undetectable levels after 1 year (p<0.0001, Figure 1B, top panel) however it did not reduce total proviral load (Figure 1B, bottom panel). In contrast, treatment initiated during acute infection resulted in both a rapid decrease of plasma viremia so that all participants had undetectable viremia at one-year post-ART (p<0.0001, Figure 1C top panel) and steady decrease of total proviral load over the same time period (p=0.0004) (Figure 1C, bottom panel). Even though treatment initiation during both chronic and acute infection resulted in complete suppression of plasma viral load after 1 year (Figure 1D, top panel), total proviral load was still detectable with the early treated group having 1.3 times lower levels of total proviral HIV DNA compared to the chronic treated group (p=0.02, Figure 1D, bottom panel). These results indicate that early treatment leads to a measurable decline in proviral DNA during the first year of treatment that is not seen when therapy is initiated during chronic infection.

Factors associated with total proviral DNA load after 1 year of suppressive ART

To further understand the impact of host, virological and immunological factors, as well as timing of treatment on the establishment and maintenance of the HIV reservoir, we analysed the associations

between viral and immune factors and HIV-1 proviral DNA after 1 year of treatment (Table 2). Analyses
for each treatment group were performed independently using multivariate regression models with HIV-
1 DNA levels as the dependent variable and other factors, specifically nadir CD4, pre-infection CD4,
baseline CD4 counts and peak viral load, as the independent predictor variables. The analysis showed
that when treatment was initiated during acute infection, only peak plasma viral load was significantly
associated with levels of HIV-1 proviral DNA after 1 year of treatment (p=0.02). However, when
treatment was initiated during chronic infection both baseline CD4 count (measured 1-3 days after
detection of HIV) (p=0.002) and peak plasma viral load (p=0.03) positively associated with HIV-1
proviral DNA levels, while there was a significant inverse association with nadir CD4 (p<0.0001).
Other factors such as protective HLA alleles and type of treatment regimen were not associated with
HIV-1 proviral DNA (data not shown). These data indicate that both host and viral characteristics
impact the establishment and maintenance of the viral reservoir.
156
Longitudinal genotypic characterisation of HIV-1 DNA
Quantification of total HIV-1 DNA by ddPCR as described above is based on the amplification of a
short 127 base pair fragment of the HIV-1 genome, and thus detects defective viruses that are incapable
of replication, thereby overestimating the size and functionality of the reservoir. To address this, we
next performed single template near full-genome sequencing to determine potential replication
competency by establishing the distribution of genome intact and genome defective latent viruses within
cells. Viral genome intactness was determined by the HIVSeqinR v2.7.1 computational bioinformatics
pipeline [33].
165
For this analysis we studied 24 participants: The chronic infection (late treatment) group (n=11)
consisted of individuals who remained untreated for over one year following infection and before
treatment initiation. Longitudinal sampling at untreated time points was available for 9 of these
individuals whereas in two individuals, samples were only available post-treatment initiation. The
acutely treated group (n=13) received treatment immediately post-detection (Figure 2A). We generated
a total of 697 sequences (GenBank accession numbers OR991333-OR991737 and MK643536-
MK643827) after sampling a median of 1.4 million PBMC (0.02-4.3 million) per sampling time point.
Genome-intact viruses (Figure 2B) accounted for 35% (247/697) of the total pool and were detected in
23 participants (12 from the early treatment group and 11 from the late treatment group), with a median
of 8 genome-intact viruses (range=1-60) per study participant. Phylogenetic analysis revealed a
significant difference in the mean pairwise distances of intact viral sequences derived from early treated
(median=0.12% (IQR, 0.07-0.21) compared to late treated participants (median=0.48% (IQR, 0.16-
1.08)) (p=0.04) (Figure 2C). Overall, 56% of the intact genomes collected in this study were obtained
from the untreated study arm, while 11% were obtained from late treated chronic infection and 33%
from acutely treated infections (Figure 2D).
181
Longitudinal studies revealed that intact proviruses accumulated rapidly during the course of HIV-1
infection with a relative contribution of 65% (450/697) to the total pool of proviral genomes detected.
The majority of defective viral genomes collected in this study were detected during untreated infection
(44% (199/450), left-most pie), while 39% (175/450) were detected during late treated chronic infection
and 17% (76/450) during acute treated infection. Defective genomes contributed 59% to the proviral
population in untreated infection, 87% in late (chronic) treated and 48% in acutely treated infection
(Figure 2D). Overall, defective genomes also accumulated quickly after onset of infection and were
detectable at a proportion of 47% within the first month irrespective of treatment status (data not
shown). Large internal deletions within viral genomes were the most common defect with relative
frequencies of 77%, 60% and 79% in untreated infection, chronic treated infection and acute treated

infection respectively among the pool of defective genomes. Overall, these gene deletions occurred
significantly more frequently between *integrase* and *envelope* in the integrase to envelope gene segment
compared to *gag* (p<0.0001–0.001), with *nef* being similar to *gag* (Supplementary Figure 1). APOBEC
induced hypermutations were the second most common defect observed in untreated (16%) and late
(chronic) treated (31%) infection. However, in acute treated infection, hypermutations were relatively
infrequent, comprising only 7% of the genome-defective pool. Premature stop codons in one of *gag,*
pol or *env* occurred at a frequency of 5%, 4% and 13% as a percentage of defective genomes in
sequences from untreated, chronic, and acute treated infections respectively. Internal inversions (1%,
1%, 1%), and 5' psi defects (2%, 4%, 0%) were other types of genome defects that were detected at
minor frequencies in untreated, chronically treated and acutely treated infections respectively.
202
To further understand the impact of ART timing on the composition, evolution, and dynamics of the
HIV-1 proviral landscape over time, we next performed a stratified analysis of the relative proportions
of viral genome sequences in each study arm over 1 year of follow up (Figure 3A-C). Genome-intact
viruses were detectable throughout the course of untreated infection while genome-defective viruses
also accumulated over this period (Figure 3A). Initiation of ART during chronic infection, at a median
of 456 days after detection of plasma viremia, resulted in a decrease in the relative proportion of
genome-intact viruses over 1 year of treatment (34% to 14%). However, genome-intact viruses were
not completely eradicated and were easily detectable after 12 months of treatment (Figure 3B).
Additionally, genomes with large deletions and hypermutations became more prominent in the chronic
treated group over 1 year of treatment. In contrast, there was a more rapid decrease in the proportion
of genome-intact viruses following ART initiation in acute infection such that these viruses were no
longer detectable at our sampling depth after 1 year of treatment (57% to 0%) (Figure 3C).
Hypermutated viruses were also less prominent before 12 months during early treated infection (Figure
3C). These data suggest that early treatment initiation facilitates faster clearance of genome-intact
viruses in the blood compared to late treatment.
218
Decay kinetics of intact and defective proviruses
Studies show that the biology and decay dynamics of genome-intact viruses within the viral reservoir
likely differ from that of the genome-defective provirus pool [12, 13]. However, the effect of ART
timing on the rate of decay of these different pools of viruses is not well known and has not been
investigated in African populations where immune responses and viral genetic heterogeneity may well
result in population-specific differences. Here we observed that the absolute proportions of both
genome-intact and -defective viruses per million PBMCs sampled decreased in both early treated and
late treated participants over the 1-year follow-up period of this study (Figure 4A and 4B).
227
To estimate potential differences in the rate of change between genome-intact and -defective viruses
within each treatment group we used a linear mixed effects regression model with random intercepts to
account for the correlation between repeated observations from the same individual. We fit a model
with log DNA copies as the response variable, and time, treatment group, and a time-group interaction
as fixed effects, with participant as a random effect.   The analysis was restricted to the first 6 months
after starting ART as regular measurements for both groups were available over this period.
234
235 Among the acute treated, genome-intact proviruses decreased by 0.308 log copies per month in the first
236 six months after starting ART, corresponding to a decline of 51% per month (p<0.001, Figure 4C). In
237 contrast, among the chronic treated, intact proviruses decreased by only 0.059 log copies per month,
238 corresponding to a decline of 13% per month; however, this decrease was not statistically significant
239 (p=0.68) (Figure 4E).

Genome-defective proviruses also decreased significantly in the acute treated group by 0.190 log copies per month in the first six months after starting ART, corresponding to a decline of 35% (p=0.01). However, in the chronic treated group the change in the number of log copies of defective provirus in the first 6 months was only 0.015 (p=0.88) corresponding to a decline of just 2% per month (Figure 4D and 4E). These results indicate that early treatment is associated with a faster decline of both genome-intact and -defective proviruses compared to late treatment.

Contribution of clonal expansion to maintenance of proviral populations

Studies show that more than 50% of the latent HIV reservoir is maintained by clonal expansion (Lui et. al., 2020). We assessed viral genome sequences to determine the extent of persistence of infected cell clones after primary infection. Viral genome sequences sharing 100% identity by FLIP-seq was used as a marker of clonal expansion of infected cells as previous studies have shown that proviral genomes that were 100% identical share the same viral integration site whereas proviruses with different integration sites do not share 100% sequence identity (Einkauf et al., 2019). At our sampling depth, we detected clonal expansion in 3/11 (27%) participants who were treated during chronic infection and 4/13 (30%) participants treated during acute infection, showing that in subtype C infection clonal expansion of infected cells occurred as early as one day post detectable viremia (Supplementary Figure 2). Defective clones were detected in two late treated participants at proportions of 6% and 13% of total proviral population, while intact clones were identified in one late treated participant at a proportion of 6% of the total proviral pool. In contrast, a higher proportion of intact clones were detected in early treated participants at 33%, 30%, 37% and 24% of the total proviral pool. Although the data is limited and needs to be interpreted with caution, this suggests that clonal expansion of intact proviral genomes is more likely to occur when treatment is initiated early, likely due to the early inhibition of viral replication that prevents the accumulation and seeding of defective viral genomes into the viral reservoir.

CTL epitope diversity in the latent reservoir

The emergence of escape mutations in viral epitopes as a mechanism to evade human leukocyte antigen (HLA) class I-restricted immune responses, specifically of CD8+ cytotoxic T lymphocytes (CTL), drives viral diversification and is a significant challenge in developing effective therapies against HIV [47-50]. We investigated the impact of late compared to early ART initiation on CTL epitope diversity and escape in the HIV reservoir by longitudinally analysing Gag, Nef and Pol CTL epitopes, [51] from single genome viral sequences (excluding only hypermutated sequences), that are restricted by HLA genotypes B*57:02, B*57:03, B*58:01, B*81:01 and A*74:01. These HLA genotypes have been associated with protection against disease progression in HIV-1 subtype C infection [52]. CTL epitope mutations were classified according to the Los Alamos HIV Molecular Immunology Database [51]. Protective HLA genotypes were present in 7/11 (64%) late treated participants and in 7/13 (54%) early treated participants. In the presence of relevant restricting HLA genotypes, mutations compared to the Clade C consensus were detected in 12% of participants with Gag, 23% with Pol and 27% with Nef targeted epitopes after 1 year of follow-up when treatment was initiated late (Figure 5A-C) in contrast to 0%, 0% and 8% respectively when treatment was initiated early (Figure 5G-I) suggesting that chronic treatment is associated with the retention of a wide spectrum of CTL escape mutations within reservoirs compared to early treatment. Escape mutations detected at baseline (up to 1 month after infection) in the presence of restricting HLA genotypes were present in 3% of participants within Gag, 19% within Pol and 23% (Figure 5A-C) within Nef targeted epitopes when treatment was initiated later compared to and 0%, 13% and 11% respectively (Figure 5G-I) with early treatment. Escape mutations observed in early treated participants were present in the earliest sequences that were derived close to the time of infection and therefore likely represent transmitted escape variants. Similar proportions of transmitted

288 escape mutations were present in participants who did not have a protective HLA genotype and
289 remained unchanged after 1 year. (Figure 5D-F and 5J-L).
290
Discussion
In this study we used a well characterised acute HIV infection longitudinal cohort of untreated, early
treated and late treated study participants, to perform an extensive quantitative and qualitative analysis
of the HIV-1 subtype C viral reservoir. Our aim was to determine whether the timing of treatment has
an impact on the viral reservoir size, genetic landscape, and decay kinetics. This was a longitudinal
study where we measured total proviral load levels by ddPCR and characterised the genetic landscape
of the viral reservoir by next generation sequencing using FLIP-Seq. Total HIV-1 DNA impacts the
pathophysiology of infection and is therefore clinically important [30, 53-55]. We found that HIV DNA
was detectable at high levels during primary infection and even in participants who were treated during
acute infection, total HIV DNA levels measured at peak viremia was very similar to untreated
participants. In contrast, we found that early but not late treatment was associated with steady decline
of total proviral load over the first year of ART. These observations confirm previous data that the viral
reservoir is seeded at the earliest stages of infection, possibly before peak viremia [2, 56-59]. However,
in contrast, studies have suggested that early ART intervention restricts the seeding of the HIV reservoir
in long-lived central memory CD4 T cells [27]. Moreover, it has recently been demonstrated that a
small fraction of deeply latent genetically intact proviruses are archived in CD4 T cells during the very
first weeks of infection [60]. However, even though early ART initiation has been associated with
continued HIV DNA reduction during long-term ART after 10 years of follow up [28], it remains
detectable in most individuals indicating that early ART alone is insufficient to achieve viral
eradication.
311
We also examined the association of clinical and virological factors with the levels of proviral DNA
after 1 year of treatment. In both early and late treatment groups, peak plasma viral load correlated
positively with the HIV DNA levels. Moreover, in the late treatment group there was a positive
association with CD4 at enrolment (baseline) and an inverse correlation with nadir CD4 counts. It is
possible that higher levels of CD4 T cells at the onset of infection (baseline) presents the availability of
more target cells for HIV-1 reservoir establishment. Similar associations with nadir CD4 have been
reported previously [61-63] suggesting that during untreated progressive HIV infection, ongoing viral
replication may drive the accumulation of long-lived latently infected cells that repopulate the immune
system by expansion during successful ART.
321
Even though total HIV DNA has been shown to be a clinically significant marker of the HIV reservoir
it does not distinguish between replication-competent and -defective viruses that contribute to the viral
reservoir. The unique design of the FRESH cohort based on frequent HIV screening and sampling
intervals of high-risk uninfected participants allowed us to examine the dynamics of the proviral
landscape from the earliest stages of infection (Fiebig I) up to 1 year of ART by near-full-length viral
genome sequencing. We performed a comparison of the proviral populations between study participants
who were treated during the acute phase of infection and those initiating ART during chronic infection.
Defective genomes accumulate rapidly after the onset of infection and contributed to almost half of the
proviral population within the first four weeks of infection irrespective of treatment status. Consistent
with other studies [33, 56, 64] the overall viral reservoir landscape in the untreated and late treated
participants was dominated by defective viruses suggesting that prolonged ongoing viral replication
before treatment initiation leads to the accumulation of defective viral genomes. Genome-intact viruses
were easily detectable throughout untreated infection but decreased after treatment. In participants who

were treated during chronic infection, genome-intact viruses were still detectable after one year of ART compared to early treated individuals where they were no longer detectable.

With our limited sampling size and depth, of a median of 1.4 million PBMC (0.02-4.3 million) per sampling time point, we cannot rule out that intact genomes may be retrieved with further sampling and investigation into tissue reservoirs of the participants who initiate ART during acute infection. However, these findings do provide further evidence that introducing ART during acute HIV infection limits the size of the HIV reservoir considerably compared to treatment during chronic infection [11, 24-28].

The intact proviral DNA assay (IPDA) which is a more scalable method that uses multiplexed ddPCR to measure individual proviruses and differentiates intact from defective proviruses without the need for long-distance PCR, has been suggested to provide more accurate quantitative information about the size and composition of the latent reservoir compared to near full genome sequence methods [65]. However, this assay has not yet been developed and optimised for quantification of subtype C HIV.

Studies have shown that during suppressive ART, intact and defective proviruses have different rates of decay that occurs in a biphasic manner [11-13]. Our analysis of decay kinetics was limited to a linear mixed effects regression model as we were unable to fit a model for biphasic decay which requires frequent viral reservoir measurements. Our analysis was further restricted to the first six months of ART as viral genomes were difficult to detect by FLIP-seq in early treated participants after this time. We found that indeed intact genomes decay faster than defective genomes in both early and late treatment groups. Cells containing intact viral genomes likely represent productively infected cells that may be preferentially targeted for clearance by the host immune response or eliminated by viral cytopathic effects [12, 13]. Moreover, early treatment results in a faster decline of both intact and defective genomes compared to treatment initiated during chronic infection and is suggestive of a more effective immune clearance mechanism from preserved immune function. Despite this, it is estimated that 226 years of effective ART is necessary to decrease intact proviral DNA levels by 4 $\log_{10}$ [12]. This further indicates that early ART in combination with novel interventional strategies will be needed to achieve a faster viral eradication.

Several studies show that clonal expansion of HIV-infected cells plays an important role in maintaining the HIV reservoir further contributing to the challenge of eradicating HIV [66-69]. Our findings suggest that clonal expansion of intact viral genomes detected predominantly in early treated infection may contribute to the maintenance of the HIV reservoir in these study participants. Further studies that extend beyond 1 year of treatment will help elucidate whether these clones expand further after several years of treatment. Our analysis of CTL epitope escape mutations, known to drive viral diversification, revealed that early treatment minimises the emergence of CTL escape in Gag, Pol and Nef epitopes despite these participants having protective HLA alleles. In contrast CTL escape was detected when treatment was initiated during chronic infection specifically in the well characterised TW10 gag epitope restricted by HLA B57/58. Transmitted CTL escape mutations detected within the first few weeks of infection were common in both treatment groups confirming previous data from this study population [70]. Moreover, the rapid rate of clearance of viral genomes observed with early treatment could be attributed to the lower proportion of cytotoxic T lymphocyte escape mutations in the reservoir of these participants compared with those treated later.

To our knowledge this is the first study in an African population, dominated by subtype C HIV infection,

that examined the impact of the timing of ART initiation on HIV reservoirs in a longitudinal setting. Moreover, our data focused on women who are underrepresented in reservoir and cure studies globally. Our data showed that early ART initiation does not blunt proviral DNA seeding in immune reservoirs, but it nevertheless results in a more rapid decay of intact viral genomes, decreases genetic complexity and immune escape and preserves innate immune factors. Although early ART alone may not be sufficient to eradicate the persisting viral reservoir, our results suggest that when combined with interventional strategies, it is more likely to achieve an effective HIV cure.

Materials and Methods

Ethics Statement
The Biomedical Research Ethics Committee of the University of KwaZulu-Natal and the Institutional Review Board of Massachusetts General Hospital approved the study. All participants provided written informed consent.

Study Design and Participants
This was a longitudinal study of the Females Rising through Education, Support, and Health (FRESH) cohort, a prospective, observational study of 18–23-year-old HIV uninfected women at high risk for HIV acquisition, established in Umlazi, Durban, South Africa [45, 46]. Finger prick blood draws were collected from FRESH study participants twice a week and subjected to HIV-1 RNA testing, with the aim of detecting acute HIV infection during Fiebig stage I. The study included a socioeconomic intervention program and HIV prevention interventions including PrEP that coincided with study visits to address challenges faced by the young women that likely contribute to the increased risk of HIV acquisition in this setting. If a participant acquired HIV-1 infection while on the study, blood samples were collected weekly for a month, then monthly until 3 months post infection, then monthly for one year and every 3 months thereafter. Days post onset of viremia (DPOPV) was calculated as the interval between the first positive HIV test and the date of sample collection. Unique participant identifier numbers were assigned to the participants and are only known to the research group.

Study participants recruited during the first 19 months of the study did not receive antiretroviral (ARV) treatment immediately after detection of acute HIV infection but were monitored and referred for treatment when they became eligible according to national treatment guidelines at the time. The South African national treatment eligibility criteria subsequently changed allowing the immediate initiation of ART for all people living with HIV (PLWH), including those with acute HIV infection as recommended under the World Health Organisation's universal test and treat policy [71]. The treatment schedule was a three-drug daily oral regimen of 300 mg tenofovir disoproxil fumarate, 200 mg emtricitabine, and 600 mg efavirenz. Additionally, raltegravir (400 mg twice a day) was introduced as a fourth drug and was continued for 90 days after viral suppression (<20 copies per mL).

For this study participants were categorized into 3 groups (untreated, late (chronic) treated, and early (acute) treated) where, 11 remained untreated during acute infection and later started ART during chronic infection at a median of 456 (297-1203) days post onset of viremia (DPOV), while 24 started ART at a median of 1 (1-3) DPOV. Participants were studied at 0-, 1-, 3-, 6-, 9-, 12- and 24-months post onset of viremia and up to 12 months post treatment.

Quantification of total HIV-1 DNA
Measurement of total HIV-1 DNA was performed as previously described [72]. Total DNA was extracted from total PBMC samples using DNeasy Blood & Tissue Kits (QIAGEN). Droplet digital

PCR (ddPCR) (Bio-Rad) was used to measure total HIV-1 DNA and host cell concentration with primers and probes covering HIV-1 5′ LTR-*gag* HXB2 coordinates 684–810 (forward primer 5′-TCTCGACGCAGGACTCG-3′, reverse primer 5′-TACTGACGCTCTCGCACC-3′ probe/56-FAM/CTCTCTCCT/ZEN/TCTAGCCTC/ 31ABkFQ/, and human RPP30 gene38 forward primer 5′-GATTTGGACCTGCGAGCG-3′, reverse primer 5′-GCGGCTGTCTCCACAAGT-3′, probe/56-FAM/CTGACCTGA/ZEN/AGGCTCT/31AbkFQ/). Thermocycling conditions for ddPCR were: 95 °C for 10 min, 45 cycles of 94 °C for 30 s and 60 °C for 1 min, 72 °C for 1 min. Thereafter droplets from each sample were analyzed on the Bio-Rad QX200 Droplet Reader and data were analysed using QuantaSoft software (Bio-Rad).

Single genome amplification and deep sequencing of near-full length HIV-1 DNA
Near full-length proviral sequences were generated as previously described [72]. Total HIV-DNA copy number as determined by ddPCR was used to calculate the DNA sample dilution to achieve one PCR-positive reaction in every three reactions. This method of limiting dilution gives a Poisson probability of 85.7% that each PCR amplicon originated from a single HIV-DNA template [73]. A nested PCR approach was used to amplify the near-full genome using one unit of Platinum™ Taq DNA Polymerase High Fidelity (Invitrogen) with the following primers: first round PCR: forward primer 5′-AAATCTCTAGCAGTGGCGCCCGAACAG-3′, reverse primer 5′-TGAGGGATCTCTAGTTACCAGAGTC-3′; second round PCR: forward primer 5′-GCGCCCGAACAGGGACYTGAAARCGAAAG-3′, reverse primer 5′-GCACTCAAGGCAAGCTT TATTGAGGCTTA-3′ (HXB2 coordinates 638-9632, 8994 bp). The 20 μL reaction mix contained 1× reaction buffer, 2 mM MgSO4, 0.2mM dNTP, 0.4 μM each of forward and reverse primers. The thermal cycling conditions were 2 min at 92 °C, 10 cycles [10 s at 92 °C, 30 s at 60 °C, 10 min at 68 °C], 20 cycles [10 s at 92 °C, 30 s at 55 °C, 10 min at 68 °C], 10 min at 68 °C, 4 °C infinite hold.

Illumina Mi-Seq and Bioinformatics Analysis
All PCR amplicons detectable by gel electrophoresis were subjected to Illumina MiSeq sequencing and thereafter the resulting small reads were *de novo* assembled using in-house UltraCycler v1.0. (Brian Seed and Huajun Wang, unpublished) [72]. Viral genome-intactness was inferred by the computational bioinformatics pipeline HIVSeqinR v2.7.1 [33].

Statistical methods
GraphPad Prism 10 was used to perform summary statistical analyses and comparisons among study groups using Fishers' Exact, Mann-Whitney and Kruskal-Wallis and multiple linear regression analysis.

Acknowledgements
The study cohort and sample collection were supported in part by grants from the Bill and Melinda Gates Foundation (OPP1066973 and OPP1146433), Gilead Sciences, Inc. (Grant ID #00406), the International AIDS Vaccine Initiative (IAVI) (UKZNRSA1001), the NIAID (R37AI067073), the Witten Family Foundation, the Dan and Marjorie Sullivan Foundation, the Mark and Lisa Schwartz Foundation, Ursula Brunner, the AIDS Healthcare Foundation, and the Harvard University Center for AIDS Research (CFAR, P30 AI060354, which is supported by the following institutes and centers co-funded by and participating with the US National Institutes of Health: NIAID, NCI, NICHD, NHLBI, NIDA, NIMH, NIA, FIC, and OAR.). Raltegravir used for immediate treatment was donated by Merck & Co., Inc. This work was also partially supported through the Sub-Saharan African Network for TB/HIV Research Excellence (SANTHE) which is funded by the Science for Africa Foundation to the Developing Excellence in Leadership, Training and Science in Africa (DELTAS Africa) programme [Del-22-007] with support from Wellcome Trust and the UK Foreign, Commonwealth & Development

Office and is part of the EDCPT2 programme supported by the European Union; the Bill & Melinda
Gates Foundation [INV-033558]; and Gilead Sciences Inc., [19275]. All content contained within is
that of the authors and does not necessarily reflect positions or policies of any SANTHE funder. For
the purpose of open access, the author has applied a CC BY public copyright licence to any Author
Accepted Manuscript version arising from this submission. The authors thank all participants in the
FRESH cohort who have made this study possible. The authors thank the Massachusetts General
Hospital Center for Computational & Integrative Biology DNA Core, specifically Dr. Nicole Stange-
Thomann, Dr. Amy Avery, Ms. Kristina Belanger, and Mr. Huajun Wang, for providing them with the
Illumina MiSeq deep sequencing service used in this manuscript.

487
Author Contributions

489 The study was conceptualized and designed by K.R., T.N., G.Q.L., M.L. and X.G.Y. and PBMC
490 samples and clinical/demographical data were collected by K.L.D., B.D.W., K.R., and T.N. HIV-1
491 genotyping laboratory work was done by K.R., G.Q.L., N.R., and T.J.B.C. Results were analyzed by
492 K.R., G.Q.L., and T.N. K.R and T.N and G.Q.L. wrote the manuscript; all authors contributed to and
493 approved the manuscript. T.N. supervised the study.

Table 1: Characteristics of study participants

Characteristics	Chronic Treated (n=11)	Acute Treated (n=24)
Age (years)	21 (19-24)	21 (18-24)
Sex		
Female, n (%)	11 (100%)	24 (100%)
Male, n (%)	0 (0%)	0 (0%)
Race/ethnicity, n (%)		
Black	11 (100%)	24 (100%)
Fiebig Stage I at detection, n (%)	10 (91%)	21 (88%)
Treatment Initiation (DPOPV)	456 (297-1203)	1 (1-3)
Time to suppression (days)	104 (30-215)	16 (6-116)
CD4 nadir (cells/µl)	383 (204-502)	561 (258-859)
CD4 pre-infection (cells/µl)	991 (395-1377)	872 (573-1612)
CD4 at study enrollment (baseline) (cells/µl)	716 (204-1377)	863 (421-2075)
Peak plasma viral load (log copies/ml)	7.04 (5.89-7.80)	4.21 (2-7.30)
*Protective HLA Allele, n (%)	6 (55%)	11 (46%)
Treatment Regimen Containing, n (%)		
FDC	11 (100%)	24 (100%)
Raltegravir	0 (0%)	16 (67%)

* HLA-B74:01, HLA-B57:02, HLA-B57:03, HLA-B58:01, HLA-B81:01

Figure 1: **Plasma viral load and total HIV DNA in acute treated and chronic treated individuals.** A) peak viral load and total HIV DNA measured at peak viral load in untreated (pre-therapy) and acute treated individuals B) longitudinal viral load and total HIV DNA in untreated acute infection and after 1 year of treatment C) longitudinal viral load and total HIV DNA in acute treated individuals D) viral load and total HIV DNA after 1 year of treatment in chronic and acute treated individuals.

Table 2: Multivariate analysis of factors that predict total HIV-1 proviral DNA load after 1 year of treatment.

Stage at Treatment Initiation	Variables	Co-efficient	Standard error	t	P value	P value summary	95% confidence interval
Acute Infection	Nadir CD4	-0,0007424	0,00102	0,7277	0,4773	ns	-0,002905 to 0,001420
	Pre-Infection CD4	-0,0001861	0,0004233	0,4395	0,6661	ns	-0,001083 to 0,0007113
	Baseline* CD4	0,0002574	0,0003727	0,6906	0,4997	ns	-0,0005326 to 0,001047
	Peak VL	0,1972	0,07938	2,485	0,0244	*	0,02895 to 0,3655
Chronic Infection	Nadir CD4	-0,006633	0,0007088	9,358	<0,0001	****	-0,008367 to -0,004898
	Pre-Infection CD4	0,0002514	0,0003896	0,6454	0,5425	ns	-0,0007019 to 0,001205
	Baseline* CD4	0,001501	0,0002905	5,166	0,0021	**	0,0007899 to 0,002211
	Peak VL	0,2658	0,09698	2,741	0,0337	*	0,02848 to 0,5031

*at study enrolment

Figure 2: Genotypic characterisation of HIV-DNA sequences. A) PBMC sequencing timepoints in untreated (red), chronic treated (green) and early treated (blue) study participants where each dot represents a sampling time point. Time of treatment initiation is shown by the vertical grey bar. B) Approximately-maximum-likelihood phylogenetic tree of intact HIV-1 DNA genomes constructed using FastTree2. This method was chosen to resolve full-viral-genome sequences with extreme homology; branch lengths were likely inflated. Viral genomes derived from acute treated participants are marked with (*). C) Comparison of intraparticipant mean pairwise distances between early and late treated participants. D) Spectrum of HIV genome sequences detected during untreated acute infection, late treated chronic infection, and acute treated infection.

Figure 3: Evolution of the proviral genetic landscape. Relative proportions of intact and defective viral genomes measured longitudinally in A) untreated acute infection for 2 years B) late (chronic) treated infection for 1 year and C) early (acute) treated infection for 1 year.

Figure 4: **Decay kinetics of intact and defective proviruses.** Absolute frequencies of intact and defective HIV-1 DNA sequences per million PBMCs during the 1st year of infection following treatment during A) acute infection and B) chronic infection. Longitudinal analysis of the change in (C) intact and (D) defective provirus copies in the 6 months after ART initiation, comparing the acute treated (blue) and chronic treated (green) groups. Dots represent a measurement from a given participant; solid lines are slopes estimated from linear mixed effect model. (E) Comparison of the monthly rate of decay of intact and defective proviruses in acute and chronic treated infection.

Figure 5: **Comparison of CTL epitope diversity in late compared to early treated participants.** Proportion of participants with wildtype, variant and CTL escape at baseline (within 1 month of infection) and up to 1 year of infection in Gag (A, D, G, J), Pol (B, E, H, K) and Nef (C, F, I, L) epitopes in participants with protective HLA genotypes (A, B, C, G, H, I) and without protective HLA genotypes (D, E, F, J, K, L).

Supplementary Table 1: Clinical and biological characteristics of 35 study participants

Identifier	HLA-A1	HLA-A2	HLA-B1	HLA-B2	HLA-C1	HLA-C2	Fiebig Stage at Detection	Timing of Treatment Initiation (DPOPV)	Peak Viral Load	ddPCR	FLIP Seq
FRESH_002	02:01	30:01	44:03	58:02	04:01	06:02	I	724	5,70E+07	✔	✔
FRESH_003	02:01	30:01	15:10	42:02	08:04	17	I	310	8,20E+06	✔	✔
FRESH_005	24:02	29:02	07:02	44:03	07:01	07:02	I	809	5,20E+07	✔	✔
FRESH_004	03:01	74:01	15:03	58:02	02:10	06:02	I	346	3,00E+07	✔	✔
FRESH_010	23:01	30:01	15:10	58:01	03:02	16:01	III	668	5,70E+07	✔	✔
FRESH_011	23:01	68:02	08:01	58:01	03:04	07:01	I	427	1,10E+07	✔	✔
FRESH_007	23:01	74:01	35:01	58:01	04:01	06:02	I	456	4,00E+06	✔	✔
FRESH_013	01:01	66:01	39:10	81:01	12:03	18	I	1202	7,70E+05	✔	✔
FRESH_009	43:01	43:01	07:02	15:03	02:10	18	I	416	9,90E+06	✔	✔
FRESH_008	02:05	02:05	58:01	58:01	07:01	07:01	I	297	9,50E+06	✔	✔
FRESH_012	23:01	29:02	42:01	53:01	03:04	17	I	1059	1,20E+07	✔	✔
FRESH_015	30:02	34:02	08:01	44:03	04:01	07:01	I	1	2,30E+04	✔	✗
FRESH_018	68:02	74:01	15:03	57:02	02:10	18	V	2	1,50E+02	✔	✔
FRESH_016	01:23	30:02	42:01	58:01	06:02	17	I	1	3,40E+04	✔	✔
FRESH_029	30:01	30:01	39:10	42:02	12:03	17:01	I	1	1,40E+04	✔	✗
FRESH_019	02:01	03:01	58:01	58:02	03:02	06:02	I	1	3,60E+05	✔	✔
FRESH_033	68:02	68:02	07:02	15:10	03:04	07:02	I	1	4,40E+02	✔	✔
FRESH_022	30:01	30:02	18:01	53:01	04:01	07:04	I	1	2,10E+05	✔	✔
FRESH_025	01:01	30:01	42:01	47:01	06:02	17	I	3	8,90E+04	✔	✔
FRESH_024	01:01	29:01	44:03	81:01	08:04	18	I	1	5,90E+03	✔	✔
FRESH_030	02:05	29:02	42:01	58:01	07:01	17:01	III	1	2,00E+07	✔	✗
FRESH_023	02:05	23:01	08:01	14:01	07:01	08:04	I	3	4,10E+04	✔	✔
FRESH_036	02:05	68:01	35:01	58:01	04:01	07:01	I	1	1,90E+04	✔	✗
FRESH_031	01:01	68:02	14:01	81:01	08:02	18	I	1	1,70E+04	✔	✔
FRESH_038	30:01	30:02	08:01	15:03	02:10	07:01	I	1	4,60E+02	✔	✗
FRESH_028	03:01	29:02	44:03	58:02	06:02	07:01	I	2	2,60E+06	✔	✔
FRESH_035	29:02	33:03	07:02	44:03	07:01	07:02	I	1	7,70E+02	✔	✗
FRESH_032	30:01	74:01	15:03	18:01	02:10	07:04	I	1	7,30E+03	✔	✔
FRESH_034	30:02	33:03	53:01	58:02	04:01	06:02	I	2	8,60E+04	✔	✔
FRESH_039	02:01	02:05	08:01	58:02	06:02	07:01	I	1	2,50E+03	✔	✗
FRESH_048	30:01:00	30:02	15:03	57:03	02:10	07:01	I	1	1,50E+04	✔	✔
FRESH_041	29:02	74:01	44:03	57:03	07:01	07:01	I	1	1,40E+03	✔	✗
FRESH_046	23:01	30:01	15:10	18:01	07:04	16:01	I	1	2,50E+03	✔	✗
FRESH_042	29:02	30:02	07:02	44:03	07:01	07:02	III	1	1,50E+04	✔	✗
FRESH_044	01:01	02:05	42:02	81:01	17:01	18	I	1	6,50E+04	✔	✗

Supplementary Figure 1: In this cohort of HIV-1 subtype C, genome deletions were most frequently observed between between Integrase and Envelope relative to Gag (p<0.0001–0.001).

Supplementary Figure 2: Clonal expansion of infected cells was detected in both defective (orange) and intact (blue) genomes in late and early treated study participants.

References

1. Finzi, D., J. Blankson, J.D. Siliciano, J.B. Margolick, K. Chadwick, T. Pierson, K. Smith, J. Lisziewicz, F. Lori, C. Flexner, T.C. Quinn, R.E. Chaisson, E. Rosenberg, B. Walker, S. Gange, J. Gallant, and R.F. Siliciano, *Latent infection of CD4+ T cells provides a mechanism for lifelong persistence of HIV-1, even in patients on effective combination therapy.* Nat Med, 1999. **5**(5): p. 512-7.

2. Siliciano, J.D., J. Kajdas, D. Finzi, T.C. Quinn, K. Chadwick, J.B. Margolick, C. Kovacs, S.J. Gange, and R.F. Siliciano, *Long-term follow-up studies confirm the stability of the latent reservoir for HIV-1 in resting CD4+ T cells.* Nat Med, 2003. **9**(6): p. 727-8.

3. Wong, J.K., M. Hezareh, H.F. Gunthard, D.V. Havlir, C.C. Ignacio, C.A. Spina, and D.D. Richman, *Recovery of replication-competent HIV despite prolonged suppression of plasma viremia.* Science, 1997. **278**(5341): p. 1291-5.

4. Deeks, S.G., N. Archin, P. Cannon, S. Collins, R.B. Jones, M. de Jong, O. Lambotte, R. Lamplough, T. Ndung'u, J. Sugarman, C.T. Tiemessen, L. Vandekerckhove, S.R. Lewin, and A.S.G.S.S.w.g. International, *Research priorities for an HIV cure: International AIDS Society Global Scientific Strategy 2021.* Nat Med, 2021. **27**(12): p. 2085-2098.

5. Chomont, N., M. El-Far, P. Ancuta, L. Trautmann, F.A. Procopio, B. Yassine-Diab, G. Boucher, M.R. Boulassel, G. Ghattas, J.M. Brenchley, T.W. Schacker, B.J. Hill, D.C. Douek, J.P. Routy, E.K. Haddad, and R.P. Sekaly, *HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation.* Nat Med, 2009. **15**(8): p. 893-900.

6. Vandergeeten, C., R. Fromentin, S. DaFonseca, M.B. Lawani, I. Sereti, M.M. Lederman, M. Ramgopal, J.P. Routy, R.P. Sekaly, and N. Chomont, *Interleukin-7 promotes HIV persistence during antiretroviral therapy.* Blood, 2013. **121**(21): p. 4321-9.

7. Mendoza, P., J.R. Jackson, T.Y. Oliveira, C. Gaebler, V. Ramos, M. Caskey, M. Jankovic, M.C. Nussenzweig, and L.B. Cohn, *Antigen-responsive CD4+ T cell clones contribute to the HIV-1 latent reservoir.* J Exp Med, 2020. **217**(7).

8. Jiang, C., X. Lian, C. Gao, X. Sun, K.B. Einkauf, J.M. Chevalier, S.M.Y. Chen, S. Hua, B. Rhee, K. Chang, J.E. Blackmer, M. Osborn, M.J. Peluso, R. Hoh, M. Somsouk, J. Milush, L.N. Bertagnolli, S.E. Sweet, J.A. Varriale, P.D. Burbelo, T.W. Chun, G.M. Laird, E. Serrao, A.N. Engelman, M. Carrington, R.F. Siliciano, J.M. Siliciano, S.G. Deeks, B.D. Walker, M. Lichterfeld, and X.G. Yu, *Distinct viral reservoirs in individuals with spontaneous control of HIV-1.* Nature, 2020. **585**(7824): p. 261-267.

9. Maldarelli, F., X. Wu, L. Su, F.R. Simonetti, W. Shao, S. Hill, J. Spindler, A.L. Ferris, J.W. Mellors, M.F. Kearney, J.M. Coffin, and S.H. Hughes, *HIV latency. Specific HIV integration sites are linked to clonal expansion and persistence of infected cells.* Science, 2014. **345**(6193): p. 179-83.

10. Wagner, T.A., S. McLaughlin, K. Garg, C.Y. Cheung, B.B. Larsen, S. Styrchak, H.C. Huang, P.T. Edlefsen, J.I. Mullins, and L.M. Frenkel, *HIV latency. Proliferation of cells with HIV integrated into cancer genes contributes to persistent infection.* Science, 2014. **345**(6196): p. 570-3.

11. Massanella, M., R.A.B. Ignacio, J.R. Lama, A. Pagliuzza, S. Dasgupta, R. Alfaro, J. Rios, C. Ganoza, D. Pinto-Santini, T. Gilada, A. Duerr, and N. Chomont, *Long-term effects of early antiretroviral initiation on HIV reservoir markers: a longitudinal analysis of the MERLIN clinical study.* Lancet Microbe, 2021. **2**(5): p. e198-e209.

12. Peluso, M.J., P. Bacchetti, K.D. Ritter, S. Beg, J. Lai, J.N. Martin, P.W. Hunt, T.J. Henrich, J.D. Siliciano, R.F. Siliciano, G.M. Laird, and S.G. Deeks, *Differential decay of intact and defective proviral DNA in HIV-1-infected individuals on suppressive antiretroviral therapy.* JCI Insight, 2020. **5**(4).

13. White, J.A., F.R. Simonetti, S. Beg, N.F. McMyn, W. Dai, N. Bachmann, J. Lai, W.C. Ford, C. Bunch, J.L. Jones, R.M. Ribeiro, A.S. Perelson, J.D. Siliciano, and R.F. Siliciano, *Complex decay dynamics of HIV virions, intact and defective proviruses, and 2LTR circles following initiation of antiretroviral therapy.* Proc Natl Acad Sci U S A, 2022. **119**(6).

14. Andrade, V.M., C. Mavian, D. Babic, T. Cordeiro, M. Sharkey, L. Barrios, C. Brander, J. Martinez-Picado, J. Dalmau, A. Llano, J.Z. Li, J. Jacobson, C.L. Lavine, M.S. Seaman, M. Salemi, and M. Stevenson, *A minor population of macrophage-tropic HIV-1 variants is identified in recrudescing viremia following analytic treatment interruption.* Proc Natl Acad Sci U S A, 2020. **117**(18): p. 9981-9990.

15. Hiener, B., B.A. Horsburgh, J.S. Eden, K. Barton, T.E. Schlub, E. Lee, S. von Stockenstrom, L. Odevall, J.M. Milush, T. Liegler, E. Sinclair, R. Hoh, E.A. Boritz, D. Douek, R. Fromentin, N. Chomont, S.G. Deeks, F.M. Hecht, and S. Palmer, *Identification of Genetically Intact HIV-1 Proviruses in Specific CD4(+) T Cells from Effectively Treated Participants.* Cell Rep, 2017. **21**(3): p. 813-822.

16. Honeycutt, J.B., A. Wahl, C. Baker, R.A. Spagnuolo, J. Foster, O. Zakharova, S. Wietgrefe, C. Caro-Vegas, V. Madden, G. Sharpe, A.T. Haase, J.J. Eron, and J.V. Garcia, *Macrophages sustain HIV replication in vivo independently of T cells.* J Clin Invest, 2016. **126**(4): p. 1353-66.

17. Koppensteiner, H., R. Brack-Werner, and M. Schindler, *Macrophages and their relevance in Human Immunodeficiency Virus Type I infection.* Retrovirology, 2012. **9**: p. 82.

18. Venanzi Rullo, E., L. Cannon, M.R. Pinzone, M. Ceccarelli, G. Nunnari, and U. O'Doherty, *Genetic Evidence That Naive T Cells Can Contribute Significantly to the Human Immunodeficiency Virus Intact Reservoir: Time to Re-evaluate Their Role.* Clin Infect Dis, 2019. **69**(12): p. 2236-2237.

19. Wightman, F., A. Solomon, G. Khoury, J.A. Green, L. Gray, P.R. Gorry, Y.S. Ho, N.K. Saksena, J. Hoy, S.M. Crowe, P.U. Cameron, and S.R. Lewin, *Both CD31(+) and CD31(-) naive CD4(+) T cells are persistent HIV type 1-infected reservoirs in individuals receiving antiretroviral therapy.* J Infect Dis, 2010. **202**(11): p. 1738-48.

20. Banga, R., F.A. Procopio, E. Lana, G.T. Gladkov, I. Roseto, E.M. Parsons, X. Lian, M. Armani-Tourret, M. Bellefroid, C. Gao, A. Kauzlaric, M. Foglierini, O. Alfageme-Abello, S.H.M. Sluka, O. Munoz, A. Mastrangelo, C. Fenwick, Y. Muller, C.G. Mkindi, C. Daubenberger, M. Cavassini, R. Trunfio, S. Deglise, J.M. Corpataux, M. Delorenzi, M. Lichterfeld, G. Pantaleo, and M. Perreau, *Lymph node dendritic cells harbor inducible replication-competent HIV despite years of suppressive ART.* Cell Host Microbe, 2023. **31**(10): p. 1714-1731 e9.

21. Keele, B.F., L. Tazi, S. Gartner, Y. Liu, T.B. Burgon, J.D. Estes, T.C. Thacker, K.A. Crandall, J.C. McArthur, and G.F. Burton, *Characterization of the follicular dendritic cell reservoir of human immunodeficiency virus type 1.* J Virol, 2008. **82**(11): p. 5548-61.

22. Olivetta, E., C. Chiozzini, C. Arenaccio, F. Manfredi, F. Ferrantelli, and M. Federico, *Extracellular vesicle-mediated intercellular communication in HIV-1 infection and its role in the reservoir maintenance.* Cytokine Growth Factor Rev, 2020. **51**: p. 40-48.

23. Ollerton, M.T., E.A. Berger, E. Connick, and G.F. Burton, *HIV-1-Specific Chimeric Antigen Receptor T Cells Fail To Recognize and Eliminate the Follicular Dendritic Cell HIV Reservoir In Vitro.* J Virol, 2020. **94**(10).

24. Namazi, G., J.M. Fajnzylber, E. Aga, R.J. Bosch, E.P. Acosta, R. Sharaf, W. Hartogensis, J.M. Jacobson, E. Connick, P. Volberding, D. Skiest, D. Margolis, M.C. Sneller, S.J. Little, S. Gianella, D.M. Smith, D.R. Kuritzkes, R.M. Gulick, J.W. Mellors, V. Mehraj, R.T. Gandhi, R. Mitsuyasu, R.T. Schooley, K. Henry, P. Tebas, S.G. Deeks, T.W. Chun, A.C. Collier, J.P. Routy, F.M. Hecht, B.D. Walker, and J.Z. Li, *The Control of HIV After Antiretroviral*

Medication Pause (CHAMP) Study: Posttreatment Controllers Identified From 14 Clinical Studies. J Infect Dis, 2018. **218**(12): p. 1954-1963.

25. Saez-Cirion, A., C. Bacchus, L. Hocqueloux, V. Avettand-Fenoel, I. Girault, C. Lecuroux, V. Potard, P. Versmisse, A. Melard, T. Prazuck, B. Descours, J. Guergnon, J.P. Viard, F. Boufassa, O. Lambotte, C. Goujard, L. Meyer, D. Costagliola, A. Venet, G. Pancino, B. Autran, C. Rouzioux, and A.V.S. Group, *Post-treatment HIV-1 controllers with a long-term virological remission after the interruption of early initiated antiretroviral therapy ANRS VISCONTI Study.* PLoS Pathog, 2013. **9**(3): p. e1003211.

26. Ananworanich, J., N. Chomont, L.A. Eller, E. Kroon, S. Tovanabutra, M. Bose, M. Nau, J.L.K. Fletcher, S. Tipsuk, C. Vandergeeten, R.J. O'Connell, S. Pinyakorn, N. Michael, N. Phanuphak, M.L. Robb, Rv, and R.S.s. groups, *HIV DNA Set Point is Rapidly Established in Acute HIV Infection and Dramatically Reduced by Early ART.* EBioMedicine, 2016. **11**: p. 68-72.

27. Ananworanich, J., A. Schuetz, C. Vandergeeten, I. Sereti, M. de Souza, R. Rerknimitr, R. Dewar, M. Marovich, F. van Griensven, R. Sekaly, S. Pinyakorn, N. Phanuphak, R. Trichavaroj, W. Rutvisuttinunt, N. Chomchey, R. Paris, S. Peel, V. Valcour, F. Maldarelli, N. Chomont, N. Michael, P. Phanuphak, J.H. Kim, and R.S.S. Group, *Impact of multi-targeted antiretroviral treatment on gut T cell depletion and HIV reservoir seeding during acute HIV infection.* PLoS One, 2012. **7**(3): p. e33948.

28. Buzon, M.J., E. Martin-Gayo, F. Pereyra, Z. Ouyang, H. Sun, J.Z. Li, M. Piovoso, A. Shaw, J. Dalmau, N. Zangger, J. Martinez-Picado, R. Zurakowski, X.G. Yu, A. Telenti, B.D. Walker, E.S. Rosenberg, and M. Lichterfeld, *Long-term antiretroviral treatment initiated at primary HIV-1 infection affects the size, composition, and decay kinetics of the reservoir of HIV-1-infected CD4 T cells.* J Virol, 2014. **88**(17): p. 10056-65.

29. Li, J.Z., B. Etemad, H. Ahmed, E. Aga, R.J. Bosch, J.W. Mellors, D.R. Kuritzkes, M.M. Lederman, M. Para, and R.T. Gandhi, *The size of the expressed HIV reservoir predicts timing of viral rebound after treatment interruption.* AIDS, 2016. **30**(3): p. 343-53.

30. Williams, J.P., J. Hurst, W. Stohr, N. Robinson, H. Brown, M. Fisher, S. Kinloch, D. Cooper, M. Schechter, G. Tambussi, S. Fidler, M. Carrington, A. Babiker, J. Weber, K.K. Koelsch, A.D. Kelleher, R.E. Phillips, J. Frater, and S.P. Investigators, *HIV-1 DNA predicts disease progression and post-treatment virological control.* Elife, 2014. **3**: p. e03821.

31. Jeewanraj, N., T. Mandizvo, T. Mulaudzi, N. Gumede, Z. Ndhlovu, T. Ndung'u, K. Gounder, and J. Mann, *Partial compartmentalisation of HIV-1 subtype C between lymph nodes, peripheral blood mononuclear cells and plasma.* Virology, 2023. **582**: p. 62-70.

32. Kloverpris, H.N., S.W. Kazer, J. Mjosberg, J.M. Mabuka, A. Wellmann, Z. Ndhlovu, M.C. Yadon, S. Nhamoyebonde, M. Muenchhoff, Y. Simoni, F. Andersson, W. Kuhn, N. Garrett, W.A. Burgers, P. Kamya, K. Pretorius, K. Dong, A. Moodley, E.W. Newell, V. Kasprowicz, S.S. Abdool Karim, P. Goulder, A.K. Shalek, B.D. Walker, T. Ndung'u, and A. Leslie, *Innate Lymphoid Cells Are Depleted Irreversibly during Acute HIV-1 Infection in the Absence of Viral Suppression.* Immunity, 2016. **44**(2): p. 391-405.

33. Lee, G.Q., K. Reddy, K.B. Einkauf, K. Gounder, J.M. Chevalier, K.L. Dong, B.D. Walker, X.G. Yu, T. Ndung'u, and M. Lichterfeld, *HIV-1 DNA sequence diversity and evolution during acute subtype C infection.* Nat Commun, 2019. **10**(1): p. 2737.

34. Mabuka, J.M., A.S. Dugast, D.M. Muema, T. Reddy, Y. Ramlakhan, Z. Euler, N. Ismail, A. Moodley, K.L. Dong, L. Morris, B.D. Walker, G. Alter, and T. Ndung'u, *Plasma CXCL13 but Not B Cell Frequencies in Acute HIV Infection Predicts Emergence of Cross-Neutralizing Antibodies.* Front Immunol, 2017. **8**: p. 1104.

35. Moir, S., C.M. Buckner, J. Ho, W. Wang, J. Chen, A.J. Waldner, J.G. Posada, L. Kardava, M.A. O'Shea, S. Kottilil, T.W. Chun, M.A. Proschan, and A.S. Fauci, *B cells in early and chronic*

HIV infection: evidence for preservation of immune function associated with early initiation of antiretroviral therapy. Blood, 2010. **116**(25): p. 5571-9.

36. Muema, D.M., N.A. Akilimali, O.C. Ndumnego, S.S. Rasehlo, R. Durgiah, D.B.A. Ojwach, N. Ismail, M. Dong, A. Moodley, K.L. Dong, Z.M. Ndhlovu, J.M. Mabuka, B.D. Walker, J.K. Mann, and T. Ndung'u, *Association between the cytokine storm, immune cell dynamics, and viral replicative capacity in hyperacute HIV infection.* BMC Med, 2020. **18**(1): p. 81.

37. Naidoo, K.K., A.J. Highton, O.O. Baiyegunhi, S.P. Bhengu, K.L. Dong, M.J. Bunders, M. Altfeld, and T. Ndung'u, *Early initiation of antiretroviral therapy preserves the metabolic function of CD4+ T-cells in subtype C HIV-1 infection.* J Infect Dis, 2023.

38. Naidoo, K.K., O.C. Ndumnego, N. Ismail, K.L. Dong, and T. Ndung'u, *Antigen Presenting Cells Contribute to Persistent Immune Activation Despite Antiretroviral Therapy Initiation During Hyperacute HIV-1 Infection.* Front Immunol, 2021. **12**: p. 738743.

39. Ndhlovu, Z.M., S.W. Kazer, T. Nkosi, F. Ogunshola, D.M. Muema, G. Anmole, S.A. Swann, A. Moodley, K. Dong, T. Reddy, M.A. Brockman, A.K. Shalek, T. Ndung'u, and B.D. Walker, *Augmentation of HIV-specific T cell function by immediate treatment of hyperacute HIV-1 infection.* Sci Transl Med, 2019. **11**(493).

40. Rosenberg, E.S., M. Altfeld, S.H. Poon, M.N. Phillips, B.M. Wilkes, R.L. Eldridge, G.K. Robbins, R.T. D'Aquila, P.J. Goulder, and B.D. Walker, *Immune control of HIV-1 after early treatment of acute infection.* Nature, 2000. **407**(6803): p. 523-6.

41. Rosenberg, E.S., J.M. Billingsley, A.M. Caliendo, S.L. Boswell, P.E. Sax, S.A. Kalams, and B.D. Walker, *Vigorous HIV-1-specific CD4+ T cell responses associated with control of viremia.* Science, 1997. **278**(5342): p. 1447-50.

42. Chang, J.J., M. Woods, R.J. Lindsay, E.H. Doyle, M. Griesbeck, E.S. Chan, G.K. Robbins, R.J. Bosch, and M. Altfeld, *Higher expression of several interferon-stimulated genes in HIV-1-infected females after adjusting for the level of viral replication.* J Infect Dis, 2013. **208**(5): p. 830-8.

43. Meier, A., J.J. Chang, E.S. Chan, R.B. Pollard, H.K. Sidhu, S. Kulkarni, T.F. Wen, R.J. Lindsay, L. Orellana, D. Mildvan, S. Bazner, H. Streeck, G. Alter, J.D. Lifson, M. Carrington, R.J. Bosch, G.K. Robbins, and M. Altfeld, *Sex differences in the Toll-like receptor-mediated response of plasmacytoid dendritic cells to HIV-1.* Nat Med, 2009. **15**(8): p. 955-9.

44. Sterling, T.R., D. Vlahov, J. Astemborski, D.R. Hoover, J.B. Margolick, and T.C. Quinn, *Initial plasma HIV-1 RNA levels and progression to AIDS in women and men.* N Engl J Med, 2001. **344**(10): p. 720-5.

45. Dong, K.L., A. Moodley, D.S. Kwon, M.S. Ghebremichael, M. Dong, N. Ismail, Z.M. Ndhlovu, J.M. Mabuka, D.M. Muema, K. Pretorius, N. Lin, B.D. Walker, and T. Ndung'u, *Detection and treatment of Fiebig stage I HIV-1 infection in young at-risk women in South Africa: a prospective cohort study.* Lancet HIV, 2018. **5**(1): p. e35-e44.

46. Ndung'u, T., K.L. Dong, D.S. Kwon, and B.D. Walker, *A FRESH approach: Combining basic science and social good.* Sci Immunol, 2018. **3**(27).

47. Borrow, P., H. Lewicki, X. Wei, M.S. Horwitz, N. Peffer, H. Meyers, J.A. Nelson, J.E. Gairin, B.H. Hahn, M.B. Oldstone, and G.M. Shaw, *Antiviral pressure exerted by HIV-1-specific cytotoxic T lymphocytes (CTLs) during primary infection demonstrated by rapid selection of CTL escape virus.* Nat Med, 1997. **3**(2): p. 205-11.

48. Goonetilleke, N., M.K. Liu, J.F. Salazar-Gonzalez, G. Ferrari, E. Giorgi, V.V. Ganusov, B.F. Keele, G.H. Learn, E.L. Turnbull, M.G. Salazar, K.J. Weinhold, S. Moore, C.C.C. B, N. Letvin, B.F. Haynes, M.S. Cohen, P. Hraber, T. Bhattacharya, P. Borrow, A.S. Perelson, B.H. Hahn, G.M. Shaw, B.T. Korber, and A.J. McMichael, *The first T cell response to transmitted/founder*

virus contributes to the control of acute viremia in HIV-1 infection. J Exp Med, 2009. **206**(6): p. 1253-72.

49. Goulder, P.J., A. Edwards, R.E. Phillips, and A.J. McMichael, *Identification of a novel HLA-A24-restricted cytotoxic T-lymphocyte epitope within HIV-1 Nef.* AIDS, 1997. **11**(15): p. 1883-4.

50. Koup, R.A. and D.D. Ho, *Shutting down HIV.* Nature, 1994. **370**(6489): p. 416.

51. Mamrosh, J.L., David-Fung E.S., Korber, B.T.M., Brander, C., Barouch, D., de Boer, R., Haynes, B.F., Klasse, P.J., Koup, R., Peters, B., Seaman, M., and Walker, B.D., *HIV Molecular Immunology 2022.* 2023, Los Alamos National Laboratory, Theoretical Biology and Biophysics, Los Alamos: New Mexico.

52. Goulder, P.J. and B.D. Walker, *HIV and HLA class I: an evolving relationship.* Immunity, 2012. **37**(3): p. 426-40.

53. Avettand-Fenoel, V., L. Hocqueloux, J. Ghosn, A. Cheret, P. Frange, A. Melard, J.P. Viard, and C. Rouzioux, *Total HIV-1 DNA, a Marker of Viral Reservoir Dynamics with Clinical Implications.* Clin Microbiol Rev, 2016. **29**(4): p. 859-80.

54. Rouzioux, C. and V. Avettand-Fenoel, *Total HIV DNA: a global marker of HIV persistence.* Retrovirology, 2018. **15**(1): p. 30.

55. Rouzioux, C., P. Tremeaux, and V. Avettand-Fenoel, *HIV DNA: a clinical marker of HIV reservoirs.* Curr Opin HIV AIDS, 2018. **13**(5): p. 389-394.

56. Bruner, K.M., A.J. Murray, R.A. Pollack, M.G. Soliman, S.B. Laskey, A.A. Capoferri, J. Lai, M.C. Strain, S.M. Lada, R. Hoh, Y.C. Ho, D.D. Richman, S.G. Deeks, J.D. Siliciano, and R.F. Siliciano, *Defective proviruses rapidly accumulate during acute HIV-1 infection.* Nat Med, 2016. **22**(9): p. 1043-9.

57. Chun, T.W., L. Stuyver, S.B. Mizell, L.A. Ehler, J.A. Mican, M. Baseler, A.L. Lloyd, M.A. Nowak, and A.S. Fauci, *Presence of an inducible HIV-1 latent reservoir during highly active antiretroviral therapy.* Proc Natl Acad Sci U S A, 1997. **94**(24): p. 13193-7.

58. Colby, D.J., L. Trautmann, S. Pinyakorn, L. Leyre, A. Pagliuzza, E. Kroon, M. Rolland, H. Takata, S. Buranapraditkun, J. Intasan, N. Chomchey, R. Muir, E.K. Haddad, S. Tovanabutra, S. Ubolyam, D.L. Bolton, B.A. Fullmer, R.J. Gorelick, L. Fox, T.A. Crowell, R. Trichavaroj, R. O'Connell, N. Chomont, J.H. Kim, N.L. Michael, M.L. Robb, N. Phanuphak, J. Ananworanich, and R.V.s. group, *Rapid HIV RNA rebound after antiretroviral treatment interruption in persons durably suppressed in Fiebig I acute HIV infection.* Nat Med, 2018. **24**(7): p. 923-926.

59. Whitney, J.B., A.L. Hill, S. Sanisetty, P. Penaloza-MacMaster, J. Liu, M. Shetty, L. Parenteau, C. Cabral, J. Shields, S. Blackmore, J.Y. Smith, A.L. Brinkman, L.E. Peter, S.I. Mathew, K.M. Smith, E.N. Borducchi, D.I. Rosenbloom, M.G. Lewis, J. Hattersley, B. Li, J. Hesselgesser, R. Geleziunas, M.L. Robb, J.H. Kim, N.L. Michael, and D.H. Barouch, *Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys.* Nature, 2014. **512**(7512): p. 74-7.

60. Gantner, P., S. Buranapraditkun, A. Pagliuzza, C. Dufour, M. Pardons, J.L. Mitchell, E. Kroon, C. Sacdalan, N. Tulmethakaan, S. Pinyakorn, M.L. Robb, N. Phanuphak, J. Ananworanich, D. Hsu, S. Vasan, L. Trautmann, R. Fromentin, and N. Chomont, *HIV rapidly targets a diverse pool of CD4(+) T cells to establish productive and latent infections.* Immunity, 2023. **56**(3): p. 653-668 e5.

61. Boulassel, M.R., N. Chomont, N.P. Pai, N. Gilmore, R.P. Sekaly, and J.P. Routy, *CD4 T cell nadir independently predicts the magnitude of the HIV reservoir after prolonged suppressive antiretroviral therapy.* J Clin Virol, 2012. **53**(1): p. 29-32.

62. Burgard, M., F. Boufassa, J.P. Viard, I. Garrigue, A. Ruffault, J. Izopet, A. Vabret, D. Descamps, P. Colson, J.M. Seigneurin, C. Rouzioux, and A.A.W. Group, *Factors influencing*

peripheral blood mononuclear cell-associated HIV-1 DNA level after long-term suppressive antiretroviral therapy in 236 patients. AIDS, 2009. **23**(16): p. 2165-71.

63. Fourati, S., P. Flandre, R. Calin, G. Carcelain, C. Soulie, S. Lambert-Niclot, A. Maiga, Z. Ait-Arkoub, R. Tubiana, M.A. Valantin, B. Autran, C. Katlama, V. Calvez, and A.G. Marcelin, *Factors associated with a low HIV reservoir in patients with prolonged suppressive antiretroviral therapy.* J Antimicrob Chemother, 2014. **69**(3): p. 753-6.

64. Pinzone, M.R., D.J. VanBelzen, S. Weissman, M.P. Bertuccio, L. Cannon, E. Venanzi-Rullo, S. Migueles, R.B. Jones, T. Mota, S.B. Joseph, K. Groen, A.O. Pasternak, W.T. Hwang, B. Sherman, A. Vourekas, G. Nunnari, and U. O'Doherty, *Longitudinal HIV sequencing reveals reservoir expression leading to decay which is obscured by clonal expansion.* Nat Commun, 2019. **10**(1): p. 728.

65. White, J.A., J.T. Kufera, N. Bachmann, W. Dai, F.R. Simonetti, C. Armstrong, J. Lai, S. Beg, J.D. Siliciano, and R.F. Siliciano, *Measuring the latent reservoir for HIV-1: Quantification bias in near full-length genome sequencing methods.* PLoS Pathog, 2022. **18**(9): p. e1010845.

66. Coffin, J.M., D.W. Wells, J.M. Zerbato, J.D. Kuruc, S. Guo, B.T. Luke, J.J. Eron, M. Bale, J. Spindler, F.R. Simonetti, S. Hill, M.F. Kearney, F. Maldarelli, X. Wu, J.W. Mellors, and S.H. Hughes, *Clones of infected cells arise early in HIV-infected individuals.* JCI Insight, 2019. **4**(12).

67. Cohn, L.B., I.T. Silva, T.Y. Oliveira, R.A. Rosales, E.H. Parrish, G.H. Learn, B.H. Hahn, J.L. Czartoski, M.J. McElrath, C. Lehmann, F. Klein, M. Caskey, B.D. Walker, J.D. Siliciano, R.F. Siliciano, M. Jankovic, and M.C. Nussenzweig, *HIV-1 integration landscape during latent and active infection.* Cell, 2015. **160**(3): p. 420-32.

68. Einkauf, K.B., G.Q. Lee, C. Gao, R. Sharaf, X. Sun, S. Hua, S.M. Chen, C. Jiang, X. Lian, F.Z. Chowdhury, E.S. Rosenberg, T.W. Chun, J.Z. Li, X.G. Yu, and M. Lichterfeld, *Intact HIV-1 proviruses accumulate at distinct chromosomal positions during prolonged antiretroviral therapy.* J Clin Invest, 2019. **129**(3): p. 988-998.

69. Ho, Y.C., L. Shan, N.N. Hosmane, J. Wang, S.B. Laskey, D.I. Rosenbloom, J. Lai, J.N. Blankson, J.D. Siliciano, and R.F. Siliciano, *Replication-competent noninduced proviruses in the latent reservoir increase barrier to HIV-1 cure.* Cell, 2013. **155**(3): p. 540-51.

70. Gounder, K., N. Padayachi, J.K. Mann, M. Radebe, M. Mokgoro, M. van der Stok, L. Mkhize, Z. Mncube, M. Jaggernath, T. Reddy, B.D. Walker, and T. Ndung'u, *High frequency of transmitted HIV-1 Gag HLA class I-driven immune escape variants but minimal immune selection over the first year of clade C infection.* PLoS One, 2015. **10**(3): p. e0119886.

71. WHO *Guideline on when to start antiretroviral therapy and on pre-exposure prophylaxis for HIV.* 2015. 76.

72. Lee, G.Q., N. Orlova-Fink, K. Einkauf, F.Z. Chowdhury, X. Sun, S. Harrington, H.H. Kuo, S. Hua, H.R. Chen, Z. Ouyang, K. Reddy, K. Dong, T. Ndung'u, B.D. Walker, E.S. Rosenberg, X.G. Yu, and M. Lichterfeld, *Clonal expansion of genome-intact HIV-1 in functionally polarized Th1 CD4+ T cells.* J Clin Invest, 2017. **127**(7): p. 2689-2696.

73. Lee, G.Q., *Chemistry and Bioinformatics Considerations in Using Next-Generation Sequencing Technologies to Inferring HIV Proviral DNA Genome-Intactness.* Viruses, 2021. **13**(9).