Title: Racial and Ethnic Disparities among Takotsubo Cardiomyopathy; A Nation-Wide Analysis

Abstract

Background: Takotsubo cardiomyopathy (TSC) is a transient left ventricular dysfunction often precipitated by stress, with unclear mechanisms ranging from coronary artery spasm to microvascular dysfunction. Its true incidence remains uncertain, estimated at 1 to 2% among suspected acute coronary syndrome cases. This condition can have a favorable prognosis but is associated with fatal complications. Evidence suggests that TSC disproportionately affects minority racial groups due to elevated stress levels, highlighting the need for healthcare equity.

Aim: Our study focused on exploring racial and ethnic disparities in-hospital outcomes and clinical characteristics for patients diagnosed with TSC.

Results: In our analysis of 32,785 TSC hospitalizations, racial and ethnic disparities were significant: 80.5% were White, 6.7% Black, and 5.8% Hispanic, with the remainder from other minorities. Black and Hispanic patients, averaging 63 years, were younger than the 67-year average, with most below the 50th percentile of income, the majority being Black patients (50%). Black patients experienced more extended hospital stays (mean 4.5 days, p < 0.001) compared to Whites (3.4 days, mean decrease 1.1 days, p = 0.002), and Native Americans had the longest stays (5.1 days, mean increase 1.75 days, p = 0.045). Asian patients faced the highest healthcare charges (mean $79,111, increase $23,542, p < 0.02). Myocardial infarction rates were highest in Native Americans (25.7%, OR 2.4, p = 0.035). Cardiogenic shock with 8.25% in Black patients (OR 1.34, p < 0.001) and 11% in Asians (OR 2, p < 0.001). Blacks had the highest risk of stroke (4.8%, OR 2.1, p = 0.003). Pulmonary embolism in Blacks was (OR 1.8, p-value <0.001). Disparities extended to inpatient mortality and cardiac arrest, with Asians and Blacks facing the highest risks (1.8% mortality in Asians, OR 1.97, p < 0.001; cardiac arrest in Blacks 3.9%, OR 2.3, p = 0.002) compared to Whites.

Conclusion: Our findings reveal significant racial and ethnic disparities in TSC hospitalizations, with minority groups experiencing adverse outcomes and higher healthcare costs. These disparities underscore the need for targeted healthcare interventions and further research to understand and address the barriers to equitable care for TSC patients.

Introduction

Takotsubo cardiomyopathy (TSC), first described in Japan in 1990, manifests as transient left ventricular dysfunction triggered by emotional or physical stress, resembling a "Takotsubo" octopus trap [1.] This condition predominantly affects postmenopausal women and mimics heart attack symptoms, though it occurs without coronary artery obstruction[1,2]. The underlying mechanisms of TSC are not fully understood. However, proposed theories include stress-induced increases in catecholamines, microvascular dysfunction, inflammation, estrogen deficiency, coronary vessel spasms, and myocardial stunning without obstructive coronary artery disease [3]. TSC can have a favorable prognosis but can be associated with fatal complications [2]. The real incidence of TSC remains unclear. It makes up for 1 to 2% of patients suspected of having acute coronary syndrome[3].

TSC cardiomyopathy demonstrates an evident correlation with physical stressors (36%) more than emotional stressors (27.7%) [6]. It's worth noting that minority race groups, who often face labor disparities and are more likely to engage in physically demanding jobs compared to white patients [15,16,] may be at increased risk of TSC due to heightened levels of physical stress [6].

While studies have elucidated various aspects of this syndrome, there remains a paucity of research investigating potential racial and ethnic disparities among affected patients. Understanding such disparities is essential for optimizing patient care and treatment strategies and addressing healthcare inequities.
Our study focused on exploring racial and ethnic disparities in-hospital outcomes for patients diagnosed with Takotsubo Cardiomyopathy (TSC). Through a systematic analysis that included patient demographics and hospitalization metrics such as length of stay and total charges, clinical outcomes, and mortality rates, we offer a comprehensive view of the disparities within the healthcare system.

Methods

Using the National Inpatient Sample (NIS) Database Registry, we analyzed data from recent years 2017 to 2020.

Database
The National Inpatient Sample (NIS) is part of the Healthcare Cost and Utilization Project (HCUP) and is maintained by the Agency for Healthcare Research and Quality (AHRQ) [4]. The NIS contains information on all inpatient stays (not individual patients) in 48 states plus the District of Columbia, representing approximately 98% of the United States population, excluding rehabilitation and long-term acute care hospitals [4]. Unweighted, it contains data from more than 7 million hospital stays each year, and weighted, it estimates more than 35 million hospitalizations nationally [4] observations in the NIS contain a primary diagnosis, up to 39 secondary diagnoses, and up to 25 procedure codes, depending on the year [4]. All discharge diagnoses were identified using the International Classification of Diseases, 10th edition (ICD-10) codes [5]. The AHRQ made these data available to the principal author via the HCUP.

Statistical Analysis
National estimates were obtained using the discharge-level weight variable (DISCWT) provided by the HCUP. Weighted data were used for all statistical analyses. Missing values were excluded from the analysis. Categorical variables were compared using the Chi-square test and are described using frequency with percentages. Continuous variables were compared using the student's t-test and are reported as mean (±SD) if their distribution was normal or compared

using the Mann–Whitney U test and reported as median (interquartile range [IQR]) if their distribution was skewed. Multivariate linear and logistic regression was used to compare hospital outcomes between groups, adjusting for potential confounders, which included age, Charlson comorbidity index, hospital bed size, teaching status, and insurance. Statistical analysis was performed using Stata version 18 (Stata Corp, College State, Texas). A p-value ≤ 0.05 was considered to indicate statistical significance for all analyses. Variables used in the regression model building were either selected from the dataset as provided variables or abstracted with the ICD-10 codes.

Inclusion criteria and study variables.
The study cohort consisted of all hospitalizations among individuals categorized as either white, Hispanic, Black, Asian, Native American Indian, and other races. Patients who had missing race or reported other race/ethnicity were included as others. We extracted the following independent variables: sociodemographic characteristics (including sex, age, mean annual income in zip code, and insurance type), hospital characteristics, comorbidities, Charlson comorbidity index, and outcomes (Tables 1 and 2). We used ICD-10 codes to identify principal/secondary diagnoses (Supplemental table). To investigate disparities, we studied baseline characteristics and outcomes for TSC hospitalizations among all races (Supplementary Table 1).

Results

Patient Demographics and Hospital Metrics:
Among 32,785 hospitalizations with a principal diagnosis of TSC, demographic distributions were as follows: White patients comprised 26,395 (80.5%), Black patients 2,155 (6.7%), Hispanic patients 1,865 (5.8%), Native American Indians 175 (0.5%), Asian patients 555 (1.7%), and others 725 (2.2%). The average age was 67, with 89.4% of patients being female. However, among Black and Hispanic patients, the average age was 63. The primary hospital location was in the Southern region for 50% of the Black patient population, 32.15% of White patients, and 43% of Hispanic patients, compared to 48% of Native American Indians and 61.2% of Asian patient populations in the Midwest. Most patients were below the 50th percentile of income: 50% of Black patients, 38% of Native American Indians, and 32% of Hispanic patients were in the 0-25th percentile, compared to 52% of White patients and 66% of Asian patients, which were above the 50th percentile (Table 1 and Figure 1).

Length of Stay (LOS) and Total Charges (TOTCHG):
The mean length of stay (LOS) among patients with TSC was 3.5 days. Black patients had a mean LOS of 4.5 days with a mean increase of 1.1 days (p-value <0.001, 95% CI 0.6 to 2.5), and Native American Indian patients had a mean LOS of 5.1 days with an increase of 1.75 days (p-value 0.045 95% CI 0.3-3.5), compared to LOS of 3.5 days for White patients with a mean decrease of 1.1 days (p-value 0.002, 95% CI -1.6 to -0.5). However, Asian patients had a mean LOS of 3.8 days with an increase of 0.4 days (p-value 0.17), and Hispanic patients had a mean LOS of 3.3 days (p-value 0.9), others with 3.5 LOS (p-value), which were not statistically significant. (Tables 1 and 3)

The mean total charge (TOTCHG) among patients with TSC was $55569. Black patients faced a mean TOTCHG of $65,635, with a mean increase of $12,019 (p = 0.02, 95% CI 1,785-22,252). Hispanic patients had a mean TOTCHG of $71,443, with a mean increase of $15,874 (p = 0.002, 95% CI 5,497-25,364). The highest mean TOTCHG was noted for Asian patients at $79,111, an increase of $23,542 (p < 0.02, 95% CI 3,377-42,339). Patients categorized as other races had a mean TOTCHG of $71,272, with an increase of $15,703 (p < 0.02, 95% CI 3,377-

35,939). Conversely, Native American Indians and White patients had mean TOTCHGs of $71,234 and $53,042, respectively. Notably, White patients experienced a decrease of $14,442 from the mean (p < 0.001, 95% CI -23,931 to -4,954). (Tables 1 and 3)

Myocardial Infarction and Cardiogenic Shock:
Myocardial infarction rates were 16.35% among White patients (OR 1.3, p-value 0.039, 95% CI 1.0 to 1.8), 25.7% among Native Americans (OR 2.4, p-value 0.035, 95% CI 1 to 5.4), compared to 12.5% among Black patients (OR 0.7, p-value 0.039, 95% CI 0.5 to 0.98). Hispanic and Asian patients had p-values of 0.896 and 0.453, respectively.

Cardiogenic shock was reported in 8.25% of Black patients (OR 1.34, p-value <0.001, 95% CI 1.16 to 1.56), 8.8% of Hispanic patients (OR 1.4, p-value <0.001, 95% CI 1.2 to 1.7), and 11% of Asian patients (OR 2, p-value <0.001, 95% CI 1.5 to 2.5), compared with Native American Indian patients at 8.59% (p-value 0.172), other races at 7.6% (p-value 0.121), and White patients at 6.3% (OR 0.7, p-value <0.001, 95% CI 0.6 to 0.8). Ventricular arrhythmia was only associated with the Black population at 5.8% (OR 1.2, p-value 0.041, 95% CI 1.1 to 1.3) while other races were not statistically significant.

Pulmonary Embolism and Stroke:
Stroke prevalence was 4.8% among Black patients (OR 2.1, p-value 0.003, 95% CI 1.2 to 3.4) and 4.14% among other races (p-value <0.001), with Native American Indian patients at 5.7% (p-value 0.218), Asian patients at 1.8% (p-value 0.776), and Hispanic patients at 2.68% (p-value 0.720) compared to 2.39% among White patients (OR 0.4, p-value 0.003, 95% CI 0.3 to 0.7). Pulmonary embolism was significant in 2.5% of Black patients (OR 1.8, p-value <0.001, 95% CI 1.4 to 2.4), other at 3.7% (OR 2.76, p-value <0.001, 95% CI 1.8-4.2), compared to 1.7% of Hispanic (p-value 0.227), 0.6% for Asian (p-value 0.094), 1% for Native American Indian (p-value 0.665), and 1.4% for White patients (OR 0.53, p-value <0.001, 95% CI 0.4-0.7).

Mortality and Cardiac Arrest:
Inpatient mortality was 1.8% for Asian patients (OR 1.97, p-value <0.001, 95% CI 1.5 to 2.5), 0.5% for Hispanic patients (OR 1.38, p-value <0.001, 95% CI 1.17 to 1.63), 4.14% for other races (OR 1.8, p-value <0.001, 95% CI 1.6 to 2.1), and 2% for Black patients (OR 1.5, p-value <0.001, 95% CI 1.3 to 1.7) compared to 1.5% for White patients (OR 0.7, p-value <0.001, 95% CI 0.6 to 0.8) and Native American Indians (p-value 0.465). Cardiac arrest was 3.9% in the Black population (OR 2.3, p-value 0.002, 95% CI 1.7 to 3.9), 2.1% for Hispanic (p-value 0.720), 3.6% for Asian (p-value 0.776), for Native American Indians (p-value 0.218), and for other (p-value 0.183) compared to 1.7% for White patients (OR 0.43, p-value 0.002, 95% CI 0.3 to 0.6). Outcome data for all races are presented in Tables 2 and 3.

Discussion

We analyzed the national inpatient sample and noted potential disparities in patient characteristics and clinical outcomes among different racial and ethnic groups hospitalized with Takotsubo cardiomyopathy (Figure 2).

We found that most TSC hospitalizations were among White patients, with Black, Hispanic, Native American Indians, and other minority groups making up a smaller portion of the total patient population. The average age was 67, and females accounted for 89% of the cases, aligning with existing research data [6]. Our analysis also revealed that Black and Hispanic patients were, on average, younger at hospitalization (age 63), suggesting potential disparities in disease manifestation or healthcare access. Notably, 50% of the Black population with TSC fell within the lowest income quartile (0-25th percentile) (Figure 1).

There were significant disparities in the length of hospital stays and hospital charges. Black and Native American patients experienced longer hospitalizations, a 2–3-day increase compared to the average. Asian patients encountered the highest total charges, while both Blacks and

Hispanics incurred higher costs than their White counterparts (Figure 3). These differences could reflect greater severity of disease presentation, delays in receiving care, or biases within healthcare facilities. The increased financial burden on Black and Hispanic patients highlights the urgent need for policy changes to ensure equitable healthcare costs.

Our findings indicated that Black, Asian, Hispanics, and Native American patients were associated with a higher prevalence of comorbidities (Figure 4). Native American (OR 2.4, p-value 0.03) and White patients (OR 1.36, p-value 0.039) had a higher risk of myocardial infarction during hospitalization compared to Blacks. Furthermore, cardiogenic shock showed a higher prevalence and risk in Black (OR 1.35 p-value <0.001) and Asian populations.

Black patients exhibited a higher risk of cardiogenic shock, even though they had a lower risk of myocardial infarction in this study. This phenomenon may be attributed to significant disparities in cardiac care for Black individuals, encompassing prevention, invasive treatments, and outcomes [7,8]. Notably, Black patients often face longer delays in receiving medications, undergoing revascularization, and obtaining surgical treatments for acute coronary syndrome, including extended door-to-balloon times [8]. Genetic factors, such as the presence of the APOL1 and ADRB2 16R alleles in Black populations, might also play a crucial role in prompting worse outcomes [9,10]. The etiology of cardiogenic shock appears to be multifactorial, suggesting a potential link between cardiogenic shock and other major cardiovascular complications prevalent among Black patients [11, 12, 13].

Pulmonary embolism was notably higher in Black patients (OR 1.8, p-value <0.001) and patients of other races (OR 2.3, p-value <0.001) compared to other populations. Stroke prevalence was the highest at 4.8% among Black hospitalized patients (OR 2.1, p-value 0.003) and 4.14% among other races (OR 1.7 p-value <0.001), significantly higher than in other races. In contrast, White patients showed no significant association (OR 0.4, p-value 0.003) with stroke. These outcomes suggest disparities may stem from underlying health conditions, differential access to preventative care, or variability in care standards.

Mortality Rates:
Disparities in mortality rates were evident, with the highest inpatient risk of mortality observed in Asian (OR 1.97, p <0.001) and other race populations (OR 1.8, p <0.001), followed by 2% of Black patients (OR 1.5, p <0.001). Black patients also demonstrated a notable association with cardiac arrests (OR 2.3, p-value 0.002) compared to White patients, who showed no significant mortality association (OR 0.6, p <0.001).

In summary, it was noted that the Black population had more adverse outcomes compared to other races, which is consistent with literature [14]. The heightened presence of comorbidities among Black individuals may be elucidated by the disparities in the labor market and employment [15,16,] which inadvertently expose them to more physically strenuous and hazardous occupations.

Additionally, there is a hypothesis that chronic medical and socioeconomic stressors, which disproportionately affect specific minority racial groups such as Black communities, may influence the severity of Takotsubo cardiomyopathy outcomes [17]. It is proposed that the worse outcomes observed in these populations could be partially explained by elevated levels of chronic stress, potentially leading to increased catecholamine production [1]. This hypothesis is supported by existing research indicating that the convergence of medical and socioeconomic

challenges is more prevalent in certain minority groups and may significantly contribute to the pathophysiology of Takotsubo cardiomyopathy [17,1].

This finding underscores the importance of addressing social determinants, such as occupational factors, in understanding and mitigating health disparities.

Limitations:
The study faced several limitations. The reliance on an administrative database dependent on ICD10-CM codes, where incorrect coding could influence findings, was a primary concern. The lack of randomization was addressed using univariate and multivariate analysis. The strength of our study lies in the power of the sample size; we utilized a large population database, thus improving the validity of our findings.

Conclusion

This study reveals insight into significant racial and ethnic disparities in Takotsubo Cardiomyopathy outcomes. Despite TSC's higher prevalence among White patients, Black, other races, and Hispanic patients face a greater risk of severe conditions like pulmonary embolism and higher mortality rates, especially among Black patients. These findings highlight the crucial impact of socioeconomic factors and healthcare access on patient outcomes.

Further research and interventions are needed to address these disparities, focusing on understanding and overcoming the socioeconomic and healthcare barriers that minority groups face.

References

1. Liauw, Liu, & Bastiany. (2023). Takotsubo cardiomyopathy. In *Biology of Women's Heart Health* (pp. 55-74). Springer International Publishing. https://doi.org/10.1007/978-3-031-39928-2_5
2. Akashi YJ, Goldstein DS, Barbaro G, Ueyama T. (2008). Takotsubo cardiomyopathy: A new form of acute, reversible heart failure. *Circulation*, 118(25), 2754-2762. https://doi.org/10.1161/CIRCULATIONAHA.108.767012
3. Ahmad SA, Brito D, Khalid N, et al. (2023). Takotsubo cardiomyopathy. In *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing. Available from: https://www.ncbi.nlm.nih.gov/books/NBK430798/
4. Healthcare Cost and Utilization Project (HCUP). (2022). NIS Database Documentation. Agency for Healthcare Research and Quality. https://www.hcup-us.ahrq.gov/db/nation/nis/nisdbdocumentation.jsp
5. World Health Organization. (2022). International classification of diseases (ICD). https://www.who.int/standards/classifications/classification-of-diseases. Accessed April 9, 2022.
6. Templin C, Ghadri JR, Diekmann J, Napp LC, Bataiosu DR, Jaguszewski M, et al. (2015). Clinical features and outcomes of Takotsubo (stress) cardiomyopathy. *The New England Journal of Medicine*, 373, 929-938. https://doi.org/10.1056/NEJMoa1406761
7. Edmund Anstey, D., Li, S., Thomas, L., Wang, T. Y., & Wiviott, S. D. (2016). Race and Sex Differences in Management and Outcomes of Patients After ST-Elevation and Non-ST-Elevation Myocardial Infarct: Results From the NCDR. *Clinical cardiology*, *39*(10), 585–595. https://doi.org/10.1002/clc.22570

8. Ya'qoub, L., et al. (2021). "Racial, Ethnic, and Sex Disparities in Patients with STEMI and Cardiogenic Shock." *JACC: Cardiovascular Interventions*, 14(6), 653-660. https://doi.org/10.1016/j.jcin.2021.01.003

9. Cornelissen, et al. (2021). "APOL1 Genetic Variants Are Associated with Increased Risk of Coronary Atherosclerotic Plaque Rupture in the Black Population." *Arteriosclerosis, Thrombosis, and Vascular Biology*, 41(7), 2201-2214. Published on 2021-07-01. https://doi.org/10.1161/ATVBAHA.120.315788

10. Cresci, S., et al. (2012). Adrenergic-pathway gene variants influence beta-blocker–related outcomes after acute coronary syndrome in a race-specific manner. *Journal of the American College of Cardiology, 60*(10), 898-907. https://doi.org/10.1016/j.jacc.2012.02.051

11. Javed Z, Sardar MR, Saeed W, Sarwar U, Williams BA, Aziz M, et al. (2021). Race, racism, and cardiovascular health: Applying a social determinants of health framework to racial/ethnic disparities in cardiovascular disease. *Circulation: Cardiovascular Quality and Outcomes*, 14, e007917. https://doi.org/10.1161/CIRCOUTCOMES.121.007917

12. Chioncel O, Mebazaa A, Harjola VP, Coats AJS, Piepoli MF, Crespo-Leiro MG, et al. (2020). Epidemiology, pathophysiology, and contemporary management of cardiogenic shock – A position statement from the Heart Failure Association of the European Society of Cardiology. *European Journal of Heart Failure*, 22(8), 1315-1341. https://doi.org/10.1002/ejhf.1922

13. Vyas A, Vedanthan R, Crown A, Tandon N, Fuster V. (2020). Cardiovascular disease burden and major adverse cardiac events in young black patients: A national analysis of 2 cohorts 10 years apart (2017 versus 2007). *Journal of the American Heart Association*, 9(19), e029895. https://doi.org/10.1161/JAHA.120.029895

14. Vriz O, Mushtaq AH, Elshaer AN, Shaik A, Landi I, Alzahrani T. (2023). Takotsubo syndrome in Black Americans: Insights from the national inpatient sample. *Texas Heart Institute Journal*, 50(5), e228055. https://doi.org/10.14503/THIJ-22-8055. PMID: 37853912; PMCID: PMC10658156.

15. Economic Policy Institute. (n.d.). Understanding black-white disparities in labor market outcomes requires models that account for persistent discrimination and unequal bargaining power. https://www.epi.org/publication/Black-white-disparities-labor-market-outcomes/

16. Andrasfay T, Goldman N. (2021). Physical work conditions and disparities in later life functioning: Potential pathways. *SSM - Population Health*, 13, 100990. https://doi.org/10.1016/j.ssmph.2021.100990

17. Zaghlol, et al. (2020). Racial differences in Takotsubo cardiomyopathy outcomes in a large nationwide sample. *ESC Heart Failure*, 7(3), 1056-1063. https://doi.org/10.1002/ehf2.12664

Authors

1. Jamal Christopher Perry (Corresponding Author)
   Email: jcperry908@bhmcny.org
   Affiliation: Brookdale University Hospital Medical Center

2. Oluwasegun Matthew Akinti

Email: drsegunakinti@bhmcny.org
Affiliation: Brookdale University Hospital Medical Center

3. Henry Alberto Becerra
   Email: habecerra@bhmcny.org
   Affiliation: Brookdale University Hospital Medical Center

4. Henry Osarumme Aiwuyo
   Email: haiwuyo@bhmcny.org
   Affiliation: Brookdale University Hospital Medical Center

5. Charles Oluwarotimi Poluyi
   Email: cpoluyi@bhmcny.org
   Affiliation: Brookdale University Hospital Medical Center

6. Abdullah Khan
   Email: abkhan@bhmcny.org
   Affiliation: Brookdale University Hospital Medical Center

Patient Characteristics

Race	White	Black	Hispanic	Asian	Native American	Other	Total	p-values
Patients n(%)	26395(82.8%)	2155(6.7%)	1865(5.8%)	555(1.7%)	175(0.5%)	725(2.2%)	32785	<0.001
Age (mean in years)	68	63	63	66	64	65	67	<0.001
Age ≥18 to <49 years	6.5%	15.6%	16.9%	14.0%	8.6%	10.3%	9.8%	<0.001
Age ≥49 to <59 years	18.3%	21.1%	21.5%	15.0%	28.6%	22.1%	17.0%	<0.001
Age ≥59 to <69 years	25.8%	25.3%	25.2%	19.0%	28.6%	27.6%	24.0%	<0.001
Age ≥69 years	49.5%	38.0%	36.5%	50.4%	34.3%	40.0%	49.0%	<0.001
Gender								
Male	16.4%	24.0%	20.8%	25.1%	17.7%	21.5%	10.6%	<0.001
Female	83.6%	76.1%	79.2%	74.9%	82.3%	78.5%	89.4%	<0.001
Hospital Region								
Northeast	20.2%	18.8%	19.0%	19.8%	5.7%	36.5%	21.0%	<0.001
Midwest	25.5%	22.0%	5.6%	11.7%	28.6%	9.7%	24.4%	<0.001
South	32.2%	49.9%	31.9%	7.2%	17.1%	29.7%	33.5%	<0.001
West	22.2%	9.3%	43.4%	61.2%	48.6%	24.1%	21.0%	<0.001
Median Household Income by ZIP Code								
0 to 25th percentile (low income)	20.3%	51.0%	32.9%	15.5%	38.2%	19.7%	25.2%	<0.001
26th to 50th percentile (median)	27.5%	20.3%	26.9%	17.3%	23.5%	19.7%	26.4%	<0.001
51st to 75th percentile	28.1%	18.0%	25.3%	20.9%	14.7%	28.2%	25.8%	<0.001
76th to 100th percentile (high income)	24.0%	10.0%	14.8%	46.4%	23.5%	32.4%	22.4%	<0.001
Insurance Type								
Medicare	64.7%	54.2%	50.4%	48.6%	57.6%	45.7%	64.4%	<0.001
Medicaid	6.9%	18.1%	20.9%	11.9%	12.1%	14.3%	11.1%	<0.001
Private Insurance	25.8%	21.0%	23.7%	33.9%	24.2%	33.6%	21.5%	<0.001
Uninsured	2.6%	6.1%	5.0%	5.5%	6.1%	6.4%	2.8%	<0.001
Hospital Location								
Urban	94.0%	96.3%	98.7%	98.1%	86.7%	98.6%	93.7%	<0.001
Rural	6.0%	3.7%	1.3%	0.9%	14.3%	1.4%	6.3%	<0.001
Type of Institution								
Teaching	74.2%	80.5%	82.0%	78.4%	77.1%	86.2%	76.1%	<0.001
Non-Teaching	25.8%	19.5%	18.0%	21.6%	22.9%	13.8%	23.9%	<0.001
Hospital Bed Size								
Small	16.8%	16.0%	16.9%	16.2%	31.4%	15.2%	17.7%	<0.001
Medium	28.3%	28.5%	30.6%	27.0%	17.1%	37.2%	27.5%	<0.001
Large	54.9%	55.5%	52.6%	56.8%	51.4%	47.6%	54.8%	<0.001
Hospital Metrics								
Length of Stay (days)	3.5	4.5	3.3	3.8	5.1	3.5	3.5	<0.001
Total Charge (USD)	53042	65635	71443	79111	71234	71272	55569	<0.001

Comorbidities	White	Black	Hispanic	Asian	Native American	Other	p-value
Congestive Heart Failure	34%	36%	29%	36%	37%	32%	<0.001
Mood Disorders	2.4%	4.9%	2.7%	1.8%	5.7%	4.1%	<0.001
Smoking	45%	45%	28%	23%	29%	30%	<0.001
Chronic Kidney Disease	10%	17%	9%	17%	11%	7%	<0.001
Hyperlipidemia	52.3%	47.6%	49.0%	53.2%	34.2%	49.0%	<0.001
Obesity	11.8%	14.1%	14.8%	5.9%	15.7%	9.6%	<0.001
Hypertension	38.3%	40.8%	43.7%	38.7%	31.4%	37.9%	<0.001
Diabetes Mellitus	19.0%	27.6%	32.7%	31.5%	31.4%	24.8%	<0.001
Charleson Comorbidity Index:							
0	25.1%	20.7%	25.1%	26.3%	26.3%	31.8%	<0.001
1	28.6%	26.9%	29.7%	27.8%	27.8%	25.0%	<0.001
2	21.4%	20.5%	20.9%	14.3%	14.3%	23.9%	<0.001
≥3 Comorbidities (Charlson)	24.9%	31.8%	24.2%	31.6%	31.6%	19.3%	<0.001

Outcomes	White	Black	Hispanic	Asian	Native American	Other	p-value
Pulmonary embolism	1.4%	2.5%	1.7%	0.6%	1.0%	3.7%	<0.001
Mortality	1.6%	2.0%	0.5%	1.8%	2.9%	4.1%	<0.001
Stroke	2.3%	4.9%	2.7%	1.8%	5.7%	4.1%	<0.001
Ventricular Arrhythmias	4.6%	5.8%	4.3%	2.7%	8.5%	7.6%	<0.001
Cardiogenic shock	6.3%	8.3%	8.8%	11.9%	8.6%	7.7%	<0.001
Myocardial Infarction	16.4%	12.5%	16.0%	18.9%	25.7%	17.9%	<0.001
Cardiac arrest	1.7%	3.9%	2.1%	3.6%	5.7%	3.5%	<0.001

Multivariate Logistic Regression

Outcomes	White		Black		Hispanic		Asian		Native American indian		Other	
	OR[a] (95% CI)	Pv	OR (95% CI)	Pv	OR[a] (95% CI)	Pv	OR[a] (95% CI)	Pv	OR[a] (95% CI)	P-value	OR[a] (95% CI)	Pv
Pulmonary embolism	0.53(0.4-0.5)	<0.001	1.9(1.4-2.4)	<0.001	1.2	0.2	0.4	0.09	0.7	0.6	2.76(1.8-4.2)	<0.00
Mortality	0.67(0.6-0.8	<0.001	1.5(1.2-1.7)	<0.001	1.4(1.1-1.6)	<0.001	1.97(1.5-2.5)	<0.001	1.2	0.4	1.8(1.4-2.3)	<0.00
Stroke	0.4(0.3-0.7)	0.003	2.1(1.3-3.4)	0.003	1.1	0.7	0.7	0.7	2.4	0.2	1.7	0.2
Ventricular Arrhythmias	0.8(0.7-0.9)	0.04	1.2(1-1.3)	0.046	0.9	0.7	1	0.8	1.2	0.4	1.3(1-1.7)	0.04
Cardiogenic shock	0.7(0.6-0.8)	<0.001	1.35(1.2-1.5)	<0.001	1.46(1.2-1.7)	<0.001	2.(1.6-2.5)	<0.001	1.4	0.17	1.2	0.12
Myocardial infarction	1.36(1- 1.8)	0.039	0.7(0.5-0.98)	0.039	0.9809	0.89	1.1	0.4	2.4(1-5.4)	0.035	1.11801	0.6
Cardiac arrest	0.43(0.2 - 0.7)	0.002	2.3(1.3-3.9)	0.002	1.2	0.56	2.1	0.7	3	0.2	2	0.18

Multivariate Linear Regression

	White		Black		Hispanic		Asian		Native American indian		Other	
Length of Stay	-1.1(-1.6 to -0.5)	<0.001	1.1(0.48-1.6)	<0.001	-0.03(-0.4-0.36)	0.9	0.43(1.8-4.2)	0.17	1.75(-0.0-3.5)	0.045	0.4.(0.6-0.7)	0.9
Total Charge	-14442(-23931-4954)	<0.001	12019(1785-22252)	<0.02	15431(5497-25364)	0.002	22858(3377-42339)	0.02	15629(11045-42303)	0.251	18950(1959-35939)	0.02

[a]OR =Odds Ratio. Pv= p-value

Comorbidities Among Racial Groups

- Congestive Heart Failure
- Mood Disorders
- Smoking
- Chronic Kidney Disease
- Hyperlipidemia
- Obesity
- Hypertension
- Diabetes Mellitus
- ≥3 Comorbidities (Charlson)