Potentially Modifiable Dementia Risk Factors in Canada: An Analysis of Canadian Longitudinal Study on Aging with a Multi-Country Comparison

Surim Son MSc[1,2]; Mark Speechley PhD[1]; Guangyong Zou PhD[1,3]; Miia Kivipelto MD PhD[4,5,6,7] ; Francesca Mangialasche MD PhD[4,6]; Howard Feldman MD[8,9]; Howard Chertkow MD[10,11]; Sylvie Belleville PhD[12,13]; Haakon Nygaard MD PhD[14]; Vladimir Hachinski MD DSc[1,3,15]; Pieruccini- Faria F PhD[2,16]; Manuel Montero-Odasso MD PhD[1,2,16*]

[1] Department of Epidemiology and Biostatistics, Schulich School of Medicine & Dentistry, University of Western Ontario, 1465 Richmond Street, London, ON, Canada, N6G 2M1

[2] Gait and Brain Lab, Parkwood Institute, Lawson Health Research Institute, 550 Wellington Road, London, ON, Canada, N6C 0A7

[3] Robarts Research Institute, University of Western Ontario, 1151 Richmond St, London, ON, Canada, N6A 3K7

[4] Division of Clinical Geriatrics, Center for Alzheimer Research, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Karolinska Vägen 37 A, 171 64 Solna, Sweden

[5] Neuroepidemiology and Ageing Research Unit, School of Public Health, Imperial College London, St Dunstan's Road, London, United Kingdom, W6 8RP

[6] Theme Inflammation and Aging, Medical Unit Aging, Karolinska University Hospital, Hälsovägen 7, Huddinge, 141 86 Stockholm, Sweden

[7] Stockholms Sjukhem, Research and Development Unit, Mariebergsgatan 22, 112 19 Stockholm, Sweden

[8] Alzheimer's Disease Cooperative Study, University of California San Diego, 9500 Gilman Drive, La Jolla, CA, USA, 92037-0948

[9] Department of Neurosciences, University of California San Diego, 9375 Gilman Dr, La Jolla, CA, USA, 92161

[10] Kimel Centre for Brain Health and Wellness and Anne & Allan Bank Centre for Clinical Research Trials, Baycrest Hospital, 3560 Bathurst St, Toronto, ON, Canada, M6A 2E1

[11] Canadian Consortium on Neurodegeneration in Aging (CCNA), Administration, 3560 Bathurst St, Toronto, ON, Canada, M6A 2E1

[12] Research Center, Institut Universitaire de Gériatrie de Montréal, 4565 Queen Mary Rd, Montréal, QB, Canada, H3W 1W5

[13] Department of Psychology, Université de Montréal, 90 avenue Vincent d'Indy, Montréal, QB, Canada, H2V 2S9

[14] Division of Neurology and Djavad Mowafaghian Centre for Brain Health, University of British Columbia, 2215 Wesbrook Mall, Vancouver, BC, Canada, V6T 1Z3

[15] Department of Clinical Neurological Sciences, Schulich School of Medicine and Dentistry, University of Western Ontario, 1151 Richmond St, London, ON, Canada, N6A 3K7

[16] Department of Medicine, Division of Geriatric, Schulich School of Medicine and Dentistry, University of Western Ontario, 550 Wellington Rd, London, ON, Canada, N7C 0A7

[*]**Corresponding author:** Manuel Montero-Odasso, Departments of Medicine (Geriatrics) and Epidemiology and Biostatistics, Schulich School of Medicine & Dentistry, University of Western Ontario, 500 Wellington Road, London, ON, Canada, N6C 0A7

mmontero@uwo.ca

Abstract

BACKGROUND
Previous estimates suggested that up to 40% of dementia cases worldwide are associated with modifiable risk factors, however, these estimates are not known in Canada. Furthermore, sleep disturbance, an emerging factor, has not been incorporated into the life-course model of dementia prevention.

Objective
To estimate the population impact of 12 modifiable risk factors in Canadian adults including sleep disturbance, by sex and age groups, and to compare with other countries.

Design
Cross-sectional analysis of Canadian Longitudinal Study on Aging Baseline data

Setting
Community

Participants
30,097 adults aged 45 years and older.

Measurements
Prevalence and Population Attributable Fraction (PAF) of less education, hearing loss, traumatic brain injury, hypertension, excessive alcohol, obesity, smoking, depression, social isolation, physical inactivity, diabetes, and sleep disturbance.

Results
The risk factors with the largest PAF were later life physical inactivity (10.2%; 95% CI, 6.8% to 13%), midlife hearing loss (6.5%; 3.7% to 9.3%), midlife obesity (6.4; 4.1% to 7.7%), and midlife hypertension (6.2%; 2.7% to 9.3%). The PAF of later life sleep disturbance was 3.0% (95% CI, 1.8% to 3.8%). The 12 risk factors accounted for 51.9% (32.2% to 68.0%) of dementia among men and 52.4% (32.5% to 68.7%) among women. Overall, the combined PAF of all risk factors was 49.2% (31.1% to 64.9%), and it increased with age.

Conclusion
Nearly up to 50% of dementia cases in Canada could be prevented by modifying the 12 risk factors across the lifespan. Canadian risk reduction strategies should prioritize targeting physical inactivity, hearing loss, obesity, and hypertension.

Key Words
Dementia, Prevention, Risk Reduction, Lifestyle, CLSA

INTRODUCTION   4193/5500 words

With rapid global population aging, the number of individuals living with dementia worldwide is expected to triple, from 57 million to 152 million, by 2050.[1] In Canada, dementia prevalence is projected to increase by 66% by 2031.[2] Dementia is a multifactorial syndrome that results from multiple pathologies including those that cause neurodegeneration as well as vascular, metabolic and inflammatory processes that are associated with potentially modifiable risk factors.[3,4] Lifestyle interventions offer a promising non-pharmacological approach to reducing dementia burden by tempering modifiable risk factors. Risk reduction can potentially be reached through individual and public health approaches, which could complement emerging disease-modifying treatments directed at the pathological processes.[4]

The 2020 Lancet Commission Report on Dementia Prevention, Intervention, and Care[5] indicated that up to 40% of dementia cases worldwide are attributable to 12 modifiable factors comprising health behaviours, illnesses, and environmental exposures across the lifespan, known as the life course model of dementia prevention. This estimation is based on the weighted population attributable fraction (PAF), which quantifies the contribution of a given risk factor to dementia by combining both prevalence and the association between risk factor and disease, such as risk ratio, while adjusting for intercorrelation among risk factors.

Following the Lancet series,[5,6] the population impact of dementia risk factors has been estimated in other countries with differences in risk factor profiles.[7-12] Given that the same analytical approach and risk ratio were used to estimate PAF in all the studies, the differences in PAF across countries are mainly driven by differences in risk factor prevalence. For instance, less education is a greater contributing factor to dementia than social isolation in low-income countries as they tend to have stronger social ties and social support but have limited access to education compared to high-income countries.[9] Despite being one of the high-income countries with a universal health care system, Canada

now more than ever needs dementia risk reduction strategies, as it is reaching the super-aged country status, where it has ≥ 20% of its population composed of older adults.[13] However, no studies have estimated the population impact of the life-course model of modifiable risk factors for dementia in Canada, which is crucial for guiding evidence-based decision-making in the development of reduction strategies.

Besides the 12 risk factors included in the life course model of dementia prevention, the Lancet Commission Reports also identified sleep and diet as emerging risk factors.[5,6] Given the number of intervention trials delivering sleep and diet interventions to improve cognition, it is timely to quantify their population impact.[14-16] Estimating the population impact of diet is challenging due to the difficulty in measuring the diet pattern. For sleep, a recent study estimated that 5.7% of dementia cases are attributable to unhealthy sleep duration, using UK Biobank data.[17] However, the population impact of a sleep disturbance encompassing other conditions has not been estimated nor included in the life course model of dementia prevention.

Although air pollution is included in the life-course model[5], focusing only on lifestyle risk factors, rather than including an environmental risk factor, would be more beneficial in guiding the dementia risk reduction program.

Therefore, to provide evidence to help inform future lifestyle interventions in Canada, we aimed to estimate the prevalence and potential population impact of modifiable risk factors, including sleep disturbance, in Canadian adults using the largest Canadian population cohort study, and to compare the contribution of modifiable risk factors with other countries. To further provide evidence for tailoring prevention strategies, we aimed to estimate these measures by age groups and sex. Since the PAF of dementia risk factors has not been estimated in Canada, this study is much needed to tailor and develop Canada's dementia risk reduction program.

METHODS

This was a cross-sectional analysis of baseline data from the Canadian Longitudinal Study on Aging (CLSA) Comprehensive cohort. We followed STROBE guidelines for a cross-sectional study and conducted analyses using the *survey* and *psy* package in R Version 4.2.0.

Data source

CLSA is a prospective cohort study following 51,388 Canadians aged 45-85 years at recruitment for 20 years.[18] Participants were recruited to either Tracking or Comprehensive cohorts, which differed in sampling methods and data availability, as described elsewhere and in Figure S1 in Appendix.[18] The comprehensive cohort was used as it included participants who had undergone detailed and comprehensive face-to-face assessments that included a neuro-cognitive battery, sensory-vision, audiometry, proprioception testing as well as physical and mobility assessments. Individuals with a diagnosis of cognitive impairment and or dementia, full-time members of the armed forces, residents of First Nations reserves, territories, or long-term care institutions (only those that provide 24 hours of nurse care), and those who could not respond in English or French at recruitment were excluded.[18] Baseline data were collected from 2012 to 2015.[18]

Risk factors

A total of 12 risk factors were identified using operational definitions from the Lancet report[5] and 2 initial PAF estimation studies for dementia.[19,20] Less education was defined as having less than secondary school graduation. Hearing loss was derived from an average hearing level of >25 dB at 500, 1000, 2000 and 4000 Hz in the better ear.[21] Traumatic brain injury was defined as having at least one head injury caused by a vehicular crash, fall, or sports-related activities that resulted in losing consciousness. An average systolic blood pressure ≥140 mmHg across six seated measurements excluding the first reading or self-reported diagnosis was used to define hypertension. The number of drinks per week was converted to the unit of alcohol using the alcohol unit conversion formula by UK

National Health Service, $\frac{Strength\ (ABV\%) \times Volume\ (mL)}{1,000}$.[22] Based on the converted unit of alcohol, >21

units was used to indicate excessive alcohol use. BMI of $\geq$30 kg/m^2 was used to categorize obesity.

Daily or occasional cigarette smoking in the past 30 days was used to represent current cigarette

smoking. Depression was categorized based on the self-reported diagnosis of clinical depression. Social

isolation was defined as having less than one social contact within a month with family, friends, or

neighbours. The level of physical activity was measured using the Physical Activity Scale for the

Elderly (PASE) questionnaire. The total hours of physical activity per week were computed using the

lowest point of each frequency and duration category with the exception that the mid-point was used

for the lowest frequency category. Based on the estimated minutes of moderate-to-vigorous physical

activity, physical inactivity was defined as <150 minutes. A self-report diagnosis of type 2 diabetes by a

physician was used for diabetes. We characterized sleep disturbance based on the definition used in the

systematic review[23] from which we used their reported relative risk for PAF calculation. Sleep

disturbance was broadly defined to encompass poor sleep quality, daytime sleepiness, insomnia

symptoms, obstructive sleep apnea symptoms and restless leg syndrome symptoms based on self-

reported questionnaires on sleep quality and behaviours. More details on sleep disturbance

classification are provided in Table S2 Appendix.

Statistical Analysis

Demographic characteristics were summarized using mean and standard deviation or frequency and

percentage, where appropriate. To build the life-course model, the prevalence of midlife and later life

risk factors was calculated for the age group 45-64 and 65-85, respectively. Early life risk factors were

calculated for all age groups. The prevalence estimates were weighted with inflation weights to account

for differences in selection probabilities.[24]

As described elsewhere,[9] PAF of each risk factor was calculated using Levin's formula and risk factor

overlap was adjusted by applying Norton's formula[20], which involves weighting by communalities from

principal component analysis (see Table S3 in Appendix for detailed description). Relative risk was taken from the Lancet report[5] for all risk factors except sleep disturbance, which was derived from a recent meta-analysis of 18 longitudinal studies with an average 9.5 years of follow up assessing the association between sleep disturbance and dementia.[23]

To compare our results with global estimates[5] and other countries,[7-11] we obtained the prevalence and PAF from other studies that utilized the Lancet[5] approach. The life-course model obtained from CLSA was qualitatively compared to the global estimate[5] and other countries – USA,[7] India,[9] Latin America,[9] China,[9] New Zealand,[8] Brazil,[11] Australia,[10] and Denmark.[12] A total of 9 risk factors were available across all the studied countries – less education, hearing loss, hypertension, obesity, smoking, depression, social isolation, physical inactivity, and diabetes – and were measured using similar definitions across studies.

To further explore differences in risk factor distribution by age groups and sex, analyses were stratified by 4 age groups (45-54, 55-64, 65-74, and 75-85) and sex and chi-square tests were conducted. A sensitivity analysis compared changes in prevalence and PAF with different risk factor definitions. Depression was re-operationalized as a 10-item Center for Epidemiologic Studies Depression Scale score of ≥ 10 to reflect depressive symptom. Excessive alcohol use was re-defined using Canadian guidance, which is drinking ≥ 7 standard drinks per week.[25] A five-point Steptoe social isolation index was derived to incorporate cohabitation, social contact and participation into social isolation with a cut-off of ≥ 3.

RESULTS

A total of 30,097 participants were included (Table 1). The mean age was 59.7 years (SD 10.3) and 52% were female. The majority were white (94%), living in urban areas (90%), and 74% were married.

Prevalence

The most prevalent risk factor was later life physical inactivity (83%) (Table 2), followed by later life sleep disturbance (40%), midlife obesity (31%), midlife hypertension (30%), and midlife hearing loss (21%). The least common risk factors were midlife smoking (6.2%) and social isolation (1.6%) in later life. The prevalence of physical inactivity was much higher than in other countries, which ranged from 15.3% in India to 82% in Australia, as shown in Table 3. Smoking and social isolation were noticeably less prevalent than in other countries.

PAF

The weighted individual and combined PAF of all risk factors in Canada are presented in Table 2 and Figure 1. Almost 50% of dementia cases in Canada were attributed to 12 risk factors (PAF 49.2%, 95% CI 31.1% to 64.9%) – nearly 12 points more than the worldwide estimate of 37.4% (Table 3 and Figure S2 in Appendix). The PAF of the nine risk factors available in other studied countries ranged between 33.2% and 55.9%. When using these nine risk factors, the combined PAF in Canada (41%) was higher than in USA (36%), Denmark (33.2%) and Australia (35.5%) but lower than in Latin America (55.9%). The combined PAF of the nine risk factors in Canada was similar to New Zealand, India, China, and Brazil, with a range from 39.5% and 42.1% (Table 3).

Of the 12 risk factors, later life physical inactivity had the largest weighted PAF in Canada, contributing 10.2% (95% CI 6.8% to 13.0%). The PAF of midlife hearing loss, obesity, and hypertension in Canada were 6.5%, 6.4%, and 6.2% respectively (95% CI 3.7% to 9.2% for hearing loss, 95% CI 4.1% to 7.7% for obesity, and 95% CI 2.7% to 9.3% for hypertension). Later life

smoking, midlife excessive alcohol use, and later life social isolation were associated with less than 2% of PAF in Canada. Compared to other countries, Canadian weighted PAFs were smaller for later life smoking and social isolation, while larger for later life physical inactivity, midlife traumatic brain injury and excessive alcohol use (Table 3 and Figure S2 in Appendix). The weighted PAFs for early life less education, midlife hypertension, and later life depression in Canada were larger than in other high-income countries, whereas the weighted PAF for early life less education and midlife hypertension was smaller than low- and middle-income countries. Moreover, compared to low- and middle-income countries, Canada had a larger weighted PAF for midlife obesity.

By Sex and Age groups

Prevalence of the most common risk factors differed across sexes (Table 4 and Figure S4 in Appendix). Among women, 80.0% had physical inactivity and 20.8% had depression, as compared to 72.8% and 11.8% in men. The prevalence of less education and sleep disturbance was similar between sexes (p>0.05). The prevalence of hearing loss, traumatic brain injury, hypertension, excessive alcohol use, diabetes, and social isolation was higher in men than women (p<0.001).

Overall, the 12 risk factors accounted for 52.4% (95% CI 32.5% to 68.7%) of dementia among women and 51.9% (95% CI 32.2% to 68.0%) among men (Table 4 and Figure S5 in Appendix). The weighted PAF of hearing loss, traumatic brain injury, and excessive alcohol use was higher in men, whereas the weighted PAF of depression and physical inactivity was higher in women.

Prevalence and weighted PAF also varied across age groups (Table 4 and Figure S6 in Appendix). From age 45-54 to 75-85 years, prevalence increased from 13.5% to 81.2% for hearing loss, 7.2% to 24.7% for less education, 21.5% to 59.8% for hypertension, 71.6% to 87.8% for physical inactivity, and 4.3% to 12.1% for diabetes. The highest prevalence of traumatic brain injury (15.3%) and smoking (13.7%) was observed at age 45-54 years, which declined after age 55 years and dropped to 8.9% and 3.9% at

age 75-85 years, respectively. Depression prevalence increased from 16.4% in age 45-54 to 20.6% in 55-64 years, then decreased to 9.0% in 75-85 years.

The combined PAF of all 12 risk factors increased with age (Figure S5 in Appendix). The combined PAF were 49.1% (95% CI 30.5% to 65.5%) in age 45-54 years, 51.1% (95% CI 31.4% to 67.6%) in 55-64 years, 63.4% (95% CI 40.7% to 79.4%) in 65-74 years, and 68.2% (95% CI 46.2% to 83.9%) in 75-85 years. Physical inactivity had the highest PAF among ages 45-54 years (10.2%, 95% CI 6.6% to 13.2%) and 55-64 years (8.8%, 95% CI 5.7% to 11.4%), while hearing loss had the largest PAF among ages 65-74 years (14.1%, 95% CI 9.0% to 17.3%) and 75-85 years (19.0%, 95% CI 13.4% to 21.6%). The estimated communality and unadjusted PAFs by age groups and sex are shown in Table S3-S4 in Appendix.

Sensitivity Analyses

Operationalizing risk factors with different definitions increased the prevalence and PAF, and also changed the rank of risk factors with the largest PAF (Table S5-S7 in Appendix). The prevalence increased to 16% for depression, 29% for excessive alcohol use, and 42% for social isolation. The largest change was observed with social isolation, and its prevalence increased from 1.6% to 42%. The combined PAF increased to 53.2% (95% CI 33.9% to 68.9%). Later life social isolation became the risk factor with the second largest PAF. In older age groups, social isolation had larger PAF than obesity.

DISCUSSION

Our results suggest that up to 49% of dementia cases in Canada could be delayed or prevented by modifying 12 risk factors including sleep disturbance. To our knowledge, this is the first study estimating the potential population impact of 12 modifiable risk factors for dementia in this country and the first to include sleep disturbance in a life-course model with participants as young as 45 years. The predominant contributors were later life physical inactivity, midlife hearing loss, midlife obesity, and midlife hypertension. In contrast, later life smoking, midlife excessive alcohol use, and later life social isolation had substantially less contribution. Finally, we observed a similar prevention potential in men than women.

Over 80% of Canadians were not meeting physical activity guidelines, and nearly 40% of Canadians had impaired sleeping. Furthermore, 1 in 3 Canadians were obese or had hypertension, and 1 in 5 Canadians showed hearing loss. The estimated prevalence of obesity, physical inactivity, diabetes, and sleep disturbance was similar to previous estimates but our results showed a higher prevalence of less education and hypertension, and a lower prevalence of smoking and hearing loss.[26,27]

Canada's national dementia strategy includes the promotion of healthy lifestyle to reduce dementia risk, such as initiatives and programs to prevent and manage cardiovascular diseases and promote physical activity.[26,28] This high prevalence of physical inactivity is found despite sustained national efforts to promote physical activity. It highlights the need to advocate and implement moderate-to-vigorous exercise programs for older adults to help them meet physical activity targets. For instance, aerobic and resistance exercises have been shown to improve cognition in older adults and can be delivered in exercise programs.[29,30] More innovative approaches which provide paid-for programs in high-trafficked community institutions such as the YMCAs or online training done in the home environment are worth exploration.

There has been extensive development of a national strategic plan for hypertension prevention and control in Canada since 1990s.[31] However, the prevalence of vascular risk factors, hypertension and obesity, we observed were still relatively higher than other risk factors. While a low prevalence of diabetes may reflect the national strategic plans, such as sodium reduction strategy[31], it could be also due to not considering blood glucose levels in its definition. The high prevalence of hypertension and obesity underscores the opportunity to improve both vascular and cognitive health.

Despite an unclear underlying mechanism between hearing loss and dementia, a meta-analysis of 8 longitudinal studies found a 19% lower risk of cognitive decline among individuals with corrected versus uncorrected hearing loss (HR 0.81, 95% CI 0.76-0.87).[32] These findings suggest that dementia prevention programs and primary health care providers should consider encouraging hearing tests and hearing aid use. Compared to other lifestyle strategies, addressing hearing loss faces challenges related to stigma and affordability of hearing aids.[33] Given its high PAF and associated barriers, there is a pressing need to enhance hearing health care besides promoting patient education.

Sleep disturbance had a substantial contribution to dementia and the high prevalence we found is noteworthy because it can be translated to large potential population health benefits were it to be successfully addressed. Non-pharmacological interventions for sleep disturbance or disorder include sleep hygiene education, physical activity, and in some cases, bright light therapy,[34] however their effectiveness in preventing cognitive impairment requires further study. Periodically screening sleep impairment in practice will help identify and intervene undiagnosed sleep impairment.

Our age-specific analysis indicated that up to 49% of dementia cases could be prevented if we can modify all 12 risk factors as early as at age 45, and the potential for dementia prevention increased to 69% at age 75. Interestingly, the prevalence of physical inactivity, obesity, hypertension, and sleep disturbance was already high at age 45-54 and all the predominant risk factors identified in the life-course model, except physical inactivity, were from midlife. These results support the recommendation

of commencing risk reduction interventions at midlife or earlier, before developing cognitive impairment.[35].

Our age-specific analysis results align with a study from Chile, which calculated the PAF of dementia risk factors, stratified by sex and 2 age groups (45-64 and ≥65). [36] Similarly, we found that physical inactivity, less education, and risk factors related to chronic diseases, such as hearing loss, hypertension, and diabetes, gradually increase with age. The observed decreasing trend for traumatic brain injury, smoking, and excessive alcohol use may imply a decline in risk-taking with aging.[37]

As with other studies ascertaining sex-specific estimates,[36,38,39] the prevalence of traumatic brain injury, hearing loss and excessive alcohol use were higher in men while depression and physical inactivity were more prevalent in women. Contrary to our results, sex differences in smoking were observed in previous studies.[36,38,39] Although we obtained a similar potential for dementia prevention in men and women, the observed difference in risk factor profile is important to inform public policy to focus on addressing early risk behaviours, particularly in men.

Importantly, the difference in the risk factor profile we observed by age groups and sex suggests the importance of tailoring dementia prevention programs and strategies within Canada. For instance, dementia prevention programs for middle-aged adults should include education on risk-taking behaviours, while the focus could shift to better management of hearing loss, hypertension, and diabetes in older adults. The programs for older adults should prioritize delivering group exercise or recreational programs. The dementia prevention efforts could be further enhanced by focusing on addressing depression in women and hearing loss in men.

The combined PAF of the nine risk factors that are available across all studies was 33.2% globally and ranged between 33.2% and 55.9% among studied countries. The proportion of potentially preventable dementia cases attributable to these nine risk factors in Canada was similar to studies from low- and

middle-income countries, but higher than those from other high-income countries. Latin America reported the highest PAF, using data from 6 Latin American countries.[9]

The risk factor with the highest PAF was less education in middle- and low-income countries, while it was physical inactivity and obesity in high-income countries, suggesting a sociocultural gradient in dementia risk factors. Interestingly, risk factor profiles varied even among high-income countries, highlighting the uniqueness of the Canadian population which could inform policymakers in Canada. For instance, despite cultural similarities between USA and Canada, the prevalence of hearing loss, depression, and alcohol use was 2 to 4 times higher in Canada while USA had 1.5 times or more higher prevalence of obesity, diabetes, hypertension, and social isolation.[7] Despite geographical distance, the risk factor profile of Canada was fairly similar to Australia and Denmark. This underpins the importance of identifying population-specific risk factor profiles to inform researchers and policymakers on developing dementia risk reduction strategies. International differences in risk factor prevalence could also reflect healthcare and policy contexts. For instance, the relatively lower prevalence of less education and smoking in high-income countries might be explained by the implementation of governmental compulsory education[40] and tobacco control policies.[41]

The primary strength of our study is the use of CLSA, the largest and well-characterized cohort study in Canada. Its large sample allowed us to explore how risk factor distributions differ from midlife to later life. Our study is the first to show dementia risk factor distributions across 4 age groups beginning at age 45.

There are several limitations. First, some risk factors were defined based on self-reported data and misclassification of risk factors may impact the estimated PAF. For example, depression and sleep disturbance were classified based on self-reported measures, although using valid instruments. Physical inactivity excluded light activities. However, because our sensitivity analyses produced higher estimates of PAF than our main analysis, our estimates probably underestimate PAF. Second, our

sample was composed by largely highly educated White Canadians, living in urban settings, which limits generalizability. Third, Levin's formula assumes no confounding and it produces a biased estimate of PAF when adjusted relative risk is used. Considering that the adjusted relative risk for dementia risk factors tends to be less than the unadjusted relative risk, our obtained PAF may be underestimated.[42] For age and sex-specific PAF calculation, the relative risk used was from life course-specific measures and may not accurately represent other age groups or sexes. We also acknowledge that the interpretation of the PAF is based on assumptions of causality and the complete eradication of risk factors, which is unlikely to be achieved in real life. However, it can still inform public health intervention strategies on which risk factors to be prioritized. Given the multifactorial nature of dementia, our approach of adjusting for risk factor inter-relationship to combine PAF may not fully account for confounding, interactions among risk factors and its multifactorial nature. Although diet is recognized as an important dementia risk factor[43] it was not included due to the complexities of analyzing diet. Lastly, while most of the risk factors had similar definitions, caution is needed in comparing the prevalence and PAF across studies.

Our results have critical implications for healthcare professionals, increasing their awareness of dementia risk factors and promoting the screening of prevalent risk factors to facilitate patient's risk reduction strategies from midlife. Furthermore, the findings could help researchers and policymakers design optimal prevention programs to ease the projected burden of dementia. Although today the β-amyloid-targeting therapies in early symptomatic Alzheimer's disease started to show promising results,[44,45] there is emerging evidence that the beneficial effect of lifestyle intervention are independent of β-amyloid pathology.[46] The FINGER trial showed that a multidomain intervention including exercise, nutritional counselling, cognitive training, social stimulation, and cardiometabolic management, improved cognition and reduced cognitive decline risk in 1,260 community dwellers at risk of dementia.[47] The FINGER model is being adapted and tested globally within the World-Wide

FINGERS network of multidomain trials for dementia risk reduction[14], with Canada being at the forefront of such efforts with the CAN-THUMBS UP project.[16] Findings from the present study can inform the refinement and development of the CAN-THUMBS UP multidomain interventions. Additionally, combining lifestyle intervention with disease-modifying therapy will offer an even better opportunity to effectively and precisely manage and prevent dementia based on patient's risk profiles, as demonstrated in diabetes management,[48] and multidomain lifestyle-based interventions combined with repurposed drugs (metformin) are now being tested in Europe.[49] Importantly, not all older adults at risk of dementia will be candidates for emerging disease-modifying therapy due to the therapies contraindications and limited generalizability of the current trial results, and therefore lifestyle interventions may be more applicable.[50] Analyses of the most prevalent combinations of risk factors and risk factor clustering would allow the development of impactful personalized prevention programs.

In conclusion, we determined that nearly up to 50% of dementia cases in Canada could be prevented by modifying 12 risk factors across the life span. The most prominent modifiable risk factors were later life physical inactivity, midlife obesity, midlife hypertension, and midlife hearing loss, whereas midlife excessive alcohol use, later life smoking, and later life social isolation had substantially less contribution.

ACKNOWLEDGEMENTS

Surim Son was awarded Alzheimer Society London and Middlesex Doctoral scholarship, Ontario Scholarship Award, and CCNA Trainee Synapse Challenge Funding.

FUNDING

This project was supported by Gait and Brain Health program via fundings of the Canadian Institute of Health and Research (MOP 211220, PJT 153100). This research was made possible using the data/biospecimens collected by the Canadian Longitudinal Study on Aging (CLSA). Funding for the CLSA is provided by the Government of Canada through the Canadian Institutes of Health Research (CIHR) under grant reference: LSA 94473 and the Canadian Foundation for Innovation as well as the following provinces, Newfoundland, Nova Scotia, Quebec, Ontario, Manitoba, Alberta, and British Columbia. This research has been conducted using the CLSA Baseline Comprehensive version 7.0, under Application Number 2109006, title "Potential modifiable risk factors for low cognition and dementia in Canadian Longitudinal Study of Aging", and was approved by the CLSA on January 28, 2022. The CLSA is led by Drs. Parminder Raina, Chrstina Wolfson, and Susan Kirkland.

The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of data, preparation, review, or approval of the manuscript and decision to submit the manuscript for publication.

REFERENCES

1. GBD Dementia Forecasting Collaborators. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. *Lancet Public Health*. Feb 2022;7(2):e105-e125. doi:10.1016/S2468-2667(21)00249-8

2. Alzheimer Society of C. Report summary Prevalence and monetary costs of dementia in Canada (2016): a report by the Alzheimer Society of Canada. *Health Promot Chronic Dis Prev Can*. Oct 2016;36(10):231-232. Note de synthese Prevalence et couts financiers des maladies cognitives au Canada : un rapport de la Societe Alzheimer du Canada (2016).

3. Baumgart M, Snyder HM, Carrillo MC, Fazio S, Kim H, Johns H. Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective. *Alzheimers Dement*. Jun 2015;11(6):718-26. doi:10.1016/j.jalz.2015.05.016

4. Kivipelto M, Mangialasche F, Ngandu T. Lifestyle interventions to prevent cognitive impairment, dementia and Alzheimer disease. *Nat Rev Neurol*. Nov 2018;14(11):653-666. doi:10.1038/s41582-018-0070-3

5. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. Aug 8 2020;396(10248):413-446. doi:10.1016/S0140-6736(20)30367-6

6. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *Lancet*. Dec 16 2017;390(10113):2673-2734. doi:10.1016/S0140-6736(17)31363-6

7. Lee M, Whitsel E, Avery C, et al. Variation in Population Attributable Fraction of Dementia Associated With Potentially Modifiable Risk Factors by Race and Ethnicity in the US. *JAMA Netw Open*. Jul 1 2022;5(7):e2219672. doi:10.1001/jamanetworkopen.2022.19672

8. Ma'u E, Cullum S, Cheung G, Livingston G, Mukadam N. Differences in the potential for dementia prevention between major ethnic groups within one country: A cross sectional analysis of population attributable fraction of potentially modifiable risk factors in New Zealand. *Lancet Reg Health West Pac*. Aug 2021;13:100191. doi:10.1016/j.lanwpc.2021.100191

9. Mukadam N, Sommerlad A, Huntley J, Livingston G. Population attributable fractions for risk factors for dementia in low-income and middle-income countries: an analysis using cross-sectional survey data. *Lancet Glob Health*. May 2019;7(5):e596-e603. doi:10.1016/S2214-109X(19)30074-9

10. See RS, Thompson F, Russell S, et al. Potentially modifiable dementia risk factors in all Australians and within population groups: an analysis using cross-sectional survey data. *Lancet Public Health*. Sep 2023;8(9):e717-e725. doi:10.1016/S2468-2667(23)00146-9

11. Suemoto CK, Mukadam N, Brucki SMD, et al. Risk factors for dementia in Brazil: Differences by region and race. *Alzheimers Dement*. May 2023;19(5):1849-1857. doi:10.1002/alz.12820

12. Jorgensen K, Nielsen TR, Nielsen A, Waldemar G. Potential for prevention of dementia in Denmark. *Alzheimers Dement*. Oct 2023;19(10):4590-4598. doi:10.1002/alz.13030

13. Flanagan A, Dunning, J., Wong, I., Brierley, A., MacDonald, BJ., Sinha, SK. *Enabling a More Promising Future for Long-Term Care in Canada*. 2023.

14. Kivipelto M, Mangialasche F, Snyder HM, et al. World-Wide FINGERS Network: A global approach to risk reduction and prevention of dementia. *Alzheimers Dement*. Jul 2020;16(7):1078-1094. doi:10.1002/alz.12123

15. Montero-Odasso M, Ismail Z, Livingston G. One third of dementia cases can be prevented within the next 25 years by tackling risk factors. The case "for" and "against". *Alzheimers Res Ther*. Jul 8 2020;12(1):81. doi:10.1186/s13195-020-00646-x

16. Feldman HH, Belleville S, Nygaard HB, et al. Protocol for the Brain Health Support Program Study of the Canadian Therapeutic Platform Trial for Multidomain Interventions to Prevent

Dementia (CAN-THUMBS UP): A Prospective 12-Month Intervention Study. *J Prev Alzheimers Dis*. 2023;10(4):875-885. doi:10.14283/jpad.2023.65

17. Chen H, Cao Y, Ma Y, Xu W, Zong G, Yuan C. Age- and sex-specific modifiable risk factor profiles of dementia: evidence from the UK Biobank. *Eur J Epidemiol*. Jan 2023;38(1):83-93. doi:10.1007/s10654-022-00952-8

18. Raina PS, Wolfson C, Kirkland SA, et al. The Canadian longitudinal study on aging (CLSA). *Can J Aging*. Sep 2009;28(3):221-9. doi:10.1017/S0714980809990055

19. Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet Neurol*. Sep 2011;10(9):819-28. doi:10.1016/S1474-4422(11)70072-2

20. Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C. Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. *Lancet Neurol*. Aug 2014;13(8):788-94. doi:10.1016/S1474-4422(14)70136-X

21. World Health Organization. *World Report on Hearing*. 2021. https://www.who.int/publications-detail-redirect/world-report-on-hearing

22. National Health Services England. Alcohl units. https://www.nhs.uk/live-well/alcohol-advice/calculating-alcohol-units/

23. Shi L, Chen SJ, Ma MY, et al. Sleep disturbances increase the risk of dementia: A systematic review and meta-analysis. *Sleep Med Rev*. Aug 2018;40:4-16. doi:10.1016/j.smrv.2017.06.010

24. Canadian Longitudinal Study on Aging. *CLSA Technical Report 2023: Sampling and Computation of Response Rates and Sample Weights for the Tracking (Telephone Interview) Participants and Comprehensive Participants*. 2023. https://www.clsa-elcv.ca/doc/5130

25. Paradis C, Butt P, Shield K, et al. *Canada's Guidance on Alcohol and Health: Final Report*. 2023.

26. Public Health Agency of Canada. *A dementia strategy for Canada: 2021 Annual report*. 2021.

27. Ramage-Morin PL., Banks R., Pineault D., Atrach M., H. G. *Hypertension associated with hearing health problems among Canadian adults aged 19 to 79 years*. 2021. *Health Reports*.

28. Public Health Agency of Canada. *A dementia strategy for Canada*. 2019.

29. Montero-Odasso M, Zou G, Speechley M, et al. Effects of Exercise Alone or Combined With Cognitive Training and Vitamin D Supplementation to Improve Cognition in Adults With Mild Cognitive Impairment: A Randomized Clinical Trial. *JAMA Netw Open*. Jul 3 2023;6(7):e2324465. doi:10.1001/jamanetworkopen.2023.24465

30. Northey JM, Cherbuin N, Pumpa KL, Smee DJ, Rattray B. Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br J Sports Med*. Feb 2018;52(3):154-160. doi:10.1136/bjsports-2016-096587

31. Schiffrin EL, Campbell NR, Feldman RD, Kaczorowski J, Lewanczuk R, Padwal R, Tobe SW. Hypertension in Canada: Past, Present, and Future. *Ann Glob Health*. Mar-Apr 2016;82(2):288-99. doi:10.1016/j.aogh.2016.02.006

32. Yeo CD, Yeom SW, You YS, Kim JS, Lee EJ. Association of sudden sensorineural hearing loss with increased risk of insomnia: a nationwide population-based cohort study. *J Clin Sleep Med*. May 1 2022;18(5):1335-1342. doi:10.5664/jcsm.9864

33. Jenstad L, Moon J. Systematic Review of Barriers and Facilitators to Hearing Aid Uptake in Older Adults. *Audiol Res*. May 10 2011;1(1):e25. doi:10.4081/audiores.2011.e25

34. Zee PC, Vitiello MV. Circadian Rhythm Sleep Disorder: Irregular Sleep Wake Rhythm Type. *Sleep Med Clin*. Jun 1 2009;4(2):213-218. doi:10.1016/j.jsmc.2009.01.009

35. Solomon A, Stephen R, Altomare D, et al. Multidomain interventions: state-of-the-art and future directions for protocols to implement precision dementia risk reduction. A user manual for Brain Health Services-part 4 of 6. *Alzheimers Res Ther*. Oct 11 2021;13(1):171. doi:10.1186/s13195-021-00875-8

36. Vergara RC, Zitko P, Slachevsky A, San Martin C, Delgado C. Population attributable fraction of modifiable risk factors for dementia in Chile. *Alzheimers Dement (Amst)*. 2022;14(1):e12273. doi:10.1002/dad2.12273

37. Rolison JJ, Hanoch Y, Wood S, Liu PJ. Risk-taking differences across the adult life span: a question of age and domain. *J Gerontol B Psychol Sci Soc Sci*. Nov 2014;69(6):870-80. doi:10.1093/geronb/gbt081

38. Borelli WV, Leotti VB, Strelow MZ, Chaves MLF, Castilhos RM. Preventable risk factors of dementia: Population attributable fractions in a Brazilian population-based study. *Lancet Reg Health Am*. Jul 2022;11:100256. doi:10.1016/j.lana.2022.100256

39. Nianogo RA, Rosenwohl-Mack A, Yaffe K, Carrasco A, Hoffmann CM, Barnes DE. Risk Factors Associated With Alzheimer Disease and Related Dementias by Sex and Race and Ethnicity in the US. *JAMA Neurol*. Jun 1 2022;79(6):584-591. doi:10.1001/jamaneurol.2022.0976

40. Besche-Truthe F. The global trajectories of compulsory education: clustering sequences of policy development. *Global Pathways to Education Global Dynamics of Social Policy*. Palgrave Macmillan, Cham. ; 2021.

41. Anderson CL, Becher H, Winkler V. Tobacco Control Progress in Low and Middle Income Countries in Comparison to High Income Countries. *Int J Environ Res Public Health*. Oct 24 2016;13(10)doi:10.3390/ijerph13101039

42. Darrow LA, Steenland NK. Confounding and bias in the attributable fraction. *Epidemiology*. Jan 2011;22(1):53-8. doi:10.1097/EDE.0b013e3181fce49b

43. World Health Organization. *Risk reduction of cognitive decline and dementia: WHO guidelines*. 2019.

44. Sims JR, Zimmer JA, Evans CD, et al. Donanemab in Early Symptomatic Alzheimer Disease: The TRAILBLAZER-ALZ 2 Randomized Clinical Trial. *JAMA*. Jul 17 2023;doi:10.1001/jama.2023.13239

45. van Dyck CH, Swanson CJ, Aisen P, et al. Lecanemab in Early Alzheimer's Disease. *N Engl J Med*. Jan 5 2023;388(1):9-21. doi:10.1056/NEJMoa2212948

46. Dhana K, Agarwal P, James BD, et al. Healthy Lifestyle and Cognition in Older Adults With Common Neuropathologies of Dementia. *JAMA Neurol*. Feb 5 2024;doi:10.1001/jamaneurol.2023.5491

47. Ngandu T, Lehtisalo J, Solomon A, et al. A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. *Lancet*. Jun 6 2015;385(9984):2255-63. doi:10.1016/S0140-6736(15)60461-5

48. Torgerson JS, Hauptman J, Boldrin MN, Sjostrom L. XENical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care*. Jan 2004;27(1):155-61. doi:10.2337/diacare.27.1.155

49. Barbera M, Lehtisalo J, Perera D, et al. A multimodal precision-prevention approach combining lifestyle intervention with metformin repurposing to prevent cognitive impairment and disability: the MET-FINGER randomised controlled trial protocol. *Alzheimers Res Ther*. Jan 31 2024;16(1):23. doi:10.1186/s13195-023-01355-x

50. Manly JJ, Deters KD. Donanemab for Alzheimer Disease-Who Benefits and Who Is Harmed? *JAMA*. Aug 8 2023;330(6):510-511. doi:10.1001/jama.2023.11704

Authors contribution

Son had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Son, Speechley, Montero-Odasso

Acquisition, analysis, or interpretation of data: Son, Speechley, Zou, Montero-Odasso

Drafting of the manuscript: Son

Revised the manuscript for important intellectual content: All authors

Statistical analysis: Son

Administrative, technical, or material support: Son, Speechley, Montero-Odasso

Supervision: Speechley, Zou, Montero-Odasso

Conflicts of Interest

Dr. Montero-Odasso reports grants from the CIHR including the Institute of Aging (MOP211220, PJT153100), the CCNA (FRN CAN 137794), Weston Foundation (BH210118), and Ontario Neurodegenerative Disease Research Initiative (OBI34739). He is on advisory boards for the Canadian Geriatrics Society (CGS) (serving as president) and the CIHR Institute of Aging, and the Research Executive Council of the CCNA. Prof. Kivipelto and Dr. Mangialasche are supported by FORTE grant 2023-01125, and are both part of the WW-FINGERS Network Global Scientific Coordinating Center, which is supported by the Alzheimer Disease Data Initiative. Dr. Feldman reports grant funding from CIHR to the CCNA (CNA- 163902) and Annovis (QR Pharma), Vivoryon (Probiodrug), AC Immune, and LuMind; service agreements for consulting activities with LuMind, Genentech (DSMB), Roche/Banner (DMC), Tau Consortium (SAB), Samus Therapeutics, Biosplice Therapeutics, Axon Neurosciences, Novo Nordisk Inc., Janssen Research & Development LLC; and travel funding from

World Events Forum (ADDF) with no personal funds received and all payments to UCSD. He also reports a philanthropic donation from the Epstein Family Alzheimer's Disease Collaboration for therapeutic research in AD with no personal funds received and all payments to UCSD. Dr. Belleville reports grant funding to CCNA from CIHR and ASC. She also reports consulting fees and participation on an advisory board for Lucilab and an unpaid role of scientific advisor for the Quebec Federation of Alzheimer's Societies. Dr. Nygaard reports consulting fees from Hoffman-La Roche and Biogen. As Scientific Director of CCNA, Dr. Chertkow reports grants from CIHR, Baycrest Health Sciences Foundation, Women's Brain Health Initiative, Alzheimer Society of Canada, Brain Canada, Saskatchewan Health Research Foundation, Alberta Innovates, and Weston Foundation (Weston Brain Institute). As a site investigator for clinical trials, he also receives support from Roche, Lilly, Anavex, and Alector. He reports being a member of the Ministerial Advisory Board on Dementia (2019-2022) and part of the membership organizing committee for Canadian Conference on Dementia. Drs. Montero-Odasso, Chertkow, Belleville, Feldman, and Nygaard are co-principal investigators of CAN-THUMB UP. Dr. Speechley is part of the steering committee for CAN-THUMBS UP. Son, Drs. Zou, Hachinski, and Peruccini-Faria have no competing interests to declare.

Statement of Informed Consent

All participants of CLSA study provided informed consent.

Table 1. Participant characteristics (Weighted N = 3,812,085)

	Overall (n=30,097)
Age, years	59.74 (10.30)
Age groups	
45 - 54	7,595 (39%)
55 - 64	9,856 (31%)
65 - 74	7,362 (18%)
75+	5,284 (12%)
Sex	
Women	15,320 (52%)
Men	14,777 (48%)
Ethnicity	
Non-white	1,326 (6.2%)
White	28,771 (94%)
Education	
< secondary school education	1,643 (17%)
Secondary school graduation	2,839 (12%)
Some post-secondary education	2,238 (9.1%)
Post-secondary degree/diploma	23,327 (62%)
Marital	
Single (never married/lived with a partner)	2,654 (8.7%)
Married/Common-law relationship	20,651 (74%)
Widowed/Divorced/Separated	6,784 (17%)
Income	
<$20,000	1,566 (6.9%)
$20,000 - $50,000	6,360 (23%)
$50,000 - $100,000	9,907 (33%)
$100,000 - $150,000	5,524 (20%)

>$150,000	4,799 (17%)
Province	
Alberta	2,957 (10%)
British Columbia	6,254 (28%)
Manitoba	3,113 (7.2%)
Newfoundland and Labrador	2,214 (2.1%)
Nova Scotia	3,078 (3.8%)
Ontario	6,418 (17%)
Quebec	6,063 (32%)
Region	
Pacific	6,254 (28%)
Prairie	6,070 (18%)
Central	12,481 (48%)
Atlantic	5,292 (5.8%)
Rurality	
Rural	2,424 (5.2%)
Urban	26,461 (90%)
Peri-urban	1,212 (4.3%)

Data shown are mean (SD) or n (%)

Table 2. PAF for 12 modifiable risk factors for dementia in Canada

	Risk factors	RR	Communality (%)	Prevalence (%)	Unweighted PAF (%)	Weighted PAF (%)
Early life	Less education	1.6 (1.3, 2.0)	56.3	14.0	7.8	3.2 (1.9, 4.3)
Midlife	Hearing loss	1.9 (1.4, 2.7)	52.3	21.0	15.9	6.5 (3.7, 9.3)
	Traumatic brain injury	1.8 (1.5, 2.2)	18.3	15.0	10.7	4.4 (3.3, 5.4)
	Hypertension	1.6 (1.2, 2.2)	55.4	30.0	15.3	6.2 (2.7, 9.3)
	Excessive alcohol	1.2 (1.1, 1.3)	43.5	11.0	2.2	0.9 (0.5, 1.1)
	Obesity	1.6 (1.3, 1.9)	57.1	31.0	15.7	6.4 (4.1, 7.7)
Later life	Smoking	1.6 (1.2, 2.2)	51.7	6.2	3.6	1.5 (0.6, 2.4)
	Depression	1.9 (1.6, 2.3)	22.3	12.0	9.8	4.0 (3.2. 4.8)
	Social isolation	1.6 (1.3, 1.9)	29.8	1.6	1.0	0.4 (0.2, 0.5)
	Physical inactivity	1.4 (1.2, 1.7)	54.1	83.0	24.9	10.2 (6.8, 13)
	Type 2 diabetes	1.5 (1.3, 1.8)	40.0	13.0	6.1	2.5 (2.4, 3.3)
	Sleep disturbance	1.2 (1.1, 1.3)	24.6	40.0	7.4	3.0 (1.8, 3.8)
	Combined PAF					49.2 (31.1, 64.9)

Abbreviations: PAF, population attributable fraction; RR, risk ratio

Table 3. Prevalence and weighted PAF of 12 potentially modifiable dementia risk factors over the life course in Canada compared to the estimates reported in other countries

	Risk factors	Canada Prev (%)	Canada wPAF (%)	Global[5] Prev (%)	Global[5] wPAF (%)	US[7] Prev (%)	US[7] wPAF (%)	New Zealand[8] Prev (%)	New Zealand[8] wPAF (%)	Australia[10] Prev (%)	Australia[10] wPAF (%)	Denmark[12] Prev (%)	Denmark[12] wPAF (%)	India[9] Prev (%)	India[9] wPAF (%)	Latin America[9] Prev (%)	Latin America[9] wPAF (%)	China[9] Prev (%)	China[9] wPAF (%)	Brazil[11] Prev (%)	Brazil[11] wPAF (%)
Early life	Less education†	14	3.2	40	7.1	10.7	2	31	4.6	10	1.9	9.5	1.9	92.2	13.6	68.8	10.9	75.9	10.8	46.6	7.7
Mid life	Hearing loss†	21	6.5	31.7	8.2	10.8	3	39.9	7.8	26.6	7	23.1	6.4	22.3	6.4	28.8	7.7	14.3	3.9	26.5	6.8
	Traumatic brain injury	15	4.4	12.1	3.4	17.1	4.1	18.2	3.7	n/a	n/a	4.7	1.4	n/a	n/a	n/a	n/a	n/a	n/a	12.1	3.1
	Hypertension†	30	6.2	8.9	1.9	42.2	6.8	33.5	4.9	17.2	3.3	22.5	4.3	19.3	4	56.6	9.3	38.1	6.4	46.4	7.6
	Excessive alcohol	11	0.9	11.8	0.8	3.6	0.2	9.8	0.6	8.2	0.5	9.4	0.6	n/a	n/a	n/a	n/a	n/a	n/a	4.3	0.3
	Obesity†	31	6.4	3.4	0.7	44	7.1	38.4	5.5	38.8	6.6	22.7	4.3	13.7	2.9	44.8	7.9	32	5.6	31.4	5.6
Later life	Smoking†	6.2	1.5	27.4	5.2	8.5	1.7	13.5	2.2	6.7	1.4	12.2	2.4	39.9	6.4	30	5.7	23	4.2	10.6	2.1
	Depression†	12	4	13.2	3.9	7.4	2.1	19.1	4.3	11.6	3.3	8	2.4	5.2	1.7	23.9	6.6	1.5	0.5	15.8	4.4
	Social isolation†	1.6	0.4	11	3.5	11.9	2.3	37.3	5.4	5.2	1.0	8.3	1.6	10.4	2.3	0.5	0.1	3.4	0.7	1.6	0.3
	Physical inactivity†	83	10.2	17.7	1.6	62.8	6.8	53.6	5.2	82	8.3	67.6	7.5	15.3	2.2	34.2	4.5	50.7	5.8	36.7	4.5
	Diabetes†	13	2.5	6.4	1.1	28.6	4.2	11.6	1.6	16.8	2.7	13.2	2.4	9.3	1.7	18.5	3.2	9.4	1.6	19.7	3.1
	Sleep disturbance	40	3	n/a	n/a	n/a	n/a	31	4.6	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Combined PAF of 9 risk factors†			40.9		33.2		36		41.5		35.5		33.2		41.2		55.9		39.5		42.1
Combined PAF of all risk factors			49.2		37.4		40.3		45.8		36.0		35.2		41.2		55.9		39.5		45.5

n/a Not available; Prev, Prevalence; wPAF, Weighted population attributable fraction

* Weighted PAF was calculated based on the data provided in Lee et al. (2022)[7]

† Risk factors available across all studies

Table 4. Weighted prevalence and PAF of 12 potentially modifiable risk factors by 4 age groups and sex in Canada

Risk factors	Overall		Age group									Sex				
			45-54		55-64		65-74		75-85			Female		Male		
	Prev (%)	wPAF (%)	Prev (%)	wPAF (%)	Prev (%)	wPAF (%)	Prev (%)	wPAF (%)	Prev (%)	wPAF (%)	p*	Prev (%)	wPAF (%)	Prev (%)	wPAF (%)	p*
Less education	12.6	2.7	7.2	1.9	12.5	2.6	16.8	3.9	24.7	5.8	<0.001	12.1	2.8	13.2	2.8	0.29
Hearing loss	33.2	8.7	13.5	4.9	28.5	7.7	54.2	14.1	81.2	19.0	<0.001	29.6	8.7	37.1	9.6	<0.001
Traumatic brain injury	13.9	3.8	15.3	5.0	14.4	3.9	13.0	4.0	8.9	3.0	<0.001	10.3	3.1	17.8	4.8	<0.001
Hypertension	37.0	6.9	21.5	5.2	38.8	7.1	53.2	10.4	59.8	11.9	<0.001	35.4	7.2	38.8	7.3	<0.001
Excessive alcohol use	11.5	0.9	10.9	1.0	12.4	0.9	12.4	1.0	9.8	0.9	<0.001	6.4	0.5	17.1	1.3	<0.001
Obesity	30.4	5.8	27.3	6.4	34.5	6.4	32.9	7.1	26.0	6.1	<0.001	30.1	6.3	30.7	6.0	0.02
Smoking	10.7	2.3	13.7	3.5	11.9	2.5	6.4	1.6	3.9	1.0	<0.001	10.7	2.5	10.7	2.3	0.04
Depression	16.5	4.9	16.4	5.9	20.6	5.9	14.3	4.9	9.0	3.4	<0.001	20.8	6.5	11.8	3.7	<0.001
Social isolation	1.6	0.4	1.9	0.5	1.4	0.3	0.9	0.2	1.8	0.5	0.003	1.0	0.2	2.2	0.5	<0.001
Physical inactivity	76.6	8.9	71.6	10.2	76.8	8.8	79.8	10.4	87.8	11.7	<0.001	80.0	10.0	72.8	8.7	<0.001
Diabetes	8.5	1.6	4.3	1.0	10.3	1.8	12.3	2.5	12.1	2.6	<0.001	7.2	1.4	10.0	1.8	<0.001
Sleep disturbance	43.0	3.0	43.0	3.6	45.3	3.1	39.7	3.2	41.7	3.5	0.004	42.2	3.2	43.7	3.1	0.1
Combined PAF		49.7		49.1		51.1		63.4		69.3			52.4		51.9	

Prev, Prevalence; wPAF, Weighted population attributable fraction

* Chi-square test comparing the prevalence of risk factor by age groups or sex

Chi-square test was weighted with the CLSA analytic weight

Figure 1. Weighted population attributable fraction for 12 potentially modifiable risk factors for dementia in Canada