Title: The role of viral interference in shaping RSV epidemics following
the 2009 H1N1 influenza pandemic

Authors: Ke Li[1]*, Deus Thindwa[1], Daniel M Weinberger[1], Virginia E Pitzer[1]*

Affiliations:

[1]Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New

Haven, CT, USA.

*Corresponding authors. ke.li.kl662@yale.edu, virginia.pitzer@yale.edu

One Sentence Summary: We demonstrated disrupted RSV activity in the United States

following the 2009 influenza pandemic by analyzing weekly positive tests for RSV and the

pandemic H1N1 virus. During the 2009/10 season, RSV experienced reduced activity, which

was negatively associated with the activity of pandemic influenza. In contrast, RSV showed

increased activity in the 2010/11 season due to the buildup of susceptible populations from the

previous season. By focusing on the dynamics of RSV following the pandemic, we found

evidence supporting interactions between the viruses at the population level. Our findings

suggest that infections with pandemic influenza could: 1) reduce host susceptibility to RSV

coinfection, 2) shorten the RSV infectious period in coinfected individuals, or 3) decrease RSV

infectivity in coinfection.

Abstract: Respiratory syncytial virus (RSV) primarily affects infants, young children, and

older adults, with seasonal outbreaks in the United States (US) peaking around December or

January. Despite the limited implementation of non-pharmaceutical interventions, disrupted

RSV activity was observed in different countries following the 2009 influenza pandemic,

suggesting possible viral interference from influenza. Although interactions between the

influenza A/H1N1 pandemic virus and RSV have been demonstrated at an individual level, it

remains unclear whether the disruption of RSV activity at the population level can be attributed

to viral interference. In this work, we first evaluated changes in the timing and intensity of RSV

activity across 10 regions of the US in the years following the 2009 influenza pandemic using

dynamic time warping. We observed a reduction in RSV activity following the pandemic,

which was associated with intensity of influenza activity in the region. We then developed an

age-stratified, two-pathogen model to examine various hypotheses regarding viral interference

mechanisms. Based on our model estimates, we identified three mechanisms through which

influenza infections could interfere with RSV: 1) reducing susceptibility to RSV coinfection;

2) shortening the RSV infectious period in coinfected individuals; and 3) reducing RSV

infectivity in coinfection. Our study offers statistical support for the occurrence of atypical

RSV seasons following the 2009 influenza pandemic. Our work also offers new insights into

the mechanisms of viral interference that contribute to disruptions in RSV epidemics and

provides a model-fitting framework that enables the analysis of new surveillance data for

studying viral interference at the population level.

Main Text:

INTRODUCTION

Respiratory syncytial virus (RSV) infections are a major public health concern for infants and

young children, causing severe lower respiratory tract infections (*1*, *2*). In 2019, an estimated

13,300 deaths were associated with RSV-induced acute lower respiratory infections in

hospitals, and there were 45,700 RSV-attributed deaths in infants aged 0-6 months worldwide

(*3*). In the United States (US) and other temperate regions, RSV activity is strongly seasonal,

typically beginning in the fall and peaking in winter (*4*). During the COVID-19 pandemic in

2020-2021, RSV disappeared for more than a year in the Northern Hemisphere then reemerged

out of the normal season (*5, 6*). This was likely due to the implementation of non-

pharmaceutical interventions (NPIs) to reduce the spread of SARS-CoV-2. However,

disruptions to RSV activity were also observed following the 2009 influenza pandemic in

different countries despite the limited implementation of NPIs (*7–12*). Viral interference from

the novel 2009 A/H1N1 influenza pandemic (pdmH1N1) virus was suggested as a cause of the

delayed RSV activity.

Viral interference refers to a phenomenon where one virus prevents or reduces infection by

another virus (*13*). Several possible mechanisms of interference have been proposed at the

within-host level (*14–16*). Direct competition between viruses for infection of susceptible cells

can result in viral interference, such that the consumption of target cells by one virus limits

infection by the other. Viral interference can also occur through the host immune response. For

example, antibody-mediated interference has been suggested as a mechanism for mitigating

infection between genetically close viruses, such as RSV and human metapneumovirus (*16*).

Host interferon responses (IFNs) have also been proposed as a mechanism leading to viral

interference. Chan et al. demonstrated that ferrets infected with pdmH1N1 influenza A virus

(IAV) could be protected from subsequent RSV infection depending upon the interval between

the two infections. They showed that the protection was transient and mediated by the IFNs

(*15*). More recently, viral interference between IAV and the SARS-CoV-2 was also studied

using human airway epithelial cultures. Cheemarla et al. showed that IAV infection could lead

to a robust IFN response that suppressed subsequent SARS-CoV-2 infection (*17*). In contrast,

infection with SARS-CoV-2, which elicited a relatively weak immune response, was unable to

suppress IAV infection, suggesting an asymmetric IFN-dependent viral interference

mechanism.

While it is evident that viral interference occurs at the host level, demonstrating the presence

of viral interference at the population level and quantifying its impact on disease transmission

is more challenging. For example, co-infections with two or more viruses in an individual can

potentially change the typical viral dynamics (e.g., time to infectiousness, symptoms and

severity) in the host, leading to changes in disease transmission patterns at the population level,

as reviewed in (*18*). To date, most studies that analyzed population-level viral interference have

primarily focused upon statistical associations between reported positive cases of different

viruses, using regression and correlation analyses (*16*, *19*). The method, lacking explicit

mechanistic formulation, was unable to distinguish true viral interaction from other

confounding factors, such as climate variables. Other techniques, including seasonal auto-

regressive integrated moving average models (*20*) and Granger causality (*21*, *22*), have also

been used to analyze time series of positive tests of viruses, aiming to identify potential

interactions between viruses. However, neither the biological mechanisms underpinning

potential viral interactions, nor the strength of interactions could be determined or quantified

using these models.

Mathematical models that explicitly depict the underlying mechanisms of viral transmission

have advantages in being able to integrate heterogeneous mechanisms and test different

hypotheses (*23–25*). Some mathematical models have been proposed to study the effect of viral

interference between RSV and influenza viruses at the population level and to quantify the

interactions by fitting the models to incidence data (*26–28*). However, these models did not

capture the natural infection history of RSV, which are characterized by intermediate immunity

lying between perfectly and imperfectly immunizing infections (*29*). In this work, we started

by analyzing laboratory-confirmed cases of RSV and the pdmH1N1 virus in different regions

in the US. We applied a dynamic time warping and hierarchical clustering method to identify

atypical RSV seasons following the 2009 influenza pandemic. We then built a mechanistic,

age-stratified mathematical model that incorporates an RSV transmission model into an

influenza transmission model, coupled with hypothesized viral interference mechanisms.

Using Latin Hypercube Sampling to explore the parameter space, we simulated various

scenarios for influenza dynamics and quantified the potential interactions between RSV and

pdmH1N1 influenza virus at the population level.

RESULTS

RSV activity following the 2009 influenza pandemic

RSV epidemics exhibit consistent seasonal timing and duration within each region of the US,

with variation in timing between regions (**Fig. 1A** and **Fig. S1**). Annual RSV epidemics start

in the fall and peak in the winter. Some regions (e.g., Regions 8 and 10 in the Upper Midwest

and Northwest) exhibit biennial patterns, with RSV tending to start and peak earlier with a

larger epidemic in even-numbered seasons (e.g., 2010/11 and 2012/13) compared to the odd-

numbered years (e.g., 2009/10 and 2011/12). The influenza pandemic (the shaded area, **Fig.**

1A and **Fig. S1**) began in April 2009, after the 2008/09 RSV season (dashed black lines). This

was followed by a second wave in most regions that started at the end of 2009, before the peak

of the 2009/10 RSV season. In our analysis, we focus on RSV activity during the 2009/10

season, when the pdmH1N1 virus was the sole influenza virus circulating in the population,

and the 2010/11 season following the pandemic.

Based on the laboratory reports of RSV-positive specimens (**Fig. 1A**), we quantified RSV

activity following the pandemic season and in other normal epidemic seasons. Regions 1 and

4 (Northeast and Southeast) exhibited consistent annual patterns of RSV activity, with a steady

onset and peak timing in different seasons, including the 2009/10 pandemic year (**Fig. 1B**).

However, we observed delayed RSV activity in 2010/11 compared with other seasons in the

two regions, as indicated by a shift of intense RSV activity to later epidemic weeks. Again,

Regions 8 and 10 showed a biennial pattern of RSV epidemics with an earlier onset and peak

timing in even-numbered years (**Fig. 1B**). In these regions, delayed RSV activity was also

evident in 2010/11 compared to other even-numbered years, and no notable shifts in the timing

of RSV activity were observed in the 2009/10 season.

Clustering RSV activity using dynamic time warping

Epidemic curves often exhibit variations in reporting intervals and temporal dynamics, making

it difficult to identify dissimilarities in disease spread patterns. To better identify and

characterize potential time shifts in RSV activity following the pandemic, we used dynamic

time warping (DTW) to compare RSV activity in the 2009/10 and the 2010/11 seasons with

other epidemic seasons. The fundamental concept of DTW is to find the optimal alignment

between multiple time series, allowing the adjustment of the timing of one of the time series

while minimizing the distance between corresponding data points (*30*). An optimal alignment

shows the indices of the elements in the query time-series that correspond to those in the

reference time series (e.g., RSV time-series in the 2009/10 and 2010/11 seasons). The

computed optimal alignment paths for RSV activity during the 2009/10 and 2010/11 seasons

with respect to other seasons are provided in **Fig. S2**. We found that the alignment paths cross

below the diagonal line for the 2010/11 season, indicating delayed RSV activity as depicted in

Fig. 1B.

The hierarchical clustering results of seasonal RSV activity are given in **Fig. 2**. In Region 1

(**Fig. 2A**) and Region 4 (**Fig. 2B**), the seasons following the pandemic (i.e., 2009/10 and

2010/11) were closer in the dendrogram, grouped into the same cluster, compared with other

seasons before or after the pandemic (as shown in the dashed box). This indicated more similar

trends of RSV activity in these years. We also observed that even within the same cluster, the

2009/10 season (highlighted in blue) was distant from the 2010/11 season (highlighted in red),

suggesting a different pattern between the two seasons following the pandemic. Notably, the

hierarchical clustering method successfully captured the biennial pattern in Region 8 (**Fig. 2C**),

Region 10 (**Fig. 2D**), and other regions in the US (**Fig. S3**), grouping even seasons and odd

seasons into different clusters. Specifically, the 2009/10 season was grouped with even-year

seasons in Region 8, indicating unusual RSV activity. While the 2010/11 season in Region 10

shares a cluster with other even-year seasons, its extended branch length in the dendrogram

indicated a distinction from the other even-year seasons.

Linking RSV activity and the 2009 influenza pandemic

We further quantified RSV activity following the pandemic seasons across 10 regions of the

US. Both the intensity and the center of gravity for each RSV epidemic season in each region

were calculated. The onset time of RSV activity for each season was also calculated. There was

a strong positive correlation (i.e., $\rho = 0.92$) between the center of gravity and onset time of

RSV activity, indicating these two methods of measuring epidemic timing agree (see **Fig. S4**).

We explored the trends of RSV activity in the 2009/10 and 2010/11 years across the US

regions, assessing changes in both intensity and center of gravity in comparison to the median

values of all other seasons in the same region (**Fig. 3**). Most regions in the US experienced

decreased RSV intensity in the 2009/10 season, with the timing of RSV activity consistent with

that of other seasons. By contrast, most regions in the US showed increased RSV intensity,

with delayed peak timing in the 2010/11 season. We then computed the intensity of the

pdmH1N1 virus, starting from April 2009 to May 2010. We found a negative association

between the intensity of RSV and the intensity of the pdmH1N1 virus (**Fig. 4A**)—enhanced

influenza intensity was associated with decreased RSV intensity in the 2009/10 season (i.e.,

$\rho = -0.38$). The negative correlations between RSV activity and pdmH1N1 activity suggested

the presence of viral interference from influenza on RSV transmission. No correlations were

found between the intensity of the pdmH1N1 virus and the intensity of RSV (**Fig. 4B**) or the

timing of RSV peak in 2010/11 (**Fig. S5**).

Transmission model analyses

To explore and examine various hypotheses on the mechanistic relationship between RSV

activity and viral interference from influenza that might explain the statistical associations

observed, we proposed a mathematical model that explicitly incorporated the transmission

dynamics of RSV into that of the pdmH1N1 virus via three hypothetical viral interaction

mechanisms (see **Materials and Methods** for detailed model description). We hypothesized

that in hosts infected with pdmH1N1 virus, the IFN response leads to: 1) a reduction of the

host's susceptibility to subsequent RSV infections (**Fig. S6A**), which was captured by a

parameter θ; 2) a reduction of the infectious period of RSV co-infection (**Fig. S6B**), captured

by a parameter η; and/or 3) a reduction of the force of infection of RSV (**Fig. S6C**), captured

by a parameter ξ. Detailed model equations are provided in the **Supplementary Materials**.

The three proposed mechanisms were studied separately as Models I-III. For each mechanism,

we first used Latin Hypercube Sampling (LHS) to generate a wide range of parameter values

that allowed us to simulate different transmission dynamics of pandemic influenza in the

2009/10 season and explore the effects of viral interference (including the possibility of no

interference when $\theta = 1$, $\eta = 1$ or $\xi = 1$). We then determined the goodness-of-fit of the models

based on the negative log-likelihood of the models fitted to the number of RSV-positive tests,

and filtered the models to the top 2% best-fitting models to identify the corresponding

parameter ranges for the viral interference mechanisms.

The best 2000 (i.e., top 2%) fitting models successfully captured the difference in relative

intensity of RSV activity during the 2009/10 and the 2010/11 seasons; however, the best-fit

models failed to capture the shift in timing of RSV activity during 2010/11. Based on the

calculated likelihood, the model without viral interference (green curves in **Fig. 5**) was not

included in the top 2% of models. The reduction of the host's susceptibility to RSV infection

(Model I, **Fig. 5A**), the RSV infectious period (Model II, **Fig. 5B**), or the force of infection of

RSV (Model III, **Fig. 5C**) led to decreased RSV activity in the 2009/10 season, followed by

increased RSV cases in the 2010/11 season. The relative intensity of RSV activity between the

two seasons could not be reproduced in the absence of viral interference mechanisms. Similar

results were observed for the top 3% (**Fig. S7**) and top 5% (**Fig. S8**) of the models, such that

the models also captured the relative intensity of RSV activity in the 2009/10 and 2010/11

seasons following the pandemic. Further, our models demonstrated that increased RSV activity

in the 2010/11 season could be explained by an increase in the proportion of the population

susceptible to RSV infection, as shown in **Fig. S9**.

The identified parameter spaces for θ, η and ξ from the top 2% best-fitting models for Region

1 provide insight into the effect of viral interference from influenza influencing RSV infection

and transmission (**Fig. 6**). The median estimate for the reduction of host susceptibility to RSV

infection when infected with influenza virus (θ) was 0.44 (95% CI: 0.23-0.60, **Fig. 6A**). In

epidemiological terms, the median estimate for θ indicates that the presence of the pdmH1N1

infection reduces the likelihood of hosts being subsequently infected with RSV by nearly 60%.

The median estimate for η implies the presence of the pdmH1N1 co-infection halves the RSV

infectious period, i.e. increases the rate of recovery by a factor 1.98 (95% CI: 1.62-2.53, **Fig.**

6B). Similarly, the median estimate for ξ was 0.46 (95% CI: 0.20-0.80, **Fig. 6C**), suggesting

the pdmH1N1 infection reduces RSV infectivity by 53%. The parameter distributions from the

top 3% and top 5% of the fitted models were similar (**figs. S10-11**). Given the best-fit parameter

distributions for the viral interference parameter excluding 1, our results suggest that viral

interference plays a role in mediating the dynamics of RSV infection following influenza

infection for all of the proposed mechanisms.

Our models were also able to capture the trends of RSV activity in Region 10 (showing a

biennial RSV pattern) and Region 4 (where seasonal RSV activity is the earliest in the US).

From the identified parameter space of the viral interaction terms, similar results were found

showing strong viral interference between RSV and pdmH1N1 virus in Region 10 (**Fig. 6D-**

F). The median estimate of the reduction in host susceptibility to RSV infection (θ) was 0.61

(95% CI: 0.42-0.74, **Fig. 6D**), the relative rate of recovery from co-infection (η) was 1.5 (95%

CI: 1.28-2.22, **Fig. 6E**), and the relative RSV infectivity (ξ) was 0.62 (95% CI: 0.42-0.75, **Fig.**

6F). By contrast, we found weak interference effects between the viruses in Region 4, such that

all estimated parameters were closed to 1 (**Figs. 6G-I**).

With the identified viral interference parameters, we further predicted the co-infection rate of

RSV and pdmH1N1 influenza virus in different age groups following the pandemic in the

presence or absence of viral interference. The rate was computed as the fraction of infections

in each age group that were co-infections. We found that the presence of viral interference

reduced the co-infection rate in all age groups (**Fig. S12**). In particular, our model showed that

the effect was even more profound at the young age groups (i.e., < 1 year old), reducing the

co-infection rate from 2% to 0.8%. Such patterns could be explained by the age-specific

likelihood of a susceptible individual coming into contact with an infectious individual.

DISCUSSION

By focusing on the dynamics of RSV following the 2009 H1N1 influenza pandemic, we found

evidence supporting the presence of interactions between the viruses at the population level

and examined the underlying mechanisms. This was accomplished using statistical analysis of

laboratory-confirmed positive tests of both viruses in 10 regions of the US and a mathematical

modeling approach. Using a dynamic time warping and hierarchical clustering method, we

identified atypical RSV activity following the 2009 influenza pandemic. The results support

and contribute to the current knowledge from several observational studies that RSV activity

was disrupted following the pandemic (*7–12*). We further showed there was a negative

correlation between the intensity of RSV and pdmH1N1 activity during the pandemic. Using a

two-pathogen, age-stratified transmission model, we assessed potential interactions between

the viruses and identified three mechanisms of viral interference that can replicate the relative

difference of RSV activity in the two epidemic seasons following the pandemic. The identified

parameter space suggested that infection with the pdmH1N1 virus could reduce either the

host's susceptibility to a subsequent RSV infection, or the infectious period of RSV infection,

or RSV infectivity.

Although the presence of viral interference between different pathogens is evident at the host

level (*15*, *31*, *32*), demonstrating the impact of this phenomenon at the population poses a

significant challenge. Mechanistic models provide a valuable approach for dissecting the casual

relationship among different components, integrating heterogeneous mechanisms and testing

various hypotheses. One of the most important applications of mechanistic models is to

estimate key parameters, as reviewed in (*33*). Parameter identifiability for data fitting is an

important but unresolved challenge in modeling work due to model complexity or limited time-

series on numerous quantities of interest (*34–36*). Waterlow et al. have previously highlighted

the importance of parameter identifiability (*26*). They conducted a simulation and back-

estimation study to evaluate the plausible parameter space of viral interaction parameters in an

RSV-influenza model. By fitting the model to a single season of simulated data, however, they

demonstrated that the inference results for the interaction parameters were often imprecise,

indicated by large credible intervals. By contrast, we specifically focused on the RSV-

pdmH1N1 pair in our study. During the 2009/10 epidemic season, the pdmH1N1 virus emerged

as the only circulating influenza virus, dominating the population. The outbreak of pdmH1N1

virus provided an opportunity to investigate viral interference between pdmH1N1 and RSV.

With the availability of regional-level data, we were able to dissect variations in the temporal

dynamics (i.e., annual/biennial patterns) of RSV activity in different regions in the US and

estimate the interaction parameters between the viruses. A key finding of the models suggests

that the strength of interaction between the viruses could be estimated from surveillance data

of both viruses in such situations, revealing that the activity of pdmH1N1 has an appreciable

impact on RSV activity.

At the individual level, the host interferon response has been shown to be the mechanism

underpinning viral interference between RSV and the pdmH1N1 virus (*15*). Depending upon

the interval between pdmH1N1 and RSV infections, the within-host viral production of RSV

could be completely suppressed, reduced, delayed, or shortened. Based on the within-host RSV

297 viral dynamics, we proposed three mechanisms through which innate immunity, stimulated by

298 the pdmH1N1 virus, could modulate the following RSV coinfection. The model fits showed

299 that each of the three mechanisms could successfully recapitulate a decrease in RSV activity,

300 followed by an enhanced RSV epidemic, in the two seasons following the pandemic. Our model

301 demonstrated that the increased intensity of the 2010/11 RSV season could be attributed to an

302 increased proportion of susceptible individuals in the population. Baker et al. similarly showed

303 that population susceptibility to influenza and RSV infections increased during the COVID-19

304 pandemic, leading to larger outbreaks following the relaxation of NPIs (*37*). More recently,

305 Lowensteyn et al. showed that the unprecedented RSV epidemic following the COVID-19

306 pandemic in the Netherlands was associated with waning immunity to RSV due to low

307 circulation of RSV during the NPIs period (*38*).

309 We note that none of the mechanisms considered in the model, including the null model (i.e.,

310 no viral interference), could capture the shift in timing of the 2010/11 RSV epidemic. One

311 possible explanation would be that we only assumed a transient viral interference interval

312 lasting up to a week, occurring in co-infected individuals and disappearing when the infection

313 resolves. The IFN response was not explicitly considered in our model, and could persist

314 slightly longer than the influenza infectious period. Although previous studies have suggested

315 that cross-protection following influenza infection against RSV could last more than two weeks

316 (*39*), this was not demonstrated in a ferret model, which shares several similarities with the

317 respiratory tracts of humans (*40*). The duration of cross-protection between RSV and the

318 pdmH1N1 virus only lasted a week in ferrets, as shown in (*15*).

Delayed RSV activity in the 2010/11 season could also be explained by the circulation of other

respiratory viruses (e.g., rhinovirus and influenza B virus), which our model did not explicitly

include. Here, our focus was not on exploring virus interactions among multiple influenza

strains and RSV in general. Instead, our specific emphasis was on studying viral interference

from the pdmH1N1 virus on RSV infection, where the interactions are evident at the host level.

This focus was guided by experimental studies indicating that the genetic strains of influenza

viruses elicit varying levels of host immunity (41). We also assumed that the RSV-influenza

interaction was unidirectional during the 2009/10 season, meaning that only the pdmH1N1

virus would exhibit interference on RSV. This assumption is justified considering that the

influenza pandemic preceded the normal 2009/10 RSV season, and our analysis focused on the

disruption in RSV activity following the 2009 influenza pandemic. Note that neither could, nor

did we intend to show the absence of viral interference from RSV against influenza viruses. To

study viral interference from RSV in shaping the transmission dynamics of influenza may

require time-series data from multiple epidemic seasons, as shown in a modeling study (27).

Besides potential viral interference, other factors can also change disease transmission patterns,

leading to the variations between model predictions and the surveillance data. Behavioral

changes can have significant impacts on viral circulation and transmission patterns, as observed

during the COVID-19 pandemic (42). After the implementation of NPIs, there was a sharp

decline in the number of positive RSV tests, and RSV activity remained disrupted in the

following seasons (5, 43). It is not clear how much of a role, if any, viral interference from

SARS-CoV-2 played in disrupting RSV activity. Understanding the interplay of the effects of

NPIs and viral interference is important to evaluate the role of behavioral and immunological

factors more accurately on disease transmission. Additionally, environmental factors such as

temperature and humidity (29) and vaccination coverage (44) can also play roles in shaping the

trajectory of disease spread. Incorporation of these factors into the dynamic model can be

improved in future work upon data availability.

Dynamic time warping (DTW) is a widely used statistical algorithm (*30, 45*), but its application

in identifying various disease transmission patterns has been limited. Recently, multiple studies

have used DTW to analyze the trajectories of COVID-19 in different countries, aiming to

identify, cluster and predict future trends in disease transmission (*46–49*). Here, we utilized

DTW to examine and visualize similarities of RSV time-series, yielding clusters of RSV

activity before and after the 2009 influenza pandemic in the ten different regions of the US.

The method successfully identified the biennial pattern of RSV epidemics in certain regions of

the US and atypical RSV seasons following the pandemic. The graphical representation of

clusters based on DTW provides an accessible and interpretable method for comparing both

temporal and spatial time-series of incidence data, enhancing our understanding of disease

transmission patterns longitudinally or geographically. The DTW method also has a promising

potential for detecting and identifying atypical epidemic seasons.

We identified three plausible viral interference mechanisms that could shape RSV epidemics

following the influenza pandemic. We do acknowledge that these mechanisms are not mutually

exclusive. This raises the question of what kind of data would be needed to further distinguish

among the models and examine the relative importance of each mechanism for RSV

transmission. One possible direction would be to emphasize the incidence of coinfections over

a period of time. By fitting the prevalence of coinfections to mathematical models, we could

estimate essential kinetic parameters separately, such as the force of infection or the recovery

rate from the coinfected components. The comparison between these estimated parameter

values and the baseline values (i.e., the rates from individuals infected with one virus) helps

discern different mechanisms and evaluate the relative contribution of each process.

Our study provides statistical and mathematical support for the presence of viral interference

between the pdmH1N1 influenza virus and RSV at the population level. Multiple mechanisms

mediated by the host immune response are capable of explaining RSV transmission dynamics

following the 2009 influenza pandemic. Given the experimental support for within-host

interference between RSV and influenza through IFN responses, all of these mechanisms could

contribute to shaping RSV epidemics. The results have implications for implementing and

evaluating disease control interventions. For example, mitigation measures that effectively

decrease disease transmission could decrease the epidemic size in the current season but

potentially lead to a larger outbreak in the following season, even if the measures are applied

for only a week (i.e., transient protection). Hence, it is imperative to deliberate upon the optimal

degree of disease suppression and the duration for which interventions should be applied prior

to their implementation, as discussed in (*50*).

There are some limitations to our study. First, we did not have age information on the positive

tests for RSV and influenza over time. Therefore, we assumed a well-mixed population and did

not account for varying levels of immunity across different age groups beyond the age-specific

contract matrices. It is possible that influenza virus stimulates a weaker innate immune

response in young children compared to other age groups (*51*). In our model, the estimated

interference parameters are interpreted as the average effects of viral interaction across the

entire population. Although our model cannot be used to assess the strength of viral interference

in each age group, our results are still sufficient to demonstrate the presence of viral

interference between pdmH1N1 and RSV within the population in general. Another limitation

of our analyses is that our model did not take into account other circulating pathogens (e.g.,

rhinovirus or human parainfluenza virus) due to the added complexity and additional

parameters required. For example, interference between rhinovirus and influenza viruses was

recognized by epidemiological observations (*31*), and the delayed 2009 influenza pandemic in

Europe was attributed to the prevalence of rhinovirus (*7*). The analysis of viral interference

between the pandemic influenza virus and other viruses will be left for future work.

Additionally, our model estimates did not show strong viral interference effects between the

viruses in Region 4. The difference in the estimates of viral interference for Region 4 is likely

attributable to the different pdmH1N1 activity in this region. In Regions 1 and 10, there was a

strong second wave of pdmH1N1 before the 2009/10 RSV season, whereas the second

pdmH1N1 wave was not observed in Region 4. It is not clear why the pandemic influenza

exhibited different activity in these regions.

Our findings, which indicate an association between the incidence of RSV infections and

pdmH1N1 infections, also have implications for enhanced surveillance of disease transmission

of other viruses. As other respiratory viruses (e.g., seasonal influenza viruses and SARS-CoV-

2) are expected to co-circulate in the upcoming epidemic seasons, our study provides a

framework for studying viral interactions and understanding transmission dynamics.

Additional information on the frequency of RSV and other viruses, along with coinfections,

would enable us to further validate our results. The mechanistic model proposed in this work

is flexible to incorporate the effects of vaccination in preventing disease, such as introducing a

model compartment that is resistant to infection and becomes susceptible over time. The

extended model can be used for the evaluation of various vaccination scenarios, assessing the

impact of vaccination coverage on long-term disease patterns while considering the presence

of viral interference. Through systematic analysis of these scenarios, our model can provide

valuable insights into the dynamic interplay between vaccination strategies and the patterns of

disease transmission, contributing to informed decision-making in public health interventions.

MATERIALS AND METHODS

Laboratory reporting of RSV and influenza

Weekly data on laboratory reporting of RSV tests in ten Health and Human Services (HHS)

regions in the US from June 2007 to July 2019 were obtained from The National Respiratory

and   Enteric   Virus   Surveillance   System   (https://data.cdc.gov/Laboratory-

Surveillance/Respiratory-Syncytial-Virus-Laboratory-Data-NREVSS/52kb-ccu2/about_data).

The regional map can be found on the https://www.hhs.gov/about/agencies/iea/regional-

offices/index.html website. Positive RSV tests were detected using three diagnostic methods:

1) antigen detection; 2) reverse transcription polymerase chain reaction (RT-PCR); and 3) viral

culture. Correspondingly, the data on laboratory reporting of influenza tests in the US from the

same period were obtained from the Center for Disease Control and Prevention (CDC) website

https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html.

The raw laboratory data were rescaled based on the number of positive tests to account for

variations in testing practices over time (*29*). We first calculated a one-year moving average of

the weekly number of RSV or influenza tests (both positive and negative tests) in each region

centered on each week. We then calculated a weekly scaling factor for each region equal to the

average number of RSV or influenza tests during the entire period of reporting (i.e., 12

epidemic seasons from 2007-2019) divided by the one-year moving average. The rescaled

number of RSV or influenza-positive tests for each region was then calculated as the reported

number of positive tests multiplied by the weekly scaling factor. The rescaled data for RSV

and influenza are shown in **Fig. 1** and **S1.**

Demographic Data

Information about population size in each age group was obtained from the US Census

Bureau's American Community Survey. Birth rates varied between regions and over time

based on the crude annual birth rate for each HSS region from 1990 to 2019. These were

obtained from https://wonder.cdc.gov/controller/datarequest/D66. To capture aging among

infants and children more accurately in our mathematical model, we divided the <1 year and

1-4 years age class into 12-month age groups. The remaining population was divided into 5

classes: 5-9 years, 10-19 years, 20-39 years, 39-60 years and >60 years old. Individuals were

assumed to age exponentially into the next age class, with the rate of aging equal to the

multiplicative inverse of the width of the age class.

The onset timing, center of gravity, and intensity of RSV activity

To determine the onset timing (measured by week) of each seasonal RSV epidemic for each

region, we initially fitted a p-spline curve to the RSV incidence data. We then calculated the

first and second derivatives of the fitted curve. The onset timing corresponds to the time point

at which the second derivative of the fitted p-spline curve reaches its maximum value during

the increasing segment of the first derivative (*52*). The center of gravity of RSV activity for

each season in each region ($G_{s,r}$) was measured as the mean epidemic week, with each week

weighted by the number of positive tests, such that $G_{s,r} = \sum_{w \in [1:52]} w \times Y_{s,r,w} /$

$\sum_{w \in [1:52]} Y_{s,r,w}$, where w is an index for the week of each epidemic year, and $Y_{s,r,w}$ is the

number of rescaled positive RSV tests in region r during epidemic season s and week w. To

determine the RSV intensity for each season and region, we also used the fitted p-spline curve.

The epidemic peak timing was determined at the point when RSV activity reaches its

maximum. The intensity of RSV was calculated as the fraction of positive tests before the

epidemic peak timing, corresponding to the integral of the positive first derivative of the log-

transformed fitted p-spline curve.

Dynamic time warping and hierarchical clustering

We used a dynamic time warping (DTW) (see a review for details (*53*)) to calculate the

pairwise non-linear alignment of the 12 RSV time-series (corresponding to the 12 seasons) in

each region of the US (**Fig. S2**) and quantify dissimilarity between those time-series. The DTW

algorithm computed the optimum warping path between two series under certain constraints,

including monotonicity, continuity, warping window, and boundary (*53*). The R packages *dtw*

and *dtwclust* facilitated the implementation of the algorithm and optimization (*54*).

After calculating distances, a local cost matrix (*lcm*) was generated with dimensions of $n \times m$,

where n and m represent the lengths of the pairwise time series. Considering the input time

series Q and S, for each element (i, j) of the *lcm*, the distance between Q_i and R_j was computed,

such that $lcm(i, j) = (\sum |Q_i - R_j|^p)^{1/p}$. The DTW algorithm thereby identifies the path that

minimizes the alignment between pairwise time-series Q and R by iteratively stepping through

the *lcm*, starting at $lcm(1,1)$ and finishing at $lcm(n, m)$, while aggregating the cost. At each

step, the algorithm determines the direction in which the cost increases the least under the given

constraints.

To limit the area of the *lcm* that the DTW algorithm must traverse, we implemented the Sakoe-

Chiba window as a global constraint (*54*). This constraint confines the allowed region along

the diagonal of the *lcm*. To select an optimal window size for hierarchical clustering, we

evaluated clustering using the modified Davies-Bouldin (DB) internal cluster validity index

(CVI), iterating across different values of window size from 1 to 52, corresponding to 1 to 52

weeks, while keeping cluster size as 3. For each window size, the DB CVI calculated distances

from computed cluster centroids

We subsequently performed hierarchical clustering on the distances of aligned time-series

using DTW. We utilized the Ward D2 clustering method, which minimizes the sum of squared

differences from the centroid during the merging of clusters. This hierarchical clustering of the

12 RSV epidemics in each region created a hierarchy of groups. As the level in the hierarchy

increased, clusters were formed by merging clusters from the next lower level, resulting in an

ordered sequence of groupings.

Transmission dynamic models

We used an age-stratified Susceptible-Infected-Susceptible (SIS) model, taking into account

repeat infections, to describe the transmission dynamics of RSV. The model was initially

proposed by Pitzer et al. (*29*) to study the environmental drivers of the spatiotemporal dynamics

of RSV in the US. The model assumed individuals were born with protective maternal

immunity, which waned exponentially, leaving the infants susceptible to infection. We

assumed a progressive build-up of immunity following up to four previous infections.

Following infection with RSV, individuals developed partial immunity, reducing the rate of

subsequent infection and relative infectiousness of the following infections. We also assumed

subsequent infections had a shortened recovery time compared to primary infections.

Transmission-relevant contact patterns were assumed to be frequency-dependent and were

consistent with the previous work (*29*). The model was able to reproduce the seasonal annual

or biennial patterns of RSV transmission in different regions of the US. To model influenza

transmission dynamics, we used a Susceptible-Infected-Recovered-Susceptible (SIRS) model.

We assumed waning immunity for recovered individuals, allowing influenza infection to recur

following the influenza pandemic in the 2009/10 and 2010/11 seasons.

The RSV and influenza transmission models were coupled through three hypothetical viral

interference mechanisms. The first mechanism assumed influenza infection reduced the host's

susceptibility to subsequent RSV coinfections, modulating the infection rate of susceptible

individuals (**Fig. S6A**), captured by a parameter θ, i.e. $dX_{si}/dt = -\theta\lambda_1 X_{si} + \lambda_2 X_{ss} - \gamma_2 X_{si}$

(Eq. (10) in **Supplementary Materials**). The second mechanism assumed influenza infection

reduced the infectious period (i.e. increased the rate of recovery) of subsequent RSV

coinfections (**Fig. S6B**), captured by a parameter η, i.e. $dX_{i1i}/dt = -\eta\gamma_1 X_{ii} + \lambda_1 X_{si} +$

$\lambda_2 X_{is} - \gamma_2 X_{ii}$ (Eq. (11) in **Supplementary Materials**). The third mechanism assumed

influenza infection reduced the force of infection of RSV (**Fig. S6C**), captured by a parameter

ξ, i.e. $\lambda_1 = \xi\beta_1(t)X_{ai}$, where $X_{ai,\ a = \{i1,i2,i3,i4\}}$ represents coinfection terms. The model was

described by a system of ordinary differential equations (ODEs); see **Supplementary**

Materials for details.

Model calibration

To calibrate the model parameters, we first fit the RSV dynamic model to the laboratory reports

of positive RSV specimens from 2007 to 2019. We estimated the seasonal amplitude (α_1),

seasonal offset (ϕ_1) and reporting fraction (f) for each region, respectively, using maximum

likelihood estimation. The likelihood of the data given the model was calculated by assuming

the number of positive cases in each week was Poisson-distributed with a mean equal to the

model-predicted cases times the reporting fraction. Other parameter values for the model were

adopted from (*29*), and are provided in Table 1 in the **Supplementary Materials**. For the

influenza model, we assumed the mean infectious period for primary and secondary influenza

infections is 8 days (*55*), the duration of waning immunity is 40 weeks, and the seasonal

amplitude is equal to that estimated for RSV. We started by simulating only the RSV epidemic

model, seeding the model with one RSV-infected individual in each age group except the <1

year-old age group. We used a burn-in period of 60 or 61 years, depending upon the region that

exhibits either an annual or biennial RSV pattern, to ensure the RSV model reached an

equilibrium quasi-steady state. Influenza infection was introduced to the population after the

model reached the equilibrium quasi-steady state.

To probe the effects of viral interference from pdmH1N1 infection on shaping RSV epidemics,

we applied Latin Hypercube Sampling (LHS) to generate representative samples from a wide

range of values for the parameter space $\Phi = (\Theta, \beta_2, \phi_2, \tau)$, where $\Theta = (\theta, \xi, \eta)$ represents

interference parameters; β_2 is the transmission rate of the pdmH1N1 virus; ϕ_2 is the seasonal

phase offset of influenza dynamics, and τ is the time point when influenza infection is seeded

in the population. We generated 100,000 samples from a uniform distribution $U(0,1)$ for the

parameter θ and ξ, respectively, and from a uniform distribution $U(1,3)$ for the parameter η.

We also sampled 100,000 values from a uniform distribution $U(100,110)$ for the parameter τ,

mimicking the onset time of the second H1N1 pandemic wave during the winter of 2009.

Additionally, 100,000 samples were obtained from a uniform distribution $U(2,3)$ for the

parameter β_2, based on the estimates of the basic reproduction number of the H1N1 pandemic

virus in the United States (*56*). 100,000 values of seasonal offset of influenza ϕ_2 were sampled

from a uniform distribution $U(-1,1)$.

Next, we explored the parameter space for each viral interference mechanism separately. We

generated forward simulations using the sampled 100,000 parameter sets and calculated the

likelihood to evaluate the "goodness-of-fit" of the model. The data that we used was the weekly

RSV positive tests in the 2009/10 season in Region 1 (CT: Connecticut, ME: Maine, MA:

Massachusetts, NH: New Hampshire, RI: Rhode Island, and VT: Vermont), Region 4 (AL:

Alabama, FL: Florida, GA: Georgia, KY: Kentucky, MS: Mississippi, NC: North Carolina, SC:

South Carolina, and TN: Tennessee) or Region 10 (AK: Alaska, ID: Idaho, OR: Oregon, and

WA: Washington), respectively. We focused on the 2009/10 season data because that was the

season when only the pdmH1N1 virus circulated in the population. We used the following

2010/11 season for model validation. For each set of parameters, the likelihood of the data

given the model was calculated by assuming the number of positive cases in each week was

Poisson-distributed with a mean equal to the model-predicted cases divided by corresponding

weekly scaling factors. We ranked the negative likelihood in an ascending order and

determined the distributions of parameter values (θ, ξ, η) based on the top 2% (i.e., the first

2000) models.

List of Supplementary Materials

Supplementary Figures: Fig. S1 to S12 (pdf file)

Supplementary Materials (pdf file)

Reference

1. M. Tin Tin Htar, M. S. Yerramalla, J. C. Moïsi, D. L. Swerdlow, The burden of respiratory syncytial virus in adults: a systematic review and meta-analysis. *Epidemiol. Infect.* **148**, e48 (2020).

2. T. Shi, D. A. McAllister, K. L. O'Brien, E. A. F. Simoes, S. A. Madhi, B. D. Gessner, F. P. Polack, E. Balsells, S. Acacio, C. Aguayo, I. Alassani, A. Ali, M. Antonio, S. Awasthi, J. O. Awori, E. Azziz-Baumgartner, H. C. Baggett, V. L. Baillie, A. Balmaseda, A. Barahona, S. Basnet, Q. Bassat, W. Basualdo, G. Bigogo, L. Bont, R. F. Breiman, W. A. Brooks, S. Broor, N. Bruce, D. Bruden, P. Buchy, S. Campbell, P. Carosone-Link, M. Chadha, J. Chipeta, M. Chou, W. Clara, C. Cohen, E. de Cuellar, D.-A. Dang, B. Dash-Yandag, M. Deloria-Knoll, M. Dherani, T. Eap, B. E. Ebruke, M. Echavarria, C. C. de Freitas Lázaro Emediato, R. A. Fasce, D. R. Feikin, L. Feng, A. Gentile, A. Gordon, D. Goswami, S. Goyet, M. Groome, N. Halasa, S. Hirve, N. Homaira, S. R. C. Howie, J. Jara, I. Jroundi, C. B. Kartasasmita, N. Khuri-Bulos, K. L. Kotloff, A. Krishnan, R. Libster, O. Lopez, M. G. Lucero, F. Lucion, S. P. Lupisan, D. N. Marcone, J. P. McCracken, M. Mejia, J. C. Moisi, J. M. Montgomery, D. P. Moore, C. Moraleda, J. Moyes, P. Munywoki, K. Mutyara, M. P. Nicol, D. J. Nokes, P. Nymadawa, M. T. da Costa Oliveira, H. Oshitani, N. Pandey, G. Paranhos-Baccalà, L. N. Phillips, V. S. Picot, M. Rahman, M. Rakoto-Andrianarivelo, Z. A. Rasmussen, B. A. Rath, A. Robinson, C. Romero, G. Russomando, V. Salimi, P. Sawatwong, N. Scheltema, B. Schweiger, J. A. G. Scott, P. Seidenberg, K. Shen, R. Singleton, V. Sotomayor, T. A. Strand, A. Sutanto, M. Sylla, M. D. Tapia, S. Thamthitiwat, E. D. Thomas, R. Tokarz, C. Turner, M. Venter, S. Waicharoen, J. Wang, W. Watthanaworawit, L.-M. Yoshida, H. Yu, H. J. Zar, H. Campbell, H. Nair, RSV Global Epidemiology Network, Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in young children in 2015: a systematic review and modelling study. *Lancet.* **390**, 946–958 (2017).

3. Y. Li, X. Wang, D. M. Blau, M. T. Caballero, D. R. Feikin, C. J. Gill, S. A. Madhi, S. B. Omer, E. A. F. Simões, H. Campbell, A. B. Pariente, D. Bardach, Q. Bassat, J.-S. Casalegno, G. Chakhunashvili, N. Crawford, D. Danilenko, L. A. H. Do, M. Echavarria, A. Gentile, A. Gordon, T. Heikkinen, Q. S. Huang, S. Jullien, A. Krishnan, E. L. Lopez, J. Markić, A. Mira-Iglesias, H. C. Moore, J. Moyes, L. Mwananyanda, D. J. Nokes, F. Noordeen, E. Obodai, N. Palani, C. Romero, V. Salimi, A. Satav, E. Seo, Z. Shchomak, R. Singleton, K. Stolyarov, S. K. Stoszek, A. von Gottberg, D. Wurzel, L.-M. Yoshida, C. F. Yung, H. J. Zar, Respiratory Virus Global Epidemiology Network, H. Nair, RESCEU investigators, Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in children younger than 5 years in 2019: a systematic analysis. *Lancet.* **399**, 2047–2064 (2022).

4. M. L. Boron, L. Edelman, J. R. Groothuis, F. J. Malinoski, A novel active respiratory syncytial virus surveillance system in the United States: variability in the local and regional incidence of infection. *Pediatr. Infect. Dis. J.* **27**, 1095–1098 (2008).

5. I. Kuitunen, M. Renko, Lessons to learn from the current pandemic for future non-pharmaceutical interventions against the respiratory syncytial virus – nationwide register-study in Finland. *Infect. Dis.* **53**, 476–478 (2021).

6. Z. Zheng, V. E. Pitzer, E. D. Shapiro, L. J. Bont, D. M. Weinberger, Estimation of the Timing and Intensity of Reemergence of Respiratory Syncytial Virus Following the COVID-19 Pandemic in the US. *JAMA Netw Open.* **4**, e2141779 (2021).

7. J. S. Casalegno, M. Ottmann, M. Bouscambert-Duchamp, M. Valette, F. Morfin, B. Lina,

Impact of the 2009 influenza A(H1N1) pandemic wave on the pattern of hibernal
respiratory virus epidemics, France, 2009. *Euro Surveill.* **15** (2010) (available at
https://www.ncbi.nlm.nih.gov/pubmed/20158981).

8.   T. Meningher, M. Hindiyeh, L. Regev, H. Sherbany, E. Mendelson, M. Mandelboim,
Relationships between A(H1N1)pdm09 influenza infection and infections with other
respiratory viruses. *Influenza Other Respi. Viruses.* **8**, 422–430 (2014).

9.   S. Hirsh, M. Hindiyeh, L. Kolet, L. Regev, H. Sherbany, K. Yaary, E. Mendelson, M.
Mandelboim, Epidemiological changes of respiratory syncytial virus (RSV) infections in
Israel. *PLoS One.* **9**, e90515 (2014).

10. N. Schnepf, M. Resche-Rigon, A. Chaillon, A. Scemla, G. Gras, O. Semoun, P.
Taboulet, J.-M. Molina, F. Simon, A. Goudeau, J. LeGoff, High burden of non-influenza
viruses in influenza-like illness in the early weeks of H1N1v epidemic in France. *PLoS
One.* **6**, e23514 (2011).

11. G. C. Mak, A. H. Wong, W. Y. Y. Ho, W. Lim, The impact of pandemic influenza A
(H1N1) 2009 on the circulation of respiratory viruses 2009-2011. *Influenza Other Respi.
Viruses.* **6**, e6–10 (2012).

12. L. Yang, K. H. Chan, L. K. P. Suen, K. P. Chan, X. Wang, P. Cao, D. He, J. S. M. Peiris,
C. M. Wong, Impact of the 2009 H1N1 Pandemic on Age-Specific Epidemic Curves of
Other Respiratory Viruses: A Comparison of Pre-Pandemic, Pandemic and Post-
Pandemic Periods in a Subtropical City. *PLoS One.* **10**, e0125447 (2015).

13. J. Piret, G. Boivin, Viral Interference between Respiratory Viruses. *Emerg. Infect. Dis.*
28, 273–281 (2022).

14. A. J. Gonzalez, E. C. Ijezie, O. B. Balemba, T. A. Miura, Attenuation of Influenza A Virus
Disease Severity by Viral Coinfection in a Mouse Model. *J. Virol.* **92** (2018),
doi:10.1128/JVI.00881-18.

15. K. F. Chan, L. A. Carolan, D. Korenkov, J. Druce, J. McCaw, P. C. Reading, I. G. Barr,
K. L. Laurie, Investigating Viral Interference Between Influenza A Virus and Human
Respiratory Syncytial Virus in a Ferret Model of Infection. *J. Infect. Dis.* **218**, 406–417
(2018).

16. S. Bhattacharyya, P. H. Gesteland, K. Korgenski, O. N. Bjørnstad, F. R. Adler, Cross-
immunity between strains explains the dynamical pattern of paramyxoviruses. *Proc.
Natl. Acad. Sci. U. S. A.* **112**, 13396–13400 (2015).

17. N. R. Cheemarla, T. A. Watkins, V. T. Mihaylova, E. F. Foxman, Viral Interference
During Influenza A–SARS-CoV-2 Coinfection of the Human Airway Epithelium and
Reversal by Oseltamivir. *J. Infect. Dis.*, jiad402 (2023).

18. L. Opatowski, M. Baguelin, R. M. Eggo, Influenza interaction with cocirculating
pathogens and its impact on surveillance, pathogenesis, and epidemic profile: A key
role for mathematical modelling. *PLoS Pathog.* **14**, e1006770 (2018).

19. S. Nickbakhsh, C. Mair, L. Matthews, R. Reeve, P. C. D. Johnson, F. Thorburn, B. von
Wissmann, A. Reynolds, J. McMenamin, R. N. Gunson, P. R. Murcia, Virus–virus
interactions impact the population dynamics of influenza and the common cold.
Proceedings of the National Academy of Sciences. **116**, 27142–27150 (2019).

20. W. Hendriks, H. Boshuizen, A. Dekkers, M. Knol, G. A. Donker, A. van der Ende, H.

Korthals Altes, Temporal cross-correlation between influenza-like illnesses and invasive
pneumococcal disease in The Netherlands. *Influenza Other Respi. Viruses*. **11**, 130–
137 (2017).

21.   R. E. G. Upshur, R. Moineddin, E. J. Crighton, M. Mamdani, Interactions of viral
pathogens on hospital admissions for pneumonia, croup and chronic obstructive
pulmonary diseases: results of a multivariate time-series analysis. *Epidemiol. Infect.*
134, 1174–1178 (2006).

22.   K. P. Tao, M. K. C. Chong, K. Y. Y. Chan, J. C. S. Pun, J. G. S. Tsun, S. M. W. Chow,
C. S. H. Ng, M. H. T. Wang, P. K. S. Chan, A. M. Li, R. W. Y. Chan, Suppression of
influenza virus infection by rhinovirus interference - at the population, individual and
cellular levels. *Curr. Res. Microb. Sci.* **3**, 100147 (2022).

23.   N. C. Grassly, C. Fraser, Mathematical models of infectious disease transmission. *Nat.*
Rev. Microbiol. **6**, 477–487 (2008).

24.   F. Brauer, C. Castillo-Chavez, Z. Feng, *Mathematical Models in Epidemiology* (Springer
New York).

25.   L. Pinky, H. M. Dobrovolny, Epidemiological Consequences of Viral Interference: A
Mathematical Modeling Study of Two Interacting Viruses. *Front. Microbiol.* **13**, 830423
(2022).

26.   N. R. Waterlow, S. Flasche, A. Minter, R. M. Eggo, Competition between RSV and
influenza: Limits of modelling inference from surveillance data. *Epidemics*. **35**, 100460
(2021).

27.   N. R. Waterlow, M. Toizumi, E. van Leeuwen, H.-A. Thi Nguyen, L. Myint-Yoshida, R.
M. Eggo, S. Flasche, Evidence for influenza and RSV interaction from 10 years of
enhanced surveillance in Nha Trang, Vietnam, a modelling study. *PLoS Comput. Biol.*
18, e1010234 (2022).

28.   S. C. Kramer, J.-S. Casalegno, M. Domenech de Cellès, Characterizing the
epidemiological interactions between influenza and respiratory syncytial viruses and
their implications for epidemic control. *bioRxiv* (2022), ,
doi:10.1101/2022.11.29.22282872.

29.   V. E. Pitzer, C. Viboud, W. J. Alonso, T. Wilcox, C. J. Metcalf, C. A. Steiner, A. K.
Haynes, B. T. Grenfell, Environmental drivers of the spatiotemporal dynamics of
respiratory syncytial virus in the United States. *PLoS Pathog.* **11**, e1004591 (2015).

30.   H. Sakoe, S. Chiba, Dynamic programming algorithm optimization for spoken word
recognition. *IEEE Trans. Acoust.* **26**, 43–49 (1978).

31.   A. Wu, V. T. Mihaylova, M. L. Landry, E. F. Foxman, Interference between rhinovirus
and influenza A virus: a clinical data analysis and experimental infection study. *Lancet*
Microbe. **1**, e254–e262 (2020).

32.   M. C. Swets, C. D. Russell, E. M. Harrison, A. B. Docherty, N. Lone, M. Girvan, H. E.
Hardwick, L. G. Visser, P. J. M. Openshaw, G. H. Groeneveld, M. G. Semple, J. K.
Baillie, SARS-CoV-2 co-infection with influenza viruses, respiratory syncytial virus, or
adenoviruses. *Lancet*. **399**, 1463–1464 (2022).

33.   A. Handel, L. E. Liao, C. A. A. Beauchemin, Progress and trends in mathematical
modelling of influenza A virus infections. *Current Opinion in Systems Biology*. **12**, 30–36

(2018).

34. N. Tuncer, H. Gulbudak, V. L. Cannataro, M. Martcheva, Structural and Practical
Identifiability Issues of Immuno-Epidemiological Vector-Host Models with Application to
Rift Valley Fever. *Bull. Math. Biol.* **78**, 1796–1827 (2016).

35. N. Tuncer, M. Marctheva, B. LaBarre, S. Payoute, Structural and Practical Identifiability
Analysis of Zika Epidemiological Models. *Bull. Math. Biol.* **80**, 2209–2241 (2018).

36. Y.-H. Kao, M. C. Eisenberg, Practical unidentifiability of a simple vector-borne disease
model: Implications for parameter estimation and intervention assessment. *Epidemics.*
25, 89–100 (2018).

37. R. E. Baker, S. W. Park, W. Yang, G. A. Vecchi, C. J. E. Metcalf, B. T. Grenfell, The
impact of COVID-19 nonpharmaceutical interventions on the future dynamics of
endemic infections. *Proceedings of the National Academy of Sciences.* **117**, 30547–
30553 (2020).

38. Y. N. Löwensteyn, Z. Zheng, N. Rave, M. A. G. E. Bannier, M.-N. Billard, J.-S.
Casalegno, V. E. Pitzer, J. G. Wildenbeest, D. M. Weinberger, L. Bont, Surveillance of
Pediatric REspiratory Admissions in Dutch hospitals (SPREAD) Study Group, Year-
Round Respiratory Syncytial Virus Transmission in The Netherlands Following the
COVID-19 Pandemic: A Prospective Nationwide Observational and Modeling Study. *J.*
Infect. Dis. **228**, 1394–1399 (2023).

39. J. R. Hamilton, D. Sachs, J. K. Lim, R. A. Langlois, P. Palese, N. S. Heaton, Club cells
surviving influenza A virus infection induce temporary nonspecific antiviral immunity.
Proc. Natl. Acad. Sci. U. S. A. **113**, 3861–3866 (2016).

40. C. A. Johnson-Delaney, S. E. Orosz, *Vet. Clin. North Am. Exot. Anim. Pract.*, in press.

41. Y. Cao, Y. Huang, K. Xu, Y. Liu, X. Li, Y. Xu, W. Zhong, P. Hao, Differential responses
of innate immunity triggered by different subtypes of influenza a viruses in human and
avian hosts. *BMC Med. Genomics.* **10**, 70 (2017).

42. J. van Summeren, A. Meijer, G. Aspelund, J. S. Casalegno, G. Erna, U. Hoang, B. Lina,
VRS study group in Lyon, S. de Lusignan, A. C. Teirlinck, V. Thors, J. Paget, Low levels
of respiratory syncytial virus activity in Europe during the 2020/21 season: what can we
expect in the coming summer and autumn/winter? *Euro Surveill.* **26** (2021),
doi:10.2807/1560-7917.ES.2021.26.29.2100639.

43. D. K. Yeoh, D. A. Foley, C. A. Minney-Smith, A. C. Martin, A. O. Mace, C. T. Sikazwe,
H. Le, A. Levy, C. C. Blyth, H. C. Moore, Impact of Coronavirus Disease 2019 Public
Health Measures on Detections of Influenza and Respiratory Syncytial Virus in Children
During the 2020 Australian Winter. *Clin. Infect. Dis.* **72**, 2199–2202 (2021).

44. J. Mereckiene, S. Cotter, J. T. Weber, A. Nicoll, F. D'Ancona, P. L. Lopalco, K.
Johansen, A. M. Wasley, P. Jorgensen, D. Lévy-Bruhl, C. Giambi, P. Stefanoff, L.
Dematte, D. O'Flanagan, VENICE project gatekeepers group, Influenza A(H1N1)pdm09
vaccination policies and coverage in Europe. *Euro Surveill.* **17** (2012),
doi:10.2807/ese.17.04.20064-en.

45. K. Hebbrecht, M. Stuivenga, T. Birkenhäger, M. Morrens, E. I. Fried, B. Sabbe, E. J.
Giltay, Understanding personalized dynamics to inform precision medicine: a dynamic
time warp analysis of 255 depressed inpatients. *BMC Med.* **18**, 400 (2020).

46.   J. Mellor, C. E. Overton, M. Fyles, L. Chawner, J. Baxter, T. Baird, T. Ward,
Understanding the leading indicators of hospital admissions from COVID-19 across
successive waves in the UK. *Epidemiol. Infect.* **151**, e172 (2023).

47.   L. Miralles-Pechuán, A. Kumar, A. L. Suárez-Cetrulo, Forecasting COVID-19 cases
using dynamic time warping and incremental machine learning methods. *Expert Syst.*
40 (2023), doi:10.1111/exsy.13237.

48.   J.-D. Morel, J.-M. Morel, L. Alvarez, Time warping between main epidemic time series in
epidemiological surveillance. *PLoS Comput. Biol.* **19**, e1011757 (2023).

49.   Z. Luo, L. Zhang, N. Liu, Y. Wu, Time series clustering of COVID-19 pandemic-related
data. *Data Science and Management*. **6**, 79–87 (2023).

50.   R. M. Anderson, H. Heesterbeek, D. Klinkenberg, T. D. Hollingsworth, How will country-
based mitigation measures influence the course of the COVID-19 epidemic? *Lancet*.
395, 931–934 (2020).

51.   S. Sakleshpur, A. L. Steed, Influenza: Toward understanding the immune response in
the young. *Front Pediatr*. **10**, 953150 (2022).

52.   R. M. Hamilton, R. E. Foster, T. J. Gibb, C. J. Johannsen, J. B. Santini, Pre-visible
Detection of Grub Feeding in Turfgrass using Remote Sensing. *Photogrammetric
Engineering & Remote Sensing*. **75**, 179–192 (2009).

53.   P. Senin, Dynamic time warping algorithm review (2008) (available at
https://www.semanticscholar.org/paper/Dynamic-Time-Warping-Algorithm-Review-
Senin/8096fe77e22ee54d829861642ac71064d866b913).

54.   T. Giorgino, Computing and Visualizing Dynamic Time Warping Alignments in R: The
dtw Package. *J. Stat. Softw.* **31**, 1–24 (2009).

55.   G. De Serres, I. Rouleau, M. E. Hamelin, C. Quach, D. Skowronski, L. Flamand, N.
Boulianne, Y. Li, J. Carbonneau, A. Bourgault, M. Couillard, H. Charest, G. Boivin,
Contagious period for pandemic (H1N1) 2009. *Emerg. Infect. Dis.* **16**, 783–788 (2010).

56.   L. F. White, J. Wallinga, L. Finelli, C. Reed, S. Riley, M. Lipsitch, M. Pagano, Estimation
of the reproductive number and the serial interval in early phase of the 2009 influenza
A/H1N1 pandemic in the USA. *Influenza Other Respi. Viruses*. **3**, 267–276 (2009).

Acknowledgments: This work was supported by a grant from the National Institutes of

Health (R01AI137093). The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

Author contribution:

Conceptualization; KL, DMW, VEP

Data curation; KL

Formal analysis; KL, DMW, VEP

Funding acquisition; DMW, VEP

Methodology; KL, DMW, VEP

Software; KL

Supervision; DMW, VEP

Visualization; KL

Writing - original draft; KL

Writing - review & editing; KL, DT, DMW, VEP

All authors read and approved the final manuscript.

Competing interests: DMW has been principal investigator on grants from Pfizer and

Merck to Yale University for work unrelated to this manuscript and has received consulting

and/or speaking fees from Pfizer, Merck, and GSK/Affinivax. The other authors declare no

competing interests.

Data and availability： All visualization was performed in R (version 4.0.2), and code to

produce all figures and data used in this analysis are available in the GitHub repository

https://github.com/keli5734/viral-interference.

Figures

Fig. 1. Laboratory-confirmed positive tests for pdmH1N1 virus and RSV following the pandemic.
(A) Positive laboratory tests for pdmH1N1 virus (red) and RSV (black) before (dashed line), during (shaded area) and following the 2009 influenza pandemic in Region 1 (CT: Connecticut, ME: Maine, MA: Massachusetts, NH: New Hampshire, RI: Rhode Island, VT: Vermont), Region 4 (AL: Alabama, FL: Florida, GA: Georgia, KY: Kentucky, MS: Mississippi, NC: North Carolina, SC: South Carolina, TN: Tennessee) Region 8 (CO: Colorado, MT: Montana, ND: North Dakota, SD: South Dakota, UT: Utah, WY: Wyoming) and Region 10 (AK: Alaska, ID: Idaho, OR: Oregon, WA: Washington) of the US. Dashed gray lines indicate the 1st of January each year. **(B)** Heatmap of RSV activity by epidemic season for the four selected regions. Epidemic seasons are defined as starting in July and ending in June of the following year. RSV activity was calculated as the fraction of positive tests among the total number of tests for each season. The resulting fraction was then normalized to a range between 0 and 1,

with red indicating high activity and blue indicating low activity. RSV activity during the pandemic 2009/10 season is highlighted in the black box.

Fig. 2. Hierarchical clusters and RSV time-series in the four selected regions.
An agglomerative clustering algorithm with a Ward variance method was used to group the 12 RSV epidemic seasons in each region based on the alignment paths (**Fig. S2**). The dendrograms (in the dashed box) give the clustering results based on the optimal alignments between time-series computed by DTW. **(A)-(D)** show the dendrogram and corresponding RSV time-series in Regions 1, 4, 8 and 10, respectively. RSV time-series following the pandemic seasons are highlighted in blue (2009/10) and red (2010/11). The clustering results for other regions are given in **Fig. S3**.

Fig. 3. The changes in RSV intensity and the center of gravity of RSV activity following the pandemic.

RSV intensity and the center of gravity of RSV activity were calculated for each epidemic season in each region of the US individually. For each region, the changes in the intensity of RSV activity in either the 2009/10 or 2010/11 season (i.e., query) were compared to the median values of all other seasons (i.e., reference), such that intensity changes = (query − reference) / reference × 100%. The changes in the center of gravity were given by the difference between the RSV-weighted weekly average for the query and the reference.

Fig. 4. Correlations between RSV intensity and pdmH1N1 virus intensity.
(A) The correlation between the intensity of RSV in the 2009/10 season and the intensity of pdmH1N1 virus during the pandemic period ($\rho = -0.38$), and **(B)** the correlation between the intensity of RSV in the 2010/11 season and the intensity of pdmH1N1 virus during the pandemic period ($\rho = 0$). The colors indicate the drop (brown) or rise (green) of RSV intensity in certain regions, corresponding to the US map in **Fig. 3**.

Fig. 5. Results of model fits for laboratory-confirmed RSV-positive tests in the 2009/10 and 2010/11 seasons in Region 1.

The observed number of RSV-positive tests in Region 1 is shown by the black line, while the top 2% of best-fitting models (based on 100,000 models generated from Latin Hypercube Sampling) is shown by the gray shaded region for **(A)** Model I—pdmH1N1 infection reduces susceptibility of RSV infection, **(B)** Model II—pdmH1N1 infection reduces RSV infectious period, and **(C)** Model III—pdmH1N1 infection reduces RSV infectivity. The median prediction, given by the median estimates of the viral interference parameter and associated other parameters, is indicated by the red curve. The green curve shows the scenario of no viral interference. Note that the models were only fitted to the RSV incidence data in the 2009/10 season, and the data from the 2010/11 season were used for model validation.

Fig. 6. Identified parameter space for viral interference parameters.
Histograms show the frequency of the values of viral interference parameters for the top 2% best-fitting models (2000 samples). The cumulative density functions (CDFs) are given by the solid lines, and the dashed red lines indicate the median estimates. **(A, D, G)** show the identified parameter distribution for the viral interference mechanism that reduces the host's susceptibility to RSV infection, θ (Model I), in Region 1, 10 and 4, respectively. **(B, E, H)** show the identified parameter distribution for the viral interference mechanism that shortens RSV infectious period, η (Model II), in Region 1, 10 and 4, respectively. **(C, F, I)** show the identified parameter distribution for the viral interference mechanism that reduces RSV infectivity, ξ (Model III), in Region 1, 10 and 4, respectively.