Predictors of good contraception attitude and practice among female students of television studies in Nigeria: a secondary analysis

Philip Adewale Adeoye,[1†*] Tolu Adeniji,[1] Hadizah Abigail Agbo[1,2]

[1]Department of Community Medicine, Jos University Teaching Hospital, Jos, Plateau State, Nigeria,

[2]Department of Community Medicine, University of Jos, Jos, Plateau State, Nigeria.

Corresponding author: Philip Adewale ADEOYE. Department of Community Medicine, Jos University Teaching Hospital, Jos, Plateau State, Nigeria. E-mail: philipadeoye@gmail.com.Mobile: +2348062081617

Abstract

Background: Globally contraceptive uptake among young people is increasing; so too is the high unmet need for family planning among this population, especially in sub-Sharan Africa. This study assessed attitude and practice item predictors of contraception among female students of Television studies in Nigeria.

Methods: This is a secondary analysis of a cross-sectional study among 227 female TV undergraduates in Nigeria; selected through a simple random technique by balloting. Data was analyzed using SPSS V.25 and qualitative data were presented as frequencies and proportions. Predictors of good attitude and practice were determined by multivariable logistic regression. P-value<0.05 was adjudged significant.

Results: most (91%) of respondents are aware of contraception. Most (94.9%) of the respondents have had unplanned pregnancies and only 42.1% had ever used contraception. Predictors of good attitude include good knowledge that female sterilization is one way to avoid pregnancy (aOR: 2.439 [95%CI: 1.245, 4.775]; p=0.009); increased risk of breast cancer with those taking OCPs (aOR: 2.188 [95%CI: 1.076, 4.448]; p=0.031); and opportunity for switching in case of side-effect (aOR: 2.113 [95%CI: 1.041, 4.288]; p=0.038). The predictor of good contraceptive practice is a poor perception of the adequacy of the current method being used (aOR: 3.236 [95%CI: 1.455, 7.196]; p=0.004).

Conclusion: This study shows that most respondents are aware of contraception. However, they show poor attitudes to and practice of contraception. There is a need to consolidate comprehensive sex education in all secondary and tertiary institutions; especially in non-science disciplines to improve the attitude and practice of contraception and ensure reproductive well-being and educational development of the girl-child. There is also the need for community action through community dialogue to improve poor contraceptive attitudes among young women.

Keywords: *knowledge, attitude, practice, contraception, female undergraduates*

Introduction

Almost 30 years ago since the 1994 International Conference on Population and Development, significant progress has been made in adolescent reproductive rights and health. Adolescents are more likely to marry later, delay sexual intercourse, use contraception and delay childbirth compared to 30 years ago. This progress has however been uneven within and between countries.[1]

Though sexual debt among adolescents has declined in sub-Saharan Africa, sexual activity is highest among adolescents compared to other regions; reflecting the higher rates of child marriages and early sexual debut within marriage in the region. Due to increased experimentation with sexual activity, sexually transmitted infections (STIs) have increased among young people; with Africa having the highest prevalence for genital herpes, and gonorrhoea- with females having a higher prevalence.[1]

The proportion of young women who were married before 18 years decreased in the last 30 years; with the highest declines reported in Northern Africa, Western and Southern Asia. Also, early childbearing has declined since the early 1990s. However, this remained highest in sub-Saharan Africa Latin America and the Caribbean.[1] Shorter interval pregnancies among young mothers pose further risks for both the mother and child.[2]

However, there has been a doubling of contraceptive use among adolescents compared to a decade ago- which currently stands at 21%. Latin America has the highest contraceptive use and a rapid rise in sub-Saharan Africa (from 4 to 15%) in the last 20 years. Contraception among unmarried sexually active adolescent girls is higher compared to their married counterparts.[2] Condoms account for 70% of the total modern contraceptive uptake among older sexually active adolescents; while pills and injectables account for a similar 70% of modern contraceptive use among married older adolescents.[3] Contraception is a way by which safe sex is ensured, unplanned pregnancies, early childbearing, STI, and maternal health risks are reduced.

With reduced age at menarche and onset of sexual activity, there is a heightened risk of unplanned sexual intercourse, unplanned pregnancy, unsafe abortion, severe puerperal illness, infertility maternal mortality, and adverse birth outcomes. Individual factors such as risk perception, fear of side-effects, opposition of partners, health system limitations, insufficient/lack of knowledge, sociocultural and religious factors are some of the barriers to contraception among female university undergraduates in sub-Saharan Africa.[4-8]

In Nigeria, contraceptive use increased from 6.0% in 19990 to 16.6% in 2018; with the south-west having the highest prevalence of 35.1% and the north-west with the lowest prevalence (6.7%). The north-central has a prevalence of 16.2%. In the north-central, plateau state has the highest prevalence (21.4%) of modern contraceptive use. contraceptive prevalence among sexually active unmarried women is 37%, and 28% for modern contraceptive use. Condom remains the commonest among unmarried sexually active women. There is a higher demand

and unmet need for contraception among sexually unmarried compared to married women in
Nigeria.[4]

However, appropriate knowledge and attitude should drive healthful and healthy reproductive
behaviour across populations. Among university female undergraduates who are older young
persons, the situation is the same; and in this population, knowledge, attitude and practice on
contraception is at best average.[9] It is during this period that adolescents and young people
begin to experiment with sexual activities while lacking requisite knowledge on reproductive
health.[10,11] This lack of knowledge on reproductive health issues, limited implementation of
comprehensive sex education in schools and poor parental knowledge has hindered
discussion on sexual health issues. This makes adolescents, who are coming from a restrictive
high school environment to a liberal university environment, to be vulnerable to harmful
sexual behaviours.

Though there have been many studies assessing knowledge, attitude and practice among
female undergraduates in Nigeria, there has not been any such study- to the best of the
knowledge of the researchers- specifically among female TV studies undergraduates in
Nigeria; as this population may be involved in mass media health education and risk
communication for community mobilization in the future. Also, a secondary item analysis
might provide insight into the levels of knowledge, attitude and practice on contraception
earlier reported in a previous study.[9]

Methods

Study design: this is a secondary analysis of a previously published cross-sectional study;[9]
which revealed the levels of good knowledge, attittude and practice of contaception to be
about average; with majority obtaining information on contraception from friends and the
internet; and the commonly used contraception being condom and oral contraceptive pills.
Further item analysis is to maximize the value and to provide insight into the items that were
of value in predicting good attitudes and practice of contraception among the study
population.

Study area: Nigeria is a sub-Saharan African country in the western region. It is the most
populous black nation on earth with an estimated population of 200million. There are about
64 million youths in Nigeria-about a third of the Nigerian population; with 51.6% female
population.[12]

Plateau State, located in the North-central region of Nigeria, covers a land area of 26,899
square kilometres with an estimated population of almost 5 million. It is a multi-ethnic state;
with English and Hausa being the commonly spoken languages. Because of its high altitude,
it has a near-temperate climate. It has 5 specialized higher educational institutions out of the
12 higher institutions awarding post-secondary school certificates.[13]

NTA TV College is a specialized higher educational institution established in 1980 to address
human resource challenges in the TV/broadcast industry. It is located in Rayfield, Jos-
metropolis, Plateau State. It is the only TV College offering diplomas and degrees in TV

journalism and production in Nigeria. It is currently an affiliate of Ahmadu Bello University,
Zaria, Nigeria.[14]

Study population: these are female NTA TV College students who consented to participate
in this study. The primary study shows that this study participants have an average age of
21.9 ($\pm$ 2.64) years; with most (91.2%) not living with a spouse or partner; Christian (91.2%);
almost two-thirds (58.1%) under the degree programme; and half (51%) with a monthly
income below the minimum wage; almost two-thirds are resident of the state and a majority
(76.7%) living off-campus when in school and 69.6% in the TV journalism course.[9]

Sample size determination: sample size was determined using Cochran's sample size
formula: $n=z^2pq/d^2$;[15] where n=minimum sample size; z=standard normal deviate at 95%
confidence interval=1.96; p=proportion of female undergraduates with good knowledge of
contraception=0.84;[16] q=alternate probability =1-0.84=0.16; d= precision at 5% error=0.05.
Therefore, the minimum sample size is 206. Correcting for 10% nonresponse makes the total
sample size=227.

Sampling technique: the female students were selected using a simple random sampling
technique by balloting for all six levels during classes in the college using the proportion-to-
size approach.

Study tool/data collection method: data collection was by the use of semi-structured self-
administered questionnaires following informed written consent by each selected participant.
The questionnaire has four sections: knowledge, attitude and practice of contraception and
sexual behaviour. It was piloted and pretested in a comparable institution (National Film
Institute) before the commencement of the study to detect errors, test for fidelity and
feasibility and participant understanding of the items of the questionnaire.

Data analysis: data was entered in and analyzed by SPSS V.25 (IBM, Armonk, NY, USA).
Qualitative variables were presented as frequencies and proportions. Simple logistic
regression was used to determine factors that are associated with good attitude and practice to
contraception; and to determine candidate variables for the multivariable analysis. Variables
with less than 10% probability of error were selected and added into the multivariable
omnibus model to determine the predictors of good attitude and practice of contraception. P-
value<0.05 was adjudged statistically significant.

Ethics approval and consent to participate: ethics approval was obtained from the JUTH
Research and Ethics Committee (JUTHH/DCS/IREC/127/XXXI/2619). Written informed
consent was obtained from each of the study participants before the commencement of the
study; which was voluntary and withdrawal was allowed at any stage of the research process.
There is no potential hazard to study participants.

158 **Results**

159 **Table 1: Knowledge of Contraception among female students at NTA TV College Jos**

Variable	Frequency (%)
Ever heard of contraception (n=214)	
Yes (good knowledge)	195 (91.1)
No/ I don't know (poor knowledge)	19 (8.9)
Pills are effective with 2-3 omissions (n=211)	
No (good knowledge)	59 (28.0)
Yes/I don't know (poor knowledge)	152 (72.0)
Female sterilization is one way to avoid pregnancy (n=213)	
Yes (good knowledge)	98 (46.0)
No/ I don't know (poor knowledge)	115 (54.0)
Health education is important for its use (n=215)	
Yes (good knowledge)	191 (88.8)
No/ I don't know (poor knowledge)	24 (11.2)
Pills guarantee 100% protection (n=214)	
No (good knowledge)	133 (62.1)
Yes/I don't know (poor knowledge)	81 (37.9)
Condoms do not prevent STI (n=214)	
No (good knowledge)	78 (36.4)
Yes/I don't know (poor knowledge)	136 (63.6)
Pill common side-effect include mood swings and weight gain (n=213)	
Yes (good knowledge)	104 (48.8)
No/ I don't know (poor knowledge)	109 (51.2)
Increased risk of breast cancer with those taking estrogen-containing contraceptives (n=217)	
Yes (good knowledge)	78 (35.9)
No/ I don't know (poor knowledge)	139 (64.1)
Those using depo must get injections every 3 months (n=213)	
Yes (good knowledge)	70 (32.9)
No/ I don't know (poor knowledge)	143 (67.1)
If there is a side-effect; switching might help (n=217)	
Yes (good knowledge)	70 (32.3)
No/ I don't know (poor knowledge)	147 (67.7)
Using both condoms ad pills is very effective (n=216)	
Yes (good knowledge)	124 (57.4)
No/ I don't know (poor knowledge)	92 (42.6)
Pills increase the risk of cervical cancer (n=217)	
Yes (good knowledge)	100 (46.1)
No/ I don't know (poor knowledge)	117 (53.9)

Table 1 above shows that the majority have heard about family planning; and know that health
education is important for its use. About two-thirds know that pills do not guarantee 100% protection;
and know that using both the condom and pills is very effective. Below half of the respondents know
that pills may increase the risk of breast and cervical cancers; know the common side-effects of pills
and know that female sterilization is a method of contraception. About one-third know that condoms

do prevent STIs; increase the risk of breast cancer among those using oral contraceptive pills; Depo
given every 3 months and a switch might be necessary if there is a side-effect. Only 28% know that
pills become ineffective with 2-3 omissions or missed intake.

Table 2: Attitude to contraception among female students at NTA TV College Jos

Variables	Frequency (percentage)
It should be used to limit my number of children (n=214)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	176 (82.2)
Good attitude (Strongly Agree/Agree)	38 (17.8)
Should be used to increase the time interval between my child's births (n=214)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	179 (83.6)
Good attitude (Strongly Agree/Agree)	35 (16.4)
Spacing will allow a child to be healthier (n=216)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	190 (88.0)
Good attitude (Strongly Agree/Agree)	26 (12.0)
Ideal age for having a first child is between 20 and 30 years of age (n=211)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	180 (85.3)
Good attitude (Strongly Agree/Agree)	31 (14.7)
Ideal number of children between 3 and 5 (n=212)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	181 (85.4)
Good attitude (Strongly Agree/Agree)	31 (14.6)
Contraceptives provide a sense of safety (n=211)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	176 (83.4)
Good attitude (Strongly Agree/Agree)	35 (16.6)
The method I am using is adequate (n=203)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	149 (73.4)
Good attitude (Strongly Agree/Agree)	54 (26.6)
Benefits males too (n=209)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	150 (71.8)
Good attitude (Strongly Agree/Agree)	59 (28.2)
Discussion about contraception with partner is embarrassing (n=209)	
Poor (Strongly Agree/Agree/Neutral)	165 (78.9)
Good (Disagree/Strongly Disagree)	44 (21.1)
Partner does not approve of my use of contraceptives (n=206)	
Poor (Strongly Agree/Agree/Neutral)	159(77.2)
Good (Disagree/Strongly Disagree)	47 (22.8)
Can protect the health of family and community (n=212)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	173 (81.6)
Good attitude (Strongly Agree/Agree)	39 (18.4)
Religious beliefs can prevent women from using contraceptives (n=213)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	169 (79.3)
Good attitude (Strongly Agree/Agree)	44 (20.7)
Cultural beliefs can prevent women from using contraceptives (n=211)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	179 (84.8)
Good attitude (Strongly Agree/Agree)	32 (15.2)
Partner's objection can prevent women from using contraceptives (n=216)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	180 (83.3)
Good attitude (Strongly Agree/Agree)	36 (16.7)
Change in male attitude can improve contraceptive use (n=216)	
Poor attitude (Neutral/Disagree/Strongly Disagree)	161 (74.5)
Good attitude (Strongly Agree/Agree)	55 (25.5)

171 *Table 2* above shows that the majority, or most, of the tertiary students have poor attitudes to issues

172 regarding contraception. For example, only a minority of respondents believe that contraception

173   provides a sense of safety; important in child spacing and child health; and the perception of the
174   adequacy of the contraception in use.

176   **Table 3: Practice of contraception among female students at NTA TV College Jos**

Variables	Frequency (percentage)
How often do you visit the healthcare centre for contraception (n=215)	
Poor practice (usually/sometimes/seldom/never)	73 (34.0)
Good practice (Always)	142 (66.0)
Uses contraceptives to prevent unplanned pregnancy (n=215)	
Poor practice (usually/sometimes/seldom/never)	103 (47.9)
Good practice (Always)	112 (52.1)
Ever had unplanned pregnancy due to contraceptive non-use (n=215)	
Poor (Always/usually/sometimes/seldom)	204 (94.9)
Good practice (Never)	11 (5.1)
Do you use contraceptives when you do not intend to get pregnant (n=215)	
Poor practice (usually/sometimes/seldom/never)	103 (47.9)
Good practice (Always)	112 (52.1)
Use different types of contraceptives (n=215)	
Poor practice (usually/sometimes/seldom/never)	64 (29.8)
Good practice (Always)	151 (70.2)
Current method changes from time to time (n=211)	
Poor (Always/usually/sometimes/seldom)	197 (93.4)
Good practice (Never)	14 (6.6)
Practice traditional/natural methods when not using any contraceptive (n=215)	
Poor practice (usually/sometimes/seldom/never)	68 (31.6)
Good practice (Always)	147 (68.4)
Ever used or currently using contraception (n=214)	
No	124 (57.9)
Yes	90 (42.1)

Table 3 above shows that more than two-thirds always visit health facilities for contraception; always
use different types of contraceptives per time; and always practice natural methods when not using
any modern method. About half always use contraceptives to prevent unplanned pregnancy or when
they do not want to get pregnant. Only about 5.1% had never had an unplanned pregnancy due to non-
use, and 6.6% had never changed their current contraceptive from time to time.

Table 4 below shows individuals who know that female sterilization is a method of contraception are
twice significantly more likely to have a good attitude towards contraception compared to those who
do not know. Those who know that there is an increased risk of breast cancer among those using
OCPs are twice significantly more likely to have a good attitude towards contraception compared to
those who do not know. Those who know that switching to another contraceptive might help with
dealing with the side-effects of a method are twice significantly more likely to have a good attitude
towards contraception compared to those who do not know.

195 **Table 4: knowledge predictors of good attitude to contraception among female students at NTA TV**
196 **College Jos**

VARIABLE	β	OR (95% CI)	*P-value*	β	aOR (95% CI)	*P-value*
Ever heard of contraception			*0.151*			*0.103*
good knowledge		0.490			0.354	
poor knowledge (ref)	-0.714	(0.185, 1.297)		- 1.024	(0.105, 1.229)	
Pills are effective with 2-3 omissions			*0.078*** \#			*0.178*
good knowledge		1.725			1.640	
poor knowledge (ref)	0.545	(0.940, 3.168)		0.495	(0.799, 3.367)	
Female sterilization is one way to avoid pregnancy			*0.000*** \#			*0.009**
good knowledge		3.356			2.439	
poor knowledge (ref)	1.211	(1.910, 5.897)		0.891	(1.245, 4.775)	
Health education is important for its use			*0.063* \#			*0.298*
good knowledge		2.403			1.775	
poor knowledge (ref)	0.877	(0.953, 6.059)		0.574	(0.602, 5.236)	
Pills guarantee 100% protection			*0.434*			-
good knowledge		0.802				
poor knowledge (ref)	-0.221	(0.461, 1.395)		-	-	
Condoms do not prevent STI			*0.424*			-
good knowledge		0.796				
poor knowledge (ref)	- 0.228	(0.455, 1.393)		-	-	
Pill common side-effect include mood swings and weight gain			*0.012*** \#			*0.609*
good knowledge		2.011			1.194	
poor knowledge (ref)	0.699	(1.165, 3.474)		0.177	(0.606, 2.352)	
Increased risk of breast cancer with those taking estrogen-containing contraceptives			*0.002*** \#			*0.031**
good knowledge		2.444			2.188	
poor knowledge (ref)	0.893	(1.383, 4.316)		0.783	(1.076, 4.448)	
Those using Depo must get injections every 3 months			*0.005*** \#			*0.130*
good knowledge		2.334			1.722	
poor knowledge (ref)	0.848	(1.299, 4.192)		0.543	(0.853, 3.476)	
If there is a side-effect; switching might help			*0.000*** \#			*0.038**
good knowledge		3.650			2.113	
poor knowledge (ref)	1.295	(1.993, 6.685)		0.748	(1.041, 4.288)	

199 **Table 5** below shows that those who have poor attitudes towards the adequacy of their current
200 contraception are three times significantly more likely to have good contraceptive practices compared
201 to those who have good attitudes.

207 **Table 5: knowledge and attitude predictors of good contraception practice among female students at NTA**
208 **TV College Jos**

VARIABLE	B	OR (90% CI)	*P-value*	β	aOR (95% CI)	*P-value*
Ever heard of contraception						
good knowledge		0.928				
poor knowledge (ref)	-0.075	(0.361, 2.384)	*0.877*	-	-	-
Pills are effective with 2-3 omissions						
good knowledge		0.740				
poor knowledge (ref)	-0.302	(0.405, 1.352)	*0.327*	-	-	-
Female sterilization is one way to avoid pregnancy						
good knowledge		1.191				
poor knowledge (ref)	0.175	(0.694, 2.043)	*0.877*	-	-	-
Health education is important for its use						
good knowledge		1.032				
poor knowledge (ref)	0.031	(0.442, 2.412)	*0.825*	-	-	-
Pills guarantee 100% protection						
good knowledge		0.720				
poor knowledge (ref)	-0.329	(0.413, 1.254)	*0.942*	-	-	-
Condoms do not prevent STI						
good knowledge		1.767			1.702	
poor knowledge (ref)	0.570	(1.006, 3.106)	*0.048**#	0.532	(0.885, 3.275)	*0.111*
Pill common side-effect include mood swings and weight gain						
good knowledge		1.868			1.520	
poor knowledge (ref)	0.625	(1.084, 3.217)	*0.042**#	0.419	(0.778, 2.970)	*0.221*
Increased risk of breast cancer with those taking estrogen-containing contraceptives						
good knowledge		1.829			1.307	
poor knowledge (ref)	0.604	(1.041, 3.212)	*0.036**#	0.268	(0.654, 2.611)	*0.448*
Those using Depo must get injections every 3 months						
good knowledge		1.701			1.152	
poor knowledge (ref)	0.532	(0.953, 3.039)	*0.073*#	0.141	(0.580, 2.287)	*0.687*
If there is a side-effect; switching might help						
good knowledge		2.077			1.333	
poor knowledge (ref)	0.731	(1.159, 3.723)	*0.014**#	0.287	(0.670, 2.654)	*0.413*
Using both condoms and pills is very effective						
good knowledge		2.410			1.650	
poor knowledge (ref)	0.880	(1.386, 4.190)	*0.000**#	0.501	(0.849, 3.206)	*0.140*
Pills increase the risk of ovarian, endometrial or cervical cancer						
good knowledge		1.279				
poor knowledge (ref)	0.246	(0.749, 2.183)	*0.002**#	- 0.368	-	-
It should be used to limit the number of children						
poor attitude		1.163				
good attitude (ref)	0.151	(0.576, 2.346)	*0.674*	-	-	-

Model characteristics: Omnibus Test of Model Coefficient (X^2; df):33.063;10; p=0.000; -2log likelihood=227.560; Cos & Snell R^2=0.161; Nagelkerke R^2=0.215; Hosmer-Lemeshow R^2 p=0.228; percentage accuracy=68.1%.
*=significance at p<0.05; #=candidate variable for multivariable regression

212   **Table 5 (contd): knowledge and attitude predictors of good contraception practice among female students**
213   **at NTA TV College Jos**

VARIABLE	B	OR (90% CI)	P-value	β	aOR (95% CI)	P-value
Should be used to increase the time interval between my child's births			*0.500*			
poor attitude		1.284				
good attitude (ref)	0.250	(0.621, 2.656)		-	-	-
Spacing will allow a child to be healthier			*0.604*			
poor attitude		1.243				
good attitude (ref)	0.217	(0.546, 2.827)		-	-	-
Ideal age for having a first child is between 20 and 30 years of age			*0.541*			
poor attitude		1.269				
good attitude (ref)	0.239	(0.591, 2.729)		-	-	-
Ideal number of children between 3 and 5			*0.012*[*#]*			
poor attitude		2.886			1.532	
good attitude (ref)	0. 060	(1.260, 6.611)		0.427	(0.555, 4.231)	*0.411*
Contraceptives provide a sense of safety			*0.282*			
poor attitude		1.494				
good attitude (ref)	0.401	(0.719, 3.106)		-	-	-
The method I am using is adequate			*0.001*[*#]*			
poor attitude		2.791			3.236	
good attitude (ref)	0.089	(1.536, 5.748)		0.174	(1.455, 7.196)	*0.004**
Benefits males too			*0.211*			
poor attitude		0.679				
good attitude (ref)	- 0.388	(0.370, 1.246)		-	-	-
Discussion about contraception with partner is embarrassing			*0.521*			
poor attitude		0.804				
good attitude (ref)	-0.219	(0.412, 1.566)		-	-	-
Partner does not approve of my use of contraceptives			*0.707*			
poor attitude		1.134				
good attitude (ref)	0.125	(0.290, 2.180)		-	-	-
Can protect the health of family and community			*0.173*			
poor attitude		1.633				
good attitude (ref)	0.490	(0.807, 3.303)		-	-	-
Religious beliefs can prevent women from using contraceptives			*0.213*			
poor attitude		0.653				
good attitude (ref)	-0.427	(0.333, 1.278)		-	-	-
Cultural beliefs can prevent women from using contraceptives			*0.168*			
poor attitude		0.580				
good attitude (ref)	-0.544	(0.268, 1.257)		-	-	-
Partner's objection can prevent women from using contraceptives			*0.092*[*#]*			
poor attitude		0.529			0.449	
good attitude (ref)	-0.637	(0.252, 1.109)		0.800	(0.183, 1.102)	*0.080*
Change in male attitude can improve contraceptive use			*0.063*[*#]*			
poor attitude		0.552			0.486	
good attitude (ref)	-0.594	(0.295, 1.033)		0.722	(0.221, 1.069)	*0.073*

Discussion

Most of the respondents, in this study, had heard of contraception. This is similar to those
observed among similar populations in Dodoma-Tanzania, states in North-Central Nigeria,
Thohoyandou-South Africa, and Suva-Fiji.[5,6,17-19] This may be due to increased sensitization
and community mobilization on contraception in most countries of the world; especially in
lower- and middle-income countries (LMICs).[20] This will need to be sustained especially in
hard-to-reach regions of the world.

Furthermore, almost half of the respondents know that female sterilization is an option to
avoid pregnancy. Lower values and similarly average proportions have been reported similar
proportion in Dodoma- Rwanda and Nigeria among similar populations.[4,5] This variation may
be due to limited practical in-depth details about contraception; especially at the grassroots.

Health education has been reported by the majority of these study respondents as important
for successful contraceptive use. This is similar to those observed in Kigali-Rwanda,
Gaborone-Botswana and Suva-Fiji among similar populations.[6,21,22] This shows a
corresponding enthusiasm that a very high awareness might have created.

Below average of respondents know that risks associated with OCP include breast, and
cervical cancers. A similar observation was reported among similar populations in Suva-Fiji.[6]
However, higher proportions were reported in Thohoyandou- South Africa.[18] This may be
due to improved reporting and access to some cancer screening services in the Southern
Africa region in Sub-Saharan Africa.[23]

About a third of respondents know that switching may help if there is a side-effect; a higher
proportion of knowledge was reported among women in Suva-Fiji.[6] Despite this, only a small
proportion of contraceptive users switch among Nigerian unmarried sexually active women
in a recent survey.[4] This might be due to improved selection of appropriate contraceptives for
women of reproductive age group which had probably minimized the side-effects associated
with contraceptives.

Most respondents of this study have poor attitude towards contraception across contraceptive
attitude. This is similar to the attitude of undergraduates in Pa Nang-Vietnam and
Malaysia.[24,25] This may be as are result of socio-cultural factors that still surround
contraception in many LMiCs.[20]

The majority of respondents do not believe that spacing will make their children healthy; and
found it unnecessary to increase birth intervals. Contrary opinion was expressed by an equal
proportion of women in Suva -Fiji.[6] A similar opinion in support of definite birth spacing was

reported in Senegal, where *Nef* (short birth interval) is frowned upon and stigmatizing; and in
line with evidence of adverse birth outcomes for the mother and child.[26] Thus, there is a need
for culture-specific tailoring of contraceptive education and mobilization; as not all caps fit
all.

Almost three-quarters do not believe that contraceptives benefit males too. This is contrary to
that which is observed in similar populations in Kigali-Rwanda and Suva-Fiji;[6,22] where more
than three-quarters have good attitude that males benefit also from contraception. This may
explain why contraception is thought to be in the main interest of the woman and some
Nigerian men will be involved in family planning programmes to show love and support for
their wives.[7]

Majority of the female undergraduates in the study site disagree that the ideal age for the
onset of childbearing should be between 20 and 30 years of age. This is, however, different
for the majority of women in Suva-Fiji.[6] This might be due to the widespread teenage
pregnancies; which are being normalized in many states in Northern Nigeria childhood and
teenage pregnancies and marriage also have the backing of cultural and religious factors in
many communities in Nigeria. The increasing numbers of out-of-school teenagers in Nigeria
might have also contributed to the normalizing of teenage pregnancies and marriages in
Nigeria.[28]

From this study, a majority do not believe that their current contraceptive is adequate. This is
contrary to the perception expressed by a similar population in Suva-Fiji.[6] This might be
responsible for the use of different types and frequent changes of contraceptives from time to
time among the study population which was not seen among similar populations in Suva-Fiji.
Therefore, there may be a need for increased confidence in contraception via community
mobilization and participation in contraception activities.

This study shows that many of the study participants visit the healthcare centres for
contraceptive uptake. This might be due to the freely available contraceptives at these public
healthcare facilities in the state in which the study was done.[29] This is however contrary to
those by contrary to those reported by similar populations in Dodoma-Tanzania, Suva-Fiji
and women of reproductive age in many LMiCs.[4,5,7] This might also be due to cultural and
social factors surrounding procurement of contraceptives by young people from the public
sector   which might be ebbing in places where contraceptives are procured from public
places. Procuring contraceptives by this population from sites outside the public sector may
affect accurate reporting of contraceptive use as these sites are not routinely captured in
national data.[30]
Most of the study participants have had unplanned pregnancies due to the non-use of
contraception. This is, however, contrary to a similar population in Suva-Fiji whose majority
had never had unplanned pregnancy.[6] This might be responsible for the use and intention to
use. Contraception, subsequently, by more than half of the respondents. There is therefore the
need to strengthen the Federal Ministry of Education's family life and HIV/AIDS Education
early in the education sector to prevent the adverse effects of teenage pregnancies.[31] There is

also the need to engage and commit families in the design and implementation of
comprehensive sex education.[8]

Majority of the study participants practised traditional/natural methods (TCM) when not
using any modern contraception. This is similar to those reported among similar populations
in Kilimanjaro-Tanzania and Suva-Fiji.[6,32] This is however contrary to those reported among
similar populations in Kigali-Rwanda, Romania, and Kano-Nigeria.[33-35] This may be as a
result of unmet needs, safety concerns, side-effect and opposition to modern contraceptive
use by partners; and the effectiveness of TCM.[35,36]

Less than half of respondents have ever used contraception among the study population. This
is similar to those reported among similar populations in Kilimanjaro- Tanzania and
Nigeria.[4,32] This is however different for similar populations in Kigali-Rwanda, Romania and
Makurdi- Nigeria.[19,22,33] This may be due to social and cultural, health systems barriers that
hinder young people from accessing contraceptive services in many LMiCs.[4,20] This might
have also explained the higher number of unplanned pregnancies experienced by this study
population.[20,23]

This study shows that knowing that female sterilization is one way to prevent pregnancy is a
prediction of good contraceptive attitude. Though studies on female sterilization are generally
low;[37] it can be hypothesized that a high knowledge of female sterilization is correlated with
a high level of knowledge of other contraceptive methods because knowledge about female
sterilization is lower compared to other methods in many populations. Increased knowledge
of female sterilization is often associated with increased acceptance and uptake, all things
being equal it has been shown compared to other methods, to improve mood and sex life.[22, 37]
Generally, it has been said to be the most common family planning method globally among
women of reproductive group.[38]

The knowledge of an increased risk of breast cancer with those taking oral contraceptive pills
is a significant predictor of good contraceptive attitude. This might be due to the awareness of
side-effects and options available to reduce such side effects. Thus, improved knowledge of
the appropriate contraceptive options in the presence of risks boosts the client's confidence
and self-efficacy on the safety and use of contraception; while awareness breeds fear which
often leads to non-use or discontinuation from misinformation, disinformation or incomplete
information.[20]
The knowledge that switching might help in the occurrence of side effects is a significant
predictor of good contraceptive attitude. This may increase improved sense of safety; self-
efficacy and perception that clients have control over their reproductive health. The need to
conceive in the near future, fewer side-effects and the expensive nature of long-acting
contraceptives are some of the reasons why long-acting reversible contraceptives have low
patronage compared to the short-acting contraceptive methods in many LMiCs.[20]
The doubt that the method clients are using is adequate is a significant predictor of good
contraceptive use. This might be due to past unplanned pregnancies reported in this study
population and frequent changes and use of different types of contraceptives among this

population. This might have improved attendance at follow-ups after contraceptive uptake where information on correct use will be consolidated and prompt report of side-effects and cross-confirmation of any contraceptive information shared might have been responsible for good practice despite less confidence in the method being used. However, the health system should continue to boost self-confidence in this population to ensure sustainability of good contraceptive use.

This study cannot be generalized to the total female undergraduate population in other higher institutions since the population is from a specific discipline that may be involved in health communication and education via mass media in the future. The use of questionnaires might bias participant's responses as sensitive questions are included in the items. However, respondents were provided a private setting, adequate preamble, use of anonymous research assistants and ethical conduct of the interview made respondents relaxed to answer the questions to the best of their abilities. Also, due to the cross-sectional nature of the study, temporality cannot be established. Our findings have been one of the few among similar populations in literature and have huge policies for health promotion; now and in the future. This is a cross-sectional study and therefore causality cannot be established

Conclusion

This study showed that though most respondents are aware of contraception, they generally show poor attitude to contraception. Majority had a history of unplanned pregnancies and few had ever used contraception; though with a willingness to do so. There is, therefore, the need to consolidate early and appropriate family life and HIV/AIDS education in all levels of formal education to improve family planning knowledge, attitude and practice in many young people. There is also a need for community dialogue to improve the poor contraceptive attitude and the need to involve families and communities in the design and implementation of contraception activities. This will ensure reproductive health and all-round development of the girl-child.

HIGHLIGHTS

- Poor attitude and practice of contraception, has been one of the main driven driving up unsustainable population growth in low- and middle-income countries.
- Most of the study participants have had past history of unplanned pregnancy
- Less than half have ever used contraception
- Knowledge of female sterilization, risk of breast cancer and opportunity of switching predicts good attitude
- Perception about the adequacy of current method is a significant predictor of good contraceptive practice

Acknowledgements

We are grateful to the students of NTA TV College of Nigeria for their willingness to participate in
the study. We also grateful for the research assistants that patiently carried out the data collection.

Funding

No funding received for this study

Conflict of interest

Authors declare no conflict of interest

Contributions

Conceptualization and design: PAA, HAO;

Data analysis and interpretation: PAA;

Drafting and critical review: HAO, PAA, TA;

Final approval: HAO, PAA, TA.

References

1.   Liang M, Simelane S, Fortuny Fillo G, Chalasani S, Weny K, Salazar Canelos P, et al. The
State of Adolescent Sexual and Reproductive Health. Journal of Adolescent Health.
2019;65(6):S3-S15.

2.   Kozuki N, Lee A, Silveira M, Victora C, Adair L, Humphrey JN, Robert, et al. The
associations of birth intervals with small-for-gestational-age, preterm, and neonatal and infant
mortality: a meta-analysis. BMC Public Health. 2013;13(Suppl 3):S3.

3.   Loaiza E, Luchsinger G, Liang M. Universal Access to Reproductive Health: Progress and
Challenges. Report. New York: UNFPA; 2016.

4.   Nigeria Population Commission. Nigeria demographic and health survey 2018. Abuja,
Nigeria: NPC, ICF; 2019.

5.   Kara WSK, Benedicto M, Mao J. Knowledge, Attitude, and Practice of Contraception
Methods Among Female Undergraduates in Dodoma, Tanzania. Cureus. 2019;11(4):e4362.

6.   Lincoln J, Mohammadnezhad M, Khan S. Knowledge, attitudes, and practices of family
planning among women of reproductive age in Suva, Fiji in 2017. Journal of Women's Health
Care. 2018;7(3):431.

7. Bradley SEK, Shiras T. Where Women Access Contraception in 36 Low- and Middle-Income
Countries and Why It Matters. Global Health: Science and Practice. 2022;10(3):e2100525.

8. Wekesah FM, Nyakangi V, Onguss M, Njagi J, Bangha M. Comprehensive Sexuality
Education in Sub-Saharan Africa. Nairobi, Kenya: African Population and Health Research
Center; 2019.

9. Agbo HA, Adeoye PA, Yilzung DR, Mangut JS, Ogbada PF. Levels and predictors of
knowledge, attitude and practice of contraception among female TV undergraduates in
Nigeria: a cross-sectional study. medRxiv. 2024.

10. Azzopardi PS, Hearps SJC, Francis KL, Kennedy EC, Mokdad AH, Kassebaum NJ, et al.
Progress in adolescent health and wellbeing: tracking 12 headline indicators for 195 countries
and territories, 1990-2016. Lancet. 2019;393(10176):1101–18.

11. Weiss HA, Ferrand RA. Improving adolescent health: an evidence-based call to action.
Lancet. 2019;393(10176):1073-5.

12. National Buerau of Statistics, Federal Ministry of Youth Development. National Baseline
Youth Survey 2012. Abuja, Nigeria: National Bureau of Statistics; 2012.

13. Plateau State ICT Development Agency. At a Glance: About Plateau State Jos, Nigeria:
Plateau State Government; 2020 [Available from: https://www.plateaustate.gov.ng/plateau/at-
a-glance.

14. NTA Television College. About Us Jos, Nigeria: NTA Television College,; 2021 [Available
from: https://ntatvc.edu.ng/?page_id=2876.

15. Bolarinwa OA. Sample size estimation for health and social science researchers: The
principles and considerations for different study designs. Nigerian Postgraduate Medical
Journal. 2020;27(2):67-75.

16. Eniojukan J, Ijeoma O, Okinedo P. Knowledge, perception and practice of contraception
among staff and students in a University Community in Delta State, Nigeria. Pharmaceutical
and Biosciences Journal. 2015;4(1):71-81.

17. Abdul I, Imohagene AO, L , Adesina K, Oguntoye M, Popoola G. Awareness, knowledge and
contraceptive use among female students in University of Ilorin, Nigeria. Journal of
Epidemiological Society of Nigeria. 2018;2(2):1-8.

18. Raselekoane RN, Chinyakata R, Gwatimba L. An investigation of the knowledge, attitudes
and use of contraceptives by youth development female students at the University of Venda. J
Hum Ecol. 2018;61(1-3):1-8.

19. Utoo B, Eka P, Utoo P, Unazi E, Maanongun M. Sexual Activity and Contraceptive use
among University Female Undergraduates in Makurdi, North-Central Nigeria. Western
Journal of Medical and Biomedical Sciences. 2020;1(2):132-8.

20. Sedgh G, Ashford LS, Hussain R. Unmet need for contraception in developing countries:
examining women's reasons for not using a method. New York: Guttmacher Institute; 2016.

21. Hoque ME, Ntsipe T, Mokgatle-Nthabu M. Awareness and practices of contraceptive use
among university students in Botswana. SAHARA J. 2013;10(2):83-8.

22. Sitini B, Okova R, Nizeyimana F, Rutayisire E. Knowledge, Attitude and Practices towards
contraceptive Use among Unmarried Females Students at Mount Kenya University Rwanda.
Journal of Clinical Review & Case Reports. 2020;5(12):520-7.

23.   Adeoye PA. Epidemiology of Breast Cancer in Sub-Saharan Africa. In: Selim Sözen and
Seyfi Emir, editors. Breast Cancer Updates. London, United Kingdom: IntechOpen Limited;
2023.

24.   Ngoc TA, Sen HTN. Knowledge and Attitude Towards Contraceptive Methods Among First-
Year Nursing Students in Da Nang, Vietnam. Ramathibodi Medical Journal. 2022;45(3):23-
32.

25.   Oo MS, Ismail NBM, Ean WR, Hamid HA, Affendi NR. Knowledge, attitude and perception
of contraception among medical students in Universiti Putra Malaysia. Malaysian Journal of
Public Health Medicine. 2019;19(2):11-9.

26.   Duclos D, Cavallaro FL, Ndoye T, Faye SL, Diallo I, Lynch CA, et al. Critical insights on the
demographic concept of "birth spacing": locating Nef in family well-being, bodies, and
relationships in Senegal. Sexual and Reproductive Health Matters. 2019;27(1):136-45.

27.   Adelekan A, Omoregie P, Edoni E. Male involvement in family planning: challenges and way
forward. International Journal of Population Research. 2014;2014(Article ID 416457):9.

28.   Joel M. Understanding Nigeria's teenage pregnancy burden Johannesburg: Africa Check;
2021 [Available from: https://africacheck.org/fact-checks/factsheets/factsheet-understanding-
nigerias-teenage-pregnancy-burden.

29.   McGuire C, Calhoun LA-O, Mumuni T, Maytan-Joneydi A, Odeku M, Speizer IS.
Government stakeholders' perspectives on the family planning environment in three Nigerian
cities: qualitative findings from the Nigerian Urban Reproductive Health Initiative (NURHI)
Sustainability Study. Global health action. 2020;13(1):1847821.

30.   Cunningham M, Bock A, Brown N, Sacher S, Hatch B, Inglis A, et al. Estimating
Contraceptive Prevalence Using Logistics Data for Short-Acting Methods: Analysis Across
30 Countries. Global Health: Science and Practice. 2015;3(3):462-81.

31.   Shiffman J, Kunnuji M, Shawar YR, Robinson RS. International norms and the politics of
sexuality education in Nigeria. Globalization and health. 2014;14(1):63.

32.   Sweya MN, Msuya SE, Mahande MJ, Manongi R. Contraceptive knowledge, sexual behavior,
and factors associated with contraceptive use among female undergraduate university students
in Kilimanjaro region in Tanzania. Adolescent health, medicine and therapeutics. 2016;7:109-
15.

33.   Blidaru IE, Furau G, Socolov D. Female Romanian university students' attitudes and
perceptions about contraception and motherhood. The European Journal of Contraception &
Reproductive Health Care. 2016;21(1):39-48.

34.   Mbarushimana V, Ntaganira J. Knowledge and attitude to female condom use among
undergraduates of Kigali Health Institute. Rwanda Journal of Health Sciences. 2013;2(1):16-
25.

35.   Rabiu A, Rufa'i AA. The role of traditional contraceptive methods in family planning among
women attending primary health care centers in Kano. Annals of African medicine.
2018;17(4):189–95.

36.   Bertrand JT, Ross JA, Sauter SR. Trends in contraceptive method mix among adolescents and
youth aged 15-24 in low- and middle-income countries. Frontiers in Global Women's Health.
2022;3:1061648.

37.   Kohls F, Kuehnle E, Brodowski L, Staboulidou I. Female sterilization as a method of
490   contraception: women's acceptance and knowledge-a review. Medical Research Archives.
491   2017;5(9).

492   38.   Sung S, Abramovitz A. Tubal Ligation. Treasure Island (FL): StatPearls Publishing; 2019.

493