Association analysis between an epigenetic alcohol risk score and blood pressure

Helena Bui, BA[1,2]*; Amena Keshawarz, PhD, MPH[1,2] *; Mengyao Wang, MS[3]*; Mikyeong Lee, PhD[4]; Scott M. Ratliff, MS[5]; Lisha Lin, MPH[5]; Kira S. Birditt, PhD[6]; Jessica D. Faul, PhD[6]; Annette Peters, PhD[7,8,9] Christian Gieger, PhD[9,10]; Thomas Delerue, PhD[10]; Sharon L. R. Kardia, PhD[5]; Wei Zhao[5], PhD; Xiuqing Guo, PhD[11]; Jie Yao, MS[11]; Jerome I. Rotter, MD[11]; Yi Li[3]; Xue Liu[3]; Dan Liu, PhD[12]; Juliana F. Tavares, MS[12]; Gökhan Pehlivan, MD[12]; Monique M.B. Breteler, MD, PhD[12,13]; Irma Karabegovic, DSc[14]; Carolina Ochoa-Rosales, PhD[14,15]; Trudy Voortman, PhD[14,16]; Mohsen Ghanbari, PhD[14]; Joyce B.J. van Meurs, PhD[17]; Mohamed Kamal Nasr, MS[18,19]; Marcus Dörr, MD[19,20]; Hans J. Grabe, MD[21,22]; Stephanie J. London, MD, DrPH[4]; Alexander Teumer, PhD[18,19,23]; Melanie Waldenberger, PhD[9,10]; David R. Weir, PhD[6]; Jennifer A. Smith, PhD[5,6]; Daniel Levy, MD[1,2] †; Jiantao Ma, PhD[24] †; Chunyu Liu, PhD[2,3] †

1. Population Sciences Branch, Division of Intramural Research, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA
2. Framingham Heart Study, Framingham, MA, USA
3. Department of Biostatistics, Boston University School of Public Health, Boston, MA
4. Epidemiology Branch, National Institute of Environmental Health Sciences, National Institutes of Health, Department of Health and Human Services, Research Triangle Park, NC, USA
5. Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, MI
6. Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, MI
7. Institute of Epidemiology, Helmholtz Munich, German Research Center for Environmental Health, German
8. Institute for Medical Informatics, Biometrics and Epidemiology, Ludwig-Maximilians-Universität München, Munich, Germany
9. German Center for Cardiovascular Research (DZHK), Partner Site Munich Heart Alliance, Munich, Germany
10. Research Unit Molecular Epidemiology, Institute of Epidemiology, Helmholtz Munich, German Research Center for Environmental Health, Bavaria, Germany
11. The Institute for Translational Genomics and Populations, Department of Pediatrics, Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA 90502, USA
12. Population Health Sciences, German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany
13. Institute for Medical Biometry, Informatics and Epidemiology (IMBIE), Faculty of Medicine, University of Bonn, Bonn, Germany
14. Department of Epidemiology, Erasmus MC University Medical Center, Rotterdam, The Netherlands
15. Centro de Vida Saludable de la Universidad de Concepción, Concepción, Chile
16. Division of Human Nutrition and Health, Wageningen University and Research, Wageningen, The Netherlands
17. Department of Internal Medicine, Erasmus MC University Medical Center, Rotterdam, The Netherlands
18. Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany
19. German Center for Cardiovascular Research (DZHK), Partner Site Greifswald, Greifswald, Germany
20. Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany
21. Department of Psychiatry and Psychotherapy, University Medicine, Greifswald, Germany
22. German Center of Neurodegenerative Diseases (DZNE), Rostock/Greifswald, site Greifswald, Germany
23. Department of Population Medicine and Lifestyle Diseases Prevention, Medical University of Bialystok, Bialystok, Poland
24. Gerald J. and Dorothy R. Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA, USA

* Denotes shared first authorship
† Denotes multiple corresponding authors

Corresponding Author(s):

Daniel Levy, MD
Framingham Heart Study
73 Mt. Wayte Avenue
Framingham, MA 01702
levyd@nhlbi.nih.gov
508-935-3458

Jiantao Ma
Gerald J. and Dorothy R. Friedman School of Nutrition Science and Policy, Tufts University
150 Harrison Avenue
Boston, MA 02111
jiantao.ma@tufts.edu

Chunyu Liu
Boston University School of Public Health, Boston University
715 Albany Street
Boston, MA 02118
liuc@bu.edu
(617) 638-5104

Abstract

Background

Epigenome-wide association studies have revealed multiple DNA methylation sites (CpGs) associated with alcohol consumption, an important lifestyle risk factor for cardiovascular diseases.

Results

We generated an alcohol consumption epigenetic risk score (ERS) based on previously reported 144 alcohol-associated CpGs and examined the association of the ERS with systolic blood pressure (SBP), diastolic blood pressure (DBP), and hypertension (HTN) in 3,898 Framingham Heart Study (FHS) participants. We found an association of alcohol intake with the ERS in the meta-analysis with 0.09 units higher ERS per drink consumed per day ($p < 0.0001$). Cross-sectional analyses in FHS revealed that a one-unit increment of the ERS was associated with 1.93 mm Hg higher SBP ($p = 4.64E-07$), 0.68 mm Hg higher DBP ($p = 0.006$), and an odds ratio of 1.78 for HTN ($p < 2E-16$). Meta-analysis of the cross-sectional association of the ERS with BP traits in eight independent external cohorts (n = 11,544) showed similar relationships with blood pressure levels, i.e., a one-unit increase in ERS was associated with 0.74 ($p = 0.002$) and 0.50 ($p = 0.0006$) mm Hg higher SBP and DBP, but could not confirm the association with hypertension. Longitudinal analyses in FHS (n = 3,260) and five independent external cohorts (n = 4,021) showed that the baseline ERS was not associated with a change in blood pressure over time or with incident HTN.

Conclusions

Our findings provide proof-of-concept that utilizing an ERS is a useful approach to capture the recent health consequences of lifestyle behaviors such as alcohol consumption.

Introduction

Approximately 178,307 people die annually from alcohol-related causes, making alcohol

consumption one of leading preventable causes of death in the United States[1]. Alcohol has

complex effects on multiple biological processes and systems, including the cardiovascular

system. Several studies suggest that habitual, heavy alcohol use can lead to cardiovascular

sequelae such as dilated cardiomyopathy and heart failure[2]. However, the benefits and

potential arms of moderate drinking have been a subject of controversy. A few studies have

indicated that the association presented a J-shaped curve between alcohol consumption and a

lower risk of cardiovascular disease[3–6]. While studies with Mendelian randomization method

suggested a non-linear and increased risk of cardiovascular risks with any dose of alcohol

intake[3,5]. Additionally, multiple studies support a causal relationship of alcohol consumption

with elevated blood pressure and the risk of hypertension (HTN)[2,7]. Furthermore, high blood

pressure is one of the leading risk factors for cardiovascular diseases (CVDs)[8,9]. Therefore,

understanding the molecular changes underlying alcohol consumption is crucial to comprehend

the relationship between alcohol consumption, high blood pressure, and CVD.

One of the most studied epigenetic modifications, DNA methylation, regulates gene

expression through the transfer of a methyl group onto DNA cytosine-phosphate-guanine (CpG)

sites. The extent of DNA methylation at certain CpG sites is associated with phenotypic variation

in numerous CVD-related traits including body mass index (BMI)[10], blood lipids[11],

glycemic traits[12], blood pressure[13], and inflammatory biomarkers[14]. DNA methylation has

also been linked to lifestyle behaviors such as alcohol consumption. A large-scale meta-analysis

of data from thirteen population-based cohorts including the Framingham Heart Study (FHS)

identified 144 differentially methylated CpG sites associated with heavy alcohol intake[15].

A standardized, biomarker of alcohol consumption may correct for limitations of self-

reported alcohol consumption, such as impression management bias[16] or faulty recall of

drinking history[17,18], and reveal alcohol-related disease risks that otherwise might not be

apparent. In this study, we used 144 alcohol-related, differentially-methylated CpGs[15] to

generate an alcohol consumption epigenetic risk score (ERS) and examine the association of the

ERS with blood pressure traits in cross-sectional and longitudinal analyses. We hypothesized

that a DNA methylation-based alcohol consumption ERS would be associated with blood

pressure, cross-sectionally and longitudinally. We tested our hypothesis by analyzing the

association of our alcohol-associated ERS with blood pressure traits, including systolic blood

pressure (SBP), diastolic blood pressure (DBP), and HTN in 3,898 FHS participants. In addition,

we carried out replication analyses of these findings in eight independent cohorts using meta-

analysis (**Figure 1**). The alcohol consumption ERS provides an opportunity to investigate the

relations of alcohol intake to health outcomes in situations where self-reported intake data is

unavailable or unreliable.

METHODS

Study population

Data from nine population-based cohort studies were used in the analysis. In addition to the

FHS[19], our investigation included the Agricultural Lung Health Study (ALHS)[20], the

Cooperative Health Research in the Region Augsburg (KORA)[21], the Genetic Epidemiology

Network on Arteriopathy (GENOA) Study[22], the Health and Retirement Study (HRS)[23], the

Multi-Ethnic Study of Atherosclerosis (MESA) Study[24], the Rhineland Study[25], the

Rotterdam Study[26], and the Study of Health in Pomerania (SHIP)[27]. Institutional review

committees of all cohorts approved this study, and all study participants provided written

informed consent. In each cohort, participants with prevalent CVD (angina pectoris, coronary

insufficiency, cerebrovascular accident, atherothrombotic infarction of the brain, transient

ischemic attack, cerebral embolism, intracerebral hemorrhage, subarachnoid hemorrhage, or

intermittent claudication), prevalent heart failure, and prevalent atrial fibrillation were excluded.

We also excluded participants without DNA methylation data at baseline examination at which

blood samples were collected for DNA methylation profiling. After exclusions, 3,898

participants in FHS and 11,544 participants in eight independent external cohorts were included

in cross-sectional association analyses, while 3,260 participants in FHS and 3,910 participants in

five external cohorts were included in longitudinal association analyses (**Figure 1**).

Clinical and behavioral data collection

Overall, clinical data for traits such as age, BMI, SBP, DBP, and the use of antihypertensive

medication were collected at in-person examinations. HTN was defined as SBP $\geq$ 140 mm Hg,

DBP $\geq$ 90 mm Hg, or use of antihypertensive medication for treating hypertension at the

examination. We added 15 mm Hg and 10 mm Hg to a participant's measured SBP and DBP

values, respectively, for participants currently using antihypertensive medication.

Participants' cigarette smoking status was determined based on self-reported smoking

behavior. Current smokers were defined as participants who smoked on average at least one

cigarette per day in the past year; former smokers were defined as participants who previously

smoked on average at least one cigarette per day but stopped smoking for at least one year; never

smokers were defined as participants who never smoked. Self-reported alcohol intake was

captured via questionnaires wherein the participants reported the frequency with which they

consumed various alcoholic beverages (i.e., beer, liquor, or wine). This study included nine

population-based cohorts, and therefore, we focused on habitual alcohol consumption in general

populations rather than examining specifically for alcohol disorder. Study-specific methods for

clinical data collection are presented in the **Supplemental Text** and **Supplemental Table 1**.

DNA methylation data collection and processing

Whole blood samples were assayed for DNA methylation via the Infinium Human Methylation

450 BeadChip platform or Infinium MethylationEPIC platform (San Diego, CA) (**Supplemental**

Text). The methylated probe intensity and total probe intensities were extracted using the

Illumina Genome Studio (version 2011.1) with the methylation module (version 1.9.0).

Preprocessing of the methylated (M) signal and unmethylated signal (U) was conducted;

methylation beta-value (β_M) was defined as $\beta = \frac{M}{U+M}$. Further information regarding DNA

extraction and processing has been outlined[15]. Further information regarding the collection of

DNA samples and assay details about each external cohort are presented in the Supplemental

Materials.

Epigenetic risk score calculation

To develop the ERS for this investigation, we used 144 CpGs identified in a previous EWAS

meta-analysis[15]. The score was calculated by multiplying individual-level methylation values

at each CpG site by the CpG's corresponding Least Absolute Shrinkage and Selection Operator

(LASSO) β estimate. LASSO is a regression analysis method that performs both variable

selection and regularization to enhance the prediction accuracy and interpretability of the

resulting statistical model by shrinking the coefficients of some variables to exactly zero[28].

The weighted individual values were summed across all 144 CpGs sites to create an ERS for

each participant, representing weighted DNA methylation levels in response to alcohol

consumption. This methodology was applied to all participating cohorts' methylation data to

generate the ERS. Across five cohorts, one drink of alcohol consumption was associated with

0.09 higher unit of ERS (**Supplemental Table 2**). Methods for calculating the ERS for cohorts

missing certain CpG methylation values can be found in the **Supplemental Text**.

Analysis of the epigenetic risk score with blood pressure traits in FHS (discovery)

We performed both cross-sectional and longitudinal regression analyses in FHS to examine the

association between the ERS (independent variable) and blood pressure traits: SBP (continuous),

DBP (continuous), and HTN (dichotomous) (dependent variables). Linear mixed regression

models were used to evaluate the association of the ERS with the two continuous blood pressure

traits. Generalized estimating equations (GEE) were used to evaluate the association of the ERS

with dichotomous HTN. A total of 3,898 participants were included in the cross-sectional

analysis from the FHS Offspring cohort (n = 2,393; examination 8) and FHS Third Generation

cohort (n = 1,505; examination 2) participants. All models were adjusted for age, age-squared,

sex, BMI, familial correlation (for family data), and current smoking status.

Longitudinal analyses of all blood pressure traits included FHS Offspring cohort

participants (n = 1,932) who attended both examinations 8 and 9 and Third Generation

participants (n = 1,328) who attended both examinations 2 and 3. Our linear mixed regression

models evaluated the association of change in blood pressure over time (i.e.**,** ΔSBP and ΔDBP)

with the baseline ERS after adjusting for baseline age, baseline age-squared, sex, baseline BMI,

baseline smoking status, baseline SBP/DBP (i.e., if the model's outcome was ΔDBP, we adjusted

for baseline DBP), and time between baseline and the follow-up examination. Our GEE models

evaluated the association of incident HTN with the baseline ERS after adjusting for baseline age,

baseline age-squared, baseline BMI, baseline smoking status, and time between baseline and

follow-up examination; in addition, these GEE models excluded all participants with HTN at

baseline examination. In the sensitivity analysis, we defined participants with stage 1

hypertension using the 2017 guideline (i.e., ≥ 130/80 SBP/DBP mm Hg or with antihypertension

treatment)[29]. We performed cross-sectional and longitudinal GEE models to investigate the

associations of ERS with prevalent and incident hypertension using the new definition.

Meta-analysis (replication)

For replication, independent external cohorts (n = 11,544) were used in an inverse-variance

weighted, fixed-effects meta-analysis by assuming that there is one true effect between ERS and

a BP trait. Eight studies were used in cross-sectional meta-analysis for replication while five

studies were utilized in the longitudinal meta-analysis (n = 4,021).

Analysis of the epigenetic risk score with blood pressure traits in participants without

antihypertension medication

To minimize the possible effects of antihypertension medication on DNA methylation, we

conducted a sensitivity-analysis among participants without antihypertension medication in five

cohorts (i.e., FHS, GENOA, HRS, Rhineland Study, and SHIP). Similar to the primary analysis,

we conducted the cross-sectional analysis using linear mixed effects model with ERS as the

independent variable and blood pressure traits as the dependent variables in each cohort.

Analysis of the epigenetic risk score with alcohol consumption

We used a linear mixed regression model to test the cross-sectional association between the ERS

(outcome) and self-reported alcohol intake (exposure) in each of the five cohorts (i.e., FHS,

GENOA, HRS, Rhineland Study, and SHIP). The change in the ERS associated with one drink

of alcohol consumption per day was calculated with adjustment for age, age-squared, sex, BMI,

current smoking status, and familial correlation.

Association analysis of alcohol consumption with blood pressure traits

To compare the association of blood pressure traits with ERS and questionnaire-based alcohol

consumption, we performed cross-sectional (i.e., FHS, GENOA, HRS, Rhineland Study, and

SHIP) and longitudinal (i.e., FHS, GENOA, and SHIP) analyses between blood pressure traits

and alcohol consumption. We used linear mixed or GEE models to quantify the associations

between SBP/DBP/HTN (outcome variables) and alcohol consumption (predictor). Covariates

included age, age-squared, sex, BMI, current smoking status, and family structure.

Association of epigenetic risk score with biological biomarkers of alcohol intake

We tested the association of the ERS with two established biomarkers of chronic alcohol

consumption: aspartate amino transferase (AST) and alanine aminotransferase (ALT)

concentrations. Separate linear mixed regression models were used with each enzyme as the

dependent variable. Serum AST and ALT were measured on fasting morning samples using the

kinetic method (Beckman Liquid-State Reagent Kit)[30]. Model 1 (i.e., the reduced model)

quantified the association between the self-reported alcohol intake and liver enzyme

concentrations after adjusting for age, sex, BMI, and smoking status. Model 2 (i.e., the full

model) further adjusted for the ERS. In order to compare the two models, we also performed a

likelihood ratio test (LRT) to gauge whether the addition of the ERS significantly improved

model fit.

Analysis of individual alcohol-associated CpGs with blood pressure traits in FHS

We examined the cross-sectional association of 144 DNA methylation probes from ERS with

blood pressure traits in the FHS. We applied the linear mixed effect model to account for the

pedigree with each CpG probe as the predictor variable and SBP/DBP as the outcome variable.

Covariates included age, age-squared, sex, BMI, and current smoking status.

All statistical analyses were conducted using the R (version 4.0.3) software package[31].

Meta-analyses was conducted with the *metafor* package (version 3.0.2)[32]. LRT was performed

using the 'lrtest' function in the R package *lmtest* in R (version 0.9.39)[33]. Statistical

significance was defined as two-sided $p<0.05$.

RESULTS

Participant characteristics

This present study included discovery and replication analyses (**Figure 1**). The discovery

association analysis was performed in the FHS. The replication association analysis and meta-

analysis was performed in up to eight cohorts (**Figure 1**). At the baseline examination, FHS

participants (n = 3,898) were, on average, 58 years old (SD = 13 years) and consisted of slightly

more women (55%) than men (45%) (**Supplemental Table 3**). In addition, approximately 42%

of FHS participants had hypertension at baseline. Furthermore, women consumed less alcohol

compared to men (mean alcohol intake 0.3 drinks/day vs. 0.7 drinks/day). The FHS participants

(n = 3,260) were followed up for six years and used for longitudinal association analyses with

blood pressure traits (**Supplemental Table 2**).

Overall, mean age of participants in the eight independent cohorts (n_{total} = 11,544) ranged

from of 49 years (SHIP) to 68 years (HRS) (**Supplemental Tables 3-5**). Similar to what was

observed in FHS, women reported a lower average amount of alcohol consumed per day

compared to men. DNA methylation was measured using blood samples collected at the same

time when alcohol consumption data were assessed in all nine cohorts. Blood pressure traits

measured contemporaneously were used in cross-sectional analysis and those measured six to ten

years after alcohol consumption measurement were used for longitudinal studies (**Supplemental**

Table 1). Additionally, the mean values of the ERS ranged from -15.35 (SD = 0.74) to -3.85 (SD

= 0.61) across all cohorts at the baseline examination (**Supplemental Table 6**).

Epigenetic risk score and blood pressure: cross-sectional and longitudinal analysis in FHS

The alcohol intake showed a significant association with ALT (p = 2.9E-09) and AST (p = 1.3E-

10), but borderline for the AST/ALT ratio (p = 0.054) in Model 1. The addition of the ERS in

Model 2 (i.e., full model) improved model fit with respect to cross-sectional ALT (p = 2.8E-07),

AST (p = 2.3E-12), and the AST/ALT ratio (p = 0.0076) (**Supplemental Table 7**).

Cross-sectional analyses in FHS participants revealed significant association of the ERS

with SBP, DBP, and HTN. A one-unit increment of the ERS was associated with a 1.98 mm Hg

higher SBP (SE = 0.39, p = 4.6E-07), a 0.68 mm Hg higher DBP (SE = 0.25, p = 0.006), and an

odds ratio of 1.78 for HTN (95% CI = [1.55, 2.04], p < 2E-16) (**Table 1**). In contrast,

longitudinal analyses did not reveal significant associations of the ERS with ΔSBP, ΔDBP, or

incident HTN (p > 0.3 for all) (**Table 1**).

In the sensitivity analysis with the updated definition for stage 1 hypertension (i.e., ≥

130/80 mm Hg or with the antihypertension treatment), we observed consistent results as the

stage 2 hypertension definition (i.e., ≥ 140/90 mm Hg or with the antihypertension treatment).

One-unit higher ERS was positively associated with the prevalent hypertension (OR = 1.70, 95%

CI = [1.54, 1.98], p < 2E-16) but not significantly associated with the incident hypertension (p =

0.98) (**Supplementary Table 8**).

Replication meta-analysis

Meta-analysis of eight independent external cohorts (n = 11,544) revealed significant cross-

sectional associations. A one-unit greater ERS was associated with a 0.74 (95% CI = [0.26,

1.22], p = 0.002) mm Hg higher SBP (**Figure 2**) and a 0.50 (95% CI = [0.21, 0.78], p = 0.0006)

mm Hg higher DBP (**Figure 3**). As can be seen in the forest plots, there was heterogeneity in the

meta-analysis results for SBP (Q = 17.27, p = 0.008), but not for DBP (Q = 5.16, p = 0.52). No

significant association was observed between the ERS and HTN in meta-analysis (OR = 1.02,

95% CI = [0.83, 1.24], p = 0.10; **Supplemental Figure 1**). As a sensitivity analysis for cross-

sectional HTN, meta-analysis was replicated excluding the Rhineland Study, which accounted

for 73% of the pooled effect size, but did not change results significantly (**Supplemental Figure**

2).

No significant associations of the ERS with blood pressure traits in longitudinal meta-

analysis were observed (ΔSBP β = 0.63, 95% CI = [-0.12, 1.38], p = 0.10; ΔDBP β = 0.10. 95%

CI = [-0.30, 0.50], p = 0.61; incident HTN β (log OR) = 0.003, 95% CI [-0.05, 0.06], p = 0.92).

These analyses included five cohorts (GENOA, KORA, MESA, the Rotterdam Study, and SHIP)

that had follow-up examination data available (for ΔSBP, ΔDBP total sample size n = 3,910; for

incident HTN, n = 3,228). **Supplemental Figures 3, 4, and 5** display the meta-analysis results

for ΔSBP, ΔDBP, and incident HTN, respectively. Cross-sectional and longitudinal analysis

results for individual cohort are shown in **Supplemental Tables 9 and 10**, respectively.

Sensitivity analyses

Association of epigenetic risk score with blood pressure traits in participants without

antihypertension medication

We conducted sensitivity analyses between ERS and SBP/DBP in 2577 FHS participants (66.1%

of the entire study sample) who were not receiving antihypertension medication. We observed

stronger results in the participants without antihypertension treatment compared to the results in

all participants (**Supplementary Table 11**). One-unit higher ERS was associated with a 2.6 mm

Hg higher SBP (p = 1.5E-7) and a 1.54 mm Hg higher DBP (p = 1.1E-6), which were stronger

than the estimates obtained from all FHS participants (SBP: β =1.98, p = 4.6E-7; DBP: β = 0.68,

p = 0.006). However, associations were not significant between ERS and SBP/DBP among

untreated participants (p > 0.05) in the four external independent cohorts (**Supplementary Table**

11), except a marginally significant association between ERS and DBP (β = 0.40, p = 0.09) in

Rhineland untreated participants. Of note, most of the cross-sectional associations were

nonsignificant in these four cohorts between ERS and blood pressure traits before treated

participants were removed (**Supplementary Table 9**).

Association of epigenetic risk score with alcohol consumption

We conducted association between ERS and alcohol consumption in FHS. Each drink of self-

reported alcohol consumption per day was associated with a 0.25-units higher ERS (p < 0.0001)

in FHS. Consistent positive associations were also observed in three independent cohorts,

GENOA, HRS, and Rhineland Study (**Supplemental Table 2**). One drink of self-reported

alcohol consumption per day was associated with a 0.32-unit higher ERS in GENOA (p <

0.0001), with a 0.26-unit higher ERS in HRS (p < 0.0001), and with a 0.01-unit higher ERS in

the Rhineland Study (p = 0.02). No significant association was observed in SHIP (**Supplemental**

Table 2).

Association of alcohol consumption with blood pressure traits

Alcohol consumption was associated with SBP in both cross-sectional and longitudinal analyses

in the FHS (**Supplemental Tables 12-13**). One additional drink/day alcohol consumption was

associated with a 0.88 mm Hg higher SBP (p = 3.7e-4) in cross-sectional analysis and with a

0.92 mm Hg increase in SBP (p = 2.6e-4) between two exams in longitudinal analysis. We also

found that one additional drink/day alcohol consumption was associated with 1.13 times of odds

of being hypertension (95% CI = 1.04, 1.22; p = 0.006) in cross-sectional analysis. Associations

of alcohol consumption with DBP were not significant ($p > 0.05$ in both cross-sectional and

longitudinal analyses) in FHS (**Supplemental Tables 12-13**). In Rhineland and SHIP, we

observed that higher alcohol consumption was significantly associated with SBP (Rhineland: β =

0.38, p = 0.006; SHIP: β = 2.94, p = 0.008) and DBP (Rhineland: β = 0.38, p = 1.6e-6; SHIP: β =

2.06, p = 0.002) (**Supplemental Table 12**). However, prevalent hypertension was not associated

with alcohol consumption in either cohort. Whereas in HRS, higher level of DBP (β = 0.72, p =

0.04) and higher odds of hypertension (β = 0.17, p = 0.02) were associated with higher alcohol

intake in cross-sectional analysis. No longitudinal significant association was observed between

alcohol consumption and any of the blood pressure traits in GENOA or SHIP participants

(**Supplemental Table 13**).

Additionally, we investigated whether the DNA methylation CpGs that were used for

constructing the ERS were associated with blood pressure traits in cross-sectional analysis in

FHS. When applying a relaxed threshold by false discovery rate (FDR) < 0.05, 26 and 17 of 144

CpG probes were significantly associated with SBP and DBP, respectively (**Supplemental**

Figures 6a-c, Supplemental Table 14-15).

DISCUSSION

This investigation builds upon our previous work that identified 144 CpGs that discriminated

current heavy alcohol drinkers from non- and light-drinkers[15]. Using these CpGs, we

constructed an alcohol consumption ERS that was associated with self-reported alcohol intake

and explained additional variance (model R^2 = 0.11 vs. 0.07 for AST and 0.14 vs. 0.11 for ALT)

in hepatic biomarkers associated with alcohol intake. The risk score was associated with systolic

and diastolic blood pressure in cross-sectional analyses both in discovery and replication

analyses. In contrast, the risk score was not associated with longitudinal changes in blood

pressure or incident hypertension either in discovery or replication studies.

The relationship between alcohol and hypertension is fairly well-defined with a

comprehensive body of cross-sectional and longitudinal epidemiological studies, revealing a

significant association between the two[34]. In addition, intervention studies and Mendelian

randomization studies have suggested that the alcohol-blood pressure relationship is causal[7].

There are multiple mechanisms by which alcohol consumption can affect blood pressure. For

example, chronic alcohol consumption has been reported to interfere with nitrous oxide (NO)

production from endothelial cells, which affects various blood pressure regulatory

mechanisms[35].

DNA methylation is an epigenetic mechanism that involves the transfer of a methyl

group onto the fifth carbon position of the DNA building block, cytosine, to form 5-

methylcytosine. This modification regulates gene expression by recruiting proteins to primarily

inhibit the binding of transcription factor(s) to DNA. Therefore, certain cells may develop a

stable and unique DNA methylation pattern that regulates tissue-specific gene expression. As

previously mentioned, a prior study from our team[15] aimed to address the lack of reliable

measures of alcohol intake by examining DNA methylation as a novel biomarker of alcohol use.

As noted, DNA methylation provides a reliable measure of heavy alcohol intake and addresses a

critical need.

While our analyses reveal that a risk score comprised of 144 CpGs reveals an association

of current alcohol intake with blood pressure, it does not indicate an association of the risk score

with longitudinal change in blood pressure. Several potential reasons may explain this finding.

As follow-up examinations may be years apart for several of these participating cohorts,

participants may change their lifestyle behaviors, which may affect methylation patterns. For

example, participants may change their diet or initiate hypertensive medication during follow-up;

participants who were moderate-to-heavy drinker may also reduce their alcohol consumption

over time with aging. A similar finding was reported with longitudinal smoking traits and DNA

methylation, where the majority of the differentially methylated CpG sites observed in analysis

of current versus never smokers returned to the level of never-smokers within five years of

smoking cessation[36]. The relationship between alcohol consumption and the degree of

methylation changes over various periods of time is still being explored. A recent longitudinal

study identified that 1414 CpGs were significantly associated with alcohol consumption in cross-

sectional analysis while about a third of CpGs (n = 513) displayed associations between the

changes in the methylation levels of these CpGs and the change of alcohol consumption between

two exams[37]. This study indicated, as a dynamic measurement, DNA methylation may show

stronger associations with cross-sectional behaviors and clinical phenotypes than associations

with changes in traits during the follow-up, which was consistent with the findings in our study.

In addition, cross-sectional meta-analysis for SBP revealed significant heterogeneity. Our

project performed a fixed-effects, inverse-variance weighted meta-analysis which assumes all

independent cohorts are estimating the same underlying effect where the variation between

estimates is attributed to random error. This heterogeneity may be attributable to a number of

factors such as participants' country or origin, ethnicity, age, sex, diet, and differences in sample

collection methodology[38]. For example, the GENOA recruited participants who had at least 2

siblings diagnosed with essential hypertension before the 60-year-old[22]. Therefore, the

GENOA participants may be more susceptible to hypertension than the general population due to

common pathogenic genes and shared behavioral habits with their siblings. The SHIP recruited

subjects from Northeast Germany, where the population had the lowest life expectancy in

Germany at the time of recruitment[27]. These different recruitment criteria may distinguish

participants in GENOA and SHIP from the general population, which may influence the relation

between alcohol consumption, DNA methylation, and blood pressure traits. Furthermore,

measurement error and a much higher SBP and DBP in GENOA and Rotterdam Study may

partly explained differential associations between the ERS and alcohol consumption in these two

studies, compared to results in other seven cohorts. Nevertheless, there was still a significant

effect for this outcome, and our meta-analysis results support our findings in FHS.

Another key finding of our study was the improved model fit after the addition of the risk

score in a cross-sectional, linear mixed regression model examining the relationship between

self-reported alcohol intake and liver enzymes, ALT and AST, in FHS. While the association

between self-reported alcohol intake and these liver enzymes was significant, the risk score was

able to capture additional interindividual variations of the elevated liver enzymes (**Supplemental**

Table 7). Whether this observation is due to the risk score as a tool that more accurately gauge

degrees of alcohol consumption as survey or because the risk score represents other biological or

environmental factors warrants future analyses.

This study has several limitations. Most of the participants are of European ancestry,

which may render the findings of our project not applicable to other racial/ethnic populations.

Secondly, we validated the utility of the risk score through association analysis with two

enzymes associated with liver function, ALT and AST. While elevated levels of these enzymes

are useful, common clinical indicators of chronic alcohol consumption, the levels of these

enzymes can be modified by other means. Many prescription drugs such as cholesterol-lowering

agents, anti-tuberculosis drugs, and non-steroidal anti-inflammatory drugs including aspirin are

known to cause mildly elevated AST and ALT[39] which were not adjusted for. These enzyme

levels can also be altered due to conditions such as autoimmune hepatitis: because of lacking

validated data, FHS participants with such conditions were not excluded from analysis but cases

of such conditions are supposed to be very few in the FHS. In addition, at the time of this study,

the FHS did not have other biomarkers available for validation analysis such as gamma-glutamyl

transferase which is another enzyme marker for heavy alcohol usage[40]. Our study also has

several strengths. Our analyses showed that the risk score was significantly associated with self-

reported alcohol intake data and with clinically-useful biomarkers of alcohol consumption. Our

findings regarding the relationship between the risk score and blood pressure traits in FHS were

replicated in meta-analysis of eight independent external cohorts.

CONCLUSION

Our study developed an epigenetic risk score of alcohol consumption based on 144 alcohol-

associated CpG probes identified in a large meta-analysis. This score showed an association with

cross-sectional, but not longitudinal, blood pressure and prevalent hypertension among middle-

aged and older participants. Our findings also provide a proof-of-concept that a DNA

methylation-based epigenetic score capturing alcohol consumption can provide insights into

alcohol-related disease risks, and further help to assess specific lifestyle factors that contribute to

an individual's health profile. This is especially pertinent in situations where self-reported

behavioral data (e.g., alcohol consumption) are unavailable, susceptible to recall bias, or subject

to significant data loss.

Abbreviations:
ERS: epigenetic risk score
SBP: systolic blood pressure
DBP: diastolic blood pressure
HTN: hypertension
AST: aspartate amino transferase
ALT: alanine aminotransferase
8
9   AVAILABILITY OF DATA AND MATERIALS
10   Data and analytical codes that support our findings are available from the corresponding authors
11   upon request.
12
13   ACKNOWLEDGEMENTS
14   We thank all participants in the cohorts for their dedication in research.
15
ETHICS DECLARATIONS
Ethics approval and consent to participate
Institutional review committees of all cohorts approved this study. All study participants
provided written informed consent. The FHS was reviewed and approved by The Boston
Medical Center Institutional Review Boards. The ALHS was approved by the Institutional
Review Board at National Institute of Environmental Health Sciences. The KORA cohort ethical
approval was granted by the ethics committee of the Bavarian Medical Association (REC
reference numbers: F4: #06068, FF4: #06068) and all were carried out in accordance with the
principles of the Declaration of Helsinki. The GENOA was approved by Institutional Review
Boards at the University of Michigan, University of Mississippi Medical Center, and Mayo
Clinic. The HRS was approved by the University of Michigan. The MESA was approved by the
Institutional Review Board of each study site, and written informed consent was obtained from
all participants (IRB: #00009029). The Rhineland Study was approved by the ethics committee
of the University of Bonn, Medical Faculty. The Rotterdam Study was approved by the medical
ethics committee of the Erasmus University Medical Center, Rotterdam, the Netherlands. The
SHIP was approved by the medical ethics committee of the University of Greifswald.
Consent for publication
Not applicable.
Competing Interests
All first and corresponding authors of this manuscript have no conflicts of interest to disclose.
36

FUNDING
The FHS is supported by NIH contract N01-HC-25195. The analytical component of this project
was funded by the Division of Intramural Research, National Heart, Lung, and Blood Institute,
National Institutes of Health, Bethesda, MD (D. Levy, Principal Investigator). J. Ma is supported
by NIH grant R01AA028263; C. Liu is supported by R01AA028263 and R01HL15569 grants.
The ALHS is supported by the Intramural Research Program of the National Institutes of Health,
the National Institute of Environmental Health Sciences (Z01-ES102385, Z01-ES049030, Z01-
ES043012) and the National Cancer Institute (Z01-CP010119). The KORA study was initiated
and financed by the Helmholtz Zentrum München –German Research Center for Environmental
Health, which is funded by the German Federal Ministry of Education and Research (BMBF)

and by the State of Bavaria. The GENOA is supported by the National Heart, Lung and Blood Institute (U10 HL054457, RC1 HL100185, R01 HL087660, R01 HL119443, R01 HL141292, and R01 HL133221). The HRS is supported by the National Institute on Aging (NIA U01AG009740). The MESA is supported by contracts 75N92020D00001, HHSN268201500003I, N01-HC-95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003, N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, UL1-TR-001420, UL1TR001881, DK063491, and R01HL105756. The Rhineland Study is supported by the Diet-Body-Brain Competence Cluster in Nutrition Research funded by the Federal Ministry of Education and Research (grant numbers 01EA1410C and FKZ:01EA1809C). The Rotterdam Study is supported by the Erasmus MC University Medical Center and Erasmus University Rotterdam; The Netherlands Organization for Scientific Research (NWO); The Netherlands Organization for Health Research and Development (ZonMw); the Research Institute for Diseases in the Elderly (RIDE); The Netherlands Genomics Initiative (NGI); the Ministry of Education, Culture and Science; the Ministry of Health, Welfare and Sports; the European Commission (DG XII); and the Municipality of Rotterdam. The SHIP is supported by the Federal Ministry of Education and Research (grants no. 01ZZ9603, 01ZZ0103, and 01ZZ0403), the Ministry of Cultural Affairs as well as the Social Ministry of the Federal State of Mecklenburg-West Pomerania, and the network 'Greifswald Approach to Individualized Medicine (GANI_MED)' funded by the Federal Ministry of Education and Research (grant 03IS2061A). DNA methylation data have been supported by the DZHK (grant 81X3400104).

AUTHOR INFORMATION

Helena Bui, Amena Keshawarz, and Mengyao Wang contributed equally to this work.

Authors and affiliations

Population Sciences Branch, Division of Intramural Research, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA
Helena Bui, Amena Keshawarz, Daniel Levy

Framingham Heart Study, Framingham, MA, USA
Helena Bui, Amena Keshawarz, Daniel Levy, Chunyu Liu

Department of Biostatistics, Boston University School of Public Health, Boston, MA
Mengyao Wang, Yi Li, Xue Liu, Chunyu Liu

Epidemiology Branch, National Institute of Environmental Health Sciences, National Institutes of Health, Department of Health and Human Services, Research Triangle Park, NC, USA
Mikyeong Lee, Stephanie J. London

Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, MI
Scott M. Ratliff, Lisha Lin, Sharon L. R. Kardia, Wei Zhao, Jennifer A. Smith

Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, MI
Kira S. Birditt, Jessica D. Faul, David R. Weir, Jennifer A. Smith

Institute of Epidemiology, Helmholtz Munich, German Research Center for Environmental Health, German
Annette Peters

Institute for Medical Informatics, Biometrics and Epidemiology, Ludwig-Maximilians-Universität München, Munich, Germany
Annette Peters

German Center for Cardiovascular Research (DZHK), Partner Site Munich Heart Alliance, Munich, Germany
Annette Peters, Christian Gieger, Melanie Waldenberger
Research Unit Molecular Epidemiology, Institute of Epidemiology, Helmholtz Munich, German Research Center for Environmental Health, Bavaria, Germany
Christian Gieger, Thomas Delerue, Melanie Waldenberger
The Institute for Translational Genomics and Populations, Department of Pediatrics, Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA 90502, USA
Xiuqing Guo, Jie Yao, Jerome I. Rotter
Population Health Sciences, German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany
Dan Liu, Juliana F. Tavares, Gökhan Pehlivan, Monique M. B. Breteler
Institute for Medical Biometry, Informatics and Epidemiology (IMBIE), Faculty of Medicine, University of Bonn, Bonn, Germany
Monique M. B. Breteler
Department of Epidemiology, Erasmus MC University Medical Center, Rotterdam, The Netherlands
Irma Karabegovic, Carolina Ochoa-Rosales, Trudy Voortman, Mohsen Ghanbari
Centro de Vida Saludable de la Universidad de Concepción, Concepción, Chile
Irma Karabegovic, Carolina Ochoa-Rosales
Division of Human Nutrition and Health, Wageningen University and Research, Wageningen, The Netherlands
Trudy Voortman
Department of Internal Medicine, Erasmus MC University Medical Center, Rotterdam, The Netherlands
Joyce B.J. van Meurs
Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany
Mohamed Kamal Nars, Alexander Teumer
German Center for Cardiovascular Research (DZHK), Partner Site Greifswald, Greifswald, Germany
Mohamed Kamal Nars, Marcus Dörr, Alexander Teumer
Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany
Marcus Dörr
Department of Psychiatry and Psychotherapy, University Medicine, Greifswald, Germany
Hans J. Grabe
German Center for Neurodegenerative Diseases (DZNE), Rostock/Greifswald, site Greifswald, Germany
Hans J. Grabe
Department of Population Medicine and Lifestyle Diseases Prevention, Medical University of Bialystok, Bialystok, Poland
Alexander Teumer
Gerald J. and Dorothy R. Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA, USA
Jiantao Ma

Contributions

D Levy, JM, and CL designed the study. HB, AK, M Wang, ML, SMR, LL, KSB, JDF, AP, CG, TD, SLRK, WZ, XG, JY, JIR, YL, XL, D Liu, JFT, GP, MMBB, IK, COR, TV, MG, JBJvM, MKN, MD, HJG, SJL, AT, M Waldenberger, DRW, JAS, D Levy, JM, and CL were involved in the data collection, data preparation, and statistical analyses. HB, AK, and M Wang drafted the manuscript. D Levy, JM, and CL supervised the study and provided critical review. All authors read and approved this manuscript.

Corresponding author

Correspondence to Daniel Levy, Jiantao Ma, Chunyu Liu.

Table 1. Cross-sectional and longitudinal association analyses of epigenetic risk score and blood pressure traits in the Framingham Heart Study

Continuous variable	β	SE	p
SBP	1.98	0.39	4.64E-07
DBP	0.68	0.25	0.006
ΔSBP	0.44	0.44	0.32
ΔDBP	-0.0012	0.26	0.99
Binary variable	**OR**	**95% CI**	**p**
Hypertension	1.78	1.55, 2.04	< 2E-16
Incident hypertension	1.02	0.83, 1.24	0.85

SBP/DBP, systolic/diastolic blood pressure; ΔSBP/ΔDBP, longitudinal change in systolic/diastolic blood pressure; SE, standard error; OR, Odds ratio; 95% CI, 95% confidence interval. The cross-sectional and longitudinal analyses were performed on 3,898 and 3,260, respectively, Framingham Heart Study (FHS) participants.

Figure 1. Study design.

Figure 2. Meta-analysis of cross-sectional association analyses of ERS in relation to systolic blood pressure in eight independent external cohorts (n = 11,544). ALHS, Agricultural Lung Health Study; GENOA, Genetic Epidemiology Network of Arteriopathy; HRS, Health and Retirement Study; KORA, Cooperative Health Research in the Region Augsburg; MESA, Multi-Ethnic Study of Atherosclerosis; SHIP, Study of Health in Pomerania; FE, Fixed Effect; 95% CI, Confidence Interval.

Figure 3. Cross-sectional meta-analysis of ERS in relation to diastolic blood pressure in eight independent external cohorts (n = 11,544). ALHS, Agricultural Lung Health Study; GENOA, Genetic Epidemiology Network of Arteriopathy; HRS, Health and Retirement Study; KORA, Cooperative Health Research in the Region Augsburg; MESA, Multi-Ethnic Study of Atherosclerosis; SHIP, Study of Health in Pomerania; FE, Fixed Effect; 95% CI, Confidence Interval.

References

1. Centers for Disease Control and Prevention (CDC). Alcohol Related Disease Impact (ARDI) application [Internet]. 2024 [cited 2024 Apr 2]. Available from: www.cdc.gov/ARDI

2. Piano MR. Alcohol's Effects on the Cardiovascular System. Alcohol Res. 2017;38:219–41.

3. Biddinger KJ, Emdin CA, Haas ME, Wang M, Hindy G, Ellinor PT, et al. Association of Habitual Alcohol Intake With Risk of Cardiovascular Disease. JAMA Netw Open. 2022;5:e223849.

4. Zhang X, Liu Y, Li S, Lichtenstein AH, Chen S, Na M, et al. Alcohol consumption and risk of cardiovascular disease, cancer and mortality: a prospective cohort study. Nutr J. 2021;20:13.

5. Fernández-Solà J. Cardiovascular risks and benefits of moderate and heavy alcohol consumption. Nat Rev Cardiol. 2015;12:576–87.

6. Krittanawong C, Isath A, Rosenson RS, Khawaja M, Wang Z, Fogg SE, et al. Alcohol Consumption and Cardiovascular Health. Am J Med. 2022;135:1213-1230.e3.

7. Puddey IB, Mori TA, Barden AE, Beilin LJ. Alcohol and Hypertension-New Insights and Lingering Controversies. Curr Hypertens Rep. 2019;21:79.

8. Vasan RS, O'Donnell CJ. Impact of High-Normal Blood Pressure on the Risk of Cardiovascular Disease. The New England Journal of Medicine. 2001;

9. Malik R, Georgakis MK, Vujkovic M, Damrauer SM, Elliott P, Karhunen V, et al. Relationship Between Blood Pressure and Incident Cardiovascular Disease: Linear and Nonlinear Mendelian Randomization Analyses. Hypertension. 2021;77:2004–13.

10. Samblas M, Milagro FI, Martínez A. DNA methylation markers in obesity, metabolic syndrome, and weight loss. Epigenetics. 2019;14:421–44.

11. Płatek T, Polus A, Góralska J, Raźny U, Gruca A, Kieć-Wilk B, et al. DNA methylation microarrays identify epigenetically regulated lipid related genes in obese patients with hypercholesterolemia. Mol Med. 2020;26:93.

12. Liu J, Carnero-Montoro E, van Dongen J, Lent S, Nedeljkovic I, Ligthart S, et al. An integrative cross-omics analysis of DNA methylation sites of glucose and insulin homeostasis. Nat Commun. 2019;10:2581.

13. Richard MA, Huan T, Ligthart S, Gondalia R, Jhun MA, Brody JA, et al. DNA Methylation Analysis Identifies Loci for Blood Pressure Regulation. Am J Hum Genet. 2017;101:888–902.

14. Myte R, Sundkvist A, Van Guelpen B, Harlid S. Circulating levels of inflammatory markers and DNA methylation, an analysis of repeated samples from a population based cohort. Epigenetics. 2019;14:649–59.

15. Liu C, Marioni RE, Hedman ÅK, Pfeiffer L, Tsai P-C, Reynolds LM, et al. A DNA methylation biomarker of alcohol consumption. Mol Psychiatry. 2018;23:422–33.

16. Davis CG, Thake J, Vilhena N. Social desirability biases in self-reported alcohol consumption and harms. Addict Behav. 2010;35:302–11.

17. Ekholm O. Influence of the recall period on self-reported alcohol intake. Eur J Clin Nutr. 2004;58:60–3.

18. Stockwell T, Donath S, Cooper-Stanbury M, Chikritzhs T, Catalano P, Mateo C. Under-reporting of alcohol consumption in household surveys: a comparison of quantity-frequency, graduated-frequency and recent recall. Addiction. 2004;99:1024–33.

19. Mahmood SS, Levy D, Vasan RS, Wang TJ. The Framingham Heart Study and the epidemiology of cardiovascular disease: a historical perspective. Lancet. 2014;383:999–1008.

20. Hoang TT, Qi C, Paul KC, Lee M, White JD, Richards M, et al. Epigenome-Wide DNA Methylation and Pesticide Use in the Agricultural Lung Health Study. Environ Health Perspect. 2021;129:97008.

21. Holle R, Happich M, Löwel H, Wichmann HE, MONICA/KORA Study Group. KORA--a research platform for population based health research. Gesundheitswesen. 2005;67 Suppl 1:S19-25.

22. FBPP Investigators. Multi-center genetic study of hypertension: The Family Blood Pressure Program (FBPP). Hypertension. 2002;39:3–9.

23. Sonnega A, Faul JD, Ofstedal MB, Langa KM, Phillips JWR, Weir DR. Cohort Profile: the Health and Retirement Study (HRS). Int J Epidemiol. 2014;43:576–85.

24. Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. Am J Epidemiol. 2002;156:871–81.

25. Breteler MMB, Wolf H. THE RHINELAND STUDY: A NOVEL PLATFORM FOR EPIDEMIOLOGIC RESEARCH INTO ALZHEIMER DISEASE AND RELATED DISORDERS. Alzheimer's & Dementia [Internet]. 2014 [cited 2024 Jan 28];10. Available from: https://alz-journals.onlinelibrary.wiley.com/doi/10.1016/j.jalz.2014.05.810

26. Ikram MA, Brusselle G, Ghanbari M, Goedegebure A, Ikram MK, Kavousi M, et al. Objectives, design and main findings until 2020 from the Rotterdam Study. Eur J Epidemiol. 2020;35:483–517.

27. Völzke H, Schössow J, Schmidt CO, Jürgens C, Richter A, Werner A, et al. Cohort Profile Update: The Study of Health in Pomerania (SHIP). Int J Epidemiol. 2022;51:e372–83.

28. Ye Z, Zhu Y, Coffman DL. Variable selection for causal mediation analysis using LASSO-based methods. Stat Methods Med Res. 2021;30:1413–27.

29. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Journal of the American College of Cardiology. 2018;71:e127–248.

30. Loomba R, Hwang S-J, O'Donnell CJ, Ellison RC, Vasan RS, D'Agostino RB, et al. Parental obesity and offspring serum alanine and aspartate aminotransferase levels: the Framingham heart study. Gastroenterology. 2008;134:953–9.

31. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2021. Available from: https://www.R-project.org/

32. Viechtbauer W. Conducting Meta-Analyses in *R* with the **metafor** Package. J Stat Soft [Internet]. 2010 [cited 2024 Jan 28];36. Available from: http://www.jstatsoft.org/v36/i03/

33. Zeileis A, Hothorn T. Diagnostic Checking in Regression Relationships. R News. 2002;2.

34. Fuchs FD, Fuchs SC. The Effect of Alcohol on Blood Pressure and Hypertension. Curr Hypertens Rep. 2021;23:42.

35. Husain K, Ansari RA, Ferder L. Alcohol-induced hypertension: Mechanism and prevention. WJC. 2014;6:245.

36. Joehanes R, Just AC, Marioni RE, Pilling LC, Reynolds LM, Mandaviya PR, et al. Epigenetic Signatures of Cigarette Smoking. Circ Cardiovasc Genet. 2016;9:436–47.

37. Dugué P-A, Wilson R, Lehne B, Jayasekara H, Wang X, Jung C-H, et al. Alcohol consumption is associated with widespread changes in blood DNA methylation: Analysis of cross-sectional and longitudinal data. Addict Biol. 2021;26:e12855.

38. Ruppar T. Meta-analysis: How to quantify and explain heterogeneity? Eur J Cardiovasc Nurs. 2020;19:646–52.

39. Yin LK, Tong KS. Elevated Alt and Ast in an Asymptomatic Person: What the primary care doctor should do? Malays Fam Physician. 2009;4:98–9.

40. van Beek JHDA, de Moor MHM, Geels LM, Sinke MRT, de Geus EJC, Lubke GH, et al. The association of alcohol intake with γ-glutamyl transferase (GGT) levels: evidence for correlated genetic effects. Drug Alcohol Depend. 2014;134:99–105.