

**Title:**

Influenza mortality as an indicator of the effectiveness of public health interventions and their impact on United States COVID mortality

**Author:** Robert Morris, MD, PhD

**Affiliation:** Research Now Institute

**Keywords:** COVID, public health, vaccination, mortality, influenza

## Abstract

The synergistic nature of non-pharmaceutical interventions (NPIs) for control of COVID makes it difficult to assess the efficacy of any individual strategy. This study uses influenza mortality reduction (IMR) during the pandemic as an indicator of the aggregate efficacy of NPIs to assess their impact on COVID mortality.

Age-adjusted COVID mortality for US states were modeled as a function of four variables: mortality prior to the introduction of NPIs, vaccination rates, IMR relative to historical averages, and population density.

A simple linear model with only these variables explained 69% of the state-to-state variability in age adjusted COVID mortality. The resulting model suggests that NPIs alone prevented 840,000 COVID related deaths in the United States over the course of the pandemic. These results demonstrate the utility of IMRs as an indicator of the aggregate impact of NPIs for controlling transmission of respiratory infections, including COVID.



## Background

Since the outset of the COVID pandemic, non-pharmaceutical interventions (NPIs) to protect public health have come under heavy criticism for their impact on everything from the economy<sup>1</sup> to mental health<sup>2-4</sup> to education<sup>5,6</sup>. Furthermore, almost every intervention has, at some point, been declared ineffective, including masking,<sup>7,8</sup> routine testing,<sup>9</sup> school closures,<sup>10</sup> and business “lockdowns”.<sup>11,12</sup>

Meta-analyses of studies of the efficacy of individual NPIs have tended to find beneficial effects<sup>13-15</sup> with a few prominent exceptions.<sup>7,8</sup> Closures of businesses and schools, limits on social gathering, travel restrictions, social distancing rules, masking mandates, and other NPI’s act in concert to reduce the transmission of respiratory infections. Some protect the individual from the infection in the community, some protect the community from the infected individual, and some do both. Also, the effectiveness of NPI’s depends on compliance, which is difficult to quantify. How, then, do we evaluate the overall impact of these interventions on the transmission of COVID?

In an ideal natural experiment, we would have two isolated regions experiencing epidemic conditions that are identical in every way except for fully quantified and controlled differences in NPIs. Alternatively, we might have historical data for a particular disease and could examine changes in incidence and mortality after interventions were imposed. No such natural experiment occurred and, because COVID is a new human disease, we have no historical data. All of this makes the aggregate impact of NPI’s on COVID difficult to assess directly.

However, the impact of these NPIs was not limited to COVID. Interventions designed to stop one respiratory pathogen will stop others as well. Therefore, the extent to which these NPIs halted the spread of respiratory pathogens with similar patterns of airborne transmission may provide a surrogate for the efficacy of NPIs for COVID. By far the best characterized of these diseases is influenza.

The marked seasonal patterns of influenza incidence and mortality have been measured for decades. As a result, the expected influenza mortality and the variability in that mortality are well established. Multiple studies have noted the dramatic, global decline in influenza incidence and mortality<sup>16–18</sup> during the COVID pandemic and suggested an association with NPIs. In the United States, influenza mortality rates for the first two complete flu seasons (2020-22) were 80% below historic rates,<sup>19</sup> as can be seen in Figure 1. The sharpness of this decline and the fact that flu mortality rose back to pre-pandemic levels once precautions were lifted in 2022-23 makes it unlikely that the drop was due to any change in the prevalent virus or treatment options. Influenza vaccination rates did rise 7% above historical averages during the pandemic,<sup>20</sup> probably due to concurrent vaccination with COVID, but this cannot explain an 80% drop in influenza mortality.

It appears that something changed during the pandemic that resulted in a dramatic drop in respiratory disease transmission. By far the most likely explanation of this is COVID NPI's. That suggests that the extent to which influenza mortality decreased from expected levels represents a drop in respiratory disease transmission and may provide an indicator of the effectiveness of COVID NPI's.

The current study explores the association between the influenza mortality reduction (IMR) and COVID mortality at the state level. Other factors considered in the analysis were COVID mortality during the first month of the pandemic and vaccination rates. Because the effectiveness of NPIs can be influenced by population density, it was also included in the model as an interaction term.

## **Methods**

Weekly counts of influenza deaths for the period from 2016 through 2023 were abstracted from the CDC FluView System<sup>21</sup> for each state. Average annual influenza mortality rates for each state were calculated for the pre-COVID period 2016 through 2019 and for the two flu seasons during the pandemic, 2020-21 and 2021-22. The decrease in average flu season mortality for each state during the pandemic as compared to average mortality rates prior to the pandemic were calculated for each state to determine the Influenza Transmission Control.

State specific, weekly COVID mortality data were obtained from the CDC COVID Data Tracker and used to calculate mortality rates for March of 2020 to determine initial COVID mortality rates.<sup>22</sup> Vaccination rates at the end of 2023 were obtained from the same site with vaccination defined as receipt of two initial doses.<sup>22</sup> Age adjusted COVID mortality was determined based on age specific COVID mortality rates from CDC through September, 2023, the last date for which they were available.<sup>23</sup> State age distributions and population density for 2020 was obtained from the United States Census Bureau.<sup>24</sup>

Two multiple linear regression models were evaluated with state specific, age adjusted COVID mortality as the outcome variable. Model 1 included only four predictor variables, IMR, COVID mortality, vaccination rates, and population density. Model 2 added an interaction term for IMR and population density based on the assumption that the effect of NPI's as represented by IMR would depend on population density. All statistical analyses were conducted using the STATA statistical package.

The resulting models were used to evaluate the counterfactual cases of non-intervention by setting the relevant variable to zero for each state, determining predicted deaths for each state, and summing the results. Since this effectively removed IMR as a variable, the interaction term also became meaningless, so Model 1 with no interaction term was used. Monte Carlo simulations with 10,000 repetitions were used to estimate the distribution of the aggregated mortality predictions and extract confidence intervals. The three cases considered involved setting IMR, vaccination rate, and both variables to zero.

## Results

As listed in Table 1, state-level influenza mortality rates were an average of 78% lower during the two full flu seasons of the pandemic, 2020-21 and 2021-22, as compared to the three full flu seasons prior to the pandemic, 2016-17, 2017-18, and 2018-19. The decrease in mortality ranged from 49% (North Dakota) to 94% (Washington). This radical difference in influenza mortality during the pandemic at the state level is highly unlikely to reflect simple seasonal variation in the flu strain or vaccine effectiveness ( $p < 0.0001$  by simple ANOVA).

COVID mortality had a strong negative correlation with IMR and vaccination rates ( $p < 0.001$ ). As shown in Figure 1, IMR explains almost a third of the variability in COVID mortality. IMR is also strongly correlated with vaccination rates (Table 2).

Multiple linear regression model results are provided in Table 3. Model 1, with only four predictor variables, IMR, COVID mortality, vaccination rates, and population density, predicts COVID mortality at the state level with an  $r^2$  to 0.65. Introducing an interaction term for IMR and population density in Model 2 improves the adjusted model  $r^2$  to 0.69. This interaction had a positive coefficient, suggesting the rate of reduction in COVID mortality associated with IMR was diminished in more densely populated states. Also, introducing the interaction to the model converted the direction of the effect of population density on COVID mortality from positive to negative. The close fit of the model to actual age-adjusted state COVID mortality rates can be seen in Figure 2.

With IMR set to zero, the model yields an estimated COVID mortality of 1.99 million (95% CI of (1.32 – 2.66 million) suggesting that NPIs saved 840,000 lives. If we set both IMR and vaccination to zero, the COVID mortality estimate rises to 3.39 million (95% CI of (2.93 – 3.86 million) suggesting that preventative measures saved 2,250,000 lives.

## Discussion

The current study provides strong evidence that NPIs played a key role in limiting the impact of the pandemic preventing an estimated 830,000 COVID deaths respectively.

It is conceivable that a decrease in ascertainment rather than reduced transmission could contribute to a decline in influenza mortality. Some have even suggested that COVID deaths are actually influenza deaths.<sup>25</sup> Several observations allow us to dismiss these alternatives. First, failure to diagnose a fatal case of the flu correctly, even during the pandemic, seems unlikely given the well-established surveillance system and diagnostic tools for influenza. Second, the sharp drop in influenza incidence during the pandemic was observed in data from the Seattle Flu Study, which was an active surveillance program that demonstrated pandemic-related decreases

for a broad range of respiratory infections.<sup>26,27</sup> All respiratory infections dropped sharply including influenza, respiratory syncytial virus, and non-COVID corona viruses. Finally, if flu deaths were being misdiagnosed as COVID, we would expect the reduction in influenza mortality to have a strong positive correlation with COVID mortality rates, not the strong negative correlation observed in these data.

It is notable that the regression coefficient for population density changes sign when interaction with IMR is included in the model. This may reflect the fact that population density is a two-edged sword with respect to COVID mortality, inferring a higher transmission risk but providing better access to life-saving medical care. Also of note is the negative association between the pre-intervention COVID mortality and total COVID mortality. This may reflect greater compliance with interventions in the states hardest hit at the outset of the pandemic.

One key advantage in using state IMR as a measure of NPI efficacy is that a region can serve as its own control. Comparing influenza mortality during the pandemic to historical mortality rates of influenza incidence and mortality with those that prevailed during periods when COVID NPIs largely eliminates the effect of time invariant confounders.

This model estimates that NPI's alone, as indicated by IMR, prevented 840,000 COVID related deaths and that, with no public health interventions, COVID would have killed 3.3 million Americans. This is consistent with the controversial early estimates from the Imperial College of London,<sup>28</sup> although that relatively simple model assumed a far more rapid spread of the disease.

The ability of this relatively simple model to explain 69% of the variability in state COVID mortality provides compelling evidence that IMR is a useful indicator for the effectiveness of NPIs against COVID and that the factors included in the model were the primary drivers of COVID mortality. Although IMR appears to be an excellent indicator of the effect of NPIs, it does not provide any insight into exactly which interventions were effective. Understanding the contribution of various NPI's to IMR will be critical to refining management strategies for future epidemics of respiratory infectious disease.



## References

1. Das, K., Behera, R.L., and Paital, B. (2022). CHAPTER 8 - Socio-economic impact of COVID-19. In *COVID-19 in the Environment*, D. Rawtani, C. M. Hussain, and N. Khatri, eds. (Elsevier), pp. 153–190. 10.1016/B978-0-323-90272-4.00014-2.
2. Lee, J. (2020). Mental health effects of school closures during COVID-19. *Lancet Child Adolesc. Health* 4, 421. 10.1016/S2352-4642(20)30109-7.
3. Panchal, U., Salazar de Pablo, G., Franco, M., Moreno, C., Parellada, M., Arango, C., and Fusar-Poli, P. (2023). The impact of COVID-19 lockdown on child and adolescent mental health: systematic review. *Eur. Child Adolesc. Psychiatry* 32, 1151–1177. 10.1007/s00787-021-01856-w.
4. Hawrilenko, M., Kroshus, E., Tandon, P., and Christakis, D. (2021). The Association Between School Closures and Child Mental Health During COVID-19. *JAMA Netw. Open* 4, e2124092. 10.1001/jamanetworkopen.2021.24092.
5. Board, T.E. (2023). Opinion | The Startling Evidence on Learning Loss Is In. *N. Y. Times*.
6. NAEP Blog - Performance Declines in Basic Mathematics and Reading Skills Since the COVID-19 Pandemic Are Evident Across Many Racial/Ethnic Groups [https://nces.ed.gov/nationsreportcard/blog/pandemic\\_performance\\_declines\\_across\\_racial\\_and\\_ethnic\\_groups.aspx](https://nces.ed.gov/nationsreportcard/blog/pandemic_performance_declines_across_racial_and_ethnic_groups.aspx).
7. Høeg, T.B., Haslam, A., and Prasad, V. (2023). An analysis of studies pertaining to masks in Morbidity and Mortality Weekly Report: Characteristics and quality of studies from 1978 to 2023. *Am. J. Med.* 0. 10.1016/j.amjmed.2023.08.026.
8. Jefferson, T., Dooley, L., Ferroni, E., Al-Ansary, L.A., Driel, M.L. van, Bawazeer, G.A., Jones, M.A., Hoffmann, T.C., Clark, J., Beller, E.M., et al. (2023). Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database Syst. Rev.* 2023. 10.1002/14651858.cd006207.pub6.
9. Hoeg, T.B., Gandhi, M., and Brown, L. (2021). Perspective | Widespread coronavirus surveillance testing at schools is a bad idea. *Wash. Post*.
10. Fukumoto, K., McClean, C.T., and Nakagawa, K. (2021). No causal effect of school closures in Japan on the spread of COVID-19 in spring 2020. *Nat. Med.* 27, 2111–2119. 10.1038/s41591-021-01571-8.
11. Herby, J., Jonung, L., and Hanke, S. (2022). A Literature Review and Meta-Analysis of the Effects of Lockdowns on COVID-19 Mortality. *Stud. Appl. Econ.*
12. Hoeg, T.B., Henderson, T.O., Johnson, D., and Gandhi, M. (2021). Our next national priority should be to reopen all America’s schools for full time in-person learning. *The Hill*. <https://thehill.com/opinion/healthcare/544142-our-next-national-priority-should-be-to-reopen-all-americas-schools-for/>.

13. Boulos, L., Curran, J.A., Gallant, A., Wong, H., Johnson, C., Delahunty-Pike, A., Saxinger, L., Chu, D., Comeau, J., Flynn, T., et al. (2023). Effectiveness of face masks for reducing transmission of SARS-CoV-2: a rapid systematic review. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* *381*, 20230133. [10.1098/rsta.2023.0133](https://doi.org/10.1098/rsta.2023.0133).
14. Murphy, C., Lim, W.W., Mills, C., Wong, J.Y., Chen, D., Xie, Y., Li, M., Gould, S., Xin, H., Cheung, J.K., et al. (2023). Effectiveness of social distancing measures and lockdowns for reducing transmission of COVID-19 in non-healthcare, community-based settings. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* *381*, 20230132. [10.1098/rsta.2023.0132](https://doi.org/10.1098/rsta.2023.0132).
15. Littlecott, H., Herd, C., O'Rourke, J., Chaparro, L.T., Keeling, M., James Rubin, G., and Fearon, E. (2023). Effectiveness of testing, contact tracing and isolation interventions among the general population on reducing transmission of SARS-CoV-2: a systematic review. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* *381*, 20230131. [10.1098/rsta.2023.0131](https://doi.org/10.1098/rsta.2023.0131).
16. Takeuchi, H., and Kawashima, R. (2023). Disappearance and Re-Emergence of Influenza during the COVID-19 Pandemic: Association with Infection Control Measures. *Viruses* *15*, 223. [10.3390/v15010223](https://doi.org/10.3390/v15010223).
17. Soo, R.J.J., Chiew, C.J., Ma, S., Pung, R., and Lee, V. (2020). Decreased Influenza Incidence under COVID-19 Control Measures, Singapore. *Emerg. Infect. Dis.* *26*, 1933–1935. [10.3201/eid2608.201229](https://doi.org/10.3201/eid2608.201229).
18. Groves, H.E., Papenburg, J., Mehta, K., Bettinger, J.A., Sadarangani, M., Halperin, S.A., Morris, S.K., Bancej, C., Burton, C., Embree, J., et al. (2022). The effect of the COVID-19 pandemic on influenza-related hospitalization, intensive care admission and mortality in children in Canada: A population-based study. *Lancet Reg. Health – Am.* *7*. [10.1016/j.lana.2021.100132](https://doi.org/10.1016/j.lana.2021.100132).
19. Morris, R. (2023). Influenza Mortality as an Indicator of the Efficacy of COVID-Related, Non-pharmaceutical Interventions to Reduce the Spread of Respiratory Infections. Preprint at medRxiv, [10.1101/2023.11.30.23299157](https://doi.org/10.1101/2023.11.30.23299157) [10.1101/2023.11.30.23299157](https://doi.org/10.1101/2023.11.30.23299157).
20. CDC (2024). The NIVD is preliminary influenza vaccination data updated weekly. *Cent. Dis. Control Prev.* <https://www.cdc.gov/flu/fluview/dashboard/vaccination-dashboard.html>.
21. CDC (2023). Weekly U.S. Influenza Surveillance Report. *Cent. Dis. Control Prev.* <https://www.cdc.gov/flu/weekly/index.htm>.
22. CDC (2020). COVID Data Tracker. *Cent. Dis. Control Prev.* <https://covid.cdc.gov/covid-data-tracker>.
23. CDC Provisional COVID-19 Deaths by Sex and Age | Data | Centers for Disease Control and Prevention. [https://data.cdc.gov/NCHS/Provisional-COVID-19-Deaths-by-Sex-and-Age/9bhg-hcku/about\\_data](https://data.cdc.gov/NCHS/Provisional-COVID-19-Deaths-by-Sex-and-Age/9bhg-hcku/about_data).

24. US Census Bureau (1910). Population Density of the 50 States, the District of Columbia, and Puerto Rico: 1910 to 2020. N. Y. <https://www2.census.gov/programs-surveys/decennial/2020/data/apportionment/population-density-data-table.pdf>.
25. National Citizens Inquiry (NCI, CeNC) [@Inquiry\_Canada] (2024). In Canada, Influenza cases plummeted from 55,379 to a mere 69 between 2020 - 2021. Dr. Stephen Bate raises an intriguing possibility, suggesting a potential renaming of the illness. He offers a visualization of PCR cycles and scrutinizes the discrepancies in vaccination rates for... <https://t.co/G6hgSvxB6q>. Twitter. [https://twitter.com/Inquiry\\_Canada/status/1754181624604790884](https://twitter.com/Inquiry_Canada/status/1754181624604790884).
26. McCulloch, D.J., Rogers, J.H., Wang, Y., Chow, E.J., Link, A.C., Wolf, C.R., Uyeki, T.M., Rolfes, M.A., Mosites, E., Sereewit, J., et al. (2023). Respiratory syncytial virus and other respiratory virus infections in residents of homeless shelters - King County, Washington, 2019-2021. *Influenza Other Respir. Viruses* 17, e13166. 10.1111/irv.13166.
27. Chow, E.J., Uyeki, T.M., and Chu, H.Y. (2023). The effects of the COVID-19 pandemic on community respiratory virus activity. *Nat. Rev. Microbiol.* 21, 195–210. 10.1038/s41579-022-00807-9.
28. WHO Collaborating Centre for Infectious Disease Modelling Report 9 - Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. Imp. Coll. Lond. <https://www.imperial.ac.uk/medicine/departments/school-public-health/infectious-disease-epidemiology/mrc-global-infectious-disease-analysis/disease-areas/covid-19/report-9-impact-of-npis-on-covid-19/>.

Table 1. Vaccination, population density, influenza and COVID mortality rates (deaths/100,000).

State	Pre-Covid Influenza Mortality Rate	2020-2022 Influenza Mortality Rate	Infection Mortality Reduction (IMR)	Vaccination Rate (2023)	Population Density (Per mi <sup>2</sup> )	3/2020 Covid Mortality	Age-Adjusted Covid Mortality
Alabama	2.55	0.77	71.0%	53%	99	0.52	422
Alaska	2.73	0.95	60.6%	65%	1	3.59	269
Arizona	2.35	0.51	77.8%	66%	63	0.35	408
Arkansas	4.09	0.88	77.7%	57%	58	0.86	418
California	2.51	0.28	88.8%	75%	254	0.07	296
Colorado	3.15	0.80	73.4%	74%	56	0.45	302
Connecticut	3.59	0.40	88.2%	83%	745	0.73	310
Delaware	2.97	0.34	87.7%	74%	508	2.55	316
Florida	1.88	0.67	66.4%	70%	402	0.12	318
Georgia	1.58	0.57	67.4%	58%	186	0.24	387
Hawaii	3.07	0.35	87.6%	82%	227	1.84	117
Idaho	4.34	0.87	77.2%	57%	22	1.34	321
Illinois	2.80	0.35	86.3%	72%	231	0.21	305
Indiana	3.76	0.69	80.3%	58%	189	0.38	406
Iowa	5.38	1.00	79.8%	65%	57	0.82	317
Kansas	5.25	1.07	79.0%	66%	36	0.90	346
Kentucky	4.52	0.80	81.5%	60%	114	0.58	457
Louisiana	2.52	0.60	75.7%	55%	108	0.58	397
Maine	5.16	0.68	85.6%	84%	44	1.89	190
Maryland	2.03	0.37	83.0%	80%	636	0.43	305
Massachusetts	3.27	0.79	75.3%	85%	901	0.38	281
Michigan	3.18	0.65	79.3%	63%	178	0.26	349
Minnesota	3.68	0.61	82.4%	72%	72	0.46	266
Mississippi	2.77	1.17	57.3%	54%	63	0.90	508
Missouri	4.55	0.78	81.5%	59%	90	0.43	370
Montana	4.86	1.06	76.7%	59%	8	2.33	332
Nebraska	4.77	1.16	72.9%	67%	26	1.33	300
Nevada	1.62	0.53	69.1%	64%	28	0.82	423
New Hampshire	4.04	0.82	77.1%	72%	154	1.88	202
New jersey	1.94	0.25	86.9%	79%	1263	0.28	377
New Mexico	2.79	0.76	74.7%	76%	18	1.25	413
New York	1.64	0.35	78.2%	81%	429	0.13	385
North Carolina	3.47	0.44	86.7%	67%	215	0.24	332
North Dakota	3.32	1.66	49.0%	59%	11	3.36	404
Ohio	3.46	0.49	85.1%	61%	289	0.22	398
Oklahoma	4.51	1.57	64.1%	61%	58	0.65	487
Oregon	5.55	0.34	92.8%	73%	44	0.62	206
Pennsylvania	3.44	0.73	77.5%	74%	291	0.20	356
Rhode Island	4.78	0.50	88.4%	88%	1061	2.40	328
South Carolina	3.54	0.61	82.1%	60%	170	0.49	387
South Dakota	5.29	1.69	64.6%	67%	12	2.87	369
Tennessee	3.31	1.16	65.8%	56%	168	0.37	446
Texas	2.15	0.67	68.7%	64%	112	0.09	433
Utah	2.07	0.50	74.8%	67%	40	0.77	252
Vermont	5.41	0.46	90.7%	86%	70	4.07	129
Virginia	2.36	0.47	79.1%	77%	219	0.30	281
Washington	4.61	0.26	93.6%	76%	116	0.34	207
West Virginia	5.07	1.30	73.1%	60%	75	1.49	404
Wisconsin	3.91	0.52	85.8%	68%	109	0.45	271
Wyoming	3.48	0.77	75.0%	53%	6	4.51	326
Average	3.50	0.72	78%	68%	207	1.05	337

Table 2. Pairwise correlations for model variables.

	COVID mortality	Initial COVID	IMR	Vaccination Rate	Population Density
COVID mortality	1.00				
Initial COVID	-0.29	1.00			
IMR	-0.54	-0.18	1.00		
Vaccination Rate	-0.64	0.03	0.54	1.00	
Population Density	-0.05	-0.18	0.34	0.57	1.00

Table 3. Multiple linear regression results for US state, age adjusted COVID mortality rates as a function of Influenza Mortality Reduction (IMR), vaccination rate (at least 2 doses), COVID mortality in the first month of the pandemic, and population density. Model 2 introduces a term for the interaction between IMR and population density.

	State Characteristics	Coef.	95% C.I.		P> t
Model 1 $r^2= 0.65$	Initial COVID Mortality	-20.8	-34.8	-6.9	0.004
	Population Density	0.12	0.059	0.19	<0.001
	Influenza Mortality Reduction	-336	-525	-147	0.001
	Vaccination Rate	-577	-788	-367	<0.001
	Constant	986	843	1129	<0.001
Model 2 $r^2= 0.69$	Initial COVID Mortality	-29	-44.1	-14.6	<0.001
	Population Density	-0.83	-1.58	-0.08	0.03
	Influenza Mortality Reduction	-493	-710	-276	<0.001
	Vaccination Rate	1.12	0.24	2.00	<0.001
	IMR x Population Density	-523	-726	-320	0.01
	Constant	1087.31	930.67	1243.94	<0.001

Table 4. Model estimates of lives saved by NPI's (based on IMR as an indicator) and vaccination.

Mortality data in model based on values when US COVID mortality was 1.04 million.

	<u>COVID Deaths without specified Interventions</u>			<u>Lives Saved by Specified Intervention</u>	
	Estimate	95% CI		Estimate	95% CI
NPI	1,990,000	1,320,000 - 2,660,000		850,000	180,000 - 1,520,000
Vaccination and NPI	3,390,000	2,930,000 - 3,860,000		2,350,000	1,790,000 - 2,720,000




Figure 1. CDC estimates of deaths from influenza for past 13 flu seasons. Note that the CDC did not provide an estimate for the 2020-2021 season because the mortality rates were too low for their estimation procedures, which seek to account for unreported cases, so the number provided is the actual count.




Figure 2. Actual age adjusted COVID mortality as compared to model estimates for US states. Using influenza mortality reduction as an indicator of their aggregate efficacy, this study shows that non-pharmaceutical interventions (NPI's), vaccination rates, population density, and initial pandemic impact explains 69% of state specific COVID mortality through 2022 to assess their impact on COVID mortality.