| 1 | Understanding Food Insecurity in Kinshasa During the COVID-19 Pandemic |
|----|--------------------------------------------------------------------------------------------------------------------------------------------|
| 2 | Pierre Z. Akilimali ^{1,2,*} , Benito Kazenza ² , Francis Kabasubabo ¹ , Landry Egbende ² , Steve |
| 3 | Botomba ² , Dynah M. Kayembe ¹ , Branly K. Mbunga ² , Nguyen Toan Tran ^{3,4} and Désiré K. |
| 4 | Mashinda ⁵ |
| 5 | |
| 6 | ¹ Patrick Kayembe Research Center, Kinshasa School of Public Health, University of |
| 7 | Kinshasa, Kinshasa, P.O. Box 11850, Democratic Republic of the Congo; |
| 8 | dirchkayembe@gmail.com (D.M.K.); fkabasu13@gmail.com (F.K.) |
| 9 | ² Department of Nutrition, Kinshasa School of Public Health, University of Kinshasa, |
| 10 | Kinshasa, P.O. Box 11850, Democratic Republic of the Congo; landry.egbende@unikin.ac.cd |
| 11 | (L.E.); benito.kazenza@unikin.ac.cd (B.K.); steve.botomba@unikin.ac.cd (S.B.); |
| 12 | branly.mbunga@unikin.ac.cd (B.J.M) |
| 13 | ³ Australian Centre for Public and Population Health Research, Faculty of Health, University |
| 14 | of Technology Sydney, Sydney, P.O. Box 123, NSW 2007, Australia; |
| 15 | nguyentoan.tran@uts.edu.au |
| 16 | ⁴ Faculty of Medicine, University of Geneva, Rue Michel-Servet 1, 1206 Geneva, |
| 17 | Switzerland; nguyentoan.tran@uts.edu.au |
| 18 | ⁵ Department of Biostatistics and Epidemiology, Kinshasa School of Public Health, University |
| 19 | of Kinshasa, Kinshasa, P.O. Box 11850, Democratic Republic of the Congo; |
| 20 | desire.mashinda@unikin.ac.cd |
| 21 | *Corresponding Author |
| 22 | Email: pierretulanefp@gmail.com |
| 23 | |

24 Abstract

25 **Introduction**: Food insecurity is a vital issue, especially in places like Kinshasa.

Additionally, food insecurity has been worsened by the COVID-19 pandemic, particularly in low- and middle-income countries. Thus, this study examined food insecurity in Kinshasa after the peak of the pandemic to understand food insecurity in post-pandemic recovery efforts and the possible implications for public health policies for future pandemics similar to COVID-19.

31 Methods: This study was conducted in Kinshasa with a representative sample of 2,160

32 households selected from 36 enumeration areas. We interviewed participants from different

33 areas and used a questionnaire to ask them about their food situation. Interviews were

34 conducted with the head of each household or their designated representative by 150 master's

35 students using tablets powered by the SurveyCTO application. Household food security status

36 was evaluated through the Household Food Insecurity Access Scale. A logistic regression

37 model was developed to assess household risk factors associated with food insecurity.

38 **Results:** Most people we talked to were over 40 years old, and many lived in households

39 with fewer than six people. About a third of the households were overcrowded. Factors

40 associated with food insecurity included being a household head aged over 50 years,

41 insufficient living space, lower socioeconomic status, and residing in slum areas (AOR: 1.38;

42 95% CI: 1.06–1.79). In 2022, 12,627,424 individuals faced food insecurity in Kinshasa,

43 including 8,829,820 individuals who experienced severe food insecurity.

44 **Conclusion:** Living conditions play a significant role in food insecurity. Governments need

45 to do more to help people, especially those living in crowded areas. To combat economic

46 restrictions that lead to food insecurity during crises, policymakers and implementing

47 partners should enhance food assistance programs, such as cash transfers and food supply

48 initiatives, focusing on overcrowded households and the informal job sector.

49

- 50 Keywords: Slum areas, Food security, Kinshasa, Post-pandemic recovery
- 51

52 Key Messages

53 What is already known on this topic

54 This study emphasizes the multifaceted nature of food security, defined as the continuous

55 access to sufficient, safe, and nutritious food, comprising availability, accessibility,

56 utilization, and stability. Food insecurity, resulting from unmet needs in any of these

57 dimensions, correlates with poor health outcomes and increased mortality. The global

58 COVID-19 pandemic exacerbated food insecurity, particularly in low- and middle-income

59 countries, with rates exceeding 50%. Factors such as poverty, living conditions, low income,

60 lack of livestock, large household size, and psychological factors contribute significantly.

61 While prior studies in the Democratic Republic of the Congo exist, they are limited, often

62 focusing on specific groups. This study aims to comprehensively assess household food

63 security in Kinshasa during the post-COVID-19 period, identifying associated factors for a

64 more nuanced understanding.

65 What this study adds

66 This study adds to the existing literature by investigating the prevalence and determinants of

67 food insecurity during a global health crisis, employing the Household Food Insecurity

68 Access Scale for assessment. It contributes novel insights by examining the prevalence and

69 severity of food insecurity in Kinshasa, the Democratic Republic of the Congo, offering a

vinique context for understanding the impact of a global health crisis on household food

71 security.

72 How this study might affect research, practice, or policy

73 The study recommends implementing cash transfer strategies for vulnerable households,

particularly those with informal jobs and young children, based on significant associations
between lower socioeconomic status and food insecurity during the COVID-19 pandemic.
Another recommendation is to expand food assistance programs for overcrowded households
and the informal job sector, addressing the high prevalence of food insecurity in slum areas.
Other social and structural determinants of food security, such as women's empowerment and
access to water and electricity, should be further researched.

80

81 **INTRODUCTION**

82 Food security is achieved "when all people, at all times, have physical, social, and economic 83 access to sufficient, safe, and nutritious food that meets their dietary needs and food 84 preferences for an active and healthy life" [1-3]. This broad definition emphasizes four 85 distinct dimensions of food security: availability, accessibility, and utilization of food, in 86 addition to the stability of each of these factors, which refers to the ability to withstand 87 shocks to the broader food system [1, 4]. Food insecurity occurs when at least one of these 88 domain needs is not met, during which the experience at the household level may be 89 temporary or longer [5–7]. Access to adequate food is a core social determinant of health, and 90 food insecurity is related to poor nutritional intake and higher mortality rates [2, 6]. Even 91 temporary reductions in food security can affect long-term health and cause a loss of human 92 capital, from which it can take years to recover [8].

The recent COVID-19 pandemic has increased the level of food insecurity worldwide, and low- and middle-income countries have been most affected [8]. More than 50% of households experienced food insecurity during the COVID-19 pandemic [2, 6–8], and at the household level, it remains a major issue in many developing countries, particularly those in Africa. Although food insecurity remains high in low- and middle-income countries, many factors, such as poverty, exacerbate food insecurity. Food insecurity is also a significant risk

99 factor for non-adherence to treatment such as antiretroviral therapy among HIV-infected 100 individuals [9]. In contrast, factors such as living situation, low income, lack of livestock, 101 high household size, and psychological situation (anxiety and depression) were principal 102 aspects associated with food insecurity [2, 5, 8, 10]. 103 Previous studies on food security have been conducted in the Democratic Republic of 104 the Congo (DRC) over the past decade, but they have often been limited in scope, focused on 105 specific demographic groups, or conducted on a relatively small scale. Further, these studies 106 predate the COVID-19 pandemic and are often characterized by descriptive rather than 107 analytical approaches. In the first half of 2020, Performance Monitoring for Action reported a 108 40% prevalence of food insecurity in the city of Kinshasa [11]. However, all urban residents 109 do not uniformly experience food insecurity, and marginalized cities, commonly referred to 110 as "slums," represent the most significant examples of urban poverty in developing nations 111 and are often the most impacted. Consequently, there is a pressing need to conduct a 112 comprehensive assessment of food security on a broader scale, especially following the peak 113 of the COVID-19 pandemic, and to delve into the factors contributing to food insecurity. 114 Slums are typically defined by homes that are hazardous, unhealthy, unstable, and 115 overcrowded, without access to basic infrastructure and services. Slum residents in the city 116 experience more health issues compared to non-slum residents [12]. As urban populations 117 grow in developing countries, with the increasing population in informal urban areas in 118 megacities worldwide, specific urban health intervention measures are crucial, particularly 119 during times of crisis [12]. It is crucial to research and compare food insecurity levels 120 between slum and non-slum residents, especially considering the health implications of 121 urbanization and the potential exacerbation of these differences by events like the COVID-19 122 pandemic. The informal sector has grown vital for many individuals in the DRC due to the 123 fast population expansion and a persistent shortage of legitimate job opportunities. Just 2.5%

| 124 | of employees work in the formal sector, and in 2021, the informal economy accounted for |
|-----|-----------------------------------------------------------------------------------------------|
| 125 | around 42% of the nation's GDP [13]. The lockdown during the COVID-19 pandemic |
| 126 | impacted workers in the informal sector, potentially affecting their food security. |
| 127 | Consequently, it is essential to comprehend food insecurity in post-pandemic recovery efforts |
| 128 | and its possible implications for public health policies for future pandemics similar to |
| 129 | COVID-19. Hence, the primary objective of this study is to gain insights into the status of |
| 130 | food insecurity at the household level in Kinshasa during the COVID-19 pandemic and to |
| 131 | identify the associated factors. |
| 132 | |

133 METHODS

134 Settings and conceptual framework

135 The DRC is the fourth most populous country in Africa [14]. Kinshasa, the capital of the

136 DRC, is classified as one of the world's "megacities." In 2022, the metro area population of

137 Kinshasa was comprised of 15,628,000 individuals [15]. Subsequently, Kinshasa ranks as

138 Africa's third-largest metropolis, following Lagos and Cairo, and is among the continent's

139 fastest-growing urban regions [14]. Kinshasa is segmented into 35 Health Zones, each of

140 which is further divided into Health Areas. Insufficient access to water and sanitation,

141 together with inadequate hygiene practices, malnutrition, and food insecurity, are identified

142 as some of the primary risk factors contributing to mortality and disability in the country. The

143 DRC reported its initial confirmed case of COVID-19 on March 10, 2020. As of September

144 3, 2022, there were a total of 92,942 cases, with 92,940 confirmed cases and 1,357 deaths,

145 resulting in a case-fatality rate of 1.5% [16].

146 This study aligns with the conceptual framework of food security, which encompasses 147 food availability influenced by crop production, livestock holdings, and market access; food 148 access involving off-farm income and credit access; vulnerability to food shortages due to

149 shock, livestock holding, and income levels; and utilization determined by age, sex, education

- 150 level, and household size.
- 151 Study design

152 We conducted a community-based cross-sectional study spanning July 27 to August 3, 2022.

153 The survey had a two-stage cluster sampling design. Census enumeration areas (EAs) were

154 randomly selected in the first stage using the National Statistical Institute sampling frame.

155 Data were collected in 36 EAs in Kinshasa. Each EA was divided into "segments" to

streamline the field workers' efforts. Each segment was intended to consist of approximately

157 500 households and be surveyed by a specific team. The number of segments in each EA was

158 calculated by dividing the total number of households in the EA by the average segment size

159 of 500 households. Within the selected EA or segment, a listing of all households was

160 obtained and used to randomly select 60 households (second stage). In the selected

161 households, the head of the household or their designated representative was interviewed on

162 the day of the survey. In total, 2,160 households were sampled across 36 EAs, with 60

163 households chosen from the EAs or segments. Finally, this study included 2,020 households

164 (response rate of 94%).

165 **Data collection**

166 Data collection was conducted by 150 master's students from the Kinshasa School of Public

167 Health (KSPH) using tablets with the SurveyCTO program. They were taught research tools,

168 ethics, and linguistic issues. The master's students, who acted as interviewers, also received

169 instruction on how to administer questionnaires during two training sessions conducted by the

- 170 research team as part of their training. Interviews were conducted in Lingala, the local
- 171 language in Kinshasa, or French. We employed reverse translation, with the assistance of

172 bilingual academics, to assure linguistic and conceptual equivalency while translating things

173 from French to Lingala. Information was gathered and examined anonymously. The survey

| 174 | questionnaire did not include any personal identities of the participants. The primary |
|------|-------------------------------------------------------------------------------------------------|
| 175 | respondent at the household level was either the head of the household or their designated |
| 176 | representative on the day of the survey. If a selected household was inaccessible or lacked a |
| 177 | capable person to participate in the interview, interviewers were directed to make three |
| 178 | separate visits at various times before considering the residence as absent or vacant. |
| 179 | KSPH teaching staff that act as supervisors played a vital role in overseeing the |
| 180 | interviewers and maintaining data quality in the field. The collected data were routinely |
| 181 | checked by supervisors before being transmitted to the server. This process occurred every |
| 182 | evening during the data collection phase. Supervisors conducted quality control visits to |
| 183 | verify the correctness and completeness of data. It was a method of verifying if the master's |
| 184 | student had interviewed the appropriate respondents effectively. Quality control visits were |
| 185 | conducted on 5% of the households in each EA. The quality check was conducted using a |
| 186 | brief questionnaire that solely contained questions from the Household Food Insecurity |
| 187 | Access Scale (HFIAS). |
| 188 | The master's students were trained to verify the completeness and quality of their work, |
| 189 | and the supervisors reviewed all data forms before their submissions. Forms containing |
| 190 | omissions and clear errors were sent back to the master's student through their supervisor for |
| 191 | rectification or further review. The forms were checked for errors or inconsistencies at the |
| 192 | time of data entry. Additionally, the central team conducted daily data visualization using the |
| 193 | SurveyCTO server to provide timely feedback throughout the data collection period. |
| 194 | |
| 195 | Measurements |
| 10.5 | a |

196 Sociodemographic variables

197 The sociodemographic variables included age, sex, ethnicity, size of household, annual198 household income, relationship status, employment status at the time of the survey, and

199 whether the housing was located in an informal settlement (slums) [17]. Overcrowded 200 households were defined as those with four or more persons living in one room. Conversely, 201 a household was deemed to have sufficient living space when three or fewer people were 202 living in one room. 203 Food security 204 Household food security status was estimated using the HFIAS [18], which is a validated 205 instrument that distinguishes food-insecure households from food-secure households across 206 different cultural contexts. The Nutrition Risk Screening 2002 tool comprises nine questions 207 designed to reflect the following universal domains of the experience of food insecurity: (1) 208 anxiety and uncertainty about household food supply, (2) insufficient quality (including 209 variety and preferences for the type of food consumed), and (3) insufficient food intake and 210 its physical consequences. The current results are presented in the following categorical 211 format: (1) food secure, (2) mildly food insecure, (3) moderately food insecure, and (4) 212 severely food insecure [19]; we dichotomized these categories into food insecure versus food 213 secure. The dependent variable (household food security status) was a dichotomous variable 214 that was assigned a value of 1 if the household was food secure and 0 otherwise. The 215 Cronbach's alpha was 0.94, demonstrating excellent internal consistency. The HFIAS has 216 been utilized in prior research in the DRC [9]. Our study used the version that was previously 217 translated into Lingala and the three other local languages. 218 While household food insecurity estimates provide an overall picture of a household's 219 access to food, they may not accurately reflect individual experiences within the household. 220 Factors such as intra-household distribution of resources and varying dietary needs can 221 impact individual food security. To gain a more nuanced understanding, combining 222 household-level data with individual-level information or conducting separate assessments 223 for individuals within the household may be more informative.

224 Data analysis

| 225 | All statistical analyses were conducted using STATA 17 (StataCorp, College Station, TX). |
|------|--------------------------------------------------------------------------------------------------|
| 226 | Initially, we determined an overview of participants' sociodemographic characteristics, both |
| 227 | in their entirety and categorized by food security status. This involved employing cross- |
| 228 | tabulations and chi-square tests to identify significant differences between non-slum and slum |
| 229 | households. Significant differences in food security status were evaluated using chi-square |
| 230 | tests. To assess household susceptibility to food insecurity as the primary outcome variable, a |
| 231 | multivariable logistic regression model was developed. Factors associated with food |
| 232 | insecurity in bivariate analysis were entered into a logistic regression model to obtain |
| 233 | adjusted odds ratios (AOR) and their 95% confidence intervals (95%CI). To assess how the |
| 234 | association between slum neighborhood and food insecurity might differ by living space |
| 235 | status, an interaction term between living space status and slum residence was included in the |
| 236 | multivariable model, and the log-likelihood ratio test was used to assess its significance. |
| 237 | The Breslow-day test for assessing the interaction effect was used. If it was found to be |
| 238 | significant at $p < 0.05$, separate multivariate regression analyses were performed by type of |
| 239 | neighborhood. The interactions between living space status and slum residence, between |
| 240 | living space status and wealth index, and between wealth index and slum residence did not |
| 241 | suggest heterogeneity between slum and non-slum residents. All the statistical analyses were |
| 242 | conducted using Stata Version 17.0. SVY procedures in Stata and were used to account for |
| 243 | the sampling design and selection weights. ORs and 95% CIs were estimated from the |
| 244 | regression parameters. Variance-inflation factors were calculated to test for multicollinearity, |
| 245 | with the highest found to be 2.65. The significance level was set at $p < 0.05$. |
| 246 | Ethical considerations |
| 0.47 | |

247 The study was conducted in accordance with the principles outlined in the Helsinki

248 Declaration. The master's students were trained to obtain informed consent. Respondents

| 249 | who could not express themselves in French were offered consent forms in their preferred |
|-----|-------------------------------------------------------------------------------------------------|
| 250 | language. Ethical clearance was obtained from the ethical review board of the KSPH (no. |
| 251 | ESP/CE/71B/2022). Each participant had the informed consent form read aloud to them and |
| 252 | provided verbal consent. To standardize the informed consent process for illiterate |
| 253 | participants, we opted for verbal consent witnessed by a third party. The third party ensured |
| 254 | that the consent was read to the subject, who then freely agreed to participate in the study. |
| 255 | The participant was provided with a signed copy of the consent form to retain. The consent |
| 256 | acquisition process was sanctioned by the ethics committee. This study did not involve any |
| 257 | individuals under the age of 18. Information was gathered and examined anonymously. The |
| 258 | survey form did not include any personal identifying of participants, and the participants were |
| 259 | notified that their involvement was optional. They have the liberty to either accept, decline |
| 260 | participation, or withdraw at any time without facing any consequences. |
| 261 | |
| 262 | RESULTS |
| 263 | Participants' characteristics |
| 264 | Overall, most individuals identified as the head of the household and were at least 40 years |
| 265 | old (72%). Almost three-quarters of the participants were men (73.6%) and married (72.5%), |
| 266 | and 60% had a schooling level between secondary school and university. Table 1 illustrates |
| 267 | that the mean household size was six persons or fewer (75%), approximately one-third of the |
| 268 | households were overcrowded (29%), and the socioeconomic status (SES) distribution among |
| 269 | households was not equal. For these characteristics, the differences between people living in |
| 270 | non-slums and people living in slums were statistically significant. |
| 271 | |

271

272 **Table 1.** Sociodemographic characteristics of the head of the household

| | Non-slum | | Slum | | All | | |
|---|----------|---|------|---|-----|---|---|
| - | n | % | n | % | Ν | % | р |

| Age | | | | | | | .555 |
|-------------------------------|------------|---------------------|------------|---------------|------------|---------------|--------|
| Younger than 30 | 39 | 7.0 | 80 | 6.3 | 119 | 6.5 | |
| years | | | | | | | |
| 30-39 | 107 | 19.2 | 275 | 21.7 | 382 | 20.9 | |
| 40–49 50 | 161 | 28.9 | 377 | 29.7 | 538 | 29.5 | |
| 50 years or older | 250 | 44.9 | 537 | 42.3 | 787 | 43.1 | |
| Total | 557 | 100.0 | 1,269 | 100.0 | 1,826 | 100.0 | 001 |
| Sex | 207 | 69 5 | 082 | 750 | 1 260 | 726 | .001 |
| Male | 387 | 68.5 31.5 | 982 314 | 75.8 24.2 | 1,369 | 73.6 | |
| Female Total | 178 565 | 100.0 | | 24.2 100.0 | 492 | 26.4 100.0 | |
| Marital status | 565 | 100.0 | 1,296 | 100.0 | 1,861 | 100.0 | .037 |
| Married | 457 | 69.9 | 1,007 | 73.7 | 1 464 | 72.5 | .037 |
| | 437 58 | 8.9 | 1,007 | 8.9 | 1,464 | 8.9 | |
| Divorced/separated Widowed | 58 80 | 8.9 12.2 | 121 | 8.9 11.8 | 179 241 | 8.9 11.9 | |
| | 80 59 | 12.2 9.0 | 161 77 | 5.6 | 241 136 | 6.7 | |
| Single Total | 59 654 | 9.0 100.0 | | 5.0 100.0 | 2,020 | 0.7 100.0 | |
| | 034 | 100.0 | 1,366 | 100.0 | 2,020 | 100.0 | < .001 |
| Schooling Nothing/primory | 78 | 11.0 | 393 | 28.8 | 471 | 23.3 | < .001 |
| Nothing/primary | | 11.9 24.2 | | | | | |
| Secondary | 224 | 34.3 | 574 | 42.0 | 798 | 39.5 | |
| University | 338 | 51.7 | 353 | 25.8 | 691 | 34.2 | |
| No answer | 14 | 2.1 | 46 | 3.4 | 60 | 3.0 | |
| Total | 654 | 100.0 | 1,366 | 100.0 | 2,020 | 100.0 | . 001 |
| Religion | 170 | 27.2 | 252 | 10.4 | 420 | 01.0 | < .001 |
| Catholic | 178 | 27.2 | 252 | 18.4 | 430 | 21.3 | |
| Protestant | 112 | 17.1 | 168 | 12.3 | 280 | 13.9 | |
| Evangelic | 261 | 39.9 | 632 | 46.3 | 893 | 44.2 | |
| Other | 103 | 15.7 | 314 | 23.0 | 417 | 20.6 | |
| Total | 654 | 100.0 | 1,366 | 100.0 | 2,020 | 100.0 | 0.01 |
| Size of household | 540 | 02.0 | 0.66 | 70 7 | 1 5 1 4 | 75.0 | < .001 |
| Six persons or fewer | 548 | 83.8 | 966 | 70.7 | 1,514 | 75.0 | |
| Seven persons or | 106 | 16.2 | 400 | 29.3 | 506 | 25.0 | |
| more | 65 A | 100.0 | 1.000 | 100.0 | 2 0 2 0 | 100.0 | |
| Total | 654 | 100.0 | 1,366 | 100.0 | 2,020 | 100.0 | |
| Sufficient living space | | | | | | | < .001 |
| (not overcrowded)* | | | | | | | |
| Sufficient living | 515 | 78.8 | 916 | 67.1 | 1,431 | 70.8 | |
| space | | | | | | | |
| Overcrowding | 139 | 21.2 | 450 | 32.9 | 589 | 29.2 | 0.04 |
| Quintiles of SES | | . – | | | | • • • | < .001 |
| Lowest | 44 | 6.7 | 360 | 26.4 | 404 | 20.0 | |
| Lower | 74 | 11.3 | 330 | 24.2 | 404 | 20.0 | |
| Middle | 124 | 19.0 | 281 | 20.6 | 405 | 20.0 | |
| Higher | 159 | 24.3 | 244 | 17.9 | 403 | 20.0 | |
| Highest | 253 | 38.7 | 151 | 11.1 | 404 | 20.0 | |
| Total | 654 | 100.0 | 1,366 | 100.0 | 2,020 | 100.0 | |

273 Note: * Three or fewer persons per living room: sufficient living space; more than three

274 persons per living room: overcrowding. SES: socioeconomic status.

275

276 Food security conditions of the household

- 277 Significant differences were observed between non-slum and slum households across all food
- 278 security measurement items. More than half the households expressed concerns related to
- food scarcity, with 66.2% of all households reporting apprehensions about preferred food
- availability (non-slum: 59.0, slum: 69.6; p < .001). Additionally, 63.7% of the households
- 281 reported that a lack of resources prevented household members from consuming their
- preferred types of food (non-slum: 52.0, slum: 69.3; p < .001). Further, in less than half of the
- 283 households, there were instances in which individuals either went to bed hungry (31.0%) or
- had to endure an entire day without eating owing to insufficient food (50.4%; Table 2).
- 285
- **Table 2.** Households by residence and components of the nine food security conditions

| Questions managering food accounts | Non | -slum | Sl | um | A | 11 | |
|---------------------------------------------------------------------------------------------------------------------------------------------------|-----|-------|-----|------|-------|------|--------|
| Questions measuring food security | n | % | n | % | Ν | % | p |
| Were you concerned that your household did not have enough food? | 386 | 59.0 | 951 | 69.6 | 1,337 | 66.2 | < .001 |
| Have you or someone in your household been unable to eat the types of food you prefer because of a lack of resources? | 340 | 52.0 | 946 | 69.3 | 1,286 | 63.7 | < .001 |
| Did you or someone in your household eat a limited variety of foods because resources were not sufficient? | 331 | 50.6 | 949 | 69.5 | 1,280 | 63.4 | < .001 |
| Have you or someone in your household eaten food that you did not want to eat because of a lack of resources to obtain other types of food? | 348 | 53.2 | 964 | 70.6 | 1,312 | 65.0 | < .001 |
| Did you or someone in your household eat a smaller meal than you would have liked because there was not enough food? | 326 | 49.8 | 930 | 68.1 | 1,256 | 62.2 | < .001 |
| Did you or someone in your household eat fewer meals a day because there was not enough food? | 310 | 47.4 | 896 | 65.6 | 1,206 | 59.7 | < .001 |
| Has it ever happened that the household was without food at all because there were no resources to buy any? | 234 | 35.8 | 762 | 55.8 | 996 | 49.3 | < .001 |
| Did you or someone in your household go to bed hungry because there was not enough food? | 203 | 31.0 | 688 | 50.4 | 891 | 44.1 | < .001 |
| Did you or someone in your household go a | 195 | 29.8 | 645 | 47.2 | 840 | 41.6 | < .001 |

whole day without eating because there was not enough food?

287

288 Food security conditions of the individuals

- Table 3 illustrates that during the COVID-19 pandemic, a similar pattern emerged for
- 290 individuals residing in both non-slum and slum areas. Approximately 10% of the participants
- 291 expressed concerns regarding insufficient food availability, limited access to preferred foods,
- dietary variety, or having fewer meals. Additionally, fewer than 5% experienced the hardship
- of going to bed hungry or enduring an entire day without food. Further, in at least 5% of the
- 294 households, a complete absence of food was reported, and the absence was primarily
- attributable to resource constraints.

296

297 **Table 3.** Food security conditions

| Questions measuring food security | Non- | slum | Sh | ım | A | 11 |
|---------------------------------------------------------------------------------------------------------------------------------------------------|------|------|-----|------|-----|------|
| Questions measuring food security | n | % | n | % | Ν | % |
| Were you concerned that your household did not have enough food? | 46 | 7.0 | 129 | 9.4 | 175 | 8.7 |
| Have you or someone in your household been unable to | | | | | | |
| eat the types of food you prefer because of a lack of resources? | 53 | 8.1 | 142 | 10.4 | 195 | 9.7 |
| Did you or someone in your household eat a limited variety of foods because resources were not sufficient? | 49 | 7.5 | 150 | 11.0 | 199 | 9.9 |
| Have you or someone in your household eaten food that you did not want to eat because of a lack of resources to obtain other types of food? | 54 | 8.3 | 157 | 11.5 | 211 | 10.4 |
| Did you or someone in your household eat a smaller meal than you would have liked because there was not enough food available? | 41 | 6.3 | 143 | 10.5 | 184 | 9.1 |
| Did you or someone in your household eat fewer meals a day because there was not enough food? | 51 | 7.8 | 126 | 9.2 | 177 | 8.8 |
| Has it ever happened that the household was without food at all because there were no resources to buy any? | 24 | 3.7 | 94 | 6.9 | 118 | 5.8 |
| Did you or someone in your household go to bed hungry because there was not enough food? | 24 | 3.7 | 81 | 5.9 | 105 | 5.2 |
| Did you or someone in your household go a whole day without eating because there was not enough food? | 22 | 3.4 | 58 | 4.2 | 80 | 4.0 |

- 300 Table 4 reveals that only 19.2% of participants were classified as food secure, leaving 80.8%
- 301 (100%–19.2%) as food insecure, with 56.5% experiencing severe food insecurity.
- 302 Extrapolating these percentages to the estimated population of 15,628,000 in Kinshasa,
- 303 12,627,424 individuals (0.808 * 15,628,000) were deemed food insecure in 2022, including
- 304 8,829,820 individuals experiencing it severely (0.565 * 15,628,000).
- 305 Notably, a marked distinction emerges when comparing residents of non-slum and slum
- 306 areas within our sample in terms of severe food insecurity. Among respondents not residing
- in slums, around a third (33.8%) experienced severe food insecurity, contrasting sharply with
- 308 approximately two-thirds (67.9%) of those living in slums.
- 309
- 310 **Table 4.** Distribution of food insecurity in the total population by accessibility pillar

| Food insecurity category | Non | Non-slum | | um | All | | |
|--------------------------|-------|----------|-------|-------|--------|-------|--|
| (access) | n | % | n | % | Ν | % | |
| Food secure | 1,058 | 32.4 | 938 | 13.2 | 1,996 | 19.2 | |
| Slight food insecurity | 224 | 6.9 | 295 | 4.2 | 519 | 5.0 | |
| Moderate food insecurity | 881 | 27.0 | 1,114 | 15.7 | 1,995 | 19.2 | |
| Severe food insecurity | 1,104 | 33.8 | 4,762 | 67.0 | 5,866 | 56.5 | |
| Total | 3,268 | 100.0 | 7,109 | 100.0 | 10,377 | 100.0 | |

311

312 Factors associated with food insecurity

313 Figure 1 illustrates the factors associated with food insecurity. Specifically, individuals aged

50 years or older exhibited a significantly greater likelihood of experiencing food insecurity

315 (AOR: 2.02; 95% confidence interval (CI): 1.23–3.31) as compared to their counterparts.

316 Conversely, having a sufficient living space was associated with a reduced likelihood of food

317 insecurity (AOR: 0.56; 95% CI: 0.44–0.77). Additionally, households characterized by lower

- 318 SES (AOR lowest: 5.36; 95% CI: 3.29–8.74; AOR low: 3.30; 95% CI: 2.19–4.98; AOR
- 319 moderate: 1.97; 95% CI: 1.37–2.84; AOR higher: 1.48; 95% CI: 1.06–2.06) and residing in
- 320 slum areas displayed a heightened risk of food insecurity (AOR: 1.38; 95% CI: 1.06–1.79) as
- 321 compared to their counterparts. Living in a slum area (AOR: 1.38; 95% CI: 1.06–1.79) was

322 associated with food insecurity.

323

324 **DISCUSSION**

325 This study assessed household food insecurity during the COVID-19 pandemic. Most 326 household heads were aged at least 40 years (72%). The mean household size was six persons 327 or fewer (75%), approximately one-third of the households were overcrowded (29%), and the 328 SES distribution among households was equal. More than half the households were affected 329 by food security conditions during the pandemic, with more than half frequently facing food 330 insecurity. The prevalence of food insecurity was notably high during the pandemic, with 331 significant disparities observed between non-slum and slum areas. Being aged 50 years or 332 older, having an insufficient living space, having a certain SES (poorest, poorer, middle, or 333 wealthier), and residing in a slum area were associated with a heightened risk of food 334 insecurity. 335 These findings align with prior research demonstrating that economic shocks can lead 336 to decreased food accessibility in vulnerable households; thus, the progression of the 337 pandemic made it possible for food security to deteriorate further [2, 4, 6, 8–10, 20, 21]. 338 Lockdown measures associated with the pandemic could have disrupted food supply chains 339 and dietary habits, potentially contributing to various forms of malnutrition, including an 340 increased risk of obesity owing to the consumption of highly processed foods and reduced 341 physical activity [2]. Therefore, it is essential to closely monitor the indirect health effects of

the COVID-19 pandemic.

Food insecurity was more prevalent in informal settlement households, in which more than three-quarters of the households reported having experienced this issue. Limited access to food resources exacerbates the adverse effects of the pandemic and obstructs infection control measures. Focused social protection actions are needed against shock, accounting for

347 household profiles [2, 3, 8]. Vulnerable households, particularly those with informal jobs and 348 young children, could benefit from cash transfer strategies to protect their food security. The 349 expansion of food assistance programs, including cash transfers and food supply initiatives, 350 should target overcrowded households and the informal job sector. 351 The findings related to the worsening trend of food insecurity emphasize the urgent 352 need for policy actions addressing broader factors contributing to food insecurity, such as 353 economic development, infrastructure, and governance. Providing and evaluating not only 354 food assistance but also cash transfer initiatives, at least for the most vulnerable households, 355 is important. A cash transfer strategy for households with informal jobs and children younger 356 than five years may help protect their food security status. Monitoring food insecurity during 357 shock events, particularly among older individuals and households with informal 358 employment, is recommended [3, 8]. Addressing inequities in accessing physical and social 359 infrastructure is crucial for food security beyond household income, as evidenced by Frayne 360 and McCordic's study in South Africa [22]. Households lacking consistent access to cooking 361 fuel, medical care, electricity, or water, in addition to cash income, demonstrated 362 significantly higher odds of being categorized as food insecure. These results imply that the 363 conditions of an urban environment might better predict and explain food security than 364 income alone. Similarly, Khan et al. highlighted the significance of resilience—such as 365 infrastructure, women empowerment, economic performance, human capital, and emergency 366 workforce—in mitigating the impact of crises, such as natural disasters, in low-income 367 countries [23]. These studies emphasize the need to consider broader social and structural 368 factors in strategies to enhance food security and reduce negative outcomes in large-scale 369 crises. 370 Moreover, addressing interconnected syndemic-like crises warrants a multifaceted

approach guided by data-driven strategies grounded in a flexible and intricate systems

framework [5]. Importantly, even as the COVID-19 pandemic recedes, its far-reaching
economic, health, and societal consequences are expected to persist over an extended period
[11].

375 Notably, compared with non-slum households, slum households reported a significantly 376 greater incidence rate of food insecurity. Food insecurity affected more than three-quarters of 377 the slum households. This finding underscores that the COVID-19 pandemic has had adverse 378 effects on food security across the socioeconomic spectrum, with particularly severe 379 consequences for those with low SES. Thus, it is imperative to establish robust mechanisms 380 for monitoring the food insecurity status of households during shock events, paying particular 381 attention to older individuals and households that rely on informal employment. A study 382 conducted in Mexico during the COVID-19 pandemic recommended monitoring for food 383 insecurity in the general population, including critical vulnerable groups such as those with 384 low- and middle-SES [2, 24, 25]. However, it is crucial to acknowledge the limitation of 385 using the SES scale, which may not capture changes in economic circumstances resulting 386 from the pandemic and would instead reflect only pre-pandemic SES. 387 This study had several strengths, including the comprehensive assessment of food 388 insecurity on a large scale during the COVID-19 pandemic, which allows for generalizability 389 across all of Kinshasa and other similar urban centers in the Democratic Republic of Congo 390 and other sub-Saharan countries characterized by the presence of slums and non-slum areas. 391 However, certain limitations must be acknowledged. First, this study did not assess the 392 broader multisectoral impacts of COVID-19 beyond its influence on food security. 393 Additionally, the pre-pandemic food security conditions of households were not evaluated, 394 thereby preventing direct comparisons of conditions before and during the pandemic. 395 Despite employing random sampling, we lack the means to confirm the 396 representativeness of our sample in relation to the population of Kinshasa. The DRC last

| 397 | conducted a census in 1984, and there is no clear indication of whether our sample accurately |
|-----|-----------------------------------------------------------------------------------------------|
| 398 | represents the population of Kinshasa. Gender considerations, albeit rarely openly addressed |
| 399 | in the DRC, could impact food security in Kinshasa. This study is also limited by the lack of |
| 400 | collected information on gender. Future research should analyze how food insecurity and the |
| 401 | factors contributing to it develop, particularly in response to the economic consequences of |
| 402 | the COVID-19 pandemic. Social and structural determinants potentially contributing to food |
| 403 | security, such as women's empowerment and access to water, electricity, and medical care, |
| 404 | should be further researched. |
| 405 | |
| 406 | CONCLUSION |
| 407 | The COVID-19 pandemic impacted household food security in Kinshasa and has been a |
| 408 | primary contributor to heightened food insecurity conditions among households, potentially |
| 409 | resulting in adverse consequences. Thus, governments must develop targeted strategies aimed |
| 410 | at mitigating household vulnerability during periods of crisis. Additionally, to combat |
| 411 | economic restrictions that lead to food insecurity during crises, policymakers and |
| 412 | implementing partners might enhance food assistance programs, such as cash transfers and |
| 413 | food supply initiatives, focusing on overcrowded households and the informal job sector. |
| 414 | |
| 415 | LIST OF ABBREVIATIONS |
| 416 | AOR: Adjusted odds ratio |
| 417 | CI: Confidence interval |

- 418 DRC: Democratic Republic of the Congo
- 419 EA: Enumeration area
- 420 HFIAS: Household Food Insecurity Access Scale
- 421 KSPH: Kinshasa School of Public Health

- 422 SES: Socioeconomic status
- 423

424 **DECLARATIONS**

425 Ethics approval and consent to participate

- 426 Ethical clearance was obtained from the ethical review board of the Kinshasa School of
- 427 Public Health (KSPH; no. ESP/CE/71B/2022). Written informed consent was obtained from
- 428 each respondent during the data collection process. Privacy and confidentiality were
- 429 maintained throughout this study.

430 **Consent for publication**

431 Informed consent was obtained from all the participants involved in this study.

432 Availability of data and materials

- 433 All the relevant data for this study are available from KSPH. The materials will be made
- 434 available by the leading author upon request. Dataset can be found also in osf:
- 435 https://osf.io/h4jt2/?view_only=ad40b39d0f584bf480fb04571ed86b70

436 **Competing interests**

- 437 All authors have completed the ICMJE uniform disclosure form at
- 438 www.icmje.org/coi_disclosure.pdf and declare no support from any organization for the
- 439 submitted work; no financial relationships with any organizations that might have an interest
- 440 in the submitted work in the previous three years; and no other relationships or activities that
- 441 could appear to have influenced the submitted work.

442 Funding

- 443 This study was conducted with financial support from the World Bank: ID Projet REDISSE
- 444 IV: P167817 IDA 64980-ZR and Don IDA D5160-ZR
- 445 Authors' contributions

- 446 Conceptualization, APZ and MDK; methodology, APZ, TNT, BS, EL, KB, MBK, KF, KDM,
- 447 and MDK; software, APZ, TNT, KF, and KDM; validation, APZ, TNT, BS, EL, KB, MBK,
- 448 KF, KDM, and MDK; formal analysis, APZ, TNT, KF, KDM, and MDK; investigation,
- 449 APZ, TNT, BS, EL, KB, MBK, KF, KDM, and MDK; resources, APZ, BS, EL, KB, MBK,
- 450 KF, KDM, and MDK; data curation, APZ, KB, KF, KDM, and MDK; writing—original draft
- 451 preparation: APZ, TNT, BS, EL, KB, MBK, KF, KDM, and MDK; writing-review and
- 452 editing: APZ, TNT, BS, EL, KB, MBK, KF, KDM, and MDK; visualization, APZ, TNT, KF,
- 453 KDM, and MDK; supervision, APZ, TNT, and MDK; project administration, APZ and
- 454 MDK; and funding acquisition, APZ and MDK. All the authors have read and agreed to the
- 455 final version of the manuscript. All authors consented to the publication of the most recent
- 456 version of the present article.

457 Acknowledgments

- 458 The authors thank the KSPH students who were involved in data collection. We also thank all
- 459 respondents for sharing their experiences with us.
- 460 Author information
- 461 Pierre Akilimali, the corresponding author, is a Professor at the University of Kinshasa
- 462 School of Public Health.
- 463

464 **REFERENCES**

- 4651Fitzpatrick KM, Harris C, Drawve G, et al. Assessing food insecurity among US
- 466 adults during the COVID-19 pandemic. J Hunger Environ Nutr 2021;16:1–18.
- 467 2 Gaitán-Rossi P, Vilar-Compte M, Teruel G, et al. Food insecurity measurement and
- 468 prevalence estimates during the COVID-19 pandemic in a repeated cross-sectional
- survey in Mexico. Public Health Nutr 2021;24:412–21.

| 470 | 3 | Owens MR, Brito-Silva F, Kirkland T, et al. Prevalence and social determinants of |
|-----|----|----------------------------------------------------------------------------------------|
| 471 | | food insecurity among college students during the COVID-19 pandemic. Nutrients |
| 472 | | 2020;12:2515. |
| 473 | 4 | Abegaz KH. Determinants of food security: Evidence from Ethiopian rural household |
| 474 | | survey (ERHS) using pooled cross-sectional study. Agric & Food Secur 2017;6:70. |
| 475 | 5 | Tomita A, Ramlall S, Naidu T, et al. Major depression and household food insecurity |
| 476 | | among individuals with multidrug-resistant tuberculosis (MDR-TB) in South Africa. |
| 477 | | Soc Psychiatry Psychiatr Epidemiol 2019;54:387–93. |
| 478 | 6 | Sultana A, Kiani A. Determinants of food security at household level in Pakistan. Afr |
| 479 | | J Bus Manage 2011;5:12972–9. |
| 480 | 7 | Ngema PZ, Sibanda M, Musemwa L. Household food security status and its |
| 481 | | determinants in Maphumulo local municipality, South Africa. Sustainability |
| 482 | | 2018;10:3307. |
| 483 | 8 | Gebeyehu DT, East L, Wark S, et al. Impact of COVID-19 on the food security and |
| 484 | | identifying the compromised food security dimension: A systematic review protocol. |
| 485 | | PLOS ONE 2022;17:e0272859. |
| 486 | 9 | Musumari PM, Wouters E, Kayembe PK, et al. Food insecurity is associated with |
| 487 | | increased risk of non-adherence to antiretroviral therapy among HIV-infected adults |
| 488 | | in the Democratic Republic of Congo: A cross-sectional study. PLoS One. 2014 Jan |
| 489 | | 15;9(1):e85327. doi: 10.1371/journal.pone.0085327. PMID: 24454841; PMCID: |
| 490 | | PMC3893174. |
| 491 | 10 | Wolfson JA, Leung CW. Food insecurity and COVID-19: Disparities in early effects |
| 492 | | for US adults. Nutrients 2020;12:1648. |
| 493 | 11 | University of Kinshasa School of Public Health, the Bill & Melinda Gates Institute for |
| 494 | | Population and Reproductive Health at the Johns Hopkins Bloomberg School of |
| | | |

| 495 | | Public Health. Jhpiego. In: Performance monitoring for action (PMA) democratic |
|-----|----|-----------------------------------------------------------------------------------|
| 496 | | republic of Congo (Kinshasa & Kongo Central) phase 1: COVID-19 female follow-up |
| 497 | | survey. 11th version, PMA2020/CD-COVID-19. Baltimore, MD: DRC 2020. |
| 498 | 12 | UN-HABITATS. The Challenge of Slums: Global Report on Human Neighborhood |
| 499 | | 2003. Earthscan Publications Ltd.; London, UK: 2003. London and Sterling. [Google |
| 500 | | Scholar] |
| 501 | 13 | Italian Institute for International Political Studies. Kinshasa looks ahead: What |
| 502 | | prospects for development? December 19, 2023. |
| 503 | | https://www.ispionline.it/en/publication/kinshasa-looks-ahead-what-prospects-for- |
| 504 | | development- |
| 505 | | 157291#:~:text=In%20other%20words%2C%20the%20informal,low%2Dincome%20 |
| 506 | | countries%20in%20Africa (Accessed on March 3, 2024). |
| 507 | 14 | United Nations. World Population Prospects: The 2015 Revision, Volume II: |
| 508 | | Demographic Profiles (ST/ESA/SER.A/380) United Nations, Department of |
| 509 | | Economic and Social Affairs, Population Division; New York, NY, USA: |
| 510 | | 2015. [Google Scholar]. |
| 511 | 15 | Macrotrends. Kinshasa, Republic of Congo metro area population 1950–2024. |
| 512 | | https://www.macrotrends.net/cities/20853/kinshasa/population#:~:text=The%20curre |
| 513 | | nt%20metro%20area%20population,a%204.4%25%20increase%20from%202021 |
| 514 | | (accessed December 1, 2023). |
| 515 | 16 | RDC CMR COVID-19, SECRETARIAT TECHNIQUE : Covid-19/ Bulletin n°636 |
| 516 | | dimanche, le 19 Décembre 2021: Situation épidémologique Covid-19. [Google |
| 517 | | Scholar]. |

- 518 17 Akilimali PZ, Tran NT, Gage AJ. Heterogeneity of modern contraceptive use among
- 519 urban slum and nonslum women in Kinshasa, DR Congo: Cross-sectional analysis. Int
- 520 J Environ Res Public Health 2021;18:9400.
- 521 18 Food and Nutrition Technical Assistance Project. Household food insecurity access
- 522 scale (HFIAS) for measurement of food access: indicator guide version 3. 2007.
- 523 https://www.fantaproject.org/monitoring-and-evaluation/household-food-insecurity-
- 524 access-scale-hfias (accessed December 1, 2023).
- 525 19 Coates J, Swindale A, Bilinsky P. Household food insecurity access scale (HFIAS)
- 526 for measurement of food access: Indicator guide. Washington, DC: Food and
- 527 Nutrition Technical Assistance Project, Academy for Educational Development 2007.
- 528 20 Régis GB, Okombi FHO, Bemba ELP, et al. Management of multidrug-resistant

529 tuberculosis in Brazzaville. Health Sci Dis 2020;21:12–6 [in French].

- 530 21 Kent K, Murray S, Penrose B, et al. Prevalence and socio-demographic predictors of
- 531 food insecurity in Australia during the COVID-19 pandemic. Nutrients 2020;12:2682.
- 532 22 Frayne B, McCordic C. Planning for food secure cities: Measuring the influence of
- 533 infrastructure and income on household food security in Southern African

534 cities. Geoforum 2015;65:1–11.

- 535 23 Khan MTI, Anwar S, Batool Z. The role of infrastructure, socio-economic
- development, and food security to mitigate the loss of natural disasters. Environ Sci
 Poll Res 2022;29:52412–37.
- 538 24 Kakaei H, Nourmoradi H, Bakhtiyari S, et al. Chapter One Effect of COVID-19 on
- 539 food security, hunger, and food crisis [Internet]. Dehghani MH, Karri RR, Roy S,
- 540 editors. ScienceDirect. Elsevier; 2022. p. 3–29. Available from:
- 541 https://www.sciencedirect.com/science/article/pii/B9780323913072000055?via%3Di
- 542 hub

- 543 25 Brown H, Mills S, Albani V. Socioeconomic risks of food insecurity during the
- 544 Covid-19 pandemic in the UK: Findings from the understanding society covid survey.
- 545 BMC Public Health 2022;22:590.
- 546

547 FIGURE CAPTIONS

- 548 Figure 1. Forecast plot presenting the associated factors of food insecurity with their adjusted
- 549 ORs

