

Investigating Ethical Tradeoffs in Crisis Standards of Care through Simulation of Ventilator Allocation Protocols

Jonathan Herington^{1,2*}, Jessica Shand^{1,3}, Jeanne Holden-Wiltse⁴, Anthony Corbett⁴,
Richard Dees², Chin-Lin Ching⁵, Margie Shaw¹, Xueya Cai⁶, Martin Zand^{4,7}.

¹ Department of Health Humanities and Bioethics, University of Rochester, Rochester, NY, USA

² Department of Philosophy, University of Rochester, Rochester, NY, USA

³ Department of Pediatrics, University of Rochester, Rochester, NY, USA

⁴ Clinical and Translational Sciences Institute, University of Rochester, Rochester, NY, USA

⁵ Department of Medicine, University of Rochester, Rochester, NY, USA

⁶ Department of Biostatistics, University of Rochester, Rochester, NY, USA

⁷ Division of Nephrology, Department of Medicine, University of Rochester, Rochester, NY, USA

* Corresponding author

Email: jonathan.herington@rochester.edu (JH)

1 Investigating Ethical Tradeoffs in Crisis Standards 2 of Care through Simulation of Ventilator Allocation 3 Protocols

4 **Abstract**

5 **Introduction:** Arguments over the appropriate Crisis Standards of Care (CSC) for public health
6 emergencies often assume that there is a tradeoff between saving the most lives, saving the most life-
7 years, and preventing racial disparities. However, these assumptions have rarely been explored
8 empirically. To quantitatively characterize possible ethical tradeoffs, we aimed to simulate the
9 implementation of five proposed CSC protocols for rationing ventilators in the context of the COVID-19
10 pandemic.

11 **Methods:** A Monte Carlo simulation was used to estimate the number of lives saved and life-years saved
12 by implementing clinical acuity-, comorbidity- and age-based CSC protocols under different shortage
13 conditions. This model was populated with patient data from 3707 adult admissions requiring ventilator
14 support in a New York hospital system between April 2020 and May 2021. To estimate lives and life-
15 years saved by each protocol, we determined survival to discharge and estimated remaining life
16 expectancy for each admission.

17 **Results:** The simulation demonstrated stronger performance for age- and comorbidity-sensitive
18 protocols. For a capacity of 1 bed per 2 patients, ranking by age bands saves approximately 28.7 lives
19 and 3408 life-years per thousand patients, while ranking by Sequential Organ Failure Assessment
20 (SOFA) bands saved the fewest lives (13.2) and life-years (416). For all protocols, we observed a positive
21 correlation between lives saved and life-years saved. For all protocols except lottery and the banded
22 SOFA, significant disparities in lives saved and life-years saved were noted between White non-Hispanic,
23 Black non-Hispanic, and Hispanic sub-populations.

24 **Conclusion:** While there is significant variance in the number of lives saved and life-years saved, we did
25 not find a tradeoff between saving the most lives and saving the most life-years. Moreover, concerns
26 about racial discrimination in triage protocols require thinking carefully about the tradeoff between
27 enforcing equality of survival rates and maximizing the lives saved in each sub-population.

28 **1.0 Introduction**

29 The allocation of scarce resources during public health emergencies is often presumed to require
30 balancing several high-level ethical goals. For scarce resource allocation, potential ethical goals may
31 include (i) saving the most lives,(1) (ii) saving the most life-years,(2,3) (iii) respecting principles of non-
32 discrimination,(4) or (iv) promoting health equity.(5) Crisis Standard of Care (CSC) protocols, which
33 allocate scarce hospital resources in the setting of public health emergencies, are one such area of
34 policymaking. Published and widely recognized protocols, such as the New York State (NYS) Ventilator
35 Allocation Guidelines, the Pennsylvania Crisis Standards of Care, and the Colorado Crisis Standards of
36 Care Plan (1,3,6) are taken to encode judgements about how to weight competing ethical values. The
37 NYS guidelines use the Sequential Organ Failure Assessment (SOFA) with the stated goal of maximizing
38 survival to discharge,(1) while the Colorado guidelines use SOFA plus quantitative adjustments for
39 comorbidities to maximize life-years saved.(6) These protocols make an implicit assumption that the
40 chosen clinical criteria will promote the intended ethical goal. Moreover, they assume that tradeoffs exist
41 between goals--for example between saving the most lives and ameliorating health inequities. Because
42 these policies are implemented in complex, dynamic systems, these assumptions may turn out to be
43 false. Modelling of COVID-19 vaccine allocation strategies, for example, suggested no tradeoff between
44 vaccination strategies that saved the most lives and life-years because of the strong association between
45 age and COVID-19 mortality.(7) Rigorous and reproducible methodologies to test ethical assumptions is
46 therefore essential if CSC protocols are to serve their intended ethical goals.

47 Recent work (8–10) has attempted to simulate the implementation of CSC protocols in the setting
48 of COVID-related scarcity. Miller et. al.(9) and Wunsch et. al.(10) assessed the predictive value and
49 potential racial bias of popular protocols, but used a convenience data set of non-COVID positive
50 individuals without simulating demand or resource scarcity. Bhavani et. al.(8) used simulation methods to

51 investigate allocation protocols in the setting of scarcity, but simulated triage only for COVID+ patients
52 and did not estimate post-discharge life expectancy. Finally, Kim et. al.(11) used a discrete event
53 simulation approach, with accompanying demand models for COVID-19 patients, to train a bespoke
54 machine learning model for prioritizing critical care beds, but they did not assess the performance of
55 proposed triage protocols. None of the previously reported simulation methods attempted to evaluate the
56 tradeoff between saving the most lives to discharge and saving the most life-years.

57 We expand on this work to develop a simulation of CSC protocols that estimates tradeoffs
58 between lives saved, life-years saved, and equal allocation of resources amongst racial groups. We
59 ground our analysis in a more realistic context than prior work by evaluating previously published and
60 adopted CSC protocols in the setting of a mixed population of COVID+ and COVID- patients. Our hope is
61 that our application of quantitative simulation to a real-world clinical dataset can be broadly adapted to
62 other bioethical debates about the appropriate mechanism for resource allocation in health systems.

63 **2.0 Methods**

64 We sought to simulate the implementation of five proposed ventilator allocation protocols: 1) lottery; 2)
65 age-banding; 3) pure SOFA 4) SOFA bands (New York), 5) SOFA plus comorbidity (Colorado), and 6) a
66 different SOFA plus comorbidity model, with age as a tie breaker (Bhavani). The age-banding protocol
67 assigns priority by 10-year age bands, with ties broken by lottery. The NY SOFA protocol assigns priority
68 tiers by SOFA score (SOFA ≤ 7 = Tier 1; SOFA 8-11 = Tier 2; SOFA ≥ 12 = Tier 3), with ties broken
69 through a lottery. The Colorado protocol assigns points by SOFA band (≤ 5 SOFA = 1 pt; 6-9 SOFA = 2
70 pts; 10-12 SOFA = 3 pts, >12 SOFA = 4 pts), and by a modified Charlson comorbidity index, with ties
71 broken by lottery.(6) The Bhavani protocol (8) assigns points by SOFA bands (<9 SOFA = 1 pt; 9-11
72 SOFA = 2 pts; 12-14 SOFA = 3 pts; >14 SOFA = 4 pts) and adds +3 points to the SOFA score of any
73 patient with a “severe” comorbidity, understood as a van Walraven acute Elixhauser score >4 (i.e. the
74 upper tertile of Elixhauser scores with a “90% predicted 1 year mortality” in (12)). Ties between patients in
75 the Bhavani protocol are then broken by course-grained age-bands (0-49, 50-69, 70-84, ≥ 85) and then by
76 lottery. To simplify this study, we ignored protocol instructions that involved re-assessment and re-
77 assignment of ventilators.

78 Our study population included all critically-ill adult patients who received mechanical ventilation
79 between April 2020 and May 2021 within the University of Rochester Medical Center system (including
80 three hospitals in the Rochester metropolitan area). The URMIC Coronavirus Ethics Response Group
81 (CERG) convened in March 2020 to adapt the 2015 NYS Ventilator Allocation Guidelines into a usable
82 triage algorithm.(13) Using the Clinical Translational Science Institute BLIS database and in collaboration
83 with hospital IT, the group created a robust data workflow that allowed for the extraction and database
84 capture of clinical information from the electronic medical record system at 30 minute intervals on all
85 inpatient that could then be used as the basis for allocation decisions. For each patient, the data included
86 their ventilator status, their sequential organ failure (SOFA) score, and their admission date. Through a
87 data broker, we received deidentified data from the database combined with additional data for COVID
88 infection status, age, sex, self-identified race and ethnicity, acute comorbidities (ICD-10 diagnosis), and
89 discharge status from electronic health records for all subjects. Missing values relevant to SOFA score
90 calculation within the first 24 hours of ventilation were assumed to be normal (rather than imputed),
91 aligning with common practice and the necessity for timely data in a resource shortage.(10,14)

92 Our study involved secondary re-use of a de-identified version of this dataset. It was deemed by
93 the University of Rochester Research Subject Review Board as exempt from review and a waiver of
94 informed consent was granted. The dataset was fully anonymized before being accessed by the study
95 team, and was provided to us on the 15th of January 2022.

96 4604 admissions involving mechanical ventilation, from 4147 unique patients, were identified
97 during the study period. Of these, 897 (19.5%) were excluded because they (i) lacked the lab results
98 required to calculate an initial SOFA score within ± 24 hours of their initial intubation (n=320), (ii) had no
99 discharge disposition, ICD-10 codes or COVID test data (n=245), or (iii) had an unclear pattern of
100 admissions and intubations (n=332). The distribution of age, sex, race and COVID positivity amongst the
101 excluded patients was similar to the included patients. There was no significant difference in survival
102 rates amongst the excluded (72%) and included (72%) encounters (Table S5). Demographic features of
103 the final modelling cohort of 3707 encounters from 3512 unique patients are described in Table 1.

Table 1: Descriptive Statistics for Mechanically Ventilated Adult Patients, March 2020 to April 2021

Feature	n	%	Survival % (CI)	Age-Adjusted Survival % (CI)
Overall	3707		72 (71-74)	84 (82-86)
Sex				
Male	2197	59	73 (71 - 75)	84 (80 - 87)
Female	1510	41	72 (69 - 74)	84 (80 - 86)
Age				
<25	104	03	93 (86 - 97)	
25-34	235	06	90 (85 - 93)	
35-44	247	07	83 (77 - 87)	
45-54	456	12	81 (77 - 85)	
55-64	868	23	76 (73 - 79)	
65-74	980	26	69 (66 - 73)	
75-84	629	17	60 (54 - 64)	
>85	188	05	50 (38 - 59)	
Race				
AAPI, non-Hispanic	50	01	78 (61 - 89)	87 (24 - 97)
AIAN, non-Hispanic	3	>01	1.0	1.0
Black, non-Hispanic	577	16	76 (72 - 80)	82 (74 - 87)
Hispanic	152	04	82 (73 - 88)	89 (78 - 94)
White, non-Hispanic	2752	74	71 (69 - 74)	85 (81 - 87)
>1 Race, non-Hispanic	16	>01	75 (36 - 93)	87 (23 - 96)
Unknown, non-Hispanic	157	04		79 (81 - 87)
COVID Status				
Negative	2431	66	73 (71 - 75)	84 (81 - 86)
Positive	1276	34	71 (68 - 74)	83 (76 - 88)
SOFA Band				
1-7	3192	86	76 (74 - 77)	87 (84 - 89)
8-11	372	10	56 (49 - 63)	58 (15 - 80)
11-24	143	04	48 (34 - 59)	75 (00 - 83)

104

105 We simulated the implementation of each protocol under different levels of scarcity using a Monte

106 Carlo method. Extending the approach of Bhavani et al.,(8) our model simulates an $n/20$ (i.e. 5%,

107 10%...95% capacity) ventilator shortage by (1) randomly sampling twenty patients $\langle A, B, \dots, T \rangle$ from our

108 dataset, and assigning each patient to one of ten decision “pairs” randomly numbered 1 through 10 (e.g.

109 1:[A, R], 2:[H, C]...10: [Q, L]) , (2) ranking the patients in each pair for priority based on the relevant

110 protocol (e.g. 1:[A > R], 2:[C > H]...10:[Q>L]), (3) allocating beds to both patients in the first $n-10$ pairs

111 (i.e. if $n=15$, then assign [A=1, R=1], [C=1, H=1] etc. to first five pairs), and not allocating beds to either

112 patient in the last $10-n$ pairs (i.e. if $n=15$, then no pairs miss out on beds), 4) allocating a bed to the

113 highest priority patients in each remaining pair (e.g. [Q=1, L=0]), and 4) repeating this process until all

114 patients in our dataset have been allocated (or not) a bed. We estimated survival in each simulation by

115 observing the actual (scarcity-free) survival to discharge of patients assigned beds. We assumed that
116 patients who were not allocated a bed in our simulation did not survive. We repeated the simulation 1,000
117 times for the 50% capacity results and 250 times for all other capacity levels.

118 For each simulation, we calculated (i) survival rate, (ii) age-adjusted survival rate, and (iii)
119 aggregate comorbidity-adjusted life expectancy (in years). Due to COVID's severe age-associated
120 mortality and the distribution of age within racial groups, survival rates for sub-populations were age-
121 adjusted, and confidence intervals for both raw and age-adjusted rates were calculated using the
122 modified Gamma method of Tiwari et al.(15,16) Raw life expectancy was calculated for each subject from
123 the corresponding National Vital Statistics System (NVSS) life tables for their age, sex and race. The
124 impact of comorbidities on life expectancy was estimated by applying the adjustments previously
125 calculated by Cho et. al for Medicare recipients without a history of cancer.(17) Subjects were placed into
126 one of three comorbidity bands (none, low/medium, high) and the corresponding age adjustment in Cho
127 et. al was applied, before re-calculating the subject's life expectancy using the NVSS tables. The
128 comorbidity adjustment for a 65yr old in Cho et. al was applied to all subjects in our cohort <65yrs (we
129 explore the limitations of this approach in the discussion).

130 We then defined "lives saved" and "years of life saved" per patient for each protocol, p , and beds-
131 per-patient capacity level, ($0 < c < 1$), by first identifying the survival rate, S_B , and expected years of life
132 lived post-discharge per patient, LY_B , of our population in the baseline no scarcity scenario. Next, for each
133 capacity level, we defined the expected survival rate ($S_B \times c$) and expected years of life per patient (LY_B
134 $\times c$) as the product of the baseline rates and the relevant capacity constraint (e.g. 0.5 beds er patient).
135 Lives saved per patient, $LS_{p,c}$, is the difference between the simulated survival rate for the protocol and
136 the expected survival rate for that capacity. Years of life saved per patient, $LYS_{p,c}$, is the difference
137 between the simulated number of life-years lived per patient for that protocol, and the expected number of
138 life-years lived for that capacity.

139 All data analysis was performed in Python v3.11 using the pandas, matplotlib, numpy, scipy,
140 seaborn and statsmodels packages. De-identified data is available at

141 <https://figshare.com/s/fb7e6ac1702b5e9a95fd> . Simulation and analysis code is available at

142 <https://github.com/jcherington/triage-sim>.

143 **3. Results**

144 Our population of 3707 ventilator encounters with 3512 unique subjects had a mean (\pm std) age of 62
145 (± 16) (Table 1). Among them, Black and Hispanic patients were younger (56 ± 17 and 53 ± 18) compared to
146 non-Hispanic White patients (64 ± 15). Black patients were more likely (rate [95% CI]) to be diagnosed with
147 COVID (.45 [.41-.49]), compared to Hispanic (.39, [.32-.47]) and non-Hispanic White patients (.33, [.31-
148 .34]). In addition, SOFA scores at the initial point of intubation were higher for White subjects (mean \pm std:
149 3.69 ± 0.07) than for both Black and Hispanic subjects (3.46 ± 0.15 and 3.14 ± 0.29).

150 Contrary to prior reported results(5,8,10) in our baseline population, we did not observe significant racial
151 disparities in age-adjusted survival amongst White, Black and Hispanic patients who received mechanical
152 ventilation (Table 2). Neither did we observe significant disparities in age-adjusted survival (rate [95% CI])
153 for patients who were COVID positive (.84 [.81-.87]) and negative (.83 [.76-.88]). In a multiple variable
154 logistic regression model, SOFA score at the point of initial intubation (OR: .896, [.878-.916]), age in
155 years (OR: .964, [.958-.969]), and Elixhauser score (OR: 1.054, [1.033-1.074]) were all predictive of
156 survival, whereas sex, race and COVID-positivity were not significantly predictive (**Table S7**).

157 **3.1 Allocation Inequalities**

158 When simulating a shortage of ventilators where only one bed is available for two patients ($c=0.5$), and
159 without controlling for other factors (i.e. clinical acuity), all protocols except Lottery exhibited significant
160 racial disparities in allocation of ventilators (Table 2). Contrary to prior findings with simulated populations
161 of COVID positive patients,(8) in our simulations with mixed populations of COVID positive and negative
162 patients, allocation of ventilators favored non-Hispanic Black and Hispanic patients when compared to
163 non-Hispanic Whites (Fig S9). Allocation disparities were most pronounced for the Age-based protocol
164 and least pronounced in the context of the NY SOFA protocol.

165 3.2 Survival and Lives Saved

166 In a simulation of moderate scarcity (0.5 beds per patient), the Age-based protocol had significantly
 167 higher overall survival rates (.3916 [.3914-.3918]) than any other protocol (Table 2). On the other hand,
 168 the NY SOFA (.3760 [.3758-.3762]) protocol resulted in lower overall survival rates than all other
 169 protocols except Lottery (.3629 [.3727-.3631]), both in overall and in age-adjusted terms for every
 170 racial/ethnic sub-population. The Bhavani, Colorado, and Pure SOFA protocols exhibited significantly
 171 lower rates of age-adjusted survival for White patients compared to both their Black and Hispanic
 172 counterparts, with the biggest racial inequalities in age-adjusted survival occurring for the Colorado
 173 protocol (White: .488, Black: .524). The Age-based and Lottery protocols exhibited significantly lower age-
 174 adjusted survival for Black patients, compared to the White or Hispanic population. Unsurprisingly, the
 175 Age-based protocol heavily skewed survival towards younger Age groups, while NY SOFA and Lottery
 176 resulted in roughly similar distributions of survival across age-groups (Table S8 and Fig S10).

Table 2: Allocation and Survival by Protocol and Race and Ethnicity.

Protocol	Overall Survival (%)	Allocation Rate by Race and Ethnicity (%)			Age-Adjusted Survival Rate (%)		
		Non-Hispanic, Black	Hispanic, All Races	Non-Hispanic, White	Non-Hispanic, Black	Hispanic, All Races	Non-Hispanic, White
Baseline (100% capacity)	72.57 (70.83-74.23)	100	100	100	82.3 (74.5-87.4)	89.1 (78.4-94.1)	85 (81.4-87.4)
Lottery	36.29 (36.27-36.31)	50.1 (50.0-50.2)	50.3 (50.1-50.5)	50.0 (49.9-50.0)	41.1 (40.9-41.3)	45.0 (44.7-45.4)	42.5 (42.3-42.6)
Age	39.16 (39.14-39.18)	59.7 (59.6-59.8)	64.7 (64.5-64.9)	46.6 (46.6-46.7)	68.2 (68.1-68.2)	73.6 (73.5-73.8)	71.2 (71.1-71.2)
Pure SOFA	37.96 (37.94-37.97)	51.8 (51.7-51.9)	55.3 (55.1-55.5)	49.7 (49.7-49.7)	51.2 (51.0-51.4)	50.4 (50.1-50.7)	48.3 (48.2-48.5)
NY SOFA	37.60 (37.58-37.62)	50.5 (50.3-50.6)	51.0 (50.8-51.3)	50.0 (50.0-50.1)	44.5 (44.3-44.7)	47.1 (46.8-47.4)	44.5 (44.3-44.7)
Colorado	37.76 (37.74-37.77)	52.0 (51.9-52.1)	55.7 (55.5-55.9)	49.6 (49.6-49.6)	52.4 (52.2-52.5)	51.2 (51.0-51.5)	48.8 (48.7-48.9)
Bhavani	38.08 (38.06-38.10)	53.3 (53.2-53.4)	58.0 (57.8-58.2)	49.1 (49.1-49.2)	55.2 (55.0-55.3)	54.8 (54.6-55.0)	52.7 (52.6-52.8)

*Mean overall survival rate and 95% CIs are reported for 1000 Monte Carlo simulations at 50% scarcity.

177 Because survival rates are heavily influenced by the distribution of survival amongst sub-populations in
 178 our underlying cohort, we also estimate the number of lives saved by each protocol: i.e. the additional

179 number of lives saved by employing a protocol instead of using a lottery (Table 3 and Fig 1). In a
 180 simulation of c=0.5 beds per patient, the Age-based protocol saved more than twice as many lives per
 181 thousand patients (28.7 [28.6-28.9]) as the nearest competing protocol, Pure SOFA (16.7 [16.5-16.9]).
 182 Again, the NY SOFA protocol (13.2 [13.0-13.4]) saved the least lives per thousand patients of any
 183 protocol, both overall and for the Black (16.5 [15.4-17.5]) and Hispanic (9.9 [7.8-12.1]) sub-populations.
 184 This effect was reversed for the White sub-population, for whom the NY SOFA protocol saved the most
 185 lives (13.6, [13.3-13.9]) and the Age protocol the least (4.3 [4.0-4.6]). All protocols except NY SOFA
 186 exhibited significantly lower numbers of lives saved for White patients compared to both their Black and
 187 Hispanic counterparts, with the biggest racial inequalities in lives saved occurring for the Age protocol
 188 (White: 4.3 [4.0-4.6], Black: 99.9 [99.0-100.7], Hispanic: 143.4 [141.7-145.1]).

Table 3: Lives Saved by Protocol and Ethnicity/Race at 50% capacity

Protocol	Lives Saved per 1000 patients (95% CI)			Overall
	Non-Hispanic, Black	Hispanic, All Races	Non-Hispanic, White	
Age	99.9 (99.0-100.7)	143.4 (141.7-145.1)	4.3 (4.0-4.6)	28.7 (28.6-28.9)
Pure SOFA	40.8 (40.0-41.7)	46.3 (44.5-48.1)	12.5 (12.2-12.7)	16.7 (16.5-16.9)
NY SOFA	16.5 (15.4-17.5)	9.9 (7.8-12.1)	13.6 (13.3-13.9)	13.2 (13.0-13.4)
Colorado	41.3 (40.4-42.2)	48.3 (46.5-50.1)	9.2 (8.9-9.5)	14.7 (14.5-14.9)
Bhavani	41.3 (40.4-42.2)	48.3 (46.5-50.1)	9.2 (8.9-9.5)	18 (17.8-18.2)

189
 190 **Fig 1. Lives saved per patient by protocol: A.** Overall increase in lives saved per patient, for each
 191 protocol, at 50% scarcity. **B.** Lives saved per patient stratified by racial group, for each protocol, at 50%
 192 scarcity.

193
 194 After simulating different capacity shortages, we noted that the magnitude of the dominance of the Age
 195 protocol was sensitive to the degree of scarcity (Table S9 & Figure 2). At 0.5 beds per patient the Age
 196 based protocol saves approximately 28.7 additional lives per 1000 patients (relative to the Lottery

197 protocol), while at 0.9 beds/patient, the protocol saves only 5.6 lives per 1000 patients. The differences
 198 between NY SOFA, Colorado and the Bhavani protocols at capacities of greater than 0.85 beds per
 199 patient are not statistically significant (Table S9 & Figure 2).

200 **Fig 2. Lives saved at different levels of scarcity, for each protocol.** The greatest differences between
 201 protocols occur at moderate levels of scarcity (i.e. ~0.5 beds per patient), and differences between
 202 protocols decline at both high and low levels of scarcity.

203

204 3.3 Life-years saved

205 With respect to life-years saved, for a scarcity level of 0.5 beds per patient, the Age-based protocol saved
 206 more than twice as many life-years per thousand patients (3408 [3402-3413]) than the nearest competing
 207 protocol, Bhavani (1454 [1447-1461]) (Table 4 and Fig 3). The NY SOFA protocol saved fewer life-years
 208 than all other protocols, both overall (416 [407-424]) and for every racial/ethnic sub-population (White:
 209 377 [366-388], Hispanic: 450 [366-534], Black: 626 [591-661]). All five non-random protocols resulted in
 210 significantly fewer life-years saved for White patients when compared to both their Black and Hispanic
 211 counterparts (Table 4 and Fig 3). After simulating different capacity shortages, we again noted that the
 212 difference between Age and other protocols was sensitive to the degree of scarcity, but less so than lives
 213 saved (Table S9). For instance, at 0.9 beds/patient, the Age protocol saves only 681 [674-688] life-years
 214 per thousand patients, but continue to outperform its nearest competing protocol, Pure SOFA (299 [290-
 215 308]), by a factor of two (Table S9).

Table 4: Life-years Saved by Protocol and Ethnicity/Race at 50% capacity

Protocol	Life-years Saved per 1000 patients (95% CI)			Overall
	Non-Hispanic, Black	Hispanic, All Races	Non-Hispanic, White	
Age	5517 (5494-5540)	8531 (8478-8584)	2553 (2545-2561)	3408 (3402-3413)
Pure SOFA	1957 (1928-1985)	2245 (2172-2317)	584 (574-592)	839 (831-846)
NY SOFA	626 (591-661)	450 (366-534)	377 (366-388)	416 (407-424)

Colorado	2226 (2198-2253)	2596 (2529-2662)	698 (689-708)	999 (992-1007)
Bhavani	2916 (2890-2942)	3748 (3688-3808)	1048 (1040-1057)	1454 (1447-1461)

216

217 **Fig 3: Life years saved per patient.** A. Overall increase in life years saved per patient, for each protocol,
 218 at 50% scarcity. B. Life years saved per patient stratified by racial group, for each protocol, at 50%
 219 scarcity.

220

221 4.0 Discussion

222 In this simulation study, we found that Age-based protocols significantly outperformed all other protocols
 223 with respect to both lives saved and life-years saved. At moderate levels of scarcity (i.e. 0.5 beds per
 224 patient), we estimate that selecting Age over Lottery could save approximately 29 lives per thousand
 225 patients requiring ventilation. When scaled to a national level, this level of effectiveness compares very
 226 favorably to classic public health interventions such as seatbelt use (14,955 lives saved per year)(18) or
 227 flu vaccination (7,200 lives saved from 36 million infections in 2020, or 0.2 lives saved per 1000
 228 infections).(19) Because age-based protocols discriminate on the basis of age, they have an uncertain
 229 legal status.. Among the other four protocols, the difference between the best (Bhavani) and worst
 230 performing (NY SOFA) protocols is less pronounced but still significant, saving approximately 5 lives and
 231 1040 life-years per thousand patients at moderate levels of scarcity.

232 Notably, the level of scarcity appears to have a dramatic influence on the effect size of choosing different
 233 protocols. While the Age based protocol dominates the other protocols for both severe (0 – 0.2 beds per
 234 patient), moderate (0.4-0.6 beds per patient) and low levels of scarcity, the size of the effect diminishes
 235 substantially at low and severe scarcity. At very low levels of scarcity (e.g. 0.9 beds per patient), choosing
 236 Age over NY SOFA still saves 2.9 lives per 1000 patients. The differences between NY SOFA, Pure
 237 SOFA, Colorado and Bhavani at that level of scarcity are not statistically significant, suggesting that other
 238 ethical considerations such as complexity of implementation, transparency, and social acceptability may
 239 be decisive.

240 One possible explanation for the poor performance of the NY SOFA protocol is that it has many fewer
241 prioritization bands (i.e. only 3) than the other protocols (Age = 8 bands, Colorado = 10 bands, Pure
242 SOFA = 23 bands, Bhavani = 28 bands). In principle, more bands add greater discriminatory power to a
243 protocol and so enhance its ability to save lives at the margins. This is borne out by the significant but
244 small improvement offered by the Pure SOFA protocol. This empirical result suggests that the value of
245 treating clinically similar cases alike (by having relatively “wide” priority bands) must be weighed against
246 the value of maximizing the overall numbers of lives saved.

247 **4.1 Tradeoffs Between Lives and Life-Years**

248 Much prior work has assumed that there is an empirical tradeoff between lives saved and numbers of life-
249 years saved, thus necessitating a difficult normative decision about prioritizing different values in the
250 context of disagreement.(2) In our simulations, that tradeoff is not evident. Not only does the Age-based
251 protocol save the most lives, it also saves the most life-years. Indeed, this result appears to be robust for
252 all protocols. While there is a high degree of variance in both survival and longevity within each protocol,
253 protocols which save more lives on average also save more life-years on average.

254 This result call into question the assumed tension between maximizing survival to discharge and
255 maximizing life-years saved, but it also illuminates the different justifications one might have for focusing
256 on the distinction between lives and life-years. For utilitarians, who aim simply to maximize the aggregate
257 good, this result suggests that the distinction between lives and life-years is empirically unimportant when
258 choosing a pandemic policy. For those who hold a “fair innings” view for the importance of allocating
259 ventilators to younger patients, this result is less important – since saving people with long life
260 expectancies is an ancillary result of prioritizing patients who have yet to progress through the different
261 stages of life. Alongside the performance of the Age protocol, this result shows that prioritizing the young
262 is at least compatible with maximizing the number of lives saved.

263 **Fig 4: Tradeoffs between life years saved and lives saved.** While there is significant variance in
264 numbers of lives and life years saved, there is a strong positive correlation between each statistic for all
265 protocols.

266

267 **4.2 Tradeoffs between Performance and Racial Disparities**

268 Of those protocols examined, the protocol that offered the most equal distribution of survival rates over
269 the three racial groups had the least favorable overall performance (excluding a simple Lottery). In our
270 simulations, at all levels of scarcity and for all protocols except the NY SOFA and Lottery, the White sub-
271 population experienced significantly lower allocation rates, lives saved, and life-years saved. This finding
272 is in *prima facie* tension with previously reported simulations of triage protocols with COVID-19 positive
273 cohorts,(8) pre-pandemic cohorts,(9,10) and with the lived experience of actual COVID mortality in
274 intensive care units.(5,20) Age-adjusted rates of survival remained significantly lower for White patients,
275 and thus it is unlikely that the observed racial disparities are solely explained by the different age
276 distributions amongst White non-Hispanic, Hispanic and Black non-Hispanic patients in our sample.
277 Rather, we suspect that hospital-level population bias may explain this difference. In particular, there are
278 two major hospital systems that offer adult critical care services in our student cohort's metropolitan area,
279 and we cannot assume that the distribution of patients between these systems is the same for each sub-
280 population. Hospitals serve particular patient populations, not national averages. This highlights that
281 implementation of protocols may result in different aggregate outcomes in different contexts (i.e. with
282 national-level results coming apart from regional or hospital level outcomes).

283 More interestingly, our results suggest a tradeoff between equalizing lives saved across racial groups,
284 and maximizing the number of lives saved for each racial group. In our results, protocols other than NY
285 SOFA greatly improve the number of lives saved for Black and Hispanic populations while making only
286 marginal differences to the number of lives saved for the White population. Thus, while NY SOFA secures
287 equality in survival rates, it would deprive Black and Hispanic populations of very large gains for little
288 benefit in lives or life-years saved for White patients. Choosing NY SOFA in such cases would thus be a
289 classic case of 'levelling down' – a situation where inequality makes little difference to the absolute
290 welfare of the worst off group, but could markedly improve the welfare of other groups.(21) Notably, this
291 tradeoff is only observable by simulating the set of alternative policies in a single population, and hence
292 discussion of the value of equality in survival rates benefits from the use of simulation methods.

293 **4.3 Limitations**

294 This study has a number of important limitations. The first is that our dataset is limited to a single hospital
295 system in a region with two major healthcare institutions. As we note above, this may partially explain our
296 surprising findings regarding racial disparities in age-adjusted survival, because Black and Hispanic
297 patients with poorer social determinants of health may have been more likely to have been admitted
298 elsewhere. In future research, we hope to extract a region-wide dataset that eliminates hospital-level
299 effects.

300 Another limitation is that we did not collect, nor adjust for, the existence of a “do-not-resuscitate” order or
301 a decision to undertake compassionate withdrawal. In principle, some patients subject to DNRs may have
302 otherwise survived their admission and, given well characterized racial differences in the uptake of DNR
303 orders, this may have artificially depressed survival rates for White patients relative to other racial sub-
304 populations.(22) While this is true for survival rates, it is less likely to distort differences in the lives saved
305 and life-years saved statistics since it accounts for variation in actual mortality (including withdrawal of
306 treatment and natural expiration) and estimates deviations from “expected” survival rates for each
307 population.

308 A final limitation of our simulation is that it does not allow for “re-allocation” of ventilators. Most existing
309 protocols (including New York’s, Maryland’s and Colorado’s) allow patients a “timed trial” of ventilation,
310 and contemplate mechanisms for removing ventilators from patients who are not improving at specified
311 re-assessment timepoints.(23) The criteria for re-allocation (and the timing of re-assessments) must
312 therefore strike a balance between giving each individual a chance to recover and preventing “wastage”
313 of ventilators on patients who will not recover.(24) In this respect, the criteria for reallocation may have
314 significant effects on the survival of individual patients and the overall numbers of lives saved.

315 Unfortunately, our existing dataset and simulation strategy cannot accommodate reallocation of
316 ventilators, since it only includes SOFA scores collected at point of intubation, and thus envisages
317 allocation until survival or natural expiration. Future work should incorporate time-series datasets of
318 patient trajectories while intubated, and simulation designs that allow for “pools” of ventilators to be
319 allocated and re-allocated. We leave this extension for future work.

320 **5.0 Conclusion**

321 In this paper we have empirically tested a number of assumptions that have been at the forefront of
322 bioethical debates over the appropriate CSC protocols. In our simulation using actual patient data, we
323 show several findings. First, there are clear performance differences between protocols, and age- or
324 comorbidity-sensitive protocols appear to save more lives to discharge than protocols which rely on SOFA
325 as an indicator of clinical acuity. Second, using comorbidity-adjusted estimates of post-discharge life
326 expectancy, we show that there is unlikely to be a tradeoff between saving lives and saving life-years in
327 the aggregate. Third, we identify that while there was tradeoff between equalizing lives saved between
328 sub-populations and overall performance, those protocols with high levels of inequality were sometimes
329 better for the least racial groups than protocols with more equal distributions of lives saved. These three
330 findings cut to the heart of important bioethical debates, and should inform both philosophical and
331 implementation work to improve CSC policy.

332 Moreover, we have reproduced and extended a methodology for prospectively and retrospectively
333 analyzing the performance of different CSC protocols, under different levels of scarcity. Building on earlier
334 work by Bhavani et al, we show not only that these simulation methods can be used to estimate survival,
335 but also that they can also provide information about estimated life expectancy, inequalities between sub-
336 populations, and the optimal parameterization of models (i.e. numbers of bands, precise threshold for
337 clinical acuity scores). Nonetheless, the work of empirically exploring CSC protocols is not complete. So
338 far, no simulation protocol has been generated that includes (i) all critical care patients, (ii) is sensitive to
339 duration of resource use, and (iii) accommodates re-allocation of resources. These components are
340 critical to many currently-existing CSC protocols, and they may yield important empirical results that alter
341 or contradict the results from the static allocation simulations conducted here and elsewhere.(8) Without a
342 generalizable methodology for either post hoc or ex ante testing of the ethical assumptions underpinning
343 re-allocation of resources, we will continue to lack evidence that future allocation policies achieve their
344 intended ethical goals. We hope that the preliminary work we present in this paper spurs others to
345 develop more robust, dynamic models that can inform CSC policy choices for the next public health
346 emergency.

347 Acknowledgements

348 The authors would like to acknowledge the efforts of Data Science Masters' students - Derek Caramella,
349 Ezgi Siir Kibris, Nefle Nesli Oruc, and Walter Burnett - for assisting with initial exploration of the dataset
350 under the supervision of JH. Final data acquisition, cleaning, analysis and simulation work was conducted
351 by JH. The authors would like to thank Michael Nabozny, William Parker, Lainie Ross and the members
352 of the URMCAfrican American Patient and Family Advisory Council for comments on earlier versions of
353 the project. The project described in this publication was partially supported by the University of
354 Rochester CTSA award number UL1 TR002001 from the National Center for Advancing Translational
355 Sciences of the National Institutes of Health. The content is solely the responsibility of the authors and
356 does not necessarily represent the official views of the National Institutes of Health.

357

358 References

- 359 1. New York State Task Force on Life and the Law. Ventilator Allocation Guidelines [Internet]. New
360 York, N.Y.: NYS Department of Health; 2015 Nov [cited 2021 Jul 4]. Available from:
361 https://www.health.ny.gov/regulations/task_force/reports_publications/docs/ventilator_guidelines.pdf
- 362 2. Emanuel EJ, Persad G, Upshur R, Thome B, Parker M, Glickman A, et al. Fair Allocation of Scarce
363 Medical Resources in the Time of Covid-19. *New England Journal of Medicine*. 2020 Mar
364 23;382(21):2049–55.
- 365 3. White DB, Lo B. A Framework for Rationing Ventilators and Critical Care Beds During the COVID-19
366 Pandemic. *JAMA*. 2020 May 12;323(18):1773.
- 367 4. Silva DS. Ventilators by Lottery: The Least Unjust Form of Allocation in the Coronavirus Disease
368 2019 Pandemic. *Chest*. 2020 Sep;158(3):890.
- 369 5. Schmidt H, Roberts D, Eneanya N. Rationing, racism and justice: advancing the debate around
370 "colourblind" COVID-19 ventilator allocation. *Journal of medical ethics* [Internet]. 2022 Feb [cited
371 2022 Mar 7];48(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/33408091/>
- 372 6. Department of Public Health & Environment. Colorado Crisis Standards of Care [Internet]. 2022 [cited
373 2023 Aug 1]. Available from: <https://cdphe.colorado.gov/colorado-crisis-standards-care>
- 374 7. Goldstein JR, Cassidy T, Wachter KW. Vaccinating the oldest against COVID-19 saves both the most
375 lives and most years of life. *Proc Natl Acad Sci U S A*. 2021 Mar 16;118(11):e2026322118.
- 376 8. Bhavani SV, Luo Y, Miller WD, Sanchez-Pinto LN, Han X, Mao C, et al. Simulation of Ventilator
377 Allocation in Critically Ill Patients with COVID-19. *Am J Respir Crit Care Med*. 2021 Nov
378 15;204(10):1224–7.

- 379 9. Miller WD, Han X, Peek ME, Charan Ashana D, Parker WF. Accuracy of the Sequential Organ
380 Failure Assessment Score for In-Hospital Mortality by Race and Relevance to Crisis Standards of
381 Care. *JAMA Network Open*. 2021 Jun 18;4(6):e2113891.
- 382 10. Wunsch H, Hill AD, Bosch N, Adhikari NKJ, Rubenfeld G, Walkey A, et al. Comparison of 2 Triage
383 Scoring Guidelines for Allocation of Mechanical Ventilators. *JAMA Network Open*. 2020 Dec
384 14;3(12):e2029250–e2029250.
- 385 11. Kim J, Lim H, Ahn JH, Lee KH, Lee KS, Koo KC. Optimal Triage for COVID-19 Patients Under
386 Limited Health Care Resources With a Parsimonious Machine Learning Prediction Model and
387 Threshold Optimization Using Discrete-Event Simulation: Development Study. *JMIR Med Inform*.
388 2021 Nov 2;9(11):e32726.
- 389 12. Snow GL, Bledsoe JR, Butler A, Wilson EL, Rea S, Majercik S, et al. Comparative evaluation of the
390 clinical laboratory-based Intermountain risk score with the Charlson and Elixhauser comorbidity
391 indices for mortality prediction. *PLoS One*. 2020;15(5):e0233495.
- 392 13. University of Rochester Coronavirus Ethics Response Group. Ventilator Allocation Guidelines
393 [Internet]. Rochester, NY: University of Rochester Medical Center; 2020 May [cited 2022 Mar 9].
394 Available from:
395 [https://www.raom.org/resources/Documents/URMedicine%20Ventilator%20Allocation%20Guidelines](https://www.raom.org/resources/Documents/URMedicine%20Ventilator%20Allocation%20Guidelines_2020.pdf)
396 [_2020.pdf](https://www.raom.org/resources/Documents/URMedicine%20Ventilator%20Allocation%20Guidelines_2020.pdf)
- 397 14. Vincent JL, de Mendonca A, Cantraine F, Moreno R, Takala J, Suter PM, et al. Use of the SOFA
398 score to assess the incidence of organ dysfunction/failure in intensive care units: Results of a
399 multicenter, prospective study. *Critical Care Medicine*. 1998 Nov;26(11):1793.
- 400 15. Tiwari RC, Clegg LX, Zou Z. Efficient interval estimation for age-adjusted cancer rates. *Stat Methods*
401 *Med Res*. 2006 Dec;15(6):547–69.
- 402 16. Fay MP, Feuer EJ. Confidence intervals for directly standardized rates: a method based on the
403 gamma distribution. *Stat Med*. 1997 Apr 15;16(7):791–801.
- 404 17. Cho H, Klabunde CN, Yabroff KR, Wang Z, Meekins A, Lansdorp-Vogelaar I, et al. Comorbidity-
405 Adjusted Life Expectancy: A New Tool to Inform Recommendations for Optimal Screening Strategies.
406 *Ann Intern Med*. 2013 Nov 19;159(10):667–76.
- 407 18. National Center for Statistics and Analysis. Lives saved in 2017 by restraint use and minimum-
408 drinking-age laws. [Internet]. Washington, DC: National Highway Traffic Safety Administration; 2019
409 Mar [cited 2024 Jan 18]. (Traffic Safety Facts Crash•Stats). Report No.: DOT HS 812 683. Available
410 from: <https://crashstats.nhtsa.dot.gov/Api/Public/ViewPublication/812683>
- 411 19. CDC. Centers for Disease Control and Prevention. 2023 [cited 2024 Jan 18]. Benefits of Flu
412 Vaccination During 2019-2020 Flu Season. Available from: [https://www.cdc.gov/flu/about/burden-](https://www.cdc.gov/flu/about/burden-prevented/2019-2020.htm)
413 [prevented/2019-2020.htm](https://www.cdc.gov/flu/about/burden-prevented/2019-2020.htm)
- 414 20. Schmidt H, Roberts DE, Eneanya ND. Sequential organ failure assessment, ventilator rationing and
415 evolving triage guidance: new evidence underlines the need to recognise and revise, unjust allocation
416 frameworks. *J Med Ethics*. 2022 Feb;48(2):136–8.
- 417 21. Temkin L. Equality, Priority, and the Levelling-Down Objection. In: Clayton M, Williams A, editors.
418 *The Ideal of Equality*. Macmillan; 2000. p. 126–61.
- 419 22. Shepardson LB, Gordon HS, Ibrahim SA, Harper DL, Rosenthal GE. Racial Variation in the Use of
420 Do-Not-Resuscitate Orders. *J Gen Intern Med*. 1999 Jan;14(1):15–20.

421 23. Truog RD. Ventilator Allocation Protocols: Sophisticated Bioethics for an Unworkable Strategy.
422 Hastings Cent Rep. 2021 Sep;51(5):56–7.

423 24. Dees RH, Herington J, Chiafery M, Shand JC, D'Angio CT, Ching CL, et al. The Ethics of
424 Implementing Emergency Resource Allocation Protocols. The Journal of Clinical Ethics. 2023
425 Mar;34(1):58–68.

426

427 **Supporting Information**

428 **Table S5: Descriptive Statistics** for INCLUDED and EXCLUDED Encounters, April 2020 to May 2021

429 **Fig S5: Violin plot of age distribution** by race and ethnicity and COVID-positivity status (AAPI, AIAN,
430 Multi-Racial and Unknown sub-populations are not shown).

431 **Table S6: Distribution of SOFA Scores**

432 **Fig S6: Histogram of SOFA score distribution.** Coloring indicates NY SOFA scoring bad, Red = first
433 priority, Yellow = second priority, Blue = third priority.

434 **Fig S7: Survival rates by initial SOFA score and age.** Single explanatory variable logistic regressions
435 and 95% CI plotted. **A.** In these simple univariable logit models (Survival ~ SOFA" and "Survival ~Age"),
436 each SOFA score point at intubation is associated with a 9.25% (7.38 – 11.09%) decrease in the
437 probability of survival to discharge (Logit coeff. = -0.0971, std err = 0.010). **B.** In our population, each year
438 of subject age is associated with a 3.65% (3.11 – 4.18%) decrease in the probability of survival to
439 discharge (Logit coeff. = -0.0372, std err = 0.003).

440 **Fig S8: Initial SOFA Score and survival by race/ethnicity. A.** Distribution of SOFA score by
441 race/ethnicity **B.** In a single explanatory variable logistic regression analyzing the relationship between
442 SOFA score and survival for each racial/ethnic sub-population, each SOFA score point at intubation is
443 associated with decreases in the probability of survival to discharge of 13.15% for non-Hispanic, Black
444 patients, 3.29% (-13.2% to +8.55%) for Hispanic patients of any race, and 8.71% for non-Hispanic, White
445 patients. Logistic regressions and 95% CI plotted.

446 **Table S7: Odds ratio of survival for whole population and racial sub-populations.** A logistic
447 regression model was used to calculate the relationship between survival and six explanatory variables:

448 race (ref: White, non-Hispanic), sex-assigned-at-birth (ref: female), SOFA score, Age, COVID positivity
449 (ref:neg) and Elixhauser comorbidity index. The model was also run for each racial/ethnic sub-population.
450 Odds-ratio are reported alongside 95% confidence intervals, significant effects ($p < 0.01$) in bold.

451 **Fig S9: Racial disparities in allocation.** For each protocol, the allocation rate for each race/ethnicity are
452 shown. All rates reported are means of the underlying rates in 1000 simulations of 3700 encounters at
453 50% scarcity.

454 **Fig S10: Survival rate by Age and Protocol.**


455 **Table S8: Survival by Age Group at 50% capacity.** Mean survival and 95% CI is reported for 1000
456 Monte Carlo simulations at the indicated level of scarcity.

457 **Table S9: Survival by Protocol at different Capacities.** Mean survival and 95% CI is reported for 250
458 Monte Carlo simulations at the indicated level of scarcity. All findings within and across capacity-levels
459 are statistically significant to $P < 0.05$.

460 **Table S10: Lives Saved by Protocol at different Capacities.** Mean survival and 95% CI is reported for
461 250 Monte Carlo simulations at the indicated level of scarcity. All findings within and across capacity-
462 levels are statistically significant to $P < 0.05$.

463 **Table S11: Life-years Saved by Protocol at different Capacities.** Mean survival and 95% CI is
464 reported for 250 Monte Carlo simulations at the indicated level of scarcity. All findings within and across
465 capacity-levels are statistically significant to $P < 0.05$.

Lives Saved By Protocol



Lives Saved By Race and Protocol




Fig 1

Lives Saved at Different Levels of Scarcity





Fig 2

Years of Life Saved per Patient by Protocol (50% Scarcity)



Years of Life Saved per Patient by Race and Protocol (50% Scarcity)




Fig 3

Lives Saved by Life Years Saved (50% Scarcity)




Fig 4