"ARVs is for HIV and cream is for HPV or precancer:" Women's Perceptions and Perceived Acceptability of Self-Administered Topical Therapies for Cervical Precancer Treatment: A Qualitative Study from Kenya

Chemtai Mungo,[1,2] Aparna Ghosh Kachoria,[3] Everlyn Adoyo,[4] Graham Zulu,[5] Supreet Kaur Goraya,

[3] Jackton Omoto,[6] Cirilus Osongo,[4] Renée M. Ferrari,[2,3] Lisa Rahangdale[1,2]

Affiliations:

1. Department of Obstetrics and Gynecology, School of Medicine, University of North Carolina at

Chapel Hill, Chapel Hill, NC 27599, USA

2. Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel

Hill, NC 27599, USA

3. Department of Maternal and Child Health, Gillings School of Global Public Health, University of

North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

4. Kenya Medical Research Institute, Kisumu, Kenya

5. Butler Institute for Families, University of Denver, Denver, CO 80210, USA

6. Department of Obstetrics and Gynecology, Maseno University School of Medicine, Kisumu,

Kenya

* Correspondence:

Chemtai Mungo, MD, MPH

Chemtai_mungo@med.unc.ed

21
22

Abstract [349/350 words]

Background: Women in low- and middle-income countries (LMICs) bear a disproportionate burden

of cervical cancer, despite being a preventable disease. Prevention efforts in LMICs are hindered in

part by lack of access to cervical precancer treatment, due to weak health infrastructure and a lack of

adequate human resources to deliver current provider-administered precancer treatments. Innovative

strategies are urgently needed to close the cervical precancer treatment gap in LMICs, including the

use of self-administered topical therapies such as 5-fluorouracil and imiquimod, for which efficacy

evidence is available from high-income settings. We investigated African women's perceptions and

perceived acceptability of these therapies for cervical precancer treatment.

Methods: Between November 2022 and April 2023, we conducted five focus group discussions

(FGDs) with women ages 25-65 years undergoing cervical cancer screening or precancer treatment in

Kisumu, Kenya. The FGDs explored women's experiences with screening and precancer treatment,

their acceptability of topical therapies for precancer treatment, and perceived barriers and facilitators

to uptake. The FGDs were moderated by local qualitative research assistants, conducted in local

languages, transcribed, coded, and analyzed using qualitative description using NVIVO software.

Results: Twenty-nine women participated, with a mean age of 35.4 years (SD 6.5). All had

undergone cervical cancer screening, and 25 (83%) had a history of precancer treatment with ablation

or excision. Multiple themes were identified related to women's perceptions of topical therapies.

Participants were highly receptive of topical treatments, with many favoring the option of self-

administration compared to provider-administration of such therapies. Self-administration of topical

therapies was felt to help address challenges associated with current treatment methods, including

difficulty in access, pain with procedures, cost, and lack of privacy with pelvic exams. Participants

had a preference for topical therapies that are used less frequently compared to those used daily.

Conclusions: Among Kenyan women with a history of cervical precancer treatment, self-

administered topical therapies for precancer are acceptable and have the potential to address barriers,

including access, privacy, and cost, that hinder precancer treatment in LMICs. If supported by

efficacy studies in LMICs, self-administered topical therapies offer a scalable approach to closing the

precancer treatment gap in LMICs.

Trial registration: *Not applicable*

Keywords: cervical precancer treatment, topical therapies, self-administered treatment,

cervical cancer elimination, low- and middle-income countries, women living with HIV,

cervical cancer, sub-Saharan Africa

Background

Women in low- and middle-income countries (LMICs) shoulder a disproportionate burden of the incidence and mortality from cervical cancer, accounting for 85 percent of cases and 90 percent of deaths in 2020[1]. Additionally, women living with HIV (WLWH), the majority of whom live in LMICs, are six times more likely to develop cervical cancer and, hence, are a priority population for prevention[2], [3]. In response to this, the World Health Organization (WHO) launched the 90/70/90 global strategy to eliminate cervical cancer[4]. This strategy, adopted by most WHO member states, calls for 90% human papillomavirus (HPV) vaccination coverage of all girls by the age 15 years, 70% of women globally receiving cervical cancer screening with a high-performance test at least twice in their lifetime, and 90% of those with a positive result adequately treated by 2030[4]. Modeling studies demonstrate that achieving the 90/70/90 targets will avert 74 million new cases of cervical cancer and 62 million deaths in LMICs alone[5].

Among unvaccinated women, cervical cancer can be prevented through screening for and treating early changes in the cervix, known as cervical precancer, caused by HPV infection. Current cervical precancer treatment options include ablation or excision procedures, both of which are performed by trained healthcare professionals[6]. Despite progress in screening, access to cervical precancer treatment following abnormal screening results in LMICs is highly limited[7], [8], [9], [10][11]. In a review of the Kenya national cervical cancer screening program in 2021, only 26% of 10,983 women who screened positive for cervical precancer received treatment [12]. Similarly, between 2011 and 2015 in Malawi, only 43.3% and 31.8% of women with cervical precancer who required ablation or excision, respectively, received treatment[13]. Challenges associated with precancer treatment in LMICs include high rates of loss-to-follow-up due to cost and transportation challenges when women

screened in rural areas are referred to central facilities where treatment is available, due to a lack of

skilled healthcare providers in rural areas where most women live [9], [10], [11], [12], [14], [15]. The

failure to treat precancerous lesions while at a curable stage in these settings results in 85% of new

global cervical cancer cases occurring in LMICs, highlighting a significant disparity. This highlights

the urgent need for innovative yet resource-appropriate approaches to address the gap in cervical

precancer treatment in LMICs. One potential strategy is the use of self-administered topical therapies.

While no topical therapies are currently approved for the treatment of cervical precancer, the use of

self- or provider-administered topical therapies for cervical precancer treatment is an area of active

investigation [16], [17], [18], [19], [20], [21], [22], [23]. The feasibility, acceptability, and efficacy of

topical therapies for cervical precancer treatment has been demonstrated by several studies in high-

income countries, including randomized trials[16], [17], [20], [24]. Several of these drugs are on the

WHO List of Essential Medications and are readily available in LMICs in generic form[25]. One

such drug is Fluorouracil (5FU) cream, which has been demonstrated to be a safe and effective

cervical precancer treatment when self-administered intravaginally[16], [17]. Compared to provider-

administered precancer treatment, which is currently inaccessible for many women in LMICs,

patient-administered therapies may be a highly scalable and cost-effective cervical precancer

treatment method in these settings.

To inform ongoing (Clinicaltrials.gov identifier NCT05362955, NCT06165614, NCT05413811) and

future studies on topical therapies for cervical precancer in LMICs, studies on their acceptability and

barriers to uptake among both women and their male partners in LMICs are needed. The objective of

this study was to assess how African women receiving cervical cancer screening and precancer

treatment perceive the use of topical therapies for cervical precancer treatment and their potential

acceptability of such therapies were they to be available.

Methods

Study design and approach: This study is part of a larger project exploring the acceptability of

topical therapies for the treatment of HPV and cervical precancer, which included in-depth interviews

and focus groups with women undergoing cervical cancer screening and male partners in Kenya, in

eastern Africa. Results of a qualitative analysis of men's perspectives have been reported elsewhere

[23]. This current analysis encompasses focus group discussions with female participants.[26] We

used a constructivist paradigm to gather perspectives of women introduced to the idea of a novel

treatment method for HPV or cervical precancer. Constructivism suggests that knowledge is

constructed through individual perceptions, experiences, and social contexts [27]. We hypothesized

that acceptability of topical therapies is based on women's experiences (e.g., prior treatment

experiences, knowledge of other women's experiences) and their social contexts (e.g., relationships

with sexual partners).

We used focus group discussions (FGDs) to gather the breadth and depth of experiences from groups

of women. A predetermined sample size of five focus groups was selected based on evidence

indicating that most themes can be captured within a range of three to six focus groups [28]. Since

the topical treatment being proposed is innovative within this study's context, we conducted an

analysis of the data using qualitative description, which is highly suitable for enhancing

comprehension in a field with limited knowledge [29]. As this method remains focused on the data

itself and involves minimal interpretation, qualitative description effectively facilitated our objective

of providing a clear and direct account of the participants' perceptions, thoughts, and experiences.

Research Team: The principal investigator (CM), a Kenyan-born Obstetrician/Gynecologist with 10

years of experience, graduate students in medicine, social work, and public health (AGK, GZ, SKG),

and a senior qualitative investigator with 20 years of experience in qualitative methods and health

services research (RMF) comprised the research team. The focus groups were facilitated and

transcribed by two qualitative research assistants from the local community.

Sampling, recruitment, and data collection: We used purposive sampling and a stepped

recruitment process to recruit FGD participants, as described previously[23] [30]. Women age 25 to

65 years undergoing cervical cancer screening or precancer treatment in public clinics in western

Kenya between November 2022 and April 2023 were included in the study. Emphasis was placed on

recruiting women with a history of positive screening results or prior precancer treatment.

Participants were recruited from HIV clinics as well as clinics serving the general population. Most

women had undergone cervical cancer screening using HPV self-collection, which was available at

most clinics at the time of recruitment. Per the WHO guidelines, women who screened positive were

offered treatment with thermal ablation or referred for excision if not eligible for ablation (6). Using

focus group discussions (FGDs), we explored the women's perceptions and hypothetical

acceptability of using proposed topical, self-administered therapies for treatment of HPV or cervical

precancer, should such therapies become available for public use.

The FGDs were conducted by two female moderators from the same community as the research

participants (EA, JO). The moderators had training in qualitative research, prior experience

conducting focus group discussions, familiarity with the local context, and fluency in the local

languages. FGDs were held at facilities near the recruiting clinics and conducted in the two most

spoken local languages (*Swahili* and *Dholuo*). Discussions were guided by several domains of

inquiry: 1) baseline knowledge of HPV and cervical cancer screening and prevention, 2) the primary

treatment experience and perceived efficacy of treatment, 3) acceptability of self-administered topical

therapies as primary or adjuvant treatment to current therapies, 4) self-perceived barriers to use of

topical therapies, and 5) perceived barriers or facilitators of male partner's support for the use of

topical therapies as adjuvant treatment. Moderators used standardized language to explain cervical

cancer screening and prevention and the potential option of topical self- or provider-administered

therapies for precancer treatment. Briefly, participants were introduced to two topical therapies for

which data are available, 5-FU and Artesunate, including details on their frequency of use (5-FU

once every other week for eight applications, Artesunate daily for five days for three cycles),

abstinence requirements (two to three days of abstinence after each 5-FU application and none for

Artesunate). Participants were told that tampon use overnight was recommended following

application of the topical, and tampons were available for illustration using a pelvic model for those

who had never used one. Each FGD included 5-8 participants and lasted approximately 90 minutes.

All FGDs were audio recorded, and recordings were transcribed verbatim, translated to English, and

crosschecked to confirm accuracy.[31]

Data Analysis: A codebook was created a priori based on the focus group guide. Two coders (GZ,

SKG) read and coded two of the five FGDs to test the code application and gain a sense of additional

topics covered in the group discussions, adding emergent codes (e.g., informational needs,

interactions with health service providers) to a final codebook. All FGD transcripts were coded using

the final codebook. To ensure agreement between coders, a random sample of transcripts was chosen,

and the codes were compared for concurrence. Any inconsistencies were addressed through

discussion and mutual agreement, and any modifications made were recorded in the codebook. The

team reviewed and summarized code reports and explored the data for patterns and themes. Content

analysis and thematic development were supported using NVIVO Version 13. Although the focus

group discussions covered multiple topics, this analysis focuses on three primary topics: 1)

participants' knowledge and awareness of cervical cancer; 2) treatment preferences and comfort with

topical therapy; and 3) perceived acceptability of topical therapy for cervical precancer treatment

(Figure 1).

Figure 1. Summary of themes regarding women's perceptions of topical, self-administered therapies

for cervical precancer treatment

Knowledge/Awareness of Cervical Cancer
- Generally, participants were aware of general symptoms associated with cervical cancer
- Participants understood that intervention for treating HPV or cervical precancer was necessary
- Given individual understanding, there were mixed feelings on sharing their diagnosis with male partners

Perceived Topical Therapy Preferences/Comfort
- Many participants expressed concerns about treatment by hospital staff/health care providers in terms of respectful care and privacy; however, there was some discordance as a few participants preferred in clinic/hospital application for provider knowledge and professional application
- Participants generally favored night time application of a topical cream, especially during menstruation
- There was mild concern about the tedious nature of self application for some treatments in terms of tampon use, anatomy, and self-confidence in applying the treatment correctly

Perceived Topical Therapy Acceptability
- Participants appreciated the potential ease of topical therapy application discretely in their homes (or in clinics)
- There was higher acceptability for the treatment among women with higher awareness of cervical cancer

Results

A total of 29 women participated in five FGDs. The mean age was 35.4 years (SD 6.5). The majority,

25 (83.3%), had a history of prior precancer treatment, including during the visit they were recruited

into the study. Analysis of the FGDs identified 15 themes related to the potential use of self-

administered topical therapies for cervical precancer treatment in the study population, summarized

in Figure 2.

Figure 2. Key findings on women's general perceptions of topical therapies

Participant Experiences with Cervical Cancer Screening and Precancer Treatment

FGD participants shared their experiences of learning their HPV or cervical precancer diagnosis

following screening. Many mentioned having symptoms of pelvic pain or bleeding during intercourse

and wanting to see a doctor for screening and treatment to learn more.

"I was suspecting something was wrong because I had some pelvic pain and also some spots

whenever I had sex, I would have some blood spotting. And I used to hear that those are some

of the symptoms suggestive of cervical cancer." -R4, FGD1

Others underwent cervical cancer screening after being advised to do so by clinic staff. One woman

shared on why she had screening.

"I cannot refuse because each person just wants good health. They tested and told me that I

would be called by somebody after some time." -R6, FGD4

While there was acknowledgment and awareness of the symptoms of cervical cancer, awareness of

HPV was less common. Following notification of a positive HPV test, some participants noted that

they were initially not aware of the difference between HPV and cervical precancer or cancer, and

they often needed to seek more information to understand the differences.

"When I was told that I was HPV positive from the screening test results, I was very afraid

from that day and all I could do was to GOOGLE about it and learn as much as I could. But

what calmed me down was that when I was being treated for the HPV, I was told that having

HPV doesn't mean that I have cancer. It can be treated early before it progresses to cancer."

- R7, FGD3

Generally, participants shared that knowledge and awareness led to greater acceptance of the

recommended treatment, particularly as it relates to the difference between receiving an HPV or

cervical cancer diagnosis.

"I was told that having HPV doesn't mean I have cancer and so when I went through

treatment, I didn't feel much pain except for the day of treatment, but for a few minutes then I

was told to abstain for 6 weeks for the cervix to heal. Then I followed that, and I feel better

now." - R7, FGD3

Some women recounted their clinic experiences, noting how they felt when they had nice providers

compared to others who had previously scared them in some way, an important factor in accepting

the news of their screening and the precancer treatment they were prescribed:

"My test results for my last precancer test are out, and they are positive, and I needed to

come for more information and treatment...[the staff] lady who called talked to me...didn't

scare me she talked nicely to me then I came [to the hospital]. I was treated and I was told

the discharge will be there for one week, 10 days. But what I felt when I was being treated, I

was counseled first, I felt some cramps for some minutes, and I even screamed a little there."

-R2, FGD3

The focus groups highlighted how participants' experiences varied in learning about their screening

results and becoming aware of the treatment options available. The discussions consistently showed

that participants' understanding of HPV and cervical cancer—from prevention through screening to

treatment—played a crucial role in overcoming stigma and pursuing treatment after their diagnosis.

Experiences with Ablative or Excisional Precancer Treatment and Perceived Advantages of

Topical Therapies

During the FGDs, participants shared their experiences with traditional precancer treatments (thermal

ablation, cryotherapy, excision) and were introduced to intravaginal topical therapies (creams or

suppositories) currently being studied that can be self- or provider-administered. Participant's views

on these topical therapies, their potential integration into their lives, and comparisons with traditional

treatments were explored.

Many FGD participants showed a greater preference for topical treatments over traditional precancer

treatments, which many had undergone, citing topical therapies perceived fewer side effects,

especially pain, compared to treatments they had received.

261  *"What I feared was the LEEP [Loop electrosurgical excision procedure, a surgical precancer*

262  *treatment method] and secondly the chemoablation [thermal ablation]. A friend who came*

263  *from treatment would tell us that it is painful, and even the doctor told us that there will be*

264  *pain during the procedure, especially during heat application. And for sure, [thermal*

265  *ablation] was painful just as labor pains."* R1, FGD5

267  *"[With ablation,] during treatment, they were removing certain things [like] cotton wool and*

268  *in addition to that my sister had also said to me that there is a chemical they will spray, which*

269  *they did and I felt abdominal pain. I felt pain during treatment, but I just persevered for the*

270  *sake of treatment so that I get well."* - R4, FGD4

272 Many were happy to hear of the potential for a self-treatment option that could be done in their own

273 home, which they felt could better fit into their daily lives, offered more privacy and less discomfort

274 compared to provider-administered treatments:

275   *"I can prefer cream because that other thermos [thermal ablation] treatment or cryo*

276   *[cryotherapy], they use strange objects in the cervix and that brings tension and discomfort*

277   *because the objects going into the cervix makes you tensed and then again that type of*

278   *treatment [cryotherapy] doesn't involve one person, you find that three or two people want to*

279   *deal with your cervix and this brings some discomfort. But this one you are alone with your*

280   *husband whom you are used to, there is no fear."* R7, FGD1

282   *"I can choose [the] cream because it has some level of confidentiality, you know women don't*

283   *like it when someone is looking at her private part. So, some people can fail to go back to the*

284   *hospital because they don't want the doctor to look at their private part and with [the] cream*

285   *you apply it yourself and you are the only one who know how your private part looks like*

286   *[laughter] so I can choose this one"* – R4, FGD1

288   *"The [treatment at] the hospital that you go to be checked by the doctor, you must at least be*

289   *seen by a person when on your way there and there is no privacy there. This cream is private,"*

290   -R4, FGD1

Others cited the convenience and accessibility of a topical therapy that can be self-administered at

home as an advantage, compared to the time and costs associated with visiting clinics for provider-

administered treatments.

"[the] cream is good due to lack of transportation all the time when going for other treatment

methods. Again, I don't have to make a queue in the hospital waiting to be treated because

once I get the cream, I will be applying it by myself at home." – R3, FGD4

Some also noted a sense of increased autonomy or empowerment with use of self-administered

topical therapies, which some felt would support compliance.

"If I can be given this cream to take home, nothing can bar me from using [the] cream, I am

just being empowered and I use it accordingly." – R7, FGD2

"I think that cream is very good. You cannot fear your own body and therefore you will insert

it very well because you want to get cured fully." – R5, FGD4

Across all focus groups, the idea of a topical treatment applied at night was embraced, citing

convenience as it meant that the day's activities would be over.

"I feel [it] is better at night because you are just resting, I don't like using it daytime because I

will be walking maybe the medicine can flow [out], and that is not good." - R5, FGD2

"Applying it at night is very good because it is a time that I am retiring to bed. Secondly during

the day, I will pass urine a lot but in the night once I have put it [applied the cream? Used the

tampon] I know it is [in place?] until morning. Since I am the one who chooses my best time

that suits me…even if I have children, they would have slept by then, even if I have a husband

we will be just the two of us. I am the one to choose one that suits me." – R4, FGD3

Although most focus group discussion participants favored self-administered treatments, some

expressed a preference for provider-administered thermal ablation, emphasizing their comfort in

trusting a doctor to accurately apply the treatment to the correct area of the cervix.

"The reason I may only like the one I was done for [thermal ablation], than this [self-

administered cream], that one was done by the doctor, when she doesn't see well, she cleans

and confirms, if not well done she does it again until it reaches where she wants, but using

cream, you are in the dark, you don't know whether you have placed it well or not." – R6,

FGD2

Similarly, participants who favored having a topical treatment applied in the clinic by a healthcare

professional compared to self-application expressed confidence in a doctor's ability to administer it

more effectively than they could themselves, particularly if they encountered side effects during the

application, which may cause them to hesitate with self-administration.

"Sometimes you can decide to try it [topical therapy] a little bit and see how it is, if you find it

itching you might stop using it. And you see with the doctor he will just go ahead and apply it,

and once he applies it is done." – R3, FGD3

Generally, the majority of focus group participants were open to using a self-administered treatment

if it were accessible. Crucially, they noted that self-administered treatments at home could shield

them from adverse interactions with healthcare providers in clinics, such as being shouted at, which

some had experienced while seeking treatment for precancer.

"Women can be free to apply it [topical therapies], they have their own time without worry of

meeting a doctor, maybe one who shouted at her last time…What I felt when I was being

treated [with non-topical treatment], I was counseled first, I felt some cramps for some

minutes, and I even screamed a little there. The health providers touched my pelvic [area] and

asked me to chill."- R2, FGD3

Participant's Preferences Between Two Proposed Topical Therapies

In the FGDs, participants were introduced to two potential self-administered topical therapies for

cervical precancer treatment: topical 5FU cream and Artesunate suppositories. The differences

between the two therapies were described, including treatment length (5FU is used once every two

weeks for 8 applications over 16 weeks, while Artesunate is used nightly for 5 days, followed by a

week off, repeated for 3 cycles over six weeks), and abstinence requirements (abstinence is required

for 2 days after 5FU use, while abstinence is not required with Artesunate use). Participants were

then asked which of the two potential therapies they would prefer, if they needed to use it, based on

these described characteristics.

Those who preferred the 5FU treatment did so because of the perceived ease of the application

regimen – once every two weeks for 16 weeks, compared to daily use for Artesunate.

"What makes it [5FU] better than the other one [Artesunate] for me is that maybe you have

traveled, so you know the one for once every two weeks, even if you apply it, even if you are on

a journey, it does not worry you." – R3, FGD2

"I think 5 FU is good especially for those are not held up in their minds, there those who are

busy all the time and since the 5 FU is not complicated, for the AS [Artesunate], it is

complicated, you can forget the days, again with the menses disruption, it is not the best. I

think 5FU is the best option." – R1, FGD4

"[The 5FU] will suit me because I don't have to put it all the time." - R2, FGD3

"...[I] am comfortable with it once a week, for the daily one you might have some occasions

like funeral and finding a place for you for application may not be easy." – R7, FGD5

Others preferred Artesunate because of its use over a significantly shorter duration – only 6 weeks

compared to 16 weeks for 5FU- and the possibility that condom use may not be required with its use

"Though the [Artesunate] is a bit tedious, you are using the medication daily, but it is a shorter

period of time then it doesn't have a lot of restrictions." R6, FGD1

"The treatment that I would prefer is [Artesunate], the one where you treat for 5 days then the

following week you rest, then you also don't use a condom and it is a shorter period of

treatment than the one that goes for 16 weeks. Though the 16 weeks also have weeks when you

are skipping but it is a long period then it has condom use for the whole treatment period. So,

for me because condom will cause conflicts in my house, I would settle for [Artesunate]." – R7,

FGD1

Participants noted that their treatment preferences were influenced by their perceptions of their male

partner's opinions of such therapies, including the requirements for condom use. Many cited that

abstinence for long periods of time could be a source of conflict with their male partner's

preferences. This was cited as a reason why therapies like Artesunate, which may not require condom

use or abstinence, may be preferable over 5FU which requires both for certain periods:

"I like where there is peace...but maybe the way the husband as we were saying, they might not

understand the abstinence part and even this condom use, they usually say that they cannot use

a condom with their partners, they feel like if you insist then there is something and this alone

can cause conflicts. So, I prefer [Artesunate] even if I am applying for 5 days in peace, it is

better because I know he is going to support me, and the medication will work well than the one

where you are fighting. And you know there are some that might even end up breaking the

rules, so peace is good." – R6, FGD1

"*For me condoms can cause conflict, most men don't like using a condom and there are those*

who have never used a condom in their life." – R7, FGD1

Participants who did not favor Artesunate pointed out the inconvenience, especially the burden of

applying it every day. Furthermore, those with irregular menstrual cycles noted that 5FU was more

manageable due to its biweekly application schedule, which is simpler to follow than Artesunate's

daily regimen, which can be interrupted by irregular periods.

Considerations for Women Living with both HIV (WLWH) diagnosed with HPV

In the FGDs, participants who were living with HIV (WLWH) who had also tested positive for HPV

or cervical precancer noted feeling an increased burden. Many expressed fears about the impact of

the dual diagnosis on their children or other family members, as well as the challenges of managing

multiple medications when treating cervical precancer alongside HIV infection.

"*It also bothered me, and it stressed me out following that I am also on HIV medication, I felt*

very bad because I also infected my baby [with HIV]. I have been taking HIV medication

from 2009 up to now. So, when I imagined getting another terminal illness, I felt sad." – R6,

FGD3

"And if I consider that I had [pre]cancer and with HIV, it was double burden. The fact that

cancer can worsen and kill you I get very bad. And if I consider that I had [pre]cancer and

with HIV, it was [a] double burden and so, I decided to clear with [pre]cancer which is

curable." - R7, FGD5

One participant believed that topical treatments for cervical precancer would be insufficient due to

their concurrent HIV and HPV diagnoses. They harbored doubts about the effectiveness of such

treatments when dealing with both conditions simultaneously.

"According to me I feel the [topical] treatment is not 100% for those who have HIV, because

of our low immunity, our system is weak. So even if we are treated, we can still just get

[cancer]." – R7, FGD3

Other participants likened the use of self-administered topical therapies among HIV-positive women

to the same way WLWH are prescribed antiretroviral therapies (ARVs), which they use at home to

treat their HIV disease. The participants drew parallels between their consistent use of ARVs at home

and their potential to similarly apply self-administered topical therapies in the same settings.

"Those who are HIV+ should go for [the topical] cream because, they go to the hospital for

ARVs refill, they should take cream and use it at home just the same way they take ARVs and

adhere to its use at home." R2, FGD4

"ARVs is for HIV and cream is for HPV or precancer, so you just take your medication and

also apply your cream because they treat different things." – R7, FGD3

Generally, participants ultimately felt that the time required to apply the topical therapies was shorter

in duration in the home setting versus returning to the clinic to be treated, which greatly influenced

their perceived acceptability and desire to use topical therapies. Regardless of HIV-seropositive

status, the participants noted that if they had the knowledge and the ability to apply the cream at

home, they would be willing to do this for the betterment of their health.

Discussion

In this qualitative study evaluating Kenyan women's perceptions of topical self- or provider-

administered therapies for cervical precancer treatment, we find that participants, many of whom had

undergone traditional cervical precancer treatment, were highly receptive to topical therapies. We

found that many participants had fears following a diagnosis of HPV or cervical precancer, which

they had to overcome in order to undergo ablation or excisional treatment procedures. When

introduced to topical therapies as a potential alternative to available precancer treatments, participants

strongly favored topical therapies, citing reduced pain, improved accessibility, and privacy, compared

to the currently available provider-administered precancer treatment methods that many had

undergone. Most study participants expressed a strong preference for self-administration of topical

therapies, with many citing the lack of privacy associated with provider-administered treatments as a

barrier that those who had received precancer treatment had to overcome and that often discourages

other women from seeking treatment. Participants ' preferences varied when given an option between

two potential topical therapies with different characteristics and requirements for use. Some favored

5FU, applied every two weeks, despite its conditions for abstinence following use and consistent

condom use. Meanwhile, others favored Artesunate, which requires more frequent applications but

may have less stringent restrictions around abstinence and condom use. Despite only having had a

brief education session, participants showed high levels of awareness and body autonomy in the

discussions by displaying keen insights into potential different trade-offs associated with the two

topical therapies discussed, including the impact of irregular menstrual cycles on the ability to adhere

to a daily topical. Given the higher incidence of cervical precancer in women living with HIV, it is

noteworthy to highlight that HIV-positive participants in our study indicated concerns about

managing their HIV disease alongside a diagnosis of HPV or cervical precancer. However, most

were confident about their ability to use a self-administered topical treatment for cervical precancer,

drawing on their experience with daily use of oral antiretroviral therapy to manage HIV infection.

To our knowledge, this is the first qualitative study to explore African women's perceptions and

perceived acceptability of self- or provider-administered topical therapies for cervical precancer

treatment. In this study of urban and peri-urban Kenyan women who had undergone cervical cancer

screening and a majority of whom had undergone ablation or excisional precancer treatment, many

expressed conflicting emotions about their treatment, explicitly highlighting the challenges they had

to overcome in terms of access, pain and lack of privacy often pointing to pain and privacy issues

when receiving provider-administered treatments. Most showed a preference for topical therapies, if

available, believing they would alleviate these challenges associated with conventional treatment

methods that often deter other women from pursuing precancer treatment. The acceptability of

thermal ablation, the most widely available precancer treatment method in LMICs that was approved

by the WHO in 2019, has been demonstrated in a few studies[28], [29]. Thermal ablation, which

involves the application of a heated probe to the cervix to destroy precancerous tissue, is performed

without local anesthesia to the cervix[6]. Studies in LMICs report that while 83.9% - 90% report no

or mild pain with thermal ablation, 2.5% - 16.1% report high or moderate pain with the procedure

[28], [29]. In our qualitative findings, some participants described thermal ablation as "*too painful*"

or "*painful just as labor pains.*" Another participant noted the need to encourage women that the

procedure is "*not very painful*" and should not deter them from presenting for treatment, as the pain

perception is thought to keep women away from presenting for treatment. Studies on whether certain

women undergoing thermal ablation may require pretreatment analgesia are needed, alongside

considerations of the feasibility of providing of doing this. If topical therapies for cervical precancer

can be shown to be equally effective as ablative or excisional procedures in low- and middle-income

countries (LMICs), they could potentially alleviate the pain-related concerns associated with ablation

or excision.

Our findings of participants noting challenges with treatment access and privacy concerns associated

with provider-administered, facility-based treatments have been demonstrated in several LMIC

studies. Facility-based precancer treatment access challenges in LMICs include lack of functional

equipment or supplies[10], [13], lack of trained providers [7], [10], [13]long distance required to

access treatment facilities [11], [30]. These factors significantly contribute to the existing precancer

treatment gaps. In a study from rural Kenya, up to 40-50% of women who screened positive and

were referred to a central facility did not make their follow-up appointment [11]. Similarly, in a

qualitative study from Malawi, women with abnormal cervical cancer screening results cited lack of

transportation to referral facilities and high associated costs as major reasons for not presenting for

treatment [30]. This is reflected in our study, where women emphasized the convenience of self-

administered topical therapies that can be used at home, highlighted ease of access to topical self-

administered therapies used at home, compared to facility-based treatments, which are associated

with high transport costs and long waiting times at the facilities as a reason they would favor topical

treatments. Similarly, our findings of increased privacy as a reason women prefer self-administered

therapies to conventional treatments have been highlighted in prior studies, which found that fear of a

violation of privacy associated with pelvic exams [31], [32], [33], [34], and especially when

performed by a male provider [33], [34], [36], are barriers to screening and precancer treatment in

sub-Saharan Africa. As noted by a study participant, during her ablation procedure, "*two or three*

people want to deal with your cervix, and this brings discomfort," stating that with a self-

administered treatment, "*you are alone with your husband whom you are used to, there is no fear*."

The use of self-administered topical therapies, which women can apply in the comfort of their own

homes, can be a scalable way to address both the access challenges and privacy concerns of African

women.

Self-administered therapies can also promote women's autonomy and sense of agency, as highlighted

by our study participants, who stated that they anticipated "*feel[ing]empowered*" and would use it

correctly, as "*you cannot fear your own body*." The use of self-administered precancer treatment, if

backed by feasibility and efficacy studies in LMICs, also aligns with a recent guideline from the

World Health Organization that advocates for self-care interventions. As stated in the guideline, these

interventions have the capacity to "increase choice and autonomy," address the global shortage of

healthcare workers, and bring us closer to achieving universal health when made "accessible,

acceptable and affordable[37]." While no studies have evaluated the acceptability of self-

administered topical cervical precancer treatment in LMICs, several studies in this setting have

demonstrated high acceptability of self-care interventions, including HIV self-testing [38] and the use

of vaginal or rectal microbicides for HIV prevention[39], [40]. Similarly, in a study on the

acceptability of rectal microbicide for HIV prevention among men who have sex with men in

Thailand, ease of use, privacy, and comfort of use at home were facilitators of uptake [41], drawing

similarities to our findings.

This study has several strengths, such as the inclusion of women who have had cervical cancer

screening, as well as a deliberate oversampling of women with a history of precancer treatment. This

approach ensures that the study represents the demographic that is most likely to benefit from topical

therapies, hence whose perceptions are important in understanding acceptability. Similarly, the use of

focus groups in the qualitative design facilitated in-depth discussion among study participants who

shared similar experiences. This enabled the identification of multiple themes that impact the

acceptability of this intervention to inform feasibility studies. The study's inclusion of women living

with HIV is a significant strength due to their higher risk of cervical cancer and current unmet need

for accessible precancer treatment. A limitation of this study is that participants expressed theoretical

acceptance of the intervention but did not actually use the topical therapies. Therefore, their views

might change with actual use, an aspect future studies should explore. Another limitation was the

limited time for focus groups; more time could have offered insights into household dynamics like

decision-making and empowerment, potentially affecting women's perceptions of the therapies.

Conclusion

Our findings from Kenya indicate that women find these therapies acceptable and that they have the

potential to address significant challenges like access, privacy, and cost that hinder precancer

treatment uptake in these regions. These results support ongoing feasibility studies and call for

efficacy studies in this population to inform whether these treatments can be made available to

women.

List of Abbreviations

5FU: Fluorouracil

FGD: Focus Group Discussions

HIV: Human immunodeficiency virus

HPV: Human papillomavirus

LEEP: Loop electrosurgical excision procedure

LMIC: low-and middle-income countries

SD: standard deviation

WHO: World Health Organization

WLWH: women living with HIV

Declarations

Ethics approval and consent to participate

Ethical Considerations: The study was approved by the ethics review boards at Maseno University

School of Medicine in Kenya and the University of North Carolina Chapel Hill in the U.S.A. All

participants provided written informed consent prior to study participation.

Consent for publication: Not applicable

Availability of data and materials

Data are available upon reasonable request. For inquiries or to request access to the data, please

contact Chemtai Mungo at Chemtai_mungo@med.unc.edu.

Competing interests

The authors declare no competing financial or non-financial interests.

Funding

This research was supported by the Eunice Kennedy Shriver National Institute of Child Health &

Human Development of the National Institutes of Health under Award Number K12HD103085, the

Victoria's Secret Global Fund for Women's Cancers Career Development Award, in Partnership with

Pelotonia Foundation and the American Association of Cancer Research (AACR), and the University

of North Carolina Center for AIDS Research under award number 5-P30-AI050410. The content is

solely the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. The study funders have no role in the research.

Authors' contributions

CM conceptualized the study and associated clinical trials. EA and JO conducted focus group

discussions with study participants . GZ, SKG, RMF, and AGK worked on qualitative methodology,

with GZ and SKG conducting data analysis and RF leading methods section in this manuscript. AGK

led manuscript writing with CM and RMF. CM, AGK. EA, GZ, SKG, JO, and RMF all read and

approved the final version of the manuscript.

Acknowledgments

The research participants and leadership of Lumumba Sub-County Hospital in Kisumu, Kenya, are

acknowledged.

References

[1] H. Sung *et al.*, "Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries," *CA Cancer J Clin*, vol. 71, no. 3, pp. 209–249, May 2021, doi: 10.3322/CAAC.21660.

[2] A. B. M. Kharsany and Q. A. Karim, "HIV Infection and AIDS in Sub-Saharan Africa: Current Status, Challenges and Opportunities.," *Open AIDS J*, vol. 10, pp. 34–48, 2016, doi: 10.2174/1874613601610010034.

[3] E. Rohner *et al.*, "Cervical cancer risk in women living with HIV across four continents: A multicohort study.," *Int J Cancer*, vol. 146, no. 3, pp. 601–609, Feb. 2020, doi: 10.1002/ijc.32260.

[4] WHO, "A Global Strategy for elimination of cervical cancer." Accessed: Mar. 26, 2020. [Online]. Available: https://www.who.int/activities/a-global-strategy-for-elimination-of-cervical-cancer

[5] K. Canfell *et al.*, "Mortality impact of achieving WHO cervical cancer elimination targets: a comparative modelling analysis in 78 low-income and lower-middle-income countries.," *Lancet*, vol. 395, no. 10224, pp. 591–603, Feb. 2020, doi: 10.1016/S0140-6736(20)30157-4.

[6] World Health Organization (WHO), "WHO guideline for screening and treatment of cervical pre-cancer lesions for cervical cancer prevention," Geneva, 2021. [Online]. Available: https://pubmed.ncbi.nlm.nih.gov/34314129/

[7] C. Mungo, S. Ibrahim, E. A. Bukusi, H. H. M. Truong, C. R. Cohen, and M. Huchko, "Scaling up cervical cancer prevention in Western Kenya: Treatment access following a community-based HPV testing approach," *International Journal of Gynecology & Obstetrics*, vol. 152, no. 1, pp. 60–67, Jan. 2021, doi: 10.1002/IJGO.13171.

[8] P. E. Castle, D. Murokora, C. Perez, M. Alvarez, S. C. Quek, and C. Campbell, "Treatment of cervical intraepithelial lesions," *International Journal of Gynecology & Obstetrics*, vol. 138, pp. 20–25, 2017, doi: 10.1002/ijgo.12191.

[9] E. Rohner *et al.*, "Mapping the cervical cancer screening cascade among women living with HIV in Johannesburg, South Africaa," *Int J Gynaecol Obstet*, vol. 152, no. 1, pp. 53–59, Jan. 2021, doi: 10.1002/IJGO.13485.

[10] H. A. Cubie and C. Campbell, "Cervical cancer screening - The challenges of complete pathways of care in low-income countries: Focus on Malawi," *Womens Health (Lond)*, vol. 16, 2020, doi: 10.1177/1745506520914804.

[11] K. Khozaim *et al.*, "Successes and challenges of establishing a cervical cancer screening and treatment program in western Kenya," *Int J Gynaecol Obstet*, vol. 124, no. 1, pp. 12–18, 2014, doi: 10.1016/J.IJGO.2013.06.035.

[12] V. Mwenda *et al.*, "Cervical cancer programme, Kenya, 2011-2020: lessons to guide
elimination as a public health problem," *Ecancermedicalscience*, vol. 16, 2022, doi:
10.3332/ECANCER.2022.1442.

[13] K. P. Msyamboza, T. Phiri, W. Sichali, W. Kwenda, and F. Kachale, "Cervical cancer
screening uptake and challenges in Malawi from 2011 to 2015: Retrospective cohort study,"
BMC Public Health, vol. 16, no. 1, pp. 1–6, 2016, doi: 10.1186/s12889-016-3530-y.

[14] K. Austad *et al.*, "Barriers to Cervical Cancer Screening and the Cervical Cancer Care
Continuum in Rural Guatemala: A Mixed-Method Analysis," *J Glob Oncol*, vol. 4, no. 4, pp.
1–10, Mar. 2018, doi: 10.1200/JGO.17.00228.

[15] D. B. Vale, J. C. Teixeira, J. F. Bragança, S. Derchain, L. O. Sarian, and L. C. Zeferino,
"Elimination of cervical cancer in low- and middle-income countries: Inequality of access and
fragile healthcare systems," *Int J Gynaecol Obstet*, vol. 152, no. 1, pp. 7–11, Jan. 2021, doi:
10.1002/IJGO.13458.

[16] M. Maiman, D. H. Watts, J. Andersen, P. Clax, M. Merino, and M. A. Kendall, "Vaginal 5-
fluorouracil for high-grade cervical dysplasia in human immunodeficiency virus infection: A
randomized trial," *Obstetrics and Gynecology*, vol. 94, no. 6, pp. 954–961, 1999, doi:
10.1016/S0029-7844(99)00407-X.

[17] L. Rahangdale, Q. K. Lippmann, K. Garcia, D. Budwit, J. S. Smith, and L. Van Le, "Topical
5-flourourcil for treamtent of cervical intraepithelial Neoplasia 2☐: a Randomized Controlled
Trial," *The American Journal of Obstetrics & Gynecology*, vol. 210, no. 4, pp. 314.e1-314.e8,
2014, doi: 10.1016/j.ajog.2013.12.042.

[18] N. Desravines *et al.*, "Feasibility of 5-fluorouracil and imiquimod for the topical treatment of
cervical intraepithelial neoplasias (CIN) 2/3," *Int J Gynaecol Obstet*, vol. 163, no. 3, Dec.
2023, doi: 10.1002/IJGO.14983.

[19] C. L. Trimble *et al.*, "A first-in-human proof-of-concept trial of intravaginal artesunate to treat
cervical intraepithelial neoplasia 2/3 (CIN2/3).," *Gynecol Oncol*, vol. 157, no. 1, pp. 188–194,
Apr. 2020, doi: 10.1016/j.ygyno.2019.12.035.

[20] B. O. Fonseca *et al.*, "Topical Imiquimod for the Treatment of High-Grade Squamous
Intraepithelial Lesions of the Cervix: A Randomized Controlled Trial," *Obstetrics and
Gynecology*, vol. 137, no. 6, pp. 1043–1053, Jun. 2021, doi:
10.1097/AOG.0000000000004384.

[21] C. T. Lin *et al.*, "Topical imiquimod treatment for human papillomavirus infection in patients
with and without cervical/vaginal intraepithelial neoplasia," *Taiwan J Obstet Gynecol*, vol. 51,
no. 4, pp. 533–538, 2012, doi: 10.1016/j.tjog.2012.09.006.

[22] C. J. De Witte, A. J. M. Van De Sande, H. J. Van Beekhuizen, M. M. Koeneman, A. J. Kruse,
and C. G. Gerestein, "Imiquimod in cervical, vaginal and vulvar intraepithelial neoplasia: a
review," *Gynecol Oncol*, vol. 139, no. 2, pp. 377–384, 2015, doi:
10.1016/J.YGYNO.2015.08.018.

[23] C. Mungo *et al.*, "'There is nothing that can prevent me from supporting her:' Men's
perspectives on their involvement and support of women's use of topical therapy for cervical
precancer treatment in Kenya," *medRxiv*, Dec. 2023, doi: 10.1101/2023.12.22.23300455.

[24] C. Grimm *et al.*, "Treatment of cervical intraepithelial neoplasia with topical imiquimod: A
randomized controlled trial," *Obstetrics and Gynecology*, vol. 120, no. 1, pp. 152–159, 2012,
doi: 10.1097/AOG.0b013e31825bc6e8.

[25] World Health Organization (WHO), "WHO Model List of Essential Medicines - 23rd List,
2023," Geneva, 2023. [Online]. Available: https://www.who.int/publications/i/item/WHO-
MHP-HPS-EML-2023.02

[26] "Female Focus Group Interview Guide: Topical precancer treatments study." Mar. 01, 2023.

[27] J. V. Appleton and L. King, "Journeying from the philosophical contemplation of
constructivism to the methodological pragmatics of health services research," *J Adv Nurs*, vol.
40, no. 6, pp. 641–648, Dec. 2002, doi: 10.1046/J.1365-2648.2002.02424.X.

[28] T. Metaxas *et al.*, "Acceptability and safety of thermal ablation to prevent cervical cancer in
sub-Saharan Africa," *BMC Cancer*, vol. 22, no. 1, Dec. 2022, doi: 10.1186/S12885-022-
09202-2.

[29] C. Mungo *et al.*, "Safety and Acceptability of Thermal Ablation for Treatment of Human
Papillomavirus Among Women Living With HIV in Western Kenya," *JCO Glob Oncol*, vol.
6, no. 6, pp. 1024–1033, Nov. 2020, doi: 10.1200/GO.20.00035.

[30] J. Chapola *et al.*, "Barriers to follow-up after an abnormal cervical cancer screening result and
the role of male partners: a qualitative study.," *BMJ Open*, vol. 11, no. 9, p. e049901, Sep.
2021, doi: 10.1136/bmjopen-2021-049901.

[31] "Knowledge and beliefs about cervical cancer screening among men in Kumasi, Ghana -
PubMed." Accessed: Mar. 02, 2024. [Online]. Available:
https://pubmed.ncbi.nlm.nih.gov/23661828/

[32] A. D. Mwaka, H. R. Wabinga, and H. Mayanja-Kizza, "Mind the gaps: a qualitative study of
perceptions of healthcare professionals on challenges and proposed remedies for cervical
cancer help-seeking in post conflict northern Uganda," *BMC Fam Pract*, vol. 14, Dec. 2013,
doi: 10.1186/1471-2296-14-193.

[33] C. W. Ngugi, H. Boga, A. W. T. Muigai, P. Wanzala, and J. N. Mbithi, "Factors affecting
uptake of cervical cancer early detection measures among women in Thika, Kenya," *Health
Care Women Int*, vol. 33, no. 7, pp. 595–613, Jul. 2012, doi: 10.1080/07399332.2011.646367.

[34] N. I. Ebu and J. K. Ogah, "Predictors of cervical cancer screening intention of HIV-positive
women in the central region of Ghana," *BMC Womens Health*, vol. 18, no. 1, Feb. 2018, doi:
10.1186/S12905-018-0534-Z.

[35]   R. Ndejjo, T. Mukama, J. Kiguli, and D. Musoke, "Knowledge, facilitators and barriers to
cervical cancer screening among women in Uganda: a qualitative study," *BMJ Open*, vol. 7,
no. 6, Jun. 2017, doi: 10.1136/BMJOPEN-2017-016282.

[36]   F. I. Modibbo *et al.*, "Qualitative study of barriers to cervical cancer screening among
Nigerian women," *BMJ Open*, vol. 6, no. 1, 2016, doi: 10.1136/BMJOPEN-2015-008533.

[37]   "WHO guideline on self-care interventions for health and well-being, 2022 revision -
PubMed." Accessed: Mar. 04, 2024. [Online]. Available:
https://pubmed.ncbi.nlm.nih.gov/35914064/

[38]   A. R. Aluisio *et al.*, "HIV self-testing acceptability among injured persons seeking emergency
care in Nairobi, Kenya," *Glob Health Action*, vol. 16, no. 1, 2023, doi:
10.1080/16549716.2022.2157540.

[39]   M. A. Abdulai *et al.*, "An assessment of the likely acceptability of vaginal microbicides for
HIV prevention among women in rural Ghana," *BMC Womens Health*, vol. 12, Nov. 2012,
doi: 10.1186/1472-6874-12-40.

[40]   A. E. Tanner, "Perceptions of acceptability and utility of microbicides in Ghana, West Africa:
A qualitative, exploratory study," *Sahara J*, vol. 5, no. 1, pp. 11–18, Apr. 2008, doi:
10.1080/17290376.2008.9724897.

[41]   P. A. Newman, S. Roungprakhon, and S. Tepjan, "A social ecology of rectal microbicide
acceptability among young men who have sex with men and transgender women in Thailand,"
J Int AIDS Soc, vol. 16, no. 1, Aug. 2013, doi: 10.7448/IAS.16.1.18476.