Dietary Factors Affect Brain Iron Accumulation and Parkinson's Disease Risk

Jonathan Ahern BS[1,2,3], Mary ET Boyle PhD[1], Leo Sugrue MD PhD[4,5], Ole Andreassen MD PhD[6], Anders Dale PhD[1,3,7,8], Wesley K. Thompson PhD[2], Chun Chieh Fan MD PhD[2,8], Robert Loughnan PhD[1,2,3,9]

1. Department of Cognitive Science, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA
2. Center for Population Neuroscience and Genetics, Laureate Institute for Brain Research, Tulsa, OK 74103, USA
3. J. Craig Venter Institute, 4120 Capricorn Ln, La Jolla, CA 92037 USA
4. Department of Radiology and Biomedical Imaging, University of California, 505 Parnassus Ave, San Francisco, CA 94143, USA
5. Department of Psychiatry, University of California, 675 18th St, San Francisco, CA 94107, USA
6. NORMENT Centre, Division of Mental Health and Addiction, Oslo University Hospital and Institute of Clinical Medicine, University of Oslo, Oslo, Norway.
7. Department of Neuroscience, University of California, San Diego School of Medicine, 9500 Gilman Drive, La Jolla, CA 92037, USA.
8. Department of Radiology, University of California, San Diego School of Medicine, 9500 Gilman Drive, La Jolla, CA 92037, USA
9. Center for Human Development, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92161, USA

Abstract Word Count: 430 words
Word Count: 4228 words

Corresponding Author: Mr. Jonathan Ahern, jahern@ucsd.edu, 1-916-847-9887

Key Points

Question: How do dietary factors impact brain iron accumulation and risk for Parkinson's Disease (PD)?

Findings: In this cohort study, we identify several dietary factors that are associated with detectable differences in brain iron accumulation and PD risk. Dietary factors related to carbohydrates are associated with decreased iron accumulation and increased PD risk, while factors related to alcohol are associated with increased iron accumulation and decreased PD risk. We find evidence that dietary preferences associated with decreased brain iron accumulation are associated with increased PD risk.

Meaning: Modifiable dietary factors may impact PD risk through changes in brain iron accumulation.

Abstract

Importance: Iron dysregulation in the brain has been associated with neurodegenerative diseases like Parkinson's Disease (PD). Iron is exclusively absorbed from food; therefore, dietary patterns may impact risk.

Objective: To understand how modifiable dietary factors are associated with brain iron accumulation and PD risk.

Design: This retrospective cohort study included participants from the UK Biobank, a population-based sample. Genotypes, MRI scans, demographics, diagnoses, and diet and lifestyle measures were collected from January 2006 to May 2021. Data analysis was conducted from September 2023 to June 2024.

Setting: population-based

Participants: Disorders analyzed included Parkinson's and Parkinsonian Disorders (PD) (International Statistical Classification of Diseases and Related Health Problems, Tenth Revision [ICD-10], codes F02.3, G20-G21), and Vascular Dementia for secondary analysis (ICD-10 codes F01).

Exposures: Dietary and lifestyle preferences based on survey data

Main Outcomes: Brain iron accumulation score (derived from T2w-MRI, an iron-sensitive imaging technique) and PD diagnosis from electronic health records

Results: This study included 228,653 UK Biobank participants (30,675 with neuroimaging; 123,303 female; ages 40-83 years, largely of northern European ancestry). Factor analysis of dietary preferences revealed 20 factors that cumulatively explained 49.77% of variance across individual preferences, with prominent factors loading on alcohol (1.88% variance explained) and high-sugar (8.20% variance explained) questions. We found strong positive associations between measures of brain iron accumulation and the alcohol factor (t=4.02, p_{FDR}=0.0003) and significant negative associations between brain iron accumulation and the high-sugar factor (t=-3.73, p_{FDR}=0.0007). These factors were found to have opposite effects on PD risk, with the alcohol factor being associated with decreased PD risk (t=-5.83, $p_{FDR}<1\times10^{-8}$) and the high-sugar

factor being associated with increased PD risk (t=6.03, $p_{FDR} < 1 \times 10^{-8}$). Instrumental variable regression of preferences revealed that dietary preferences associated with higher brain iron levels were also associated with lower risk for PD (θ=-0.247, p=0.004). Voxelwise imaging analysis of alcohol and high-sugar factors, using T2w-MRI, confirmed patterns consistent with increased and decreased iron accumulation in motor regions, respectively. Using T2w-MRI and dw-MRI, we confirmed that this signal was more likely driven by iron than vascular confounds.

Conclusions and Relevance: Contrary to some prior studies' findings of higher central iron levels linked with PD risk, we find lower central iron levels to be associated with greater PD risk. Dietary components, especially those related to alcohol and carbohydrates, are related to detectable differences in brain iron accumulation in motor regions of the brain and differences in risk of PD, suggesting the potential for lifestyle interventions to influence PD risk.

Graphical Abstract

*Previous research has established that iron overload in the brain is associated with increased risk of movement disorders[1–4].

Introduction

Parkinson's disease (PD) is the fastest-growing neurodegenerative disease in the world[5] and is characterized by progressive difficulties with movement[6] and a significant reduction in quality of life[7]. Increased iron accumulation in regions like the substantia nigra and the basal ganglia and dyshomeostasis of iron metabolism are common findings in PD patients[8,9]. While iron is an essential nutrient, iron overload can result in oxidative stress and ferroptotic mechanisms implicated in PD[10,11].

Research has indicated that low peripheral iron and hemoglobin are associated with greater PD risk[11–14]. Conversely, hemochromatosis, a genetic disorder characterized by systemic iron overload[15], has also been linked to an increased PD risk[1,3]. These previous studies suggest a U-shaped relationship between iron and PD risk, with both low and high levels being linked to increased risk. A recent study demonstrated this relationship using MRI in a large cohort[3], showing that lower central iron levels were linked to a higher risk of PD in individuals without hemochromatosis. Conversely, higher central iron levels were associated with increased PD risk in individuals carrying hemochromatosis risk alleles. Under this U-shaped model, a normative cohort—free of major iron-disrupting genotypes—lies on the descending arm, exhibiting a linear negative central iron–PD risk association ("Study Population Distribution" of graphical abstract).

PD is caused by an interaction between genetic predisposition and environmental factors that accumulate throughout life[6], with diet representing an important modifiable factor. A high-quality diet and avoidance of undernutrition are associated with a lower risk of PD[16] and improved PD outcomes[17,18]. Dietary patterns, like the Mediterranean diet[19], have been shown to improve outcomes related to PD[20,21]. Conversely, extended consumption of ultraprocessed foods is related to increased odds of nonmotor features of PD[22].

Given these findings, we aim to investigate the following hypothesis: whether diet impacts PD risk through changes in brain iron. To achieve this, we build upon work identifying a brain iron-specific biomarker of PD referred to as the *Hemochromatosis Brain PolyVoxel Score* (HB-PVS). This marker quantifies the central iron dysregulation associated with hemochromatosis, as captured by T2-weighted MRI intensities (T2w-MRI), in critical motor regions of the brain[3,4]. This score is differentially associated with a higher risk of PD in individuals with and without hemochromatosis, where individuals with hemochromatosis and very high HB-PVS show increased PD risk, whereas in individuals without hemochromatosis, a lower HB-PVS compared to other healthy subjects is associated with increased PD risk[3].

We aim to investigate how dietary factors are related to differences in this iron-specific biomarker and how this is jointly related to PD risk.

Methods

UK Biobank Measures

Genotypes, MRI scans, demographics, diagnoses, and dietary measures were obtained from the UK Biobank (URLs) under accession number 27412, excluding 206 participants who withdrew their consent. All participants provided electronically signed informed consent. The study was approved by the UK Biobank Ethics and Governance Council. The recruitment period for participants was from 2006 to 2010, and participants had to be 40-69 years old during this period to be included in the sample.

Neuroimaging Data

Image Processing

Image acquisition, pre-processing, and registration are described in the supplementary methods. The average of the b = 0 images from registered diffusion-weighted MRI intensities (dw-MRI) was used as voxelwise measures of T2w-MRI. T2w-MRI refers to the average of b=0 images, and dw-MRI refers to the full diffusion acquisition (all b directions and values). Using a restriction spectrum imaging (RSI)[23] model, we separated dw-MRI signal into three compartments: intracellular, extracellular, and unhindered free water. Briefly, this model estimates fractions of diffusion attributable to these compartments using a mixture of spherical harmonic basis functions[24]. Although prior work indicated that the HB-PVS derived from T2w-MRI is a biomarker strongly linked with iron[3,4], T2w-MRI signal can also be sensitive to other cellular changes, such as gliosis and edema[25], pathologic processes that may reflect ischemic injury or inflammation[26]. We used the free water RSI component in sensitivity analysis (see "Brain Association Maps…" section) to distinguish iron-related from non-iron, vascular-related signals for discovered associations.

The Hemochromatosis Brain PolyVoxel Score

In this analysis, the primary imaging measure is the PolyVoxel Score (PVS), derived from a classifier trained on UK Biobank T2w-MRI brain scans, which are sensitive to iron[27], to differentiate between individuals with and without C282Y homozygosity. C282Y homozygosity accounts for 95% of hemochromatosis cases[28], a disorder that can lead to iron overload[15]. As detailed by Loughnan et al. (2024)[3], the PVS quantifies iron accumulation across motor-related brain regions, including the cerebellum, thalamus, caudate, putamen, pallidum, red nucleus, substantia nigra, and subthalamic nucleus.

Preference Data

Preferences data (Category 1039) were gathered from a web-based questionnaire developed and designed to reflect preferences for many foods and activities. The questionnaire

included 150 items measured on a nine-point scale from 'extreme dislike' to 'extreme like' designed to cover a range of sensory aspects as well as variable food groups[29]. After noninformative values ('Never tried' or 'Prefer not to answer') were removed, 6 columns were removed for having greater than 10% missingness across the population (preferences for bell peppers, capers, globe artichoke, going to the gym, pollock, and soy milk), and 3,459 participants were removed for having individual missingness of greater than 10%, any remaining missing values were imputed from the median value for that variable, and data was z-score normalized across all remaining subjects.

Nutrient Data

As a secondary and confirmatory analysis to our preference analyses, we analyzed the effects of estimated nutrient intake (Category 100117; estimated from a separate questionnaire to preference data) on HB-PVS and PD risk. Details of the questionnaire and data cleaning can be found in the supplementary methods.

Disease Diagnosis

Neurological diagnoses were extracted from ICD-10 diagnosis data from electronic health records (Data-Field 41270). PD diagnosis included the following diagnosis codes: F02.3, G20, G21.0, G21.1, G21.2, G21.3, G21.4, G21.8, and G21.9. Vascular Dementia (VD) diagnoses, which were used as a positive control in our imaging analysis, included the following diagnosis codes: F01.0, F01.1, F01.2, F01.3, F01.8, and F01.9. Individuals with any one or more of these diagnoses were categorized as having been diagnosed with PD (Supplementary Table 2) or VD, respectively.

Statistical Analyses

Analyses were performed using generalized linear regressions in Python using statsmodels (version 0.14.0). All analyses were controlled for sex (Data-Field 31), the maximum reported pre-tax income category (Data-Field 738), and the top 10 genetic principal components (Data-Field 22009) to account for global genetic factors. Analyses that involve the HB-PVS additionally controlled for MRI scanner study site (Data-Field 54) to account for differences in imaging equipment. All analyses controlled for age; imaging analyses controlled for the age at the time of the scan (retrieved from Data-Field 20216), disease analyses of preferences and nutrition were corrected for age at time of survey completion (Data-Field 20750 and 105010). P-values within each analysis were corrected using the multitest correction from statsmodels using Benjamini/Hochberg FDR and Bonferroni correction methods.

Lower dimensional factors of preference data were generated in Python using Educational Testing Service's Factor_Analyzer package (version 0.5.0) with default parameters, which calculates factors by minimizing residuals[30,31]. Factors were varimax rotated to increase interpretability[32]. To calculate robust standard errors, we performed 10,000 bootstrap iterations of all our regression analyses[33].

As an exploratory analysis, we sought to test whether the effect of dietary preferences on PD risk was mediated by brain iron concentration differences. To achieve this, we performed two-sample instrumental variable mediation analysis[34,35]. In this framework, individual preferences are used as instruments to test the role of mediating traits on eventual outcomes. In our analysis, we use preference questionnaire data as instruments/exposures, PD risk as outcome, and brain iron concentration as the mediator to test the following regression:

$$\beta_i^{PD} = \alpha + \theta\beta_i^{HB-PVS} + \varepsilon_i$$

Where β_i^{HB-PVS} and β_i^{PD} are the coefficients associating the i-th food preference question with HB-PVS and PD, respectively, calculated from previous analyses. α is the intercept, ε_i is the error, and θ is the mediation effect. In supplementary results, we repeat this analysis using the nutrient questionnaire data as exposures. For both analyses, we interpret the results as suggestive of dietary factors impacting PD risk through brain iron levels, but refrain from making strong claims about causality.

Our analysis is principally focused on dietary factors, so we removed eight preference questions related to non-dietary factors (preferences for taking the stairs, working up a sweat, exercising alone, bicycling, going to a cafe, watching television, cigarette smoking, exercising with others, and going to the pub) — some of which (exercise in particular) we expected to be susceptible to reverse causality with PD risk. An analysis including non-dietary questions is included in Supplementary Figure 8.

Statistical analysis for our confirmatory nutrient analysis is largely consistent with the methods described above. Further details of statistical considerations related to our nutrient analysis can be found in the supplementary methods.

Voxelwise Associations of Dietary Factors

We generated voxelwise association maps between T2w-MRI intensities, known to be sensitive to iron[27], and dietary preference factors 2 (sweets) and 7 (alcohol). To account for the potential influence of microvascular pathologies, which we expect to be preferentially related to differences in the free water fraction of RSI, we generated association maps between normalized free water diffusion from RSI models and these dietary factors.

Association maps were generated by voxelwise univariate regression using all 28,388 individuals from our preference imaging subsample (Table 1), predicting T2w-MRI and free water signal from dietary factors 2 (sweets) and 7 (alcohol), separately. Covariates of age at scan, sex, scanner, and the top 10 principal components of genetic ancestry were corrected by pre-residualizing images before model fitting. We plotted voxelwise beta effects from these models after passing a smoothing kernel with a full width at half maximum of 2.82 voxels. Beta values were masked to those used to compute the HB-PVS.

Results

Demographics on the cohorts used in imaging and non-imaging analyses can be found in Table 1.

Table 1: Sample Demographics

	Imaging		Non Imaging		
Preferences	**Sample Size:**	28,388	**Sample Size:**		132,023
	Mean Age (SD)	64.13 (7.45)	**Mean Age (SD)**		66.65 (7.70)
	# Female (%)	14,871 (52.38)	**# Female (%)**		74,360 (56.32)
	# PD Cases (%)	45 (0.16)	**# PD Cases (%)**		456 (0.35)
	Category	**%**		**Category**	**%**
Max Income	<£18,000	6.47	Max Income	<£18,000	13.29
	£18,000-£30,999	18.86		£18,000-£30,999	23.29
	£31,000-£51,999	30.22		£31,000-£51,999	28.97
	£52,000-£100,000	33.40		£52,000-£100,000	26.23
	>£100,000	11.05		>£100,000	8.22
Nutrients	**Sample Size:**	20,477	**Sample Size:**		150,603
	Mean Age (SD)	64.06 (7.56)	**Mean Age (SD)**		58.63 (8.05)
	# Female (%)	10,545 (51.50)	**# Female (%)**		80,595 (53.51)
	# PD Cases (%)	37 (0.18)	**# PD Cases (%)**		979 (0.65)
	Category	**%**		**Category**	**%**
Max Income	<£18,000	5.97	Max Income	<£18,000	15.88
	£18,000-£30,999	18.83		£18,000-£30,999	24.79
	£31,000-£51,999	30.52		£31,000-£51,999	28.37
	£52,000-£100,000	33.31		£52,000-£100,000	23.81
	>£100,000	11.38		>£100,000	7.15

Table 1: Demographic Information. Test samples were separated into four groups based on the availability of imaging and survey data related to nutrients and preferences. We use distinct, non-overlapping imaging and non-imaging cohorts to facilitate downstream instrumental variables analyses (Depicted in Figure 1A and Supplementary Figure 4C). Non-overlapping samples provide independent estimates, which reduce bias due to shared influences. We also note, as others have, that the neuroimaging cohort is healthier than the non-imaging sample, specifically with respect to PD prevalence. This is likely to reduce our statistical power to detect effects in the neuroimaging cohort relative to a more representative sample with higher PD prevalence, thereby providing a conservative estimate of significance. Across all groups, there are a total of 228,653 unique individuals (30,675 with neuroimaging).

HB-PVS Instrument

We use the HB-PVS as an instrument to indicate regional brain iron levels. Previous work provides details of HB-PVS training[3], where contributions to the HB-PVS depend on classifier weightings and brain region size (see Supplementary Figure 1). These weights indicate that, while greater HB-PVS scores indicate higher iron concentrations for most included brain regions, some brain regions that contribute less (e.g. caudate, substantia nigra) may have lower iron concentrations for greater HB-PVS values. Overall, previous work demonstrates that

the HB-PVS correlates strongly with genes associated with iron metabolism, peripheral iron levels, and PD[3,4].

Preference Analyses

Preference Analysis

To understand how dietary preferences may be related to brain iron accumulation and PD risk, we fit a series of linear and logistic regression models associating each preference with the HB-PVS and PD risk. Here we describe general trends among our preference findings. Full preference results can be found in the Supplementary Results (Supplementary Figures 2-3, Supplementary Tables 5-6).

Increased HB-PVS was associated with preferences for alcoholic beverages (red wine: t=6.72, $p_{FDR}<1\times10^{-8}$; white wine: t=5.48, $p_{FDR}<1\times10^{-5}$; etc.) and red meats (red meat: t=4.46, $p_{FDR}<1\times10^{-3}$; lamb: t=4.30, $p_{FDR}<1\times10^{-3}$; etc.). Decreased HB-PVS was associated with preferences for cereal grains (corn flakes: t=-4.61, $p_{FDR}<1\times10^{-5}$; porridge t=-3.74, $p_{FDR}=0.004$; etc.), sweet foods (biscuits: t=-3.64, $p_{FDR}=0.004$; cake: t=-3.57, $p_{FDR}=0.004$; etc.), and exercise (bicycling: t=-3.19, $p_{FDR}=0.010$; exercising alone: t=-3.03, $p_{FDR}=0.015$; etc.).

Decreased PD risk was associated with exercise preferences (taking the stairs: t=-10.87, $p_{FDR}<1\times10^{-24}$; working up a sweat: t=-7.67, $p_{FDR}<1\times10^{-10}$; etc.), vegetables (salad leaves: t=-8.27, $p_{FDR}<1\times10^{-13}$; vegetables: t=-6.24, $p_{FDR}<1\times10^{-7}$; etc.), and alcoholic beverages (spirits: t=-6.15, $p_{FDR}<1\times10^{-7}$ and red wine: t=-5.80, $p_{FDR}<1\times10^{-6}$). Increased PD risk was associated with preferences for sweet foods (sweet coffee house drinks: t=5.15, $p_{FDR}<1\times10^{-5}$; cake icing: t=4.90, $p_{FDR}<1\times10^{-4}$; etc.).

Dietary Effects on Brain Iron Predict Parkinson's Risk

We hypothesized that the effect of specific foods on PD risk was mediated through their impact on brain iron concentration. To explore this, we performed mediation analysis, using two-sample instrumental variable estimation. Here, preferences were regarded as exposures, PD as the outcome, and brain iron accumulation (HB-PVS) as the mediator. When restricting to dietary preferences, coefficients from HB-PVS and PD display a significant negative association (Figure 1A: θ=-0.247, p=0.004) - indicating that preferences associated with higher brain-iron accumulation tended to confer lower PD risk, and vice versa. This negative β-β relationship is consistent with a model in which dietary modulation of central iron contributes to Parkinson's susceptibility. We emphasize, however, that these results are exploratory: while the strength and direction of the association support an iron-mediated pathway (Diet → Brain Iron → PD), residual direct effects of diet on PD risk and potential confounders cannot be fully excluded without further validation. This trend does not persist when including non-dietary preferences in our analysis for reasons we believe to be related to reverse causation (See Supplementary Results and Supplementary Figure 8 for further discussion).

Nutrient Analysis

Analysis of estimates of nutrient intake supported several of our findings from our preference analysis. We found that estimated intakes related to sugars (maltose: t=-3.74, p_{FDR}=0.012; lactose: t=-3.36, p_{FDR}=0.013; etc.) were associated with decreased measures of the HB-PVS, supporting preference findings. We also found estimates related to sugars (carbohydrates: t=3.46, p_{FDR}=0.012; total sugar: t=3.45, p_{FDR}=0.012; etc.) were associated with increased PD risk, and estimates related to alcohol were associated with decreased PD risk, again in line with preference findings. Instrumental variable analysis also showed a general downward trend (θ=-0.421, p=0.0006) where measures associated with decreased HB-PVS were associated with increased PD risk. The full results of our nutrient analysis can be found in our supplementary results, Supplementary Figure 4, and Supplementary Tables 3-4.

Factor Analysis of Preferences

Due to the high correlation structure in dietary preferences (Supplementary Figure 5), we performed factor analysis to identify interpretable latent factors. Twenty factors accounted for approximately half (49.77%) of the total variance in the preference space (Supplementary Figure 6). A summary of each Factor's top loadings as well as an interpretive description of each factor can be found in Table 2. For full loadings, see Supplementary Figure 7 and Supplementary Table 7.

Table 2: Factor Loadings

Factor	Top Preferences (Loading)	Interpretation	Variance Explained (%)	Cumulative Variance Explained (%)
Factor 1	red meat (0.844), beef steak (0.790), barbequed or grilled meat (0.765)	Red Meat	11.77	11.77
Factor 2	cake (0.794), sweet foods (0.756), biscuits (0.709)	Sweets	8.20	19.97
Factor 3	fruit (0.717), plums (0.672), oranges (0.577)	Fruit	4.43	24.40
Factor 4	haddock (0.743), baked steamed fish (0.735), mackerel (0.721)	Fish	4.11	28.51
Factor 5	spicy foods (0.886), chili pepper (0.790), burn of spicy foods (0.746)	Spicy Foods	2.30	30.81
Factor 6	cabbage (0.711), cauliflower (0.685), Brussel sprouts (0.641)	Brassica Vegetables	2.00	32.81
Factor 7	spirits (0.648), whisky (0.604), red wine (0.578)	Alcohol	1.88	34.68
Factor 8	avocados (0.509), aubergine (0.465), asparagus (0.443)	Savory Fruits & Vegetables	1.74	36.42
Factor 9	hard cheese (0.596), soft cheese (0.589), blue cheese (0.468)	Cheese	1.54	37.96
Factor 10	tea with sugar (-0.732), coffee with sugar (-0.731), coffee without sugar (0.541)	Unsweetened Tea and Coffee	1.42	39.38
Factor 11	working up a sweat (0.753), exercising alone (0.529), bicycling (0.497)	Exercise	1.30	40.68
Factor 12	salty foods (0.701), adding salt to foods (0.526), salty pretzels (0.463)	Salty Foods	1.24	41.93
Factor 13	green olives (0.767), black olives (0.753), gherkins (0.340)	Olives	1.23	43.15
Factor 14	whole grain breakfast cereal (0.481), brown rice (0.351), porridge (0.350)	Cereal Grains	1.14	44.30
Factor 15	potatoes (0.377), chips/French fries (0.370), pasta (0.366)	Potatoes and Carbs	1.04	45.34
Factor 16	salad dressing (0.504), mayonnaise (0.419), vinegar (0.370)	Condiments	0.99	46.33
Factor 17	whole milk (0.496), cream (0.416), dairy products (0.384)	Dairy	0.93	47.27
Factor 18	chicken (-0.337), roast chicken (-0.319), herring (0.294)	Fish and Not Chicken	0.87	48.14
Factor 19	orange juice (0.501), apple juice (0.399), grapefruit (0.253)	Juices	0.85	48.99
Factor 20	diet fizzy drinks (0.356), corn flakes (0.222), regular nondiet fizzy drinks (0.204)	Fizzy Drinks	0.78	49.77

Table 2: Summary Table of the lower-dimensional factor space generated from our 144 preference variables. Visual Factor loadings can be found in Supplementary Figure 7. A full numeric summary of all factor loadings can be found in Supplementary Table 7.

Preference Factors and Brain Iron

To formalize and confirm the results identified in the preference-level analysis, we performed linear regressions in our imaging sample (N=28,388) predicting the HB-PVS from the first 20 preference factors. All of the results from the preference-level analysis were recapitulated, namely the positive association between alcohol (factor 7: t=4.02, p_{FDR}<0.001) and red meat (factor 1: t=4.73, p_{FDR}<1e-4) and the negative associations for cereal grains (factor 14: t=-2.54, p_{FDR}=0.012), sweets (factor 2: t=-3.73, p_{FDR}<0.001), and exercise (factor 11: t=-4.31, p_{FDR}<0.001).

We found additional positive associations between the HB-PVS and savory fruit and vegetables (factor 8: t=4.86, p_{FDR}<1e-4) and salty foods (factor 12: t=3.36, p_{FDR}=0.001) and negative associations between the HB-PVS and fizzy drinks (factor 20: t=-4.00, p_{FDR}<0.001), brassica vegetables (factor 6: t=-2.74, p_{FDR}=0.008), and preference for fish and a dislike for chicken (factor 18: t=-2.36, p_{FDR}=0.018) (Figure 1B, Supplementary Table 8).

Preference Factors and Parkinson's Disease Risk

Using the same top twenty preference factors, we fit models to predict PD risk in our non-imaging sample (N_{Case}=456, $N_{Control}$=131,567). We were able to recapitulate the results from the preference-level analysis, namely increased PD risk was associated with a lower preference for exercising (factor 11: t=-7.66, p_{FDR}<1e-12), lower alcohol preference (factor 7: t=-5.82, p_{FDR}<1e-7), lower vegetable preference (factor 6: t=-6.11, p_{FDR}<1e-8), lower fruit preference (factor 3: t=-2.98, p_{FDR}=0.018), and higher sweet preference (factor 2: t=6.03, p_{FDR}<1e-8).

We also found that increased PD risk was associated with a lower preference for potato products and carbohydrates (factor 15: t=-4.03, p_{FDR}<0.001) and a higher preference for juices (factor 19: t=3.89, p_{FDR}<0.001), fizzy drinks (factor 20: t=3.23, p_{FDR}=0.002), and olives (factor 13: t=2.35, p_{FDR}=0.018) (Figure 1C, Supplementary Table 9).

Figure 1: Preference Analysis

Figure 1: (A) Scatter plot of the instrumental variable regression (i) between dietary preferences and the HB-PVS and (ii) preferences and the odds ratio of developing PD. Error bars are shown for factors that are significant along a given axis. Of those factors, the top thirteen dietary preference factors furthest from the origin are annotated. A regression line shows the correlation between the effects of a dietary preference on PD and the HB-PVS, respectively. A scatter plot of the dietary and non-dietary factors can be found in Supplementary Figure 8. (B) Forest plot showing the FDR associations between preference factors and the HB-PVS. Results below the dotted line are additionally Bonferroni Significant. A full numeric summary of the effects of all factors can be found in Supplementary Table 8. (C) Forest plot showing the FDR associations between preference factors and odds ratio of developing PD. Results below the dotted line are additionally Bonferroni Significant. A full numeric summary of the effects of all factors can be found in Supplementary Table 9.

Voxelwise Association Maps of Sweet and Alcohol Factors

We generated voxelwise association maps between T2w-MRI and dietary factors 2 (sweets) and 7 (alcohol), which were selected because they were significantly associated with both PD risk and the HB-PVS.

Consistent with earlier results of factor 2 (sweets) being associated with decreased HB-PVS, voxelwise analysis demonstrated that this factor was associated with T2w-MRI hyperintensities in movement-related regions, including basal ganglia and cerebellar regions, indicating decreased iron accumulation (Figure 2) with peak voxels in the putamen. Conversely, factor 7 (alcohol) was associated with T2w-MRI hypointensities in movement-related regions, including the putamen and cerebellum (Figure 2), largely consistent with earlier results. Interestingly, we also observe regions of hyperintensities, of modest effect size, in thalamic regions.

To inspect the degree to which our T2w-MRI associations were driven by vascular pathologies, we associated factors 2 (sweets) and 7 (alcohol) with voxelwise estimates of free water. Specifically, we were concerned that a high-sugar diet may be related to microvascular pathology[36,37]. As a positive control, we additionally associated voxelwise estimates of free water with a diagnosis of Vascular Dementia (VD)[38]. We posited that vascular disruptions would increase both T2w-MRI and free water signal, whereas iron differences would only affect the T2w-MRI signal.

Preference factors for sweets and alcohol were associated with basal ganglia and cerebellar intensities for T2w-MRI imaging but not free water estimates (compare the upper and lower panels of Figure 2). Conversely, for VD, we see higher signal intensities for both T2w-MRI and free water signals (right side of Figure 2). Previous evidence of the HB-PVS's strong coupling to iron regulation[3,4] along with these results, indicates that the alcohol and sweet imaging effects are more likely related to iron processes than being confounded by vascular disruptions. Incidental free water hyperintensities are cautiously interpreted in the supplementary results.

Figure 2: Dietary Factor Association Maps Brain

Figure 2: Voxelwise association map of T2w-MRI and free water signal intensity for (i) factor 2 (sweets), (ii) factor 7 (alcohol), and (iii) VD as a positive control. Voxels are masked to those that have non-zero weights for Hemochromatosis PVS (Supplementary Figure 1). Unmasked factor association maps can be found in Supplementary Figure 9. The hemochromatosis brain weights[3] (top, far left panels) are included for reference and indicate the relative weighting that intensities in different regions of the brain contribute to the calculation of the HB-PVS, with red regions representing positive weightings and blue regions representing negative weightings, expressed in arbitrary units (Supplementary Figure 1). In our T2w-MRI brain association maps, effect sizes are displayed. Orange regions represent increased T2w-MRI intensities, consistent with lower iron concentrations for individuals with a higher score along the spectrum of a given dietary factor, and blue regions represent decreased T2w-MRI intensities, consistent with higher iron concentrations. In the free water association maps, higher values (orange) indicate increased isotropic diffusion. The similar association patterns in both T2w-MRI and free water association maps in the VD analysis suggest that changes in free water are likely driving the observed changes in T2w-MRI signal intensity. However, the association maps of the dietary factors show distinct intensity patterns: T2w-MRI signal changes in the basal ganglia and cerebellum that are not associated with changes in free water, suggesting that our T2w-MRI signal changes related to dietary factors are more likely to be driven by iron accumulation differences than vascular differences.

Discussion

We have demonstrated that dietary preferences linked to lower brain iron are associated with increased PD risk, aligning with our previous work showing a negative association between regional brain iron and PD risk[3]. Although this may appear to contradict studies reporting a positive association between brain iron and PD risk[8–10]; we believe our findings offer a broader perspective rather than a conflicting one. Specifically, while a lower HB-PVS score - reflecting overall reduced brain iron accumulation - is associated with greater PD risk, this does not imply that iron is uniformly reduced across all brain regions included in the measure. During PVS training, regularization results in some regions contributing negatively and others contributing positively to the overall score - see Supplementary Figure 1 and previous work[3]. Given these HB-PVS weightings, our results are consistent with higher iron accumulation in regions such as the substantia nigra, pallidum, and caudate being associated with greater PD risk. Simultaneously, larger regions with greater contributions to the HB-PVS, such as the thalamus and cerebellum, show a pattern where lower iron levels are associated with greater PD risk. These regionally specific contributions help reconcile our findings with the broader literature and support a more nuanced interpretation of brain iron's role in PD.

The individual regions that make up the HB-PVS show results consistent with a misdistribution of iron, where in some regions, excessive accumulation is present, and in other larger regions, there is a relative dearth of iron. However, when these results are aggregated, the HB-PVS is linked with lower central iron levels in large areas of the brain and is associated with greater PD risk. The novelty in our approach may capture a more subtle understanding of central iron homeostasis that has been difficult to discover with traditional approaches, allowing us to distill a pathogenic, U-shaped misdistribution of iron into a central tendency that is still sensitive to dietary trends and movement disorders[3].

Sugars and Carbohydrates

We consistently found carbohydrate-related preferences (cake, ice cream, biscuits, etc.) were associated with decreases in the HB-PVS, a measure of brain iron, and an increase in PD risk. These findings were supported by our nutrient and factor-level analyses. Decreases in brain iron in regions associated with the hemochromatosis brain were also supported by T2w-MRI imaging analysis, where we found relative hyperintensity in movement-related regions of the brain, including the putamen and cerebellum, indicating reduced iron accumulation.

Experimental evidence in humans demonstrates a causal relationship between sugar on iron homeostasis, with glucose ingestion decreasing serum iron through pancreatic secretion of the iron-regulatory factor, hepcidin[39], consistent with our findings. Epidemiological research shows that glycemic dysregulation and related diseases are related to increased PD risk and worse PD outcomes[70–72], in line with our findings of preferences toward sugary foods being related to greater PD risk. Other research has shown that iron stores and regulatory molecules contribute to impaired glycemic regulation[43], possibly indicating a bidirectional coupling between iron and glucose homeostasis. Taken together, our findings indicate that consumption of sweet

and sugary foods contributes to a generally iron-depleted phenotype in the brain. This iron depletion may, then, contribute to increased PD risk. These results provide a mechanistic interpretation of high-sugar diets as a potential modifiable factor mediating PD risk.

Sugars and carbohydrates may also impact PD risk and iron accumulation at the microbiome level, which is associated with health[44,45], including PD[46,47]. Preferences for sweets and carbohydrates may elevate levels of pro-inflammatory, opportunistic pathogenic bacteria in the gut, which is linked to an increased risk of PD[48]. This dietary pattern also contributes to α-synuclein pathology in the enteric and central nervous system[49], where it contributes to ferroptosis, cell toxicity related to iron-dependent lipid peroxide accumulation, in the brain[50].

Alcohol

Preferences related to alcohol (red wine, spirits, etc.) were associated with greater brain iron, as evidenced by the HB-PVS, and decreased PD risk. These findings were supported by our nutrient and factor-level analyses. Alcohol-preference-related increases in brain iron in regions associated with the hemochromatosis brain were supported by T2w-MRI imaging, which found hypointensities in the putamen and cerebellum, consistent with higher iron accumulation in those areas associated with a greater preference for alcohol. Interestingly, we also found modest effects of hyperintense regions in the thalamus, indicating that alcohol may have differential effects on subregions of the basal ganglia.

There is extensive literature suggesting that moderate alcohol consumption is associated with decreased PD risk[51,52], with one meta-analysis suggesting a U-shaped risk profile[51].

Alcohol consumption is known to impact iron absorption and accumulation. Alcohol can lead to increased iron accumulation in the brain[53]. Alcohol can downregulate the synthesis of hepcidin, the primary iron-regulating hormone, and, with excessive consumption, can cause iron overload in otherwise typical individuals[54–57]. In individuals with iron overload disorders, alcohol consumption is broadly associated with worse health outcomes[58–60]. Conversely, in genetically typical individuals, the moderate consumption of alcohol (fewer than 2 drinks a day) is associated with decreased risk of iron deficiency and its associated anemia[57]. Anemia itself has been associated with increased PD risk in some[61–64] but not all[65] previous studies[66]. Measures of iron in the brain have also been suggested to have a U-shaped association with PD risk[3]. Our findings support a more nuanced view of the relationship between iron, alcohol, and PD, where alcohol consumption may alter iron-regulation, increasing iron absorption and decreasing anemia, which together may decrease PD risk.

See supplementary materials for a discussion of non-dietary preferences and results that were individually significant in only PVS or PD analyses.

Limitations

Our dietary measures are based on self-report data, which is susceptible to reporting bias. Nevertheless, the convergence of results from our preference, nutrient, and factor-level analyses gives us confidence that we are capturing consistent results.

The HB-PVS is a univariate measure that is specifically tuned to iron disruptions in specific brain regions that are related to motor control and predictive of PD[27] risk, including the substantia nigra, thalamus, putamen, and cerebellum[3]. Collapsing the signal across multiple brain regions into a single measure enables the analyses showcased here; however, there is an inherent loss of information when performing this dimensionality reduction. Furthermore, the PVS represents larger regions like the cerebellum with greater weight than smaller regions like the substantia nigra (Supplementary Figure 1)[3]. While the HB-PVS has been shown to be representative of iron accumulation in motor regions, it will not pick up changes in other brain regions.

The HB-PVS (a T2w-MRI derivative measure) is strongly related to iron[3], and T2w-MRI is an iron-sensitive measure. However, other factors can also affect T2w-MRI signals. We have made efforts to rule out confounding factors like gliosis[25] and edema[26,38]; however, other forms of tissue mineralization[67] or imaging features could impact these measures.

Due to the demographics of the UK, the majority of individuals within this study are of European descent, so it will be important to validate these findings in non-European individuals.

Conclusion

Dietary factors and preferences, especially those related to carbohydrates and alcohol, are associated with measurable differences in iron in motor regions of the brain and with risk for PD. In a typical population, dietary preferences and nutrients associated with decreased iron accumulation in the brain are associated with increased risk of PD. These results demonstrate that diet is associated with risk for PD and offer a potential mechanistic explanation, suggesting that changes in PD risk may be related to iron levels in critical motor regions of the brain.

UK Biobank Acknowledgment

This work uses data provided by patients and collected by the NHS as part of their care and support. UK Biobank's research resource is a major contributor in the advancement of modern medicine and treatment, enabling better understanding of the prevention, diagnosis, and treatment of a wide range of serious and life-threatening illnesses – including cancer, heart diseases, and stroke. UK Biobank is generously supported by its founding funders the Wellcome Trust and UK Medical Research Council, as well as the Department of Health, Scottish Government, the Northwest Regional Development Agency, British Heart Foundation, and Cancer Research UK. The organization has over 150 dedicated members of staff, based in multiple locations across the UK.

Data Sharing Statement: See Supplement 2.

URLs

UK Biobank: https://www.ukbiobank.ac.uk/

Statsmodels: https://www.statsmodels.org/stable/index.html
FactorAnalyser: https://factor-analyzer.readthedocs.io/en/latest/index.html

Funding

This work was supported by grants R01MH122688 and RF1MH120025, funded by the National Institute for Mental Health (NIMH), and the Lundbeck Foundation fellowship (R335-2019-2318).

Citations

1. Dekker, M. C. J. *et al.* Mutations in the hemochromatosis gene (HFE), Parkinson's disease and parkinsonism. *Neurosci. Lett.* **348**, 117–119 (2003).

2. Kruer, M. C. *et al.* Neuroimaging Features of Neurodegeneration with Brain Iron Accumulation. *Am. J. Neuroradiol.* **33**, 407–414 (2012).

3. Loughnan, R. *et al.* Hemochromatosis neural archetype reveals iron disruption in motor circuits. *Sci. Adv.* **10**, eadp4431 (2024).

4. Loughnan, R. *et al.* Association of Genetic Variant Linked to Hemochromatosis With Brain Magnetic Resonance Imaging Measures of Iron and Movement Disorders. *JAMA Neurol.* **79**, 919–928 (2022).

5. Feigin, V. L. *et al.* Global, regional, and national burden of neurological disorders during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Neurol.* **16**, 877–897 (2017).

6. Bloem, B. R., Okun, M. S. & Klein, C. Parkinson's disease. *The Lancet* **397**, 2284–2303 (2021).

7. Zhao, N. *et al.* Quality of life in Parkinson's disease: A systematic review and meta-analysis of comparative studies. *CNS Neurosci. Ther.* **27**, 270–279 (2021).

8. Belaidi, A. A. & Bush, A. I. Iron neurochemistry in Alzheimer's disease and Parkinson's disease: targets for therapeutics. *J. Neurochem.* **139**, 179–197 (2016).

9. Foley, P. B., Hare, D. J. & Double, K. L. A brief history of brain iron accumulation in Parkinson disease and related disorders. *J. Neural Transm.* **129**, 505–520 (2022).

10. Zhang, P. *et al.* Ferroptosis was more initial in cell death caused by iron overload and its underlying mechanism in Parkinson's disease. *Free Radic. Biol. Med.* **152**, 227–234 (2020).

11. Medeiros, M. S. *et al.* Iron and Oxidative Stress in Parkinson's Disease: An Observational

Study of Injury Biomarkers. *PLOS ONE* **11**, e0146129 (2016).

12. Logroscino, G., Chen, H., Wing, A. & Ascherio, A. Blood donations, iron stores, and risk of Parkinson's disease. *Mov. Disord.* **21**, 835–838 (2006).

13. Deng, Q. *et al.* Lower hemoglobin levels in patients with parkinson's disease are associated with disease severity and iron metabolism. *Brain Res.* **1655**, 145–151 (2017).

14. Pichler, I. *et al.* Serum Iron Levels and the Risk of Parkinson Disease: A Mendelian Randomization Study. *PLOS Med.* **10**, e1001462 (2013).

15. Powell, L. W., Seckington, R. C. & Deugnier, Y. Haemochromatosis. *The Lancet* **388**, 706–716 (2016).

16. Liu, Y.-H. *et al.* Diet Quality and Risk of Parkinson's Disease: A Prospective Study and Meta-Analysis. *J. Park. Dis.* **11**, 337–347 (2021).

17. Bianchi, V. E., Rizzi, L. & Somaa, F. The role of nutrition on Parkinson's disease: a systematic review. *Nutr. Neurosci.* **26**, 605–628 (2023).

18. Zhang, X., Molsberry, S. A., Schwarzschild, M. A., Ascherio, A. & Gao, X. Association of Diet and Physical Activity With All-Cause Mortality Among Adults With Parkinson Disease. *JAMA Netw. Open* **5**, e2227738 (2022).

19. Guasch-Ferré, M. & Willett, W. C. The Mediterranean diet and health: a comprehensive overview. *J. Intern. Med.* **290**, 549–566 (2021).

20. Paknahad, Z., Sheklabadi, E., Derakhshan, Y., Bagherniya, M. & Chitsaz, A. The effect of the Mediterranean diet on cognitive function in patients with Parkinson's disease: A randomized clinical controlled trial. *Complement. Ther. Med.* **50**, 102366 (2020).

21. Solch, R. J. *et al.* Mediterranean diet adherence, gut microbiota, and Alzheimer's or Parkinson's disease risk: A systematic review. *J. Neurol. Sci.* **434**, 120166 (2022).

22. Wang, P. *et al.* Long-Term Consumption of Ultraprocessed Foods and Prodromal Features of Parkinson Disease. *Neurology* **104**, e213562 (2025).

23. White, N. S. *et al.* Improved Conspicuity and Delineation of High-Grade Primary and

Metastatic Brain Tumors Using "Restriction Spectrum Imaging": Quantitative Comparison with High B-Value DWI and ADC. *Am. J. Neuroradiol.* **34**, 958–964 (2013).

24. Fan, C. C. *et al.* Multivariate genome-wide association study on tissue-sensitive diffusion metrics highlights pathways that shape the human brain. *Nat. Commun.* **13**, 2423 (2022).

25. Ho, M.-L., Rojas, R. & Eisenberg, R. L. Cerebral Edema. *Am. J. Roentgenol.* **199**, W258–W273 (2012).

26. Wu, X. *et al.* Pathogeneses and Imaging Features of Cerebral White Matter Lesions of Vascular Origins. *Aging Dis.* **12**, 2031–2051 (2021).

27. Wang, J.-Y. *et al.* Meta-analysis of brain iron levels of Parkinson's disease patients determined by postmortem and MRI measurements. *Sci. Rep.* **6**, 36669 (2016).

28. Adams, P. C. Epidemiology and diagnostic testing for hemochromatosis and iron overload. *Int. J. Lab. Hematol.* **37**, 25–30 (2015).

29. UK Biobank - Food preferences web questionnaire Version 1.1. (2020).

30. Biggs, J. factor_analyzer Documentation.

31. Persson, I. & Khojasteh, J. Python Packages for Exploratory Factor Analysis. *Struct. Equ. Model. Multidiscip. J.* **28**, 983–988 (2021).

32. Abdi, H. Factor Rotations in Factor Analyses. in *Encyclopedia for Research Methods for the Social Sciences* (2003).

33. DiCiccio, T. J. & Efron, B. Bootstrap confidence intervals. *Stat. Sci.* **11**, 189–228 (1996).

34. Pierce, B. L. & Burgess, S. Efficient Design for Mendelian Randomization Studies: Subsample and 2-Sample Instrumental Variable Estimators. *Am. J. Epidemiol.* **178**, 1177–1184 (2013).

35. Smith, G. D. & Ebrahim, S. Mendelian randomization: prospects, potentials, and limitations. *Int. J. Epidemiol.* **33**, 30–42 (2004).

36. Freeman, L. R., Haley-Zitlin, V., Rosenberger, D. S. & Granholm, A.-C. Damaging effects of a high-fat diet to the brain and cognition: A review of proposed mechanisms. *Nutr.*

Neurosci. **17**, 241–251 (2014).

37. Hirai, S. *et al.* High-sucrose diets contribute to brain angiopathy with impaired glucose uptake and psychosis-related higher brain dysfunctions in mice. *Sci. Adv.* **7**, eabl6077 (2021).

38. Frantellizzi, V. *et al.* Neuroimaging in Vascular Cognitive Impairment and Dementia: A Systematic Review. *J. Alzheimers Dis.* **73**, 1279–1294 (2020).

39. Aigner, E. *et al.* Glucose acts as a regulator of serum iron by increasing serum hepcidin concentrations. *J. Nutr. Biochem.* **24**, 112–117 (2013).

40. Haas, J., Berg, D., Bosy-Westphal, A. & Schaeffer, E. Parkinson's Disease and Sugar Intake—Reasons for and Consequences of a Still Unclear Craving. *Nutrients* **14**, 3240 (2022).

41. Komici, K., Femminella, G. D., Bencivenga, L., Rengo, G. & Pagano, G. Diabetes Mellitus and Parkinson's Disease: A Systematic Review and Meta-Analyses. *J. Park. Dis.* **11**, 1585–1596 (2021).

42. Zhong, Q. & Wang, S. Association between diabetes mellitus, prediabetes and risk, disease progression of Parkinson's disease: A systematic review and meta-analysis. *Front. Aging Neurosci.* **15**, (2023).

43. Huth, C. *et al.* Biomarkers of iron metabolism are independently associated with impaired glucose metabolism and type 2 diabetes: the KORA F4 study. *Eur. J. Endocrinol.* **173**, 643–653 (2015).

44. Gomaa, E. Z. Human gut microbiota/microbiome in health and diseases: a review. *Antonie Van Leeuwenhoek* **113**, 2019–2040 (2020).

45. Wilson, A. S. *et al.* Diet and the Human Gut Microbiome: An International Review. *Dig. Dis. Sci.* **65**, 723–740 (2020).

46. Toh, T. S. *et al.* Gut microbiome in Parkinson's disease: New insights from meta-analysis. *Parkinsonism Relat. Disord.* **94**, 1–9 (2022).

47. Romano, S. *et al.* Meta-analysis of the Parkinson's disease gut microbiome suggests alterations linked to intestinal inflammation. *Npj Park. Dis.* **7**, 1–13 (2021).

48. Wallen, Z. D. *et al.* Characterizing dysbiosis of gut microbiome in PD: evidence for overabundance of opportunistic pathogens. *Npj Park. Dis.* **6**, 1–12 (2020).

49. Mahbub, N. U., Islam, M. M., Hong, S.-T. & Chung, H.-J. Dysbiosis of the gut microbiota and its effect on α-synuclein and prion protein misfolding: consequences for neurodegeneration. *Front. Cell. Infect. Microbiol.* **14**, (2024).

50. Angelova, P. R. *et al.* Alpha synuclein aggregation drives ferroptosis: an interplay of iron, calcium and lipid peroxidation. *Cell Death Differ.* **27**, 2781–2796 (2020).

51. Shao, C. *et al.* Parkinson's Disease Risk and Alcohol Intake: A Systematic Review and Dose-Response Meta-Analysis of Prospective Studies. *Front. Nutr.* **8**, (2021).

52. Mitchell, E., Chohan, H., Bestwick, J. P. & Noyce, A. J. Alcohol and Parkinson's Disease: A Systematic Review and Meta-Analysis. *J. Park. Dis.* **12**, 2369–2381 (2022).

53. Topiwala, A. *et al.* Associations between moderate alcohol consumption, brain iron, and cognition in UK Biobank participants: Observational and mendelian randomization analyses. *PLOS Med.* **19**, e1004039 (2022).

54. Harrison-Findik, D. D. Role of alcohol in the regulation of iron metabolism. *World J. Gastroenterol. WJG* **13**, 4925–4930 (2007).

55. Grochowski, C. *et al.* Serum iron, Magnesium, Copper, and Manganese Levels in Alcoholism: A Systematic Review. *Mol. Basel Switz.* **24**, 1361 (2019).

56. Shobi, V., Adeseye, A., Billy, B. & Medhat, K. Hemochromatosis, alcoholism and unhealthy dietary fat: a case report. *J. Med. Case Reports* **15**, 84 (2021).

57. Ioannou, G. N., Dominitz, J. A., Weiss, N. S., Heagerty, P. J. & Kowdley, K. V. The effect of alcohol consumption on the prevalence of iron overload, iron deficiency, and iron deficiency anemia. *Gastroenterology* **126**, 1293–1301 (2004).

58. Milman, N. T. A Review of Nutrients and Compounds, Which Promote or Inhibit Intestinal

Iron Absorption: Making a Platform for Dietary Measures That Can Reduce Iron Uptake in Patients with Genetic Haemochromatosis. *J. Nutr. Metab.* **2020**, e7373498 (2020).

59. Milman, N. T. Managing Genetic Hemochromatosis: An Overview of Dietary Measures, Which May Reduce Intestinal Iron Absorption in Persons With Iron Overload. *Gastroenterol. Res.* **14**, 66–80 (2021).

60. Al, L. Beta-Thalassemia. *PubMed* https://pubmed.ncbi.nlm.nih.gov/20301599/ (1993).

61. Kim, J. H. *et al.* The Association between Anemia and Parkinson's Disease: A Nested Case-Control Study Using a National Health Screening Cohort. *Brain Sci.* **11**, 623 (2021).

62. Hong, C. T. *et al.* Newly Diagnosed Anemia Increases Risk of Parkinson's disease: A Population-Based Cohort Study. *Sci. Rep.* **6**, 29651 (2016).

63. Savica, R. *et al.* Anemia or low hemoglobin levels preceding Parkinson disease. *Neurology* **73**, 1381–1387 (2009).

64. Rozani, V. *et al.* Anemia in men and increased Parkinson's disease risk: A population-based large scale cohort study. *Parkinsonism Relat. Disord.* **64**, 90–96 (2019).

65. Cho, I. Y. *et al.* Anemia and the risk of Parkinson's disease in Korean older adults: A nationwide population-based study. *Sci. Rep.* **10**, 4268 (2020).

66. Meléndez-Flores, J. D. *et al.* Association of anemia with Parkinson's disease: a systematic review with meta-analysis of epidemiological studies. *Rev. Mex. Neurocienc.* **24**, 019–028 (2023).

67. Chavhan, G. B., Babyn, P. S., Thomas, B., Shroff, M. M. & Haacke, E. M. Principles, Techniques, and Applications of T2*-based MR Imaging and Its Special Applications. *Radiographics* **29**, 1433–1449 (2009).