1 **Forecasting the spatial spread of an Ebola epidemic in real-time:**
2 **comparing predictions of mathematical models and experts**
3

4 James D Munday*[†1,2,3], Alicia Rosello[†1,2], W. John Edmunds[$1,2], Sebastian Funk[$1,2]
5
6 [1] Centre for mathematical modelling of infectious diseases, London School of Hygiene and
7 Tropical Medicine, United Kingdom
8 [2] Department of infectious disease epidemiology, London School of Hygiene and Tropical
9 Medicine, United Kingdom
10 [3] Department of Biosystems Science and Engineering, ETH Zürich, Switzerland
11
12 [†] These authors contributed equally
13 [$] These authors contributed equally
14
15 *Corresponding author: james.munday@bsse.ethz.ch

17 **Keywords:** Ebola, Forecast evaluation, Expert elicitation, Mathematical modelling, Outbreak
18 response

Abstract

Ebola virus disease outbreaks can often be controlled, but require rapid response efforts frequently with profound operational complexities. Mathematical models can be used to support response planning, but it is unclear if models improve the prior understanding of experts.

We performed repeated surveys of Ebola response experts during an outbreak. From each expert we elicited the probability of cases exceeding four thresholds between two and 20 cases in a set of small geographical areas in the following calendar month. We compared the predictive performance of these forecasts to those of two mathematical models with different spatial interaction components.

An ensemble combining the forecasts of all experts performed similarly to the two models. Experts showed stronger bias than models forecasting two-case threshold exceedance. Experts and models both performed better when predicting exceedance of higher thresholds. The models also tended to be better at risk-ranking areas than experts.

Our results support the use of models in outbreak contexts, offering a convenient and scalable route to a quantified situational awareness, which can provide confidence in or to call into question existing advice of experts. There could be value in combining expert opinion and modelled forecasts to support the response to future outbreaks.

Background

Following the initial emergence in 1976 in Zaire (now the Democratic Republic of the Congo, DRC)[1], epidemics of Ebola Virus Disease (EVD) have occurred, on average, every 12 - 24 months[2]. EVD is a viral haemorrhagic fever first caused by the Ebola Zaire virus (EZV), with a case fatality rate of 25-90%[3]. A major outbreak in North-Eastern provinces of DRC between 2018-2020 resulted in over 3300 reported cases and over 2100 deaths [4](Figure 1).

Figure 1. The extent of the 2018-2020 ebola outbreak in north-eastern DRC and areas included in our study. A) Daily incidence in north-eastern DRC between August 2018 and March 2020. Grey points show days prior to the study period, coloured points show days within the study period (November 2019 - March 2020), hue indicates month. B) Shows the total number of cases of ebola recorded in each Health Zone. C) Number of cases in each month and health zone during the period covered by this study, health zones outlined in red show all health zones affected by the entire epidemic.

Transmission of EZV occurs mainly through direct contact during the symptomatic phase of infection; therefore, isolation of infected individuals with strict infection control, contact tracing, and safe burials have been key to controlling past EVD outbreaks[5], although setting

59   specific challenges can hamper containment efforts [6]. More recently, vaccination has also
60   become a tool for outbreak control, with two vaccines now licensed for use[7][8].
61
EVD outbreaks typically occur in resource-poor settings where limited communication and poor
accessibility make logistics of surveillance and vaccination campaigns challenging.
Understanding the spatial risk of future spread is therefore useful to allow response teams to
focus efforts on high-risk areas. Mathematical and statistical models have been used
extensively to forecast the spread of infectious diseases, including EVD[9]. Such models rely on
a combination of statistical inference based on epidemiological data and information about the
mechanisms underlying the dynamics of infection. However, the dynamics of EVD are
frequently governed by changing contextual factors which are challenging to forecast
quantitatively. For example, violent conflicts or flooding can seriously hinder, interrupt, or even
reverse the impact of containment efforts [6,10]. Moreover, changes in healthcare capacity
and health seeking behaviour of patients can strengthen or weaken efforts to reduce
transmission [11]. The timing and impact of these factors is notoriously difficult to predict using
mathematical models.
75
Models are used by epidemic response experts to support decision making in the field. In
addition to models, experts also make judgments as to the future spread of the virus based on
their interpretation of the current status of the outbreak combined with their knowledge of
other less tangible factors such as the geography, climate (eg. seasonal variation in accessibility
of particular areas) and soft intelligence about the escalation of conflict in areas which may as a
result, be harder to access by response teams. There are clear costs and benefits to human-
made and modelled-based forecasts. Whereas models are objectively based on observations of
the past outbreak dynamics and current case data, experts have additional knowledge of the
complex factors surrounding the outbreak response. It is therefore difficult to assess the impact
mathematical models have on decision making, how much modelled forecasts differ from those
made by experts in the field and whether either modelled or human forecasts are
systematically more accurate or useful. Moreover, the knowledge of experts in the field of EVD
epidemiology, with a good understanding of the geographical area of study may provide an
invaluable resource that is currently underused in forecasting.
90
Previous studies have aimed to establish the relative performance of humans and models in
predicting infectious disease spread in human populations, particularly in the context of acute
respiratory infections such as Influenza and SARS-CoV-2. Three studies have evaluated the
predictions of humans against models, explicitly. The first of these evaluated short-term
forecasts and season-wide predictions of reports of influenza-like-illness (ILI) in the United
States of America (USA) [12] and two studies [13,14] compared short-term forecasts of cases of
and deaths from COVID-19, firstly in Germany and Poland and secondly in the United Kingdom
(UK). All three studies found that humans tended to perform better than the mathematical and
statistical models selected for comparison when predicting cases. However, the COVID-
focussed studies found the human ensembles performed worse than the ensemble prediction
of the models when predicting deaths - These results were maintained when only self-declared
'experts' were included in the forecasts. A number of other studies recorded expert predictions

without comparison to mathematical models. A study conducted early in the COVID-19
pandemic [15] evaluated the relative ability of laypeople and experts to predict the course of
the UK epidemic over the first calendar year. The study found that both experts and laypeople
typically under-predicted the impact overall, however experts' forecasts were more accurate
and better calibrated than laypeople. A study of expert predictions in the United States of
America [16] evaluated their weekly forecasts of case incidence and total deaths in the first
year against a pooled ensemble of all predictions. The study found that the ensemble
outperformed every expert individually over the period of the study. A similar study surveyed
experts regarding the total number of cases and deaths from MPox in the USA during 2022 [17],
however these predictions are yet to be evaluated. Overall, these studies provide evidence that
human predictions can play a valuable role in epidemiological prediction, providing a
comparator and complementary method to mathematical and statistical modelling.
115
In this paper we extend the use of expert forecasters to predict spatial risk of transmission in
the context of a local outbreak. We made monthly forecasts of the geographic spread of Ebola
Virus Disease (EVD) from November 2019 to March 2020 during the declining phase of the
2018-2020 outbreak in the Democratic Republic of the Congo (DRC) using both expert
predictions collected through regular interviews and with two spatially explicit computational
transmission models with different spatial interaction assumptions: a gravity model and an
adjacency model (where transmission can only occur between contiguous regions), see the
methods section for details. Alongside supporting situational awareness, these forecasts were
motivated by an aim to inform site selection for a planned vaccine trial. The objective was to
identify areas that had seen no cases yet and thus were not already being supported by
vaccination and other interventions, but were at high risk of still becoming affected by the EVD
outbreak, thus allowing estimation of efficacy [18]. Here we evaluate the performance of the
forecasts and select ensembles of the methods in predicting continued transmission and flare-
ups of EVD in health zones (HZs) close to the affected area. We further study variation in
forecast quality against a selection of factors related to local demography, case history and
forecast implementation.

Methods

Expert elicitation

Experts in EVD epidemiology with knowledge of the local geography speaking English or French were identified originally by convenience sampling. The pool of experts was then expanded through recommendation from the identified experts (snowball sampling). This approach was best suited to capture the expertise of individuals who were most often temporarily based in the field.

A pilot study was carried out in November 2019. Subsequently, monthly interviews were held over WhatsApp in December 2019, January 2020, February 2020, and March 2020. All interviews were scripted. The main biases of this type of study (availability bias, representativeness bias, overconfidence, motivational bias, anchoring on past estimates) were briefly discussed during the first interview.

Experts also were provided with an interactive map of the outbreak area and surrounding health zones (HZ) showing the number of total cases during the outbreak and during the two preceding weeks for reference (supplementary figure S1). HZ were numbered to facilitate communication with the experts.

Experts were asked to estimate the number of reported probable and confirmed cases they would expect per HZ during the following month using the online MATCH Uncertainty Elicitation Tool [19] (supplementary figure S1). Through this platform, the experts and the researcher (AR) interacted in real time. The "roulette" (chips and bins) method was used. Experts were instructed to place a total of 20 chips over the available bins (0-1 cases, 2-4 cases, …, 48-50 cases). Therefore, each chip represented for the expert a 5% probability that the number of cases was in the bin where the chip was placed. This process aimed to capture the uncertainty surrounding the expert's estimates.

The experts were asked to estimate the number of reported cases they would expect in the HZ where there had been 1 or more cases in the 2 preceding weeks, as well as Goma (Figure 2). The experts were then asked to identify any additional HZ where they would predict 1 or more cases during the following month with >5% probability, and to also estimate the number of reported cases they would expect in these HZ. In the pilot study, carried out in October 2019, experts were asked to forecast the number of cases they expected during November 2019 in 10 HZs: Beni, Goma, Kalunguta, Katwa, Lolwa, Mabalako, Mambasa, Mandima, Nyankunde, and Oicha.

Figure 2. Health zones included in the model and Expert Elicitation survey. A shows the provinces around the affected area, and included in the transmission model, the red box shows the area detailed in panel B. B shows the health zones included at least once in the Expert Elicitation survey we conducted.

Ethics

LSHTM ethics approval was obtained for this study (reference: 17633). Signed informed consent was taken from experts willing to participate and their verbal consent was requested again at the beginning of each elicitation.

Modelling framework

In parallel with the expert elicitation programme, we developed a modelling framework to forecast spatial risk of infection. In the framework, incidence of cases is forecast in each Health Zone based on historical case reports. The model was formed of two components, the autoregressive component, and the spatial component.

$$\lambda_{it} = \sum_{s=t-(D+L)}^{t-D} \left(\gamma N_{i,s} + \alpha \sum_{j} w_{ij} N_{j,s} \right) \tag{1}$$

The auto-regressive component modelled the rate of infections in a particular health zone i, on day t, to be proportional to the number of cases in the same health zone (i) between dates t-(D+L) and t-D, where L is the estimated latent period and D is the estimated infectious period. The spatial component accounts for transmission between health zones, where rate of infection was proportional to the cases in each other Health Zone (i.e. ∀j j≠i) and moderated by a pairwise specific factor defined by a spatial kernel w_{ij} We used two spatial kernels, both of which use proximity of health zones to each other and their respective population size, P_i and P_j. Firstly, the gravity model which treats interaction in an analogous way to Newtonian gravity with population size in place of mass, such that interaction reduces distance, *d*, raised to a power, *k*.

201
$$w_{ij} = \frac{P_i P_j}{d_{ij}^k} \tag{2}$$

202
203
204 Secondly, we applied a model with adjacency-based interaction. In this model only adjacent HZs
205 can interact. The strength of interaction between HZs is proportional to the product of their
206 population sizes.
207

208
$$w_{ij} = \delta_{ij} P_i P_j \tag{3}$$

$$\delta_{ij} = \begin{cases} 1, & \text{if adjacent} \\ 0, & \text{otherwise} \end{cases} \tag{4}$$

210
211
212 Cases were modelled as Poisson distributed such that:
213

214
$$N_{i,t} = \text{Poisson}(\lambda_{i,t}) \tag{5}$$

215
To forecast cases, we fitted the spatiotemporal model to historical data from the 60 days prior
to the date the forecast was made, accounting for cases in health zones in seven regions (169
HZs) centred on the location of the epidemic; Nord-Kivu, Ituri, Tshopo, Maniema, Sud-Kivu,
Haut-Uele, Bas-Uele. We fit the model using the No U-Turn Sampling (NUTS) method for
Hamiltonian Monte Carlo with Stan [20], a probabilistic programming framework. We
estimated α and γ, which vary the contribution of within-health-zone and between-health-
zone transmission. We also estimated k, which determines how rapidly transmission rate
decays with distance in the spatial component of the model. We sampled parameters from the
resultant joint posterior distribution to simulate daily incidence in all HZs in the seven regions,
up to and including the last day of the following month. We performed 1000 iterations for each
forecast date. We then extracted the full distribution of the number of cases incident within the
calendar month of interest. Forecasts were made using data up to the last day of the month
prior to the forecast period.
229
Enslemble forecasts
Ensemble forecasts were calculated as an average of the probabilities attributed by the
members of the ensemble. For the expert ensemble the arithmetic mean was calculated across
all experts with equal weighting. Similarly the model ensemble used the unweighted mean of
the model forecasts. For the mixed (model and expert) ensemble, the mean was weighted such
that the combined weight of the experts forecasts and the combined weight of the models
forecasts were equal.
237
238 *Quantification of risk and forecast evaluation*
239 To compare the model and the expert forecasts and score them according to the eventual true
240 number of cases we calculated the probability attributed to cases over four thresholds, >=2,
241 >=6, >=10 and >=20 cases.

We evaluated the forecasts using the Brier Score, a proper scoring rule which quantifies how accurate a forecast or a group of forecasts are when compared to true data after the event. The Brier score, BS, is defined as the square of the difference between the probability of observing an event and the observation O_i status, which takes a value 1 or 0 for cases observed and none observed respectively. We calculated this for multiple (N) forecasts by taking the mean of the individual forecast scores.

$$BS = \frac{1}{N} \sum_{i=1}^{N} (p_i - o_i)^2 \tag{6}$$

We also quantified the general bias and calibration of the forecasts by considering the hazard rate predicted by each forecast, which we calculated as the sum of probabilities attributed to exceeding each threshold. This gives the number of HZs the forecast 'expected' to cross the threshold in each month. To quantify the bias of each set of predictions, we took the difference between the hazard rate and the actual number of HZs that exceeded each threshold in each month. We refer to this as the *hazard gap* (HG).

$$HG = \sum_{\forall HZ} p(c > c_{thresh}) - \sum_{\forall HZ} o(c > c_{thresh}) \tag{7}$$

Where $p(c>c_{thresh})$ isthe probability of threshold exceedance and $o(c>c_{thresh})$ is the observation status of the threshold exceedence 1 if true 0 if false.

Results

Expert panel and health zones included in survey

Over the study period, we conducted a total of 40 interviews with 15 experts, three of which took place during the pilot phase (November 2019). Figure 3 shows the timeline of the expert elicitations.

Forecast Month   a   November 2019   a   December 2019   a   January 2020   a   February 2020   a   March 2020

Figure 3. Timeline of the expert elicitation. Each point shows the date of the interview of the expert labelled to obtain forecasts for the following month. Colour indicates the month for which the forecast was made, the forecast windows are highlighted with a shaded band of the same colour.

Eight experts worked at the World Health Organization, four at the London School of Hygiene & Tropical Medicine, two for Médecins Sans Frontières, and one at the DRC Ministry of Health. Most experts (10/15) had more than five years of experience working in infectious disease epidemiology. About half of the interviews (21 of 40) were conducted with experts that were in the outbreak area (defined HZs affected by EVD or Goma, the site of the international response base) or had been there within 2 weeks of the interview. Four experts had never been in the outbreak area.

Table 1: Experts and health zones included in each round of the survey. The left part of the table details the experts interviewed (highlighted in green) the health zones included in the main survey in each month. In

addition, the right part of the table details the health zones nominated by experts and the number of experts that nominated each one.

Month forecasted	Experts interviewed (Highlighted)			Health zones in interview (HZs)	No. experts	HZ nominated
December	1	2	3	Beni	7	Oicha
	4	5	6	Goma	5	Komanda
	7	8	9	Kalunguta	2	Butembo
	10	11	12	Mabalako	2	Katwa
	13	14		Mambasa	1	Lolwa
				Mandima	1	Makiso - Kisangani
					1	Nyankunde
January	1	2	3	Beni	6	Butembo
	4	5	6	Biena	3	Katwa
	7	8	9	Goma	1	Kalunguta
	10	11	12	Mabalako	1	Mangurerdjipa
	13	14		Mandima		
				Oicha		
February	1	2	3	Beni	6	Oicha
	4	5	6	Bunia	4	Biena
	7	8	9	Butembo	2	Vuhovi
	10	11	12	Goma	1	Lolwa
	13	14		Kalunguta		
				Katwa		
				Kayna		
				Mabalako		
				Mambasa		
				Mandima		
				Musienene		
March	1	2	3	Beni	3	Mabalako
	4	5	6	Butembo		
	7	8	9	Goma		
	10	11	12	Mandima		
	13	14				

Between December 2024 and March 2025, eight to ten experts provided monthly forecasts for 4 to 11 health zones (HZs), estimating the probability that reported cases would exceed various thresholds (2, 6, and 20 cases). Expert predictions showed variable accuracy across months and zones.

Experts correctly identified Mabalako as the highest-risk HZ in December. They attributed an
average 82% probability of exceeding 2 cases; Mabalako reported 38 cases that month,
exceeding all thresholds, although the probability assigned to exceeding the higher thresholds
was similar to that of Beni (3 cases). Other zones with moderate cases—Kalanguta (5), and
Mambasa (4)—were assigned lower probabilities and generally did not exceed higher case
thresholds. Interestingly, Mandima, which had no reported cases, was assigned a relatively high
probability (72%) of exceeding 2 cases, indicating some overestimation in this zone.

In January, Beni and Mabalako again accounted for most cases (22 and 11, respectively) and
were recognized as high risk by the experts. However, experts underestimated the scale of the
outbreak in Beni, assigning 0% probability to exceeding 20 cases there, despite this threshold
being surpassed. Mandima continued to be forecasted as high risk, though no cases were
reported. Predictions for other zones such as Oicha and Biena suggested moderate risk, but no
cases were confirmed.

Of the 11 nominated HZs, only Beni reported confirmed cases in February (9), exceeding the 6
case threshold. Experts collectively assigned a 70% probability to this event. Similar
probabilities assigned to Mabalako (60%), where no cases were reported, following cases in the
prior two months. Several other zones were predicted to exceed 2 cases, yet reported none.

No cases were reported in any health zone during March. Experts broadly anticipated this, with
only one expert assigning over 50% probability of exceeding 2 cases in any HZ. Beni had the
highest average assigned risk at 33%.

328 **Figure 4. Expert elicitation results and accuracy of predictions.** Only the HZs that were rated by all experts are
329 included here. Results are shown as probabilities (vertical axes) that a given health zone (horizontal panels)
330 exceeds a given threshold (horizontal axes) according to the experts (box plots) or models (square / diamond for
331 gravity and adjacency models, respectively) across different months (vertical panels). Health zone / month
332 combinations where the given thresholds were exceeded are marked in cyan, and ones where they weren't in red.
333
Performance evaluation
We evaluated the forecasts using the Brier score. The overall scores of individual experts varied
between 0 and 0.6 across the four thresholds. Collectively, the experts scored best at the
highest threshold (20 cases) and worst for forecasts of the lowest threshold (2 cases). The
models also performed better at higher thresholds than low thresholds, but the difference was
less pronounced. Overall, the gravity model ranked best amongst all forecasts at the 2 case
threshold. It also ranked best for this threshold in the month of February and consistently in the
top half of forecasts in December and January, however performed comparatively poorly in
March, ranking higher than only one of the experts. The adjacency model also performed better
than the experts overall for the 2 case threshold. Related to this, including the models
improved the ensemble forecast. Although the gravity model performed better than the
adjacency model for higher thresholds, together the models performed similarly to the expert
ensemble forecast overall. In January and February the gravity model performed well compared
to the adjacency model and the expert ensemble, however in March both models performed
particularly badly compared to the experts for all thresholds. None of the experts performed
consistently well relative to the others, experts 3 and 10 performed best for the 2 case
threshold, whereas experts 13 and 14 did best for higher thresholds.
351

Figure 5. Evaluation of forecasts made by the experts, models, and ensembles. A shows the overall Brier Score for each expert, model and ensemble (calculated over all forecasts included in the study). In B each panel shows the Brier score across all health zones for each month (vertical) at each case threshold (Horizontal). Coloured points show each expert score, the violin plot shows their distribution. The grey hollow points show the model scores, the yellow points show the ensemble scores (circles show experts alone, squares show models alone and triangles show experts and models with 50% weight given to each). C Shows the ranking of each expert and model in terms of forecast performance

To evaluate how the different forecasts may impact decision making we ranked the health zones for each month, based on the probability of exceeding each threshold of cases forecast by each ensemble and by the model alone (supplementary figure S6). In general, the model and

the ensembles all ranked health zones that did reach the threshold highly. In some cases the
model performed better, ranking health zones that did meet the threshold higher than the
experts, specifically ranking Beni Higher than Mandima in higher thresholds (>=6 and 10 cases)
for the forecast of January, where Beni ultimately had cases and Mandima did not, in that
month. Considering the models separately, the gravity model performed better than the
adjacency model in general, with the adjacency model occasionally performing worse than the
experts when ranking the HZs. This was clearest in the forecasts of November and January.
372
Bias and calibration in forecasts
We evaluated the bias in each forecast type by considering the hazard gap between forecasts
and actual cases. We found that experts systematically forecasted higher risk of the lowest
threshold (>=2 cases) than was warranted, but tended to forecast lower risk of exceeding the
highest threshold (>=20 cases) than was borne out across all HZs (Figure 6). When calculated
across all months, this bias was present in 12 of the 15 experts. The models did not show clear
consistent bias in either direction.
380

381
382 **Figure 6 Bias and calibration of forecasts.** Panels show the Hazards gap difference between the Hazard rate
383 (expected number of exceedances across all health zones) for each threshold attributed by the forecast and the
384 actual number of Health Zones that exceeded the associated threshold. Each panel shows one forecast (expert or
385 model) in each month. The bottom row shows the same for each forecast calculated over the entire study period.
386
387 *Forecasting flare-ups*
388 In addition to the health zones presented to all experts, each expert was able to nominate
389 health zones, which they deemed at risk. Experts nominated seven further HZs to forecast in
390 December, four in January, four in February and one in March (Table 1).
391

The only two HZs not included in the default list but exceeding 2 cases were Butembo and
Katwa. These were nominated by 2 of the 10 experts (4 and 10) both attributing 50% chance of
exceeding 2 cases. In contrast, Oicha and Komanda had the most nominations with 7 and 5
each, and 6 and 4 of the 10 experts interviewed allotted greater than 5% chance of 2 or more
cases in December, with attributed probabilities ranging between 5% and 95% chance of
crossing the 2 case threshold.
398
399   In January, six of the eight experts interviewed nominated Butembo (1,2,4,5,9, and 11 with
400   probabilities of 5% to 85% of crossing the 2 case threshold) while three of them also nominated
401   Katwa (2 and 9 giving 85% and 5 giving 5%). Kalunguta and Manguredjipa were also nominated
402   by one expert each, 5 with 10% and 11 with 50% respectively. None of the nominated health
403   zones crossed the threshold in January.
404
In February six of the ten experts interviewed (2, 4, 5, 11, 12 and 14) nominated Oicha with
probabilities between 10% and 95% of crossing the 2 case threshold. Four (2, 4, 5 and 11)
nominated Biena with probabilities between 10% and 95% of exceedance. Experts 4 and 8 also
nominated Vuhovi attributing 55% and 20% probability of threshold exceedance respectively.
Expert 3 nominated Lolwa alone but gave no probability of exceeding 1 case. No HZs not
included in the interview as default crossed the 2 case threshold in February.
411
412   In March three (4, 8 and 11) of the eight experts interviewed gave probabilities of 35%, 50%
413   and 15% of exceeding the 2-case threshold respectively. No HZs not included in the interview as
414   default crossed the 2 case threshold in March.
415
To compare the model with the experts we included all HZs modelled and attributed all HZs not
nominated by experts an exceedance probability of 0%. To allow comparison, we also set all
HZs given a probability of lower than 5% to 0% for both the gravity and adjacency models.
When considering the Brier Score (Figure 7), we found that the gravity model performed
comparably to some experts when forecasting for December, and February. The adjacency
model performed worse than all the experts in every month except February. In every month
the ensemble of experts did better than the models and including the models in the ensembles
reduced performance.
424
425
426
427

Figure 7 Evaluation of forecasts made in health zones not included in the main survey. Each panel (right to left) shows the Brier score across all health zones for each month. Coloured points show each expert score, density plot shows their overall distribution. The red points show the model scores, the yellow points show the ensemble scores (squares show experts alone, crosses show experts and models with 50% weight given to each).

Discussion

We compared forecasts of the geographic spread of Ebola made by experts, with those made using a modelling framework. Since the outbreak dynamics of Ebola are highly sensitive to the changeable context in which they take place, mathematical models and expert opinions are expected to have different strengths and weaknesses, with models benefiting from objective inference from previous observations and experts able to utilise detailed knowledge about the outbreak and the changing surrounding context to make informed projections of risk. By interviewing experts and asking them to forecast risk in a structured way, we were able to compare the performance of their forecasts against those made with well-established modelling approaches in a quantifiable and robust way.

Overall, the forecasts made by the group of experts as a whole performed similarly to those of the model, with a few consistent exceptions. The model performed better than the experts when considering the lowest threshold in four of the five months covered by the survey, but performance was more comparable for the higher thresholds. The model also performed marginally better when ranking health zones by risk of ongoing transmission, indicating that use of a model may improve prioritisation of health zones when attributing resources.

We found that both methods performed better when considering higher case thresholds. This is likely to be due to a combination of bias in both forecast types towards under prediction of cases and the fact that there were few instances where the higher thresholds were reached.

Although individual experts frequently out-performed ensembles in individual instances, no
individual expert outperformed the ensembles overall. This supports the practice of considering
predictions from a range of experts over a smaller number of more specialist or experienced
experts. The models tended to perform similarly to the ensemble representing more consistent
performance across all forecasts.

Experts tended to be more biased than models, especially at low case thresholds with a
tendency to over-predict cases to a greater degree than models. This bias reduced rapidly as
the case threshold increased. This may be interpreted as over cautiousness from the experts
regarding potential for geographic spread of the virus but confidence that transmission could
be contained quickly. This trend reflects a pattern amongst previous introductions into new
health zones earlier in the pandemic, where a small number of cases were reported, but the
local outbreak was quickly stopped (Figure 1).

To our knowledge, our study is the first to record experts' assessment of geographical risk at a
local level during an epidemic and the first comparison of outbreak response experts'
predictions to those of models in real-time. Although direct comparison is not possible, our
results lead to conclusions that are broadly similar to those from previous studies [12–14],
however each of these studies found that ensemble expert forecasts performed better than the
comparison models, whereas our study found no clear performance difference. This may
suggest that experts are better at predicting simple time series than geographic distribution of
cases. However, we cannot view these findings independently from the different survey designs
or study contexts.

There are a number of important limitations to consider when interpreting our results. The
context within which we conducted the study has important implications for interpretation.
Due to the timing and logistics of setting up the questionnaire, the study only began in the
closing phase of the epidemic, whereas the relative performance of experts and models may
differ during different phases of the epidemic. For example, in the early phase where dynamics
are driven more by infectious transmission than established response practices, or during the
peak where changes in intervention strategy may be more influential. The stage of the epidemic
also meant that there was a substantial trend towards 'negative' results (i.e. no threshold
exceedance), which is likely to favour some forecasting methods over others.

Additionally, experts were not all interviewed on the same day and interviews occurred several
days before the beginning of the month they were forecasting. In some cases experts were
interviewed up to 2 weeks prior to the beginning of the month. This means that the information
available differed both between experts and with the model, which was run considering data
up to the first day of each month. This reduces our ability to compare models to experts
directly, however, it could also be argued that this is a 'built-in' factor which represents the
inherent challenge of eliciting predictions from experts. In addition, the interview process for
experts was quite taxing and required a phone call - which can cause scheduling challenges

during an epidemic response. It may be that other methods, with less arduous and more
flexible data entry would improve responses.
500
Our analysis represents the comparison of expert forecasts to only two specific forecasting
models. There are a great range of models that could have been applied in this context which
may have differed in performance to those we used. We chose these models for convenience
since we were applying them to the outbreak at the time of the interviews. It is also possible
that some of the experts involved in the study had ingested results from our model, which were
available through our online dashboard, or other models being used at the time.
507
Since our findings, like those of similar studies, suggest that models and experts perform
comparably in this context, there is an argument that models have no value in informing expert
decision making. It can be argued, however, that models remain useful in outbreak response.
Firstly, while the models performed similarly to the ensemble forecasts of the experts, there
was no individual expert that performed consistently better than the models. Secondly, models
are much more easily scaled and generalised making them simple to deploy in new contexts
and to adapt as epidemics grow. Expert interviews are time-consuming and often inconvenient,
especially in the context of outbreak response activities, which are characteristically fast paced.
Models therefore offer a more convenient route to a quantified insight, which from our results,
performs comparably to the way groups of experts may think. Finally, there are ways to
combine both methods. For example, in the event that expert forecasts can be garnered, joint
ensembles can capture information from both the expert and modelled forecasts. Further, we
suggest that models can offer a role in aiding decision making by providing confidence in or
calling into question expert advice that is being considered.
522

Conclusions
Our analysis evaluated performance of experts and models when forecasting the spatial spread
of Ebola, representing the first such study incorporating local geographic distribution and the
first to focus on an epidemic in a resource poor setting. We found that forecasts made by
experts and models performed comparably overall, but experts tended to be slightly more
biased towards predicting that a small number of cases would persist. The results support the
use of models in outbreak response and provide insight into how models and expert opinion
could be combined when tackling future epidemics.

531
Acknowledgements
We would like to thank Xavier de Radiguès, Neale Batra, Nabil Tabbal, Mathias Mossoko, Chris
Jarvis, Thibaut Jombart, Denis Ardiet, Michel Van Herp, Silimane Ngoma, Olivier le Polain,
Esther van Kleef, Noé Guinko, and Amy Gimma as well as 2 other experts who preferred to
remain anonymous, for their participation as experts in this study. We would also like to thank
David Smith, Thibaut Jombart, Chris Jarvis, Flavio Finger, and Anton Camacho for their helpful
advice in conducting this survey.

Ethics
LSHTM ethics approval was obtained for this study (reference: 17633). Signed informed consent
was taken from experts willing to participate and their verbal consent was requested again at
the beginning of each elicitation.

Code and data availability
All data and code used to process the expert interview responses can be found here:
https://github.com/epiforecasts/Ebola-Expert-Interviews. The forecasts were performed using
the EpiCastR package https://github.com/epiforecasts/EpiCastR. The code used for the analysis
and scoring of the forecasts can be found here: https://github.com/epiforecasts/Ebola-Expert-
Ellicitation

Author contributions
AR and WJE conceived and designed the interviews. JDM and SF conceived and designed the
modelling framework and the evaluation of forecasts. AR conducted the expert interviews and
prepared the interview data for comparison. JDM implemented the model and performed the
formal forecast evaluations. JDM, AR, WJE and SF interpreted the results. JDM and AR wrote
the manuscript. JDM, AR, WJE and SF edited the manuscript.

Funding statement
This study was partly funded by the Department of Health and Social Care using UK Aid funding
and is managed by the National Institute for Health and Care Research (VEEPED: PR-OD-1017-
20002; AR and WJE). This study was partly funded by the Wellcome Trust (210758/Z/18/Z : JDM
and SF). The views expressed in this publication are those of the authors and not necessarily
those of the funders.

Conflicts of interest
The authors have no conflicts of interest to declare

568
569 **References**
570

1. Report of an International Commission. 1978 Ebola haemorrhagic fever in Zaire, 1976. *Bull. World*
Health Organ. **56**, 271–293.

2. Rosello A *et al.* 2015 Ebola virus disease in the Democratic Republic of the Congo, 1976-2014. *Elife*
4. (doi:10.7554/eLife.09015)

3. In press. Ebola virus disease. See https://www.who.int/news-room/fact-sheets/detail/ebola-virus-
disease (accessed on 19 December 2023).

4. WHO Regional Office for Africa. 2020 Outbreaks and Emergencies Bulletin, Week 5: 27 January -
02 February 2020.

5. Kucharski AJ, Camacho A, Flasche S, Glover RE, Edmunds WJ, Funk S. 2015 Measuring the
impact of Ebola control measures in Sierra Leone. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 14366–14371.

6. Adongo PB, Tabong PT-N, Asampong E, Ansong J, Robalo M, Adanu RM. 2016 Preparing towards
Preventing and Containing an Ebola Virus Disease Outbreak: What Socio-cultural Practices May
Affect Containment Efforts in Ghana? *PLoS Negl. Trop. Dis.* **10**, e0004852.

7. In press. WHO prequalifies Ebola vaccine, paving the way for its use in high-risk countries. See
https://www.who.int/news/item/12-11-2019-who-prequalifies-ebola-vaccine-paving-the-way-for-its-
use-in-high-risk-countries (accessed on 20 December 2023).

8. Woolsey C, Geisbert TW. 2021 Current state of Ebola virus vaccines: A snapshot. *PLoS Pathog.* **17**,
e1010078.

9. Chowell G, Viboud C, Simonsen L, Merler S, Vespignani A. 2017 Perspectives on model forecasts
of the 2014-2015 Ebola epidemic in West Africa: lessons and the way forward. *BMC Med.* **15**, 42.

10. Wannier SR *et al.* 2019 Estimating the impact of violent events on transmission in Ebola virus
disease outbreak, Democratic Republic of the Congo, 2018-2019. *Epidemics* **28**, 100353.

11. Funk S *et al.* 2017 The impact of control strategies and behavioural changes on the elimination of
Ebola from Lofa County, Liberia. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **372**.
(doi:10.1098/rstb.2016.0302)

12. Farrow DC, Brooks LC, Hyun S, Tibshirani RJ, Burke DS, Rosenfeld R. 2017 A human judgment
approach to epidemiological forecasting. *PLoS Comput. Biol.* **13**, e1005248.

13. Bosse NI *et al.* 2022 Comparing human and model-based forecasts of COVID-19 in Germany and
Poland. *PLoS Comput. Biol.* **18**, e1010405.

14. Bosse NI, Abbott S, Bracher J, van Leeuwen E, Cori A, Funk S. 2023 Human judgement forecasting
of COVID-19 in the UK. *Wellcome Open Res.* **8**, 416.

15. Recchia G, Freeman ALJ, Spiegelhalter D. 2021 How well did experts and laypeople forecast the
size of the COVID-19 pandemic? *PLoS One* **16**, e0250935.

16. McAndrew T, Reich NG. 2022 An expert judgment model to predict early stages of the COVID-19
pandemic in the United States. *PLoS Comput. Biol.* **18**, e1010485.

17. McAndrew T *et al.* 2022 Early human judgment forecasts of human monkeypox, May 2022. *Lancet*
Digit Health **4**, e569–e571.

18. Watson-Jones D *et al.* 2022 Protocol for a phase 3 trial to evaluate the effectiveness and safety of a
heterologous, two-dose vaccine for Ebola virus disease in the Democratic Republic of the Congo.
BMJ Open **12**, e055596.

19. 2014 A web-based tool for eliciting probability distributions from experts. *Environmental Modelling*
& Software **52**, 1–4.

20. Stan Team. 2012 *Stan Modeling Language User's Guide and Reference Manual.*

614
615
616
617