Single-cell RNA sequencing unraveled the expression heterogeneity of hematopoietic stem and progenitor cells and lymphoid cell development dysregulation in childhood asthma

Danying Zhu[*,1,2], Guang Li[*,3], Lang Yuan[*,1], Zeyu Zeng[1,2], Na Dong[1], Chao Wang[1], Ming Chen[1], Lijian Xie[5], Guohui Ding[#,4], Libing Shen[#,4], Xiaoyan Dong[#,1,2]

[1] Department of Pulmonology, Shanghai Children's Hospital, School of medicine, Shanghai Jiao Tong University, Shanghai, China

[2] Institute of Pediatric Infection, Immunity, and Critical Care Medicine, Shanghai Jiao Tong University School of Medicine, Shanghai, China.

[3] Daozhi Precision Medicine Technology (Shanghai) Co., Ltd; Shanghai, China.

[4] Institute for Digital Health, International Human Phenome Institutes (Shanghai), Shanghai, China.

[5] Department of Pediatrics, Jinshan Hospital, Fudan University, Shanghai, 201508, China.

*These authors contributed equally: Danying Zhu, Guang Li, Lang Yuan

#Corresponding authors

Xiaoyan Dong, *Ph.D.*

Department of Respiratory, Shanghai Children's Hospital, School of medicine, Shanghai Jiao Tong University, Shanghai, China.

Institute of Pediatric Infection, Immunity, and Critical Care Medicine, Shanghai Jiao Tong University School of Medicine, Shanghai, China.

No.355, Luding Road, Shanghai 200062, China.

Tel: 18930734642

E-mail: dongxy@shchildren.com.cn

Libing Shen, *Ph.D*

Institute for Digital Health, International Human Phenome Institutes (Shanghai), Shanghai, China.

12/F, No.258 Guoxia Road, Yangpu District, Shanghai, China.

Tel: 13482391370

E-mail: shenlibing@ihup.org.cn

Guohui Ding, *Ph.D.*

Institute for Digital Health, International Human Phenome Institutes (Shanghai), Shanghai,

China.

12/F, No.258 Guoxia Road, Yangpu District, Shanghai, China.

Tel:17721236391

Email: dingguohui@ihup.org.cn

Abstract

Asthma is a long-term inflammatory disease affecting airways and lungs with usual

onset in childhood. Its cause is not fully understood up to now. Here, using single-cell

RNA sequencing, we profile peripheral blood mononuclear cells (PBMCs) from three

pediatric patients with onset asthma and four age-matched healthy controls to

investigate the cellular etiology of childhood asthma. The overall expression features

among three asthma patients' PBMCs demonstrate that innate immunity is commonly

upregulated while adaptive immunity is commonly downregulated in childhood

asthma, but each patient has different molecular phenotypes. The analyses of the

expression profiles of hematopoietic stem and progenitor cells (HSPCs) further show

that the HSPCs of asthma patients have heterogeneous expression backgrounds with

more specific differentially expressed genes (DEGs) in each patient than common

DEGs and a common feature of low S100 protein binding gene expression. S100A8,

S100A9, S100A12, and RETN are universally upregulated in various cell types of

asthma patients. The cell developmental trajectories in three asthma cases exhibit an

abnormal immune cell development pattern compared to that in health control. The

dysregulated lymphoid lineage development is observed in all 3 patients, but there is

no identical abnormal pattern for each patient. The pseudo-time analyses of gene

expression show that the expression dynamics of two proto-oncogenes, JUN and SPI1,

and six inflammatory response related genes (S100A8, S100A9, S100A12, IL7R,

IL32, CCL5) are relevant to abnormal immune cell development in asthma patients.

The cell-cell communication analyses reveal the contribution of incoming annexin

signal towards dendritic cells and the outgoing resistin signal from dendritic cells to

asthma heterogeneity. Interestingly, the plasma blast cells of asthma patient 3 with

severe symptoms exhibit dual cell identities of both plasma blast cells and T cells. Our

scRNA-Seq analyses for three asthma patients reveal a complex cellular etiology for

childhood asthma and provide a new research direction for the comprehensive and

systematic understanding of key molecular mechanisms of childhood asthma.

Introduction

Asthma, a persistent respiratory condition, is marked by inflammation of the airways, heightened bronchial responsiveness, and variable airway obstruction. This complex disease involves a multitude of cellular participants, including eosinophils, T and B lymphocytes, macrophages, neutrophils, and epithelial cells, which interact with environmental factors to induce chronic inflammation, reversible airflow limitations, and airway remodeling in patients[1]. The involvement of various cell types and their mediators is a key aspect of asthma's pathogenesis[2]. This diversity in asthma phenotypes and the intricacy of its pathogenesis are further complicated by the fact that respiratory infections and allergen exposures are known to trigger asthma exacerbations[1]. While asthma can affect individuals across all age groups, the manifestation, underlying pathophysiology, and response to treatment can vary significantly between children and adults. Recognizing the molecular and cellular distinctions between childhood and adult asthma is crucial for crafting age-tailored diagnostic and therapeutic approaches.

childhood asthma, in particular, presents with distinct clinical and immunological characteristics compared to its adult counterpart. Asthma in children often manifests as allergic phenotypes with a higher incidence of eosinophilic inflammation, whereas adults may display a broader range of subtypes, including non-eosinophilic variants. The immature and plastic nature of the immune system in children can lead to unique cellular responses and immune dysregulation. Therefore, there are some differences in the treatment by unique age groups[3]. Single-cell sequencing offers a powerful tool to explore age-specific alterations in immune cell populations, their activation states, and functional attributes, providing a deeper understanding of the mechanisms driving childhood asthma.

Previous research has predominantly focused on the single-cell sequencing of adult asthma[4, 5], whereas this study applies single-cell technology to childhood asthma, aiming to uncover the role of immune abnormalities in the pathogenesis of asthma in children. Through single-cell sequencing, specific cell subpopulations can be identified, and their roles in the inflammatory responses of children with asthma can be understood, as well as their responses to current treatment methods. These findings may help explain why childhood asthma exhibits differences in clinical presentation and treatment response compared to adult asthma, and they provide new perspectives for future research and clinical practice.

Method

Patients

During the initial hospital stay, before any treatment, we collected three fresh peripheral blood samples from patients experiencing acute asthma. The diagnosis of asthma was confirmed using the criteria outlined in the 2023 Global Initiative for Asthma (GINA) report, which can be accessed at https://ginasthma.org/2023-gina-main-report/. Atopic status was determined by a total IgE level of 200 IU/ml or higher. For single-cell RNA sequencing (scRNA-seq), we enrolled four healthy individuals who were participating in routine physical examinations and had no recent history of fever, infection, or immunization. Data from three of these individuals have previously been published[6]. All participants were recruited from Shanghai Children's Hospital between December 2019 and February 2023. Exclusion criteria for the study included: Participants who were undergoing immunotherapy, such as anti-IgE treatment, or had received such therapy within the last three months. Individuals with significant abnormalities identified in their complete blood count. Subjects who were currently enrolled in an asthma-related pharmaceutical or interventional study, or had been part of such a study within the last four weeks. Patients with concurrent medical conditions necessitate systemic corticosteroids or other immunomodulatory treatments. The study was approved by the Ethics Committee of Shanghai Children's Hospital (Protocol Numbers: 2019R081, 2022R029-F-01). Informed consent was obtained from all participants and their guardians.

Single-cell preparation and sequencing

A total of 2 milliliters of venous blood was collected from each participant using tubes containing EDTA as an anticoagulant. These samples were processed within four hours to preserve cell integrity. Peripheral blood mononuclear cells (PBMCs) were separated from the collected blood using Ficoll-Paque medium and density gradient centrifugation. The viability of the isolated PBMCs was assessed using trypan blue staining, and only samples with more than 90% viable cells were selected for further analysis. A calculated volume of the cell suspension, estimated to contain around 12,000 cells per sample, was prepared to ensure an adequate cell count for subsequent capture and sequencing processes. The Chromium Next GEM Single Cell

V(D)J Reagent Kits v1.1 from 10x Genomics were utilized for capturing single cells

and constructing the corresponding libraries. The cell suspension, along with

barcoded gel beads and partitioning oil, were loaded onto the $10 \times$ Genomics

Chromium Chip to create single-cell Gel Beads-in-Emulsion (GEMs). Inside each

GEM, cells were lysed, and their transcripts were barcoded through reverse

transcription. After reverse transcription, the cDNA, now carrying cell barcodes,

underwent PCR amplification to ensure a sufficient quantity for sequencing. The

construction of single-cell RNA sequencing (scRNA-seq) libraries was accomplished

using the 5' Library Kits. The prepared libraries were sequenced on an Illumina

NovaSeq platform, producing paired-end reads with a length of 2×150 base pairs.

This sequencing step generates the raw data necessary for subsequent data analysis.

scRNA-seq data analysis

The demultiplexed reads were then aligned to the GRCh38 reference genome

using the Cell Ranger software suite. This gene-barcode matrix was subsequently

analyzed using Seurat version 3.0.2 for various analytical steps, including quality

control, data normalization, dimensionality reduction, batch effect correction,

clustering, and data visualization. The specific procedures carried out with Seurat

included: Applying quality control criteria for the majority of samples, which

involved ensuring a total UMI count ranging from 200 to 6000 and limiting the

percentages of mitochondrial, hemoglobin, and ribosome genes to below 10%, 0.1%,

and 3%, respectively. Integrating samples collected during the initial phase of the

study (C1-C4) to eliminate batch effects using anchor.   Scaling the integrated matrix

and employing the principal components derived from principal component analysis

(PCA) for uniform manifold approximation and projection (UMAP). Utilizing UMAP

to visualize cells in a two-dimensional space, highlighted their gene expression

pattern similarities. Conducting shared nearest neighbor graph-based clustering on the

PCA-reduced data to identify the primary cell types within the PBMCs. Examining

the expression of canonical marker genes to refine cell cluster annotations.

Determining cell identities through multimodal reference mapping with the

SeuratDisk package.

Differential expression, functional enrichment analysis, and pseudo-time analysis

The marker genes for each cluster were calculated with the FindAllMarkers function within the Seurat package. The clusterProfiler package (version 3.16.0) was used for function over-representation analysis of the differentially expressed genes (DEGs) with a false discovery rate (FDR) threshold of <0.05. Gene Ontology (GO), KEGG pathways, and hallmark gene sets from the MSigDB database (version 7.1) were used as gene function databases to assess the enrichment of specific biological functions or pathways among the DEGs. Pseudo-time analysis of cell differentiation trajectories for each sample dataset was performed with R package Monocle 2[7]. The expression feature and inferred cell type for each sample dataset from the Seurat result were used to annotate the cell dataset for the Monocle analysis pipeline. We used the Monocle built-in approach named "dpFeature" to detect the variable genes that define a cell's differentiation. Its advantages are needing no prior biological knowledge and discovering important ordering genes from the data itself. Dimension reduction was performed with 2 max components and the "DDRTree" method. Cell-cell communication analysis was performed with the R package CellChat 1.6.1[8].

Human serum resistin Enzyme-Linked Immunosorbent Assay (ELISA) analysis

EDTA-anticoagulated whole blood was transferred to the laboratory and processed immediately after collection. Centrifuge samples for 15 minutes at 1000 x g at 4 □ within 30 minutes of collection and stored at -80 □. We followed a human resistin ELISA kit (Signalway Antibody, Maryland, USA, Catalog No: EK2351) protocol to analyze peripheral blood serums.

Results

Characteristics of the study subjects for asthma patients and health controls

In our single-cell RNA analysis study, we included a cohort of 7 childhood subjects, comprising 3 children experiencing asthma exacerbations and 4 healthy individuals without asthma. T The mean age of the participants ranges from 1 to 9 years old. An additional group of 14 asthma patients and health controls ranging from 1 to 13 years old, were included in the study. Allergic testing in asthmatic children revealed sensitization to house dust mites. Allergic testing in asthmatic children revealed sensitization to house dust mites.

Single-cell RNA profiling of PBMCs in health and asthma patients

We collected the peripheral blood samples derived from 3 patients with onset asthma (A1-A3). The patients were diagnosed according to the GINA 2023 (https://ginasthma.org/2023-gina-main-report/). For each patient, the blood sample was taken on the first days before therapy. We also collected fresh peripheral blood samples from four age-matched healthy donors as controls (C1–C4).

We used the 10× Genomics platform for scRNA-seq of PBMCs isolated from the samples. PBMCs were loaded onto the platform and about 5000 - 14000 cells per sample could be recovered from the sequencing data. The total number of detected cells passing quality control was 34,354 for 3 asthma patients and 22,398 cells from healthy controls. Based on the scRNA-seq profiles, we clustered the cells across samples with Seurat 4.0 and visualized them in two-dimensional space (Fig. 1a). The PBMCs in our study could be clustered into 12 major cell types including B cells, CD4 T cells, CD8 T cells, CD14 monocytes (CD14 mono), CD16 monocytes (CD16 mono), dendritic cells (DC), mucosa-associated invariant T cells (MAIT), gamma-delta cells (gdT), hematopoietic stem and progenitor cells (HSPC), natural killer cells (NK), plasma blast cells (Plasmablast), and platelets. Each cell type was verified with PBMC multimodal reference object and canonical gene markers (Supplementary Figure 1a and 1b). The percentage of major cell types for each sample is shown in Figure 1b. The more detailed classification of cell types excluding CD4 T and CD8 T cells shows that CD14 monocytes are significantly increased and CD16 monocytes are significantly decreased in asthma patients (Figure 1c, *P*-value < 0.05). The detailed classification of T cell types in each sample is shown in Supplementary Figure 1c.

Expression features of all cells and HSPCs in asthma patients

We first compared the gene expression level of all cells in each asthma patient to controls. The overall differentially expressed genes (DEGs) were identified for each asthma patient. There are 174 upregulated genes in A1 (Asthma1), 29 upregulated genes in A2 (Asthma2), and 246 upregulated genes in A3 (Asthma3) (Figure 2a). There are 21 common upregulated genes among three asthma patients. GO analysis shows that these commonly upregulated genes are involved in the extracellular region,

secretion, and antimicrobial function (Figure 2b). GO result also shows that innate

immunity plays a major role in childhood asthma which manifests as a defense

response to bacterium and inflammatory response. There are 57 specific upregulated

genes in A1, 3 specific upregulated genes in A2, and 128 specifically upregulated

genes in A3. GO analyses of the specifically upregulated genes in A1 and A3 show

that these two patients have different molecular phenotypes (Supplemental Tables 1

and 2). A1 exhibits the specific expression of signal sequences (usually 20-30 amino

acids long) and positive regulation of T cell migration while A3 has an excessive

expression of immunoglobulin. 3 specific upregulated genes in A2 are IGHV3-73,

CDK6 and NELL2 which are not sufficient for GO analysis. We also find 33

downregulated genes in A1 (Asthma1), 80 downregulated genes in A2 (Asthma2), and

66 downregulated genes in A3 (Asthma3) (Figure 2c). There are 9 common

upregulated genes among three asthma patients. GO analysis shows that these

common downregulated genes are involved in adaptive immunity and T cell receptor

(Figure 2d). There are 13 specific downregulated genes in A1, 37 specific

downregulated genes in A2, and 20 specific downregulated genes in A3. A1 exhibits

the specific repression of the external side of the plasma membrane and B cell

receptor signaling pathway expression (Supplemental Table 3). A2 exhibits the

specific repression of the immunoglobulin complex (Supplemental Table 4). A3

exhibits the specific repression of the T cell receptor complex (Supplemental Table 5).

We further examined the expression features of HSPCs in asthma patients, because

the other types of cells are all derived from them. There are 176 upregulated HSPC

genes in A1 (Asthma1), 211 upregulated HSPC genes in A2 (Asthma2), and 199

HSPC upregulated genes in A3 (Asthma3) (Figure 3a). There are 81 common

upregulated HSPC genes among three asthma patients. GO analysis shows that these

commonly upregulated HSPC genes are mainly involved in the cellular components

of extracellular exosome and ER to Golgi transport vesicle membrane (Figure 3b).

There are 29 specifically upregulated HSPC genes in A1, 72 specifically upregulated

HSPC genes in A2, and 55 specifically upregulated HSPC genes in A3. No significant

GO result is found for 29 specifically upregulated HSPC genes in A1. HSPCs in A2

exhibit a specific expression of extracellular exosome related genes and HSPCs in A3

exhibit a specific expression of endoplasmic reticulum related genes (Supplemental

Tables 6 and 7). There are 179 downregulated HSPC genes found in A1 (Asthma1),

203 downregulated HSPC genes found in A2 (Asthma2), and 119 HSPC
downregulated genes found in A3 (Asthma3) (Figure 3c). GO analysis shows that
these common downregulated HSPC genes are mainly involved in S100 protein
binding (Figure 3d). There are 85 specifically downregulated HSPC genes in A1, 113
specifically downregulated HSPC genes in A2, and 53 specifically downregulated
HSPC genes in A3. HSPCs in A1 exhibit a specific repression of adaptive immunity
and T cell receptor complex expression (Supplemental Table 8). HSPCs in A2 exhibit
a specific repression of immune response related genes' expression (Supplemental
Table 9). HSPCs in A3 exhibit a specific repression of extracellular exosome and
immunity relate genes' expression (Supplemental Table 10). Interestingly, we find that
T cell receptor genes are repressed in all three asthma patients, although they are not
statically significant in the HSPCs of A2 and A3 (Supplemental Tables 9 and 10).
284
285 In expression feature analyses, we find that S100A8, S100A9, S100A12, and RETN
286 are universally upregulated in various cell types of asthma patients, especially
287 S100A8 and S100A9 (Figure 4). S100A8, S100A9, and S100A12 are known for
288 participating in inflammatory response while RETN has antibacterial activity[9-11].
289

Single-cell trajectory reconstructions of PBMCs in health controls and 3 asthma
patients

The expression features of asthma patients' HSPCs demonstrate that the expression of
immunity and T cell receptor genes are abnormally repressed in them. Since HSPCs
give rise to the other cells, it is worth investigating the cell developmental states in
asthma patients. Single-cell trajectory analysis could help us reconstruct the cell
developmental paths of PBMCs based on featured variable genes. The analyses show
that the PBMCs in healthy controls can be divided into 5 developmental states while
the PBMCs in asthma patients have 7, 3, and 5 developmental states, differently
(Figure 5a, 5c, 5e, and 5g). All cell types are plotted on each trajectory in each sample
(Figure 5b, 5d, 5f, and 5h).
301

Based on the major cell types and numbers, state 1 in healthy controls can be
classified as B lineage, state 2 can be classified as monocyte lineage, state 3 is a
mixed lineage (B and T), state 4 can be classified as T lineage, and state 5 can be

classified as T-NK lineage (Figure 3b and Supplement Figure 2a). Myeloid lineage
(monocytes) and two lymphoid lineages (B cells and T cells) are clearly differentiated
in the PBMCs of healthy controls. However, the clear cell development patterns
disappear in 3 asthma patients. Each patient exhibits unique cell developmental
patterns different from healthy controls. A1 has 7 states and 4 of them are T lineages
according to the major cell numbers (state 1, 2, 3, and 6, Figure 5d and Supplement
Figure 2b). A2 has only 3 states and a part of B cells are mixed with T cells (state 1,
Figure 5f and Supplement Figure 2c). A3 has 5 states, but B cells are mixed with T
cells in state 3 (state 1, Figure 5h and Supplement Figure 2d). Although different
childhood asthma patient has a different cell developmental state, abnormal lymphoid
lineage development is observed in all 3 patients (multiple T lineage states or no clear
B lineage state). This result is consistent with the expression feature analyses showing
that adaptive immunity is repressed in asthma patients. T cells and B cells are a major
constitution of adaptive immune system and their poor development naturally leads to
insufficient the adaptive immunity.
320

Pseudo-time expression dynamic analyses of proto-oncogenes and inflammatory
response related genes
The abnormal cell developmental trajectories in asthma patients should be
accompanied by abnormal gene expression patterns. JUN gene is a transcription factor
regulated by various extracellular stimuli including peptide growth factors, pro-
inflammatory cytokines, and even UV irradiation[12]. SPI1 gene is also a transcription
factor regulating gene expression during myeloid and B-lymphoid cell development[13].
Both of them are proto-oncogenes essential for cellular differentiation[14, 15]. We further
include six inflammatory response related genes in pseudo-time expression dynamic
analyses. Their express dynamics were plotted along cell developmental trajectories
in both health control and three asthma patients.
332

333   In health controls, we observed the onset expression of JUN in the early pseudo-time
334   state and a peak expression of six inflammatory response related genes in the early-
335   middle pseudo-time state. SPI1 is expressed in the late pseudo-time state (Figure 6a).
336   In A1 and A2, the expression dynamics of JUN, SPI1, and six inflammatory response
337   related genes are almost the same (Figure 6b and 6c). JUN's onset expression is

repressed while SPI1 exhibits an onset expression in the early pseudo-time state. S100A8, S100A9, and S100A12 show an early onset expression rather than a peak expression in the early-middle pseudo-time state. IL7R, IL32, and CCL5 start to express in the early-middle pseudo-time state as health control, but they show a persistent expression level instead of an obvious expression peak. Notably, IL7R, IL32, and CCL5 all participate in the biological process of negative regulation of the T cell apoptotic process (GO:0070233). It implies that the T cells have prolonged cell survival in A1 and A2. A3 shows a different expression dynamic pattern from health control, A1, and A2 (Figure 6d). In A3, JUN is also repressed while the expression patterns of IL7R, IL32, and CCL5 are more similar to health control. S100A8, S100A9, S100A12, and SPI1 show a persistent expression level without decline. Since SPI1 activates gene expression during myeloid and B-lymphoid cell development and S100 protein family plays a role in cell growth and differentiation, their persistent expression pattern indicates an abnormal lymphoid cell development fate in A3.

DCs function as a signaling hub in cell communication and play a role in asthma heterogeneity

We performed cell communication analyses for both health control and asthma patients to investigate their cell-cell interactions. The number of interactions and interaction strength for health control and three asthma patients are shown in Figure 7 . Generally, asthma patients have a weaker interaction strength than health control and A3 has the smallest number of interactions and the weakest interaction strength among four samples. Among the major immune cell types, we find that dendritic cells have the largest number of both incoming and outgoing interaction numbers(Supplemental Figure 3a, 3b, 3c, and 3d). DCs are known to be the messenger cells communicating between innate and adaptive immune systems[16]. They are also known as the antigen presenting cells which interact with T cells and B cells to activate and regulate the adaptive immune response[17]. Although DCs don't have the strongest interaction strength among the major immune cell types (Supplemental Figure 4a, 4b, 4c, and 4d), they surely function as a signaling hub in cell communication based on cell communication analyses.

We further examined the DCs' signaling differences between health control and

asthma patients. Using health control as a signaling background, the signaling

changes were detected in three asthma patients. The DCs of A1 has a specific

incoming signal of annexin (Supplemental Figure 5a). The DCs of A2 has a specific

outgoing signal of resistin (Supplemental Figure 5b). The DCs of A3 has a specific

incoming signal of annexin and a specific outgoing signal of resistin (Supplemental

Figure 5c). Annexin A1, A2 and A5 have a role in the regulation of inflammation

activation and may serve as the potential biomarkers for asthma[18-20]. They all show an

elevated expression tendency in asthma cases (Supplemental Figure 6). Resistin also

serves a potential biomarker for asthma and is related to inflammation as well[21, 22].

Our results show that annexin and resistin are actually two opposite directional signals

for the DCs of asthma patients. One of them is a sufficient predictor of asthma and

both of them indicate a severe symptom of asthma, since A3 is an acute severe asthma

patient. Both annexin and resistin show a self-reinforce trend in the CD14 monocytes

of asthma patients (Figure 8a and 8b). The upregulated signaling interactions in

asthma are mainly centered on the molecular component of external side of plasma

membrane and cell adhesion molecules (GO:0009897 and hsa04514, Supplemental

Figure 7a, 7b, and 7c). Furthermore, we found that CLEC2B-KLRB1 and LGALS9-

CD44 signaling interactions are downregulated in all three asthma patients

(Supplemental Figure 8a, 8b, and 8c). CLEC2B-KLRB1 interaction mediates the

activation of NK cells and monocytes[23]. LGALS9-CD44 interaction enhances the

stability and function of adaptive regulatory T cells[24]. The down-regulated CLEC2B-

KLRB1 and LGALS9-CD44 interactions proposes a weakened functions of NK cells,

monocytes and T cells in asthma patients.

Plasma blast cells in A3 express T cell receptor genes

Patient A3, a severe asthma sufferer who has been admitted to the pediatric intensive

care unit, exhibits a unique group of cells not present in healthy controls A1 and A2

(Figure 1a). This group predominantly consists of plasma blast cells along with a

mixed cluster comprising plasma blast cells (35.2%), erythrocytes (26.5%), CD14

monocytes (14.4%), CD4 T cells (11.2%), CD8 T cells (7.7%), and other cells (5%).

Upon comparing the gene expression profile of A3's plasma blast cells with those of

the healthy controls, A1 and A2, we discovered an unexpected finding: the plasma

blast cells in A3 express a range of T cell receptor genes, including TRBV20-1, in

addition to immunoglobulins (Figure 9a). Further investigation into the expression
levels of classical markers for plasma blast cells (CD38, CD19, CD27), CD4 T cells
(CD8A, CD27), CD8 T cells (CD40LG, CD27), and NK cells (GNLY, NKG7)
revealed through a violin plot that all these markers are indeed expressed in A3's
plasma blast cells (Figure 9b). This suggests that the plasma blast cells in A3 likely
possess dual cellular identities, exhibiting characteristics of both plasma blast cells
and T cells, which accounts for the expression of T cell receptor genes.
413

The expression level of RETN in lymphoid linage cells
The feature plot shows that the RETN expression is not evenly distributed among
different cell clusters. CD14 mono, CD16 mono, and dendritic cells have an
especially high expression level of RETN. The violin plot further shows that RETN
expression level in these four clusters is always higher in 3 asthma cases than in
healthy controls.   RETN encodes resistin, then we tested the level of resistin in
plasma between children with asthma and healthy controls, we found that resistin in
the serum increased significantly in the asthma group than in controls (Figure 10).
422

Discussions
In individuals with allergic airway disease, such as allergic asthma, the airways are
already sensitized to specific allergens. This means that the immune system has
developed an exaggerated response to these allergens, leading to chronic
inflammation and airway hyperresponsiveness. During an asthma exacerbation, this
preexisting allergic airway disease is further aggravated by additional inflammatory
stimuli or triggers. Nevertheless, the etiology of childhood asthma is so far poorly
understood[25].
431

The analyses of PBMCs from three childhood asthma patients show that childhood
asthma is a complex disease with distinct individualities at the cellular level. The
overall expression features among three asthma patients' PBMCs demonstrate that
innate immunity is commonly upregulated while adaptive immunity is commonly
downregulated in childhood asthma, but each patient has different molecular
phenotypes. The common upregulated genes in the HSPCs of asthma patient are
related to the cellular components of extracellular exosome and ER to Golgi transport
vesicle membrane and the common downregulated genes in the HSPCs of asthma

patients are related to S100 protein binding. The specific downregulated genes in the HSPCs of each asthma patients converge on immunity and T cell receptor related biological entities. Because HSPCs give the rise to the other cells, they are a cellular source of immunity. The expression abnormality in HSPCs would surely influence their descendant cells. The downregulation of S100 protein binding in the HSPCs indicates that S100 proteins are in an active form in the HSPCs of asthma patients. The downregulation of immunity related genes in HSPCs is a sign of hypo-immune tendency started from the cellular source in asthma cases. The active S100 protein and hyo-immune tendency in the HSPCs of asthma patients seem to lead to over innate immunity and low adaptive immunity for all cells of asthma patients. Although childhood asthma has common pathophysiological features at the cellular level, there are more specific differentially expressed genes (DEGs) in each patient than common DEGs, which proposes that the genetic causes of asthma vary across different children or different allergens can trigger different immune genes in a pediatric patient's immune system. These two explanations are mutually inclusive. Future studies includes bone marrow samples might give us much clearer views for this question, since HSPCs are generated from bone marrow.

The single-cell trajectory analyses further demonstrate the cell developmental heterogeneities among asthma patients. Different asthma patient has different developmental states. The abnormal lymphoid lineage development is observed in all 3 patients. A1 has multiple T cell states while A2 and A3 have a poorly developed lymphoid state of mixed B and T cells. The expression of T cell receptor genes is repressed in the HSPCs of all three asthma patients, although this repression is not statically significant in A2 and A3. It could partly explain the abnormal cell developmental states in asthma patients and the developmental heterogeneities among them. The delayed expression of T cell receptor genes could lead to the HSPC's descendant cells having difficulty receiving the lymphoid development signals. In A1, such deficiency is responded to the development of multiple T cell states by A1's immune system, which seems to compensate for the lack of a proper T cell developmental state. In A2 and A3, the delayed expression of T cell receptor genes is less severe than that in A1, which leads to the underdevelopment of lymphoid lineage. The abnormal single-cell developmental trajectories in asthma patients are also accompanied by the key genes' aberrant expression dynamics. JUN and SPI1 are two

proto-oncogenes, both of which have multiple functions in cell fate decisions[15, 26]. The

expression of JUN is repressed in the early stage of cell trajectories for all three

asthma patients. It indicates that the cell developmental fate of asthma patients has

trouble at the very beginning, which is consistent with the expression abnormality in

the HSPCs of asthma patients. In A1 and A2, the repression of JUN seems to be

replaced by the expression of SPI1. In A3, the beginning of the expression of SPI1 is

consistent with that in health control, but the expression of SPI1 shows no decline like

in health control. It implies that the developmental fates of distal cells in A3 could be

influenced by SPI1.

The cell-cell communication analyses further revealed the possible mechanisms

underlying the etiology of asthma. Although the number of cell-cell interactions is

similar in health control and asthma cases, the strength of cell-cell interactions is

much weaker in asthma patients, especially in A3. The overall weak signaling strength

is a clear sign of cell-cell communication disorder in asthma. It is in accordance with

the abnormal cell developmental trajectories of asthma. DCs induce primary immune

responses by processing and presenting antigen material on the cell surface of T

cells[27]. They function as the signaling hub communicating between innate and

adaptive immune systems. Two aberrant signals, annexin, and resistin, are found for

the DCs of asthma, both of which play a part in inflammatory response. Inflammation

is a generic immune response and is considered as a mechanism of innate immunity. It

explains why innate immunity is hyperactive in asthma patients. In our cases, each

asthma patient has his or her aberrant signaling pattern, annexin for A1, resistin for A2,

and both annexin and resistin for A3. It further explains the heterogeneities among

asthma patients. Resistin as an outgoing signal from DCs has been reported as a

predictor of asthma[21]. Our ELISA results show that its expression can be detected in

12 enlisted childhood asthma patients.

A3 is a special case in our study for the patient with severe symptoms compared to A1

and A2 whose symptoms are mild. Severe asthma means strong inflammation in

patient's airway which further causes breathlessness and continuous cough. The

profiling of A3's PBMCs first presents the patient's severity in our study. A3 has two

extra cell clusters compared with health control, A1 and A2. These two extra cell

clusters are annotated as mixed cells and plasma blast cells. The UMAP plot shows

that the mixed cells and plasma blast cells in A3 are stemmed from CD4 T cells. The expression features of A3's plasma blast cells demonstrate that they express T cell receptor genes as well. It is noticed that SPI1 has a persistent expression dynamic in A3. That A3's plasma blast cells show a duality of T cell identity could be linked to the SPI1's aberrant expression pattern. SPI1 might influence the developmental fate of plasma blast cells in A3 and further exacerbate the patient's symptoms.

Conclusions

Our research substantiates that childhood asthma is a multifaceted disease characterized by an upregulation of the innate immune system and a downregulation of the adaptive immune response. Single-cell RNA profiling of PBMCs from asthma patients has uncovered a variety of expression patterns among HSPCs, along with aberrant cell developmental trajectories and diminished cell-cell interaction signaling. While significant heterogeneity exists among different cases of asthma, the pathological features converge on certain commonalities, particularly the abnormal development of the lymphoid lineage and an overabundance of expression in genes related to S100 protein binding and immune response. Notably, S100A8, S100A9, S100A12, and RETN, along with ANXA1, ANXA2, and ANXA5, have emerged as pivotal molecular markers in asthma, directly implicated in the inflammatory response.

Contributions

Xiaoyan Dong and Guohui Ding designed the study. Danying Zhu wrote the manuscript. Libing Shen analysis the data. Zeyu Zeng did literature search. Lang Yuan and Na Dong contributed clinical samples. Chao Wang and Ming Chen collected the clincal data. Lijian Xie and Xiaoyan Dong funded the single cell sequencing.

Declaration of interests

The authors declare no competing interests.

Acknowledgements

This study was funded by Jinshan Distinct key medical specialty project (JSZK2023A04) and Institute of Pediatric Infection, Immunity, and Critical Care

540   Medicine Cultivate Research Project.

542   **Data Sharing Statement**

543   All data generated or analyzed during this study are included in this article. Further

544   enquiries can be directed to the corresponding author.

References

1. Tang W, Li M, Teng F, Cui J, Dong J, Wang W. Single-cell RNA-sequencing in asthma research. *Front Immunol*. 2022;13:988573. doi:10.3389/fimmu.2022.988573

2. Victor JR, Lezmi G, Leite-de-Moraes M. New Insights into Asthma Inflammation: Focus on iNKT, MAIT, and gammadeltaT Cells. *Clin Rev Allergy Immunol*. Dec 2020;59(3):371-381. doi:10.1007/s12016-020-08784-8

3. Reddel HK, Bacharier LB, Bateman ED, et al. Global Initiative for Asthma Strategy 2021: Executive Summary and Rationale for Key Changes. *Am J Respir Crit Care Med*. Jan 1 2022;205(1):17-35. doi:10.1164/rccm.202109-2205PP

4. Alladina J, Smith NP, Kooistra T, et al. A human model of asthma exacerbation reveals transcriptional programs and cell circuits specific to allergic asthma. *Sci Immunol*. May 12 2023;8(83):eabq6352. doi:10.1126/sciimmunol.abq6352

5. Li H, Wang H, Sokulsky L, et al. Single-cell transcriptomic analysis reveals key immune cell phenotypes in the lungs of patients with asthma exacerbation. *J Allergy Clin Immunol*. Mar 2021;147(3):941-954. doi:10.1016/j.jaci.2020.09.032

6. Wang Z, Xie L, Ding G, et al. Single-cell RNA sequencing of peripheral blood mononuclear cells from acute Kawasaki disease patients. *Nat Commun*. Sep 14 2021;12(1):5444. doi:10.1038/s41467-021-25771-5

7. Qiu X, Mao Q, Tang Y, et al. Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods*. Oct 2017;14(10):979-982. doi:10.1038/nmeth.4402

8. Jin S, Guerrero-Juarez CF, Zhang L, et al. Inference and analysis of cell-cell communication using CellChat. *Nat Commun*. Feb 17 2021;12(1):1088. doi:10.1038/s41467-021-21246-9

9. Shah RD, Xue C, Zhang H, et al. Expression of Calgranulin Genes S100A8, S100A9 and S100A12 Is Modulated by n-3 PUFA during Inflammation in Adipose Tissue and Mononuclear Cells. *PLoS One*. 2017;12(1):e0169614. doi:10.1371/journal.pone.0169614

10. Harris TA, Gattu S, Propheter DC, et al. Resistin-like Molecule alpha Provides Vitamin-A-Dependent Antimicrobial Protection in the Skin. *Cell Host Microbe*. Jun 12 2019;25(6):777-788 e8. doi:10.1016/j.chom.2019.04.004

11. Propheter DC, Chara AL, Harris TA, Ruhn KA, Hooper LV. Resistin-like molecule beta is a bactericidal protein that promotes spatial segregation of the microbiota and the colonic epithelium.

Proc Natl Acad Sci U S A. Oct 17 2017;114(42):11027-11033. doi:10.1073/pnas.1711395114

12. Wisdom R, Johnson RS, Moore C. c-Jun regulates cell cycle progression and apoptosis by distinct

mechanisms. *Embo j*. Jan 4 1999;18(1):188-97. doi:10.1093/emboj/18.1.188

13. Burda P, Laslo P, Stopka T. The role of PU.1 and GATA-1 transcription factors during normal and

leukemogenic hematopoiesis. *Leukemia*. Jul 2010;24(7):1249-57. doi:10.1038/leu.2010.104

14. Angel P, Hattori K, Smeal T, Karin M. The jun proto-oncogene is positively autoregulated by its

product, Jun/AP-1. *Cell*. Dec 2 1988;55(5):875-85. doi:10.1016/0092-8674(88)90143-2

15. Ray D, Culine S, Tavitain A, Moreau-Gachelin F. The human homologue of the putative proto-

oncogene Spi-1: characterization and expression in tumors. *Oncogene*. May 1990;5(5):663-8.

16. Monga I, Kaur K, Dhanda SK. Revisiting hematopoiesis: applications of the bulk and single-cell

transcriptomics dissecting transcriptional heterogeneity in hematopoietic stem cells. *Brief Funct*

Genomics. May 21 2022;21(3):159-176. doi:10.1093/bfgp/elac002

17. Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature*. Mar 19

1998;392(6673):245-52. doi:10.1038/32588

18. Lee SH, Lee PH, Kim BG, et al. Annexin A1 in plasma from patients with bronchial asthma: its

association with lung function. *BMC Pulm Med*. Jan 4 2018;18(1):1. doi:10.1186/s12890-017-0557-5

19. Dallacasagrande V, Hajjar KA. Annexin A2 in Inflammation and Host Defense. *Cells*. Jun 19

2020;9(6)doi:10.3390/cells9061499

20. Lee SH, Lee PH, Kim BG, Hong J, Jang AS. Annexin A5 Protein as a Potential Biomarker for the

Diagnosis of Asthma. *Lung*. Dec 2018;196(6):681-689. doi:10.1007/s00408-018-0159-x

21. Ballantyne D, Scott H, MacDonald-Wicks L, Gibson PG, Wood LG. Resistin is a predictor of

asthma risk and resistin:adiponectin ratio is a negative predictor of lung function in asthma. *Clin Exp*

Allergy. Aug 2016;46(8):1056-65. doi:10.1111/cea.12742

22. Malyszko J, Malyszko JS, Pawlak K, Mysliwiec M. Resistin, a new adipokine, is related to

inflammation and renal function in kidney allograft recipients. *Transplant Proc*. Dec

2006;38(10):3434-6. doi:10.1016/j.transproceed.2006.10.140

23. Welte S, Kuttruff S, Waldhauer I, Steinle A. Mutual activation of natural killer cells and

monocytes mediated by NKp80-AICL interaction. *Nat Immunol*. Dec 2006;7(12):1334-42.

doi:10.1038/ni1402

24. Wu C, Thalhamer T, Franca RF, et al. Galectin-9-CD44 interaction enhances stability and function

of   adaptive   regulatory   T   cells.   *Immunity*.   Aug   21   2014;41(2):270-82.

doi:10.1016/j.immuni.2014.06.011

25.   Martinez FD. Genes, environments, development and asthma: a reappraisal. *Eur Respir J*. Jan

2007;29(1):179-84. doi:10.1183/09031936.00087906

26.   Vogt PK. Fortuitous convergences: the beginnings of JUN. *Nat Rev Cancer*. Jun 2002;2(6):465-9.

doi:10.1038/nrc818

27.   Kelly RR. Microcomputers in education: age of romance, or age of reason? *Am Ann Deaf*. Sep

1982;127(5):693-7. doi:10.1353/aad.2012.1147

Figures and figure legends

Figure 1. Single-cell profiling of PBMCs in six samples. A. The integration single-cell profiling analysis of six samples including 4 healthy controls, asthma patient 1, asthma patient 2, and asthma patient 3. B. The percentage bar chart shows the continent of different cell types in each sample. C. The percentage bar chart shows the different cell types except T cells in each sample.

Figure 2. The analysis of upregulated and downregulated genes in asthma patients. A. Venn gram of the upregulated genes in 3 asthma patients. B. GO term enrichment analysis of the specific upregulated genes in 3 asthma patients. C. Venn gram of the downregulated genes in 3 asthma patients. D. GO term enrichment analysis of the specific downregulated genes in 3 asthma patients.

Figure 3. The analysis of upregulated and downregulated genes in asthma patients'

HSPC. A. Venn gram of the upregulated genes in 3 asthma patients' HSPC. B. GO

term enrichment analysis of the specific upregulated genes in 3 asthma patients'

HSPC. C. Venn gram of the downregulated genes in 3 asthma patients' HSPC. D. GO

term enrichment analysis of the specific downregulated genes in 3 asthma patients'

HSPC.

Figure 4. The violin plot of S100A9, S100A8, S100A12 and RETN gene expression

levels across different cell types for four sample groups.

Figure 5. Pseudo-time analysis of all cells in four sample groups. A. The
differentiation trajectory of all cells in healthy controls by state. B. The differentiation
trajectory of all cells in healthy controls by cell types. C. The differentiation trajectory
of all cells in asthma patient 1 (Ast1) by state. D. The differentiation trajectory of all
cells in asthma patient 1 (Ast1) by cell types. E. The differentiation trajectory of all
cells in asthma patient 2 (Ast2) by state. F. The differentiation trajectory of all cells in
asthma patient 2 (Ast2) by cell types. G. The differentiation trajectory of all cells in
asthma patient 3 (Ast3) by state. H. The differentiation trajectory of all cells in asthma
patient 3 (Ast3) by cell types.

Figure 6. Pseudo-time analyses of expression dynamics of two cell-fate and six inflammation related genes in four sample groups. A. Expression dynamics two cell-fate and six inflammation related genes in health control. B. Expression dynamics two cell-fate and six inflammation related genes in health control in asthma patient 1 (Ast1). C. Expression dynamics two cell-fate and six inflammation related genes in health control in asthma patient 2 (Ast2). D. Expression dynamics two cell-fate and six inflammation related genes in health control in asthma patient 3 (Ast3).

658   **Figure 7.** The number of cell-cell interactions and interaction strength for four sample

659   groups.

Figure 8. The annexin and resistin signaling pathway network of four sample groups. The annexin signaling pathway network is shown for health control, asthma patient 1 (Ast1) and asthma patient 3 (Ast3). The resistin signaling pathway network is shown for health control, asthma patient 1 (Ast1) and asthma patient 2 (Ast2).

Figure 9. The dual cell identity of the plasma blast cells in asthma patient 3. A. The expression heatmap of immunoglobulin (IGLV or IGHV) genes and T cell receptor (TRBV or TRAV) genes in the plasma blast cells of four sample groups. B. The violin plot of CD38, CD19, CD27, CD8A, CD40LG, GNLY and NKG7 gene expression levels across CD4+ T cell, CD8+ T cell, NK cell and plasma blast cells among four sample groups. CD38 and CD19 for the markers for plasma blast cells. CD27 is the marker for both plasma blast cells and T cells. CD8A, CD40LG, GNLY and NKG7 are the markers for T cells.

Figure 10. The expression of RETN gene in four sample groups. The level of serum

resistin between asthma patients and health controls($***p < 0.001$).

Supplemental Figures

Supplemental Figure 1. Integrated single-cell profiling of PBMCs for four sample

groups. A. The predicted cell types for the integrated single-cell profiling of PBMCs

for four sample groups B. Expression of canonical gene markers for each cell type in

four sample groups based on integration analysis. C. The detailed classification and

proportion of T cells for all seven samples in our study.

Supplemental Figure 2. Pseudo-time analysis of all cells in four sample groups. A.

The proportion of major cell types in each state for healthy controls. B. The

proportion of major cell type in each state for asthma patient 1 (Ast1). C. The

proportion of major cell type in each state for asthma patient 2 (Ast2). D. The

proportion of major cell type in each state for asthma patient 3 (Ast3).

Supplemental Figure 3. The number of incoming and outgoing interactions for each

cell type in four sample groups. A. The number of incoming and outgoing interactions

in health control. B. The number of incoming and outgoing interactions in asthma

patient 1 (Ast1). C. The number of incoming and outgoing interactions in asthma

patient 2 (Ast2). D. The number of incoming and outgoing interactions in asthma

patient 3 (Ast3).

Supplemental Figure 4. The number of incoming and outgoing interaction strength for each cell type in four sample groups. A. The number of incoming and outgoing interaction strength in health control. B. The number of incoming and outgoing interaction strength in asthma patient 1 (Ast1). C. The number of incoming and outgoing interaction strength in asthma patient 2 (Ast2). D. The number of incoming and outgoing interaction strength in asthma patient 3 (Ast3).

Supplemental Figure 5. The signaling changes of DCs in three asthma patients

(health control as the backgroup). A. The signaling changes of DCs in asthma patient

1 (Ast1). B. The signaling changes of DCs in asthma patient 2 (Ast2). C. The

signaling changes of DCs in asthma patient 3 (Ast3).

Supplemental Figure 6. The violin plot of ANXA1, ANXA2 and ANXA2 gene

expression levels across different cell types for four sample groups.

A Up-regulated signaling in Asthma1

B Up-regulated signaling in Asthma2

C Up-regulated signaling in Asthma3

711 **Supplemental Figure 7.** The upregulated signaling from DCs to the other cell types

712 in three asthma patients. A. The upregulated signaling from DCs to the other cell

713 types in asthma patient 1 (Ast1). B. The upregulated signaling from DCs to the other

714 cell types in asthma patient 2 (Ast2). C. The upregulated signaling from DCs to the

715 other cell types in asthma patient 3 (Ast3).

A

B

C

718   **Supplemental Figure 8.** The downregulated signaling from DCs to the other cell

719   types in three asthma patients. A. The downregulated signaling from DCs to the other

720   cell types in asthma patient 1 (Ast1). B. The downregulated signaling from DCs to the

721   other cell types in asthma patient 2 (Ast2). C. The downregulated signaling from DCs

722   to the other cell types in asthma patient 3 (Ast3).