Recapitulation of Cellular Senescence, Inflammation, and Fibrosis in Human Kidney-

Derived Tubuloids by Repeated Cisplatin Treatment: A Novel Pathophysiological Model

for Sensing Nephrotoxicity and Drug Screening

Yuki Nakao[1#], Makiko Mori[1#], Yuta Sekiguchi[1], Iori Morita[1], Ryota Shindoh[1], Shintaro Mandai[1],

Tamami Fujiki[1], Hiroaki Kikuchi[1], Fumiaki Ando[1], Koichiro Susa[1], Takayasu Mori[1], Yuma

Waseda[2], Soichiro Yoshida[2], Yasuhisa Fujii[2], Eisei Sohara[1], Shinichi Uchida[1], Yutaro Mori[1*]

[1]Department of Nephrology and [2]Department of Urology, Graduate School of Medical and

Dental Sciences, Institute of Science Tokyo, Tokyo 113-8510, Japan.

[#]These authors contributed equally to this work.

Lead contact and Corresponding author:

Yutaro Mori, MD, PhD

Department of Nephrology

Graduate School of Medical and Dental Sciences

Institute of Science Tokyo

1-5-45, Yushima, Bunkyo-ku, Tokyo 113-8510, Japan

Phone: +81-3-5803-5214; Fax: +81-3-5803-5215; Email: y-mori.kid@tmd.ac.jp

Abstract

In the pursuit of pathophysiological models for assessing renal drug response, the development

of kidney organoids derived from human pluripotent stem cells represents a significant step

forward. However, recapitulating aging/senescence-associated pathophysiology remains

challenging. Here, we present an innovative approach to generate epithelial-like structures

known as "tubuloid" using primary human renal proximal tubular epithelial cells (hRPTECs)

cultured from human resected kidneys, as a refined alternative. We evaluated the efficacy of

tubuloids using cisplatin treatment. Tubuloids showed highly differentiated structures. Exposure

to cisplatin increased γH2AX, Kidney Injury Molecule-1 (KIM-1) and Cleaved Caspase-3,

markers for DNA damage response, epithelial damage and apoptosis respectively. Repeated

cisplatin administration resulted in upregulation of the cellular senescence marker p16, p21 and

SA-β-Gal. Additionally, increased secretion of inflammatory cytokines, indicating the induction

of a senescence-associated secretory phenotype (SASP) were induced Supernatant collected from

cisplatin-treated tubuloids induced myofibroblast activation, indicating the onset of renal fibrosis.

We successfully established a tubuloid-based model of cisplatin-induced kidney injury using

hRPTECs. Tubuloids can replicate cellular senescence, SASP, and fibrosis, which can

recapitulate the phenotypes of chronic kidney disease (CKD). Furthermore, tubuloids provide a

novel platform for studying the response of renal epithelial cells to toxins and therapeutics and

offer innovative strategy for drug screening in a human-based fashion.

Key words

organoid, cisplatin nephrotoxicity, fibrosis, chronic kidney disease (CKD)

Translational Statement

Recapitulating aging/senescence-associated pathophysiological reaction in kidney organoids

remains challenging. Our study reveals that tubuloids could be novel candidate for chronic

kidney disease (CKD) model.

Introduction

Chronic kidney disease (CKD) is an irreversible, progressive condition characterized by the gradual sclerosis of glomeruli, accompanied by tubular atrophy, interstitial fibrosis, and infiltration of inflammatory cells [1]. Despite its significant impact on global health, the underlying pathogenic mechanisms of CKD are poorly understood, and a definitive cure has yet to be discovered.

In recent years, a shared mechanism for inflammation and fibrosis in CKD has emerged. It is proposed that the initial trigger is DNA damage in tubular epithelial cells, which initiates a cascade involving DNA damage response, cell cycle arrest [2], cellular senescence [3, 4], and the subsequent development of the senescence-associated secretory phenotype (SASP) [5]. To investigate these processes, various models have been created, including cell and animal models, as well as kidney organoids derived from human induced pluripotent stem cells (iPSCs) [6] or human embryonic stem cells (ESCs). These models show promise in understanding the pathogenesis of CKD and investigating potential therapeutic interventions [7]. Nonetheless, one critical limitation of existing models is their inability to fully capture the aging process, particularly the phenomenon of cellular senescence, which manifests prominently in humans with a lifespan greater than 80 years. Experimental animals such as mice, with a lifespan of only two years, do not fully replicate the range of age-related diseases associated with cellular senescence found in humans.

Furthermore, despite progress, kidney organoids derived from iPSCs or ESCs have primarily focused on replicating fetal kidney development stages, falling far short of accurately mimicking age-related diseases such as CKD.

In this context, "tubuloids," a three-dimensional structure resembling renal tubules, were

first reported in 2019. These tubuloids come from adult primary human renal proximal tubular

epithelial cells (hRPTECs) [8]. Schutgens et al. pioneered the culture of primary hRPTECs

derived from adult human kidneys and urine, successfully producing tubuloids capable of

replicating viral infections in proximal tubules and hereditary diseases [9]. Building on this

work, we have further refined the tubuloid generation protocol with human resected kidneys

[10]. Our research has focused on understanding the response of tubuloids to a nephrotoxicant

and a potential therapeutic agent. Despite these advances, developing a comprehensive

pathophysiological model for aging or senescence-associated diseases using tubuloids remains a

promising area of research.

Cisplatin, a commonly used therapeutic agent for various malignancies, including lung

cancer, has a well-known side effect known as cisplatin-induced nephropathy [11, 12]. This

condition is defined by the induction of DNA damage in the renal proximal tubules [13]. The

cellular response to this damage includes cell cycle arrest during the repair phase following

cisplatin-induced injury, which eventually leads to cellular senescence and SASP. These

processes are thought to be the final detrimental mechanisms that underpin the progression from

acute kidney injury (AKI) to CKD [14, 15]. Thus, modeling cisplatin-induced nephropathy is not

only useful for studying nephrotoxicity but also has the potential to mimic the progression of

CKD.

The primary goal of this study was to assess the disease-modeling potential of tubuloids

by exposing them to cisplatin to reproduce cisplatin-induced nephropathy. Furthermore, we

wanted to investigate whether tubuloids could replicate cellular senescence and its associated

responses beyond the immediate effects of cisplatin nephrotoxicity to simulate the transition

from AKI to CKD, or even model CKD itself.

Methods

Cell culture experiment: Human kidney samples were collected from patients undergoing

clinically indicated nephrectomy at the Tokyo Medical and Dental University Hospital in Tokyo,

Japan. The protocol was approved by the Institutional Review Board of the Ethics Committee of

Tokyo Medical and Dental University (M2022-005). Human renal proximal tubular epithelial

cells (hRPTECs) were isolated from unaffected areas of kidneys removed during nephrectomy

procedures performed on patients with renal cell carcinoma or urinary tract malignances by

modifying a previously established protocol[16]. Briefly, the isolation procedure involved

mincing the human renal cortex and then digesting it in a solution of Collagenase type II (1.0

mg/mL) (Worthington Biochemical, NJ, USA). The enzyme reaction was terminated with fetal

bovine serum (FBS), and the samples were resuspended in hRPTECs culture medium containing

DMEM/F-12 (Nacalai Tesque, Kyoto, Japan), with BSA (Nacalai Tesque), Antibiotic–

Antimycotic (ThermoFisher Scientific, MA, USA), hydrocortisone (ThermoFisher Scientific,

MA, USA), ITS liquid media supplement (Sigma-Aldrich, MO, USA), and human recombinant

epidermal growth factor (EGF) (ThermoFisher Scientific, MA, USA). The epithelial cells were

cultured for 7-10 days before being used in experiments.

To make mouse primary kidney fibroblasts (MPKFs), a 12-week old male C57BL/6J

mouse was sacrificed. MPKFs were isolated by mincing the mouse kidneys and digesting them

in a solution of Collagenase type II (1.0 mg/mL). The enzyme reaction was stopped with FBS,

and the samples were resuspended in DMEM with 10% FBS, 5 ng/ml FGF-basic (ThermoFisher

Scientific), and Antibiotic-Antimycotic. The MPKFs were used after 3 to 5 passages. All cells

were maintained in a CO_2 incubator (5% CO_2) at 37°C to ensure optimal growth conditions. All

animal studies were performed in accordance with the guidelines for animal research of Tokyo

Medical and Dental University. The study protocol was approved by the Animal Care and Use

Committee of Tokyo Medical and Dental University (approval no. A2023-109C).

Human Renal Tubuloids Culture: hRPTECs were seeded onto ultralow attachment plates at a

density of 5.0×10^5 cells/well using Advanced RPMI 1640 medium (ThermoFisher Scientific),

supplemented with 5% FBS. After a 2-day incubation period, Matrigel (Corning, NY, USA) was

added to stimulate tubuloid formation. The following day, the culture medium was changed to

Advanced RPMI 1640 medium with 5% FBS, EGF, FGF2, and HGF (tubuloids media). Media

was changed once or twice weekly to maintain optimal cell growth conditions. Tubuloids were

usually ready for experimentation within 1–2 weeks of starting the culture.

Fibrosis bioassay: The tubuloids were treated with or without cisplatin on a repeated basis.

Following a 48-hour exposure period, the media was replaced with serum-free DMEM to

facilitate conditioning. After an additional 48 hours of incubation, the culture supernatants were

collected as conditioned media.

MPKFs were uniformly plated at a density of 2.0×10^4 cells/well in an 8-well chamber

slide (Nunc Lab-Tek Chamber Slide system) (ThermoFisher Scientific) with 10% FBS-DMEM.

The next day, the conditioned media harvested from the tubuloids was added to MPKFs and

incubated for 48 hours. The MPKFs on the 8-well chamber slide were then washed twice with

PBS and immunostained as described below.

Immunofluorescence staining: The tubuloids were fixed with 4% paraformaldehyde (PFA) in

PBS and permeabilized with 0.1% Triton X-100 in PBS. The tubuloids were then blocked with

3% BSA in PBS and incubated with primary antibodies for 1 hour. Tubuloids were washed

thoroughly with PBS and then incubated with secondary antibodies for 30 minutes. After another

round of PBS washing, the tubuloids were placed on glass slides. The slides were mounted with

Prolong Glass Antifade Mountant with NucBlue Stain (ThermoFisher Scientific) and cover-

slipped for fluorescence microscopy.

The following primary antibodies were used: rabbit anti-LRP2 / Megalin antibody

(1:100) (ab236244, Abcam, MA, USA), rabbit anti-γH2AX antibody (1:200) (#9718, Cell

Signaling Technology, MA, USA), mouse anti-Na$^+$/K$^+$-ATPase antibody (1:200) (05-369,

Sigma-Aldrich), goat anti-KIM-1 antibody (1:200) (AF1750, R&D systems, MN, USA), rabbit

anti-caspase-3 antibody (1:200) (#9664, Cell Signaling Technology), rabbit anti-Vimentin

antibody (1:200) (10366-1-AP, Proteintech, IL, USA), rabbit anti-p16-INK4A polyclonal

antibody (1:200) (10883-1-AP, Proteintech), rabbit anti-IL-1β polyclonal antibody (1:200) (bs-

0812R, Bioss, MA, USA), rabbit anti-Collagen type 1 antibody, rabbit anti-αSMA antibody

conjugated with Cy3 (1:400) (C6198, Sigma-Aldrich) and rabbit anti-collagen type 1 antibody

(1:250) (600-406-103, rockland) were used. To stain the brush border of the proximal tubular

epithelium, LTL, Biotinylated (B-1325-2, Vector Laboratories, CA, USA) was used instead of

primary antibodies (1:200). Streptavidin, Alexa Fluor 633 conjugate (S21375, ThermoFisher

Scientific) was then applied (1:500).

For the immunofluorescence staining of fibrosis bioassay, MPKFs plated on an 8-well

chamber slide were washed before being fixed with 4% PFA-PBS. The cells were permeabilized

with 0.1% Triton X-100 in PBS and then blocked with 3% BSA.

All images were captured using either standard or confocal microscopy (Eclipse Ti2 from

Nikon), with the NIS-Elements Advanced Research microscope system serving as the operating

system. Quantification of stained area was done using ImageJ Fiji

(https://imagej.net/software/fiji/downloads).

Quantitative reverse transcription-polymerase chain reaction (qRT-PCR): The qRT-PCR

analysis of human IL-1β, IL-6, and 18S mRNA followed established protocols. The real-time

PCR detection system utilized was the Thermal Cycler Dice Real Time System Lite TP700

(Takara Bio, Shiga, Japan) along with TB green Premix Ex Taq II (Takara Bio, Shiga, Japan)

[17]. The total RNA was extracted from frozen kidney tubuloids stored at −80°C using Sepasol-

RNAI Super G (Nacalai Tesque, Kyoto, Japan). cDNA was then synthesized from the isolated

total RNA using ReverTra Ace (TOYOBO, Tokyo, Japan) [18]. The generated cDNA was then

amplified through 40 PCR cycles, with denaturation at 95°C for 5 seconds and

annealing/extension at 60°C for 30 seconds.

The nucleotide sequences of the primers utilized for qRT-PCR analysis are as follows.

human 18S.

Forward:5′-GCAGAATCCACGCCAGTACAAG-3′

Reverse:5′-GCTTGTTGTCCAGACCATTGGC-3′

human IL-1β.

Forward: 5′- CCACAGACCTTCCAGGAGAATG -3′

Reverse: 5′- GTGCAGTTCAGTGATCGTACAGG -3′

human IL-6.

Forward 5′- AGACAGCCACTCACCTCTTCAG -3′

Reverse: 5′- TTCTGCCAGTGCCTCTTTGCTG -3′

Western blot analysis: The tubuloids were lysed, and the protein was purified. Protein bands

were visualized using Western Blue (Promega, WI, USA). Rabbit anti–ERK2 (p44/42 MAPK)

antibody (9102S, Cell Signaling Technology) (1:200) was used as a loading control. Primary

antibodies were also used, such as rabbit anti-p16-INK4A polyclonal antibody (1:200)

(Proteintech), mouse anti-p21 antibody (1:100) (sc-6247, Santa Cruz Biotechnology, TX, USA),

rabbit anti-KIM-1 antibody (1:400) (PA5-20244, Thermo Fisher Scientific, MA, USA), rabbit

anti-caspase-3 antibody (1:200) (Cell Signaling Technology) and rabbit anti-Vimentin antibody

(1:1000) (#5741, Cell Signaling Technology, MA, USA). Densitometry was performed by using

Image J Fiji (https://imagej.net/software/fiji/downloads).

Senescence-associated-β-galactosidase (SA-β-GAL) staining: The tubuloids were stained by

using Cellular Senescence Detection Kit (SA-β-Gal Staining) (CBL, CA, USA) according to

directions of the manufacturer. Four wells of organoids cultured in a 24-well plate were used for

each experimental condition, and images were captured to encompass all areas of the wells.

Flowcytometry: Tubuloids were digested in using 2.5g/l-Trypsin/1mmol/l-EDTA Solution, with

Phenol Red (Nacalai Tesque Inc., Kyoto, Japan) at 37 degree for 20 minutes. Single cells

obtained from tubuloids were fixed with 4% PFA-PBS with 5% FBS, permeabilized with 0.1%

Triton X-100-PBS, and stained with primary antibodies for 1 hour and with secondary antibodies

for 45 minutes. Rabbit anti-LRP2 / Megalin antibody (1:250) (Abcam) and rabbit anti-caspase-3

antibody (1:250) (Cell Signaling Technology) were used for primary antibodies. Cells were

analyzed by BD FACSLyric (BD Biosciences, MA, USA). Data were analyzed by FlowJo (BD

Biosciences, MA, USA).

Quantification and Statistical Analysis: Figure Legends specify the number of samples

assayed in each experiment. To determine significant differences between groups, a one-way

analysis of variance was used. A p-value of less than 0.05 was considered statistically significant.

All statistical analyses were performed with GraphPad Prism by GraphPad Software Inc. (San

Diego, CA, USA).

Results

Primary human renal proximal tubular epithelial cells (hRPTECs) form highly polarized

structure

To form tubuloids, human renal proximal tubular epithelial cells (hRPTECs) were isolated from

the cortex of human kidneys, embedded in a basement membrane gel, and cultured with various

growth factors **(Figure 1A)**. Using precise techniques that included finely mincing the cortex

portion of the human kidneys and selective culturing of epithelial cells in a serum-free medium

supplemented with epidermal growth factor (EGF), primary cultured cells, primarily proximal

tubular epithelial cells, were successfully established **(Figure 1B and 1C**).

Tubuloids were then formed over approximately a two-week protocol **(Figure 1D).**

Specifically, hRPTECs were seeded on an ultralow attachment plate **(Figure 1E),** followed by

the addition of Matrigel and media containing fetal bovine serum, EGF, fibroblast growth factor

2 (FGF2), and hepatocyte growth factor (HGF). Multiple tubuloids were formed in a single well,

with nearly uniform sizes. The epithelial cells formed tubular-like structures that encircled the

basement membrane gel **(Figure 1F)**. The three-dimensional architecture was confirmed by

manipulating the microscope along the Z-axis direction **(Figure 1G)**. Notably, the cells formed a

monolayer structure similar to tubules or renal cysts, rather than a spheroid-like structure.

Furthermore, tubuloids expressed differentiation markers associated with proximal tubular

epithelial cells, such as lotus tetragonolobus lectin (LTL) and LDL Receptor Related Protein 2

(LRP2/Megalin), confirming their highly differentiated and polarized structure **(Figure 1H)**[19].

Cisplatin Exposure Induces Tubular Structure Collapse and Elicits Acute DNA Damage

Response (DDR) in Tubuloids

To test the sensitivity of tubuloids against a nephrotoxicant, we exposed them to cisplatin.

Cisplatin causes cytotoxicity by entering proximal tubular epithelial cells primarily through

Organic Cation Transporter 2 (OCT2) [20], resulting in direct DNA damage **(Figure 2A)**.

Consistent with previous reports highlighting the activation of DNA damage response (DDR)

pathways, we found significant upregulation of γH2AX, a marker of DNA double-strand breaks,

in both *in vivo* and *in vitro* [21]. Initially, tubuloids were treated with cisplatin at three

concentrations of 0.2, 2.0, and 20.0 µg/mL to determine dose-dependent responses. Notably,

high-dose cisplatin treatment collapsed the three-dimensional tubuloid architecture, **(Figure 2B)**,

demonstrating the extent of DNA damage caused by cisplatin exposure **(Figure 2C-E).**

Furthermore, immunostaining analyses revealed a significant reduction in the expression levels

of differentiation markers such as LRP2 and Na+/K+-ATPase after cisplatin treatment. This

downregulation indicates a concomitant process of cellular dedifferentiation, possibly as an

adaptive response to cytotoxic insult **(Figures 2F-K).**

Additionally, there was an increase in the expression of a renal injury marker, Kidney

Injury Molecule-1 (KIM-1) [19], and an apoptosis marker, Cleaved Caspase-3 **(Figure 2L-Q)**,

which was consistent with previous findings [20]. These findings further support the sensitivity

of tubuloids to cisplatin-induced injury and validate their use as a model system for studying

acute responses, including DDR.

Interestingly, we found that high concentrations of cisplatin increased the expression of

Vimentin, an intermediate filament specific to mesenchymal cells **(Figure 2R-T).** This suggests

that proximal tubular epithelial cells may undergo epithelial-mesenchymal transition (EMT) as

part of a tissue repair mechanism [22, 23]. The activation of Vimentin following DDR has been

reported previously [24], and our study confirms this sequential response in cisplatin-treated

tubuloids, elucidating the molecular dynamics underlying cisplatin-induced nephrotoxicity.

Tubuloids show cellular senescence and release inflammatory cytokines following repeated

cisplatin treatment

To assess the potential for replicating chronic responses, tubuloids were subjected to prolonged

treatment of cisplatin. Initially, cisplatin was administered at concentrations of 0.2 and 2.0

µg/mL. The media was then replaced three times to reduce residual cisplatin, and standard

culture conditions were applied for one day, completing one cycle. This was considered one

cycle, and the next day, cisplatin was given again at similar concentrations **(Figure 3A)**. The

cycle was repeated five times. Although treatment with 20.0 µg/mL cisplatin caused significant

disruption of tubuloids and cell death **(Figure 2)**, this concentration was excluded from the

experimental setup to focus on long-term chronic responses.

In the repeated cisplatin administration model, we observed the expression of p16, p21

and SA-β Gal, which are well-established **(Figure 3B-G)** and consistent with previous findings

[25-27]. Additionally, we confirmed an increased expression of the inflammatory cytokine IL-1β

at a concentration of 2.0 µg/mL through immunofluorescence analyses **(Figure 3H)**.

Quantitative PCR analysis revealed a significant upregulation in mRNA levels of both IL-1β and

IL-6, ranging from 50 to 100-fold compared to controls, in the group subjected to repeated

administration of 2.0 µg/mL cisplatin (*P = 0.0167*; IL-1β and *P = 0.011*; IL-6) **(Figure 3I and**

J). These findings reveal that repetitive DNA damage induced by cisplatin caused cellular

senescence, which resulted in the acquisition of an SASP.

Secretions released from cisplatin-treated tubuloids strongly promote myofibroblast

differentiation *in vitro*

We hypothesized that prolonged exposure of tubuloids to high concentrations of cisplatin would

promote myofibroblast activation, potentially contributing to tissue fibrosis via the secretion of

paracrine factors from tubuloids, as previously reported *in vivo* [28]. To test this hypothesis, we

used a fibrosis bioassay, as described previously [10] **(Figure 4A)**. Initially, tubuloids were

divided into three groups: one receiving no cisplatin, one receiving 0.2 µg/mL cisplatin, and

another receiving 2.0 µg/mL cisplatin. Each group received repeated treatments. Following

cisplatin administrations, the medium was changed three times to remove residual cisplatin, and

the tubuloids were cultured for an additional 2 days in serum-free conditions to produce

conditioned media (CM) containing secreted paracrine factors. Subsequently, the collected CM

was administered to fibroblasts derived from the mouse renal cortex **(Figure 4B)**. The activation

of myofibroblasts was assessed by evaluating the expression of α-smooth muscle actin (α-SMA)

and Collagen type I[29].

294   CM obtained from tubuloids repeatedly treated with 2.0 µg/mL cisplatin showed the most

significant promotion of fibrosis **(Figure 4C-F).** This concentration of cisplatin, as shown in

Figure 3, corresponds to the maximal secretion levels of IL-1β and IL-6 from tubuloids,

indicating consistency. Furthermore, direct administration of cisplatin to mouse fibroblasts did

not result in myofibroblast activation, decisively ruling out cisplatin as a direct fibrosis inducer

on fibroblasts. The fibrosis bioassay confirmed that various paracrine factors secreted from

tubuloids during repeated treatment of cisplatin have a fibrogenic effect on tissues.

Discussion

In this study, we succeeded to recapitulate almost all the maladaptive responses of renal

epithelial cells against cisplatin-induced toxicity by patient-derived tubuloids, not animal models

or iPS kidney organoids. By using multiple patients-derived tubuloid, our strategy has a potential

to use tubuloids as a semi-peresonalized toxicity or drug eficaciy platform. Furthermore, as we

succeeded to induce senescence markers, SA-β-GAL, p16 and p21 expression, and SASP, our

tubuloids can be not a drug toxicity model but also a new pathophysiological model for AKI to

CKD transition or CKD model itself.

310   Our findings provide compelling evidence of the negative effects of cisplatin on tubuloids,

elucidating the cascade of events initiated by DNA damage induction. The dose-dependent

increase in γH2AX expression is a reliable indicator of DNA damage, supporting previous

findings. Upon DNA damage, tubuloids exhibit a sensitive acute phase response as evidenced by

increased expression of the kidney injury marker KIM-1 and the apoptosis marker Cleaved

Caspase-3, which is consistent with DDR activation. There was also an increase in Vimentin

expression, which indicates epithelial-mesenchymal transition (EMT).

The well-preserved three-dimensional structure of tubuloids collapsed upon exposure to

cisplatin, accompanied by a process of dedifferentiation marked by reduced expression levels of

Na^+/K^+-ATPase and LRP2/Megalin. Furthermore, repeated long-term administration of cisplatin-

increased the expression of the cellular senescence markers, SA-β-GAL, p16 and p21. Tubuloids

developed a senescence-associated secretory phenotype (SASP), releasing inflammatory

cytokines like IL-1β and IL-6, causing a chronic inflammatory response in surrounding tissues.

These secretions from tubuloids significantly increased fibrosis in the surrounding interstitial

area, as confirmed by the fibrosis bioassay.

CKD research is difficult due to the complexity of the underlying conditions. A

significant impediment to research into CKD is the lack of a reliable disease model. CKD is

inextricably linked to aging and senescence, making it particularly difficult to model in

experimental settings. Currently, the primary models for studying CKD are mice, which, despite

their utility, live only about two years. This timeframe does not adequately represent the

decades-long progression of chronic diseases such as CKD in humans. Furthermore, mouse

models are a highly homogeneous population, lacking the diversity observed in human CKD

cases. Given the variability in CKD progression among humans, which is influenced by factors

such as gender, race, age, and environmental exposures, mouse models fail to capture the

complexities of human CKD adequately. This discrepancy reveals a significant gap in our

current understanding, emphasizing the need for more robust and clinically relevant models to

advance CKD research.

In recent years, kidney organoids derived from induced pluripotent stem cells (iPSCs) or

embryonic stem cells (ESCs) have emerged as promising disease models for the study of kidney

diseases [30-32]. However, current technologies primarily produce organoids that look like fetal

kidneys, making it difficult to replicate age-related features associated with CKD in older people

[33]. Furthermore, iPSC-derived kidney organoids require complex differentiation induction

procedures and skilled techniques, complicating their application. Primary cultured cells derived

from resected human kidneys and kept in two-dimensional conditions fail to fully replicate the

cellular polarity and three-dimensional structures found in living tissues. As a result, while these

models provide valuable insights, they have limitations that prevent them from accurately

representing the complexity of CKD pathogenesis and progression.

Conventional disease models for CKD, such as mouse models, iPSC/ESC kidney

organoids, and two-dimensional primary cultured cells from human kidneys, struggle to fully

replicate aging, senescence, and ease of production. However, the human kidney-derived

tubuloids developed in our study provide significant improvements over traditional pathological

models for CKD. Tubuloids are formed by cells obtained from patients that reflect their age and

may exhibit senescence, as evidenced by the control condition without cisplatin treatment

(Figure 3B, C, E and G). This unique feature addresses the limitation of other models by

incorporating the aging process directly into the system.

The formation of tubuloids follows a remarkably simple protocol, requiring no

specialized techniques. Culture is carried out in standard 24-well ultra-low attachement plates,

and as shown in **Figure 1F**, multiple tubuloids develop within each well. During sample

collection for western blotting, qPCR, and immunofluorescence, we harvested one sample per

well as an independent condition, resulting in multiple tubuloids in each sample, effectively

reducing variability across individual tubuloids.

Expanding our study to include tubuloids from multiple patients shows promise for

capturing individual differences in kidney responses to various stimuli. Notably, our tubuloid

production protocol allows us to derive these structures from different patients simultaneously,

under uniform conditions (manuscript in preparation). These characteristics point to tubuloids'

potential as a novel and useful screening tool for renal toxicity in drug discovery efforts.

Tubuloids have a highly differentiated three-dimensional structure[34] and mimic

physiological responses to nephrotoxicants[35, 36]. Previous research has shown that these well-

differentiated tubuloids express a diverse set of channels and transporters, providing them with

greater physiological functionality than cells cultured in two dimensions [37]. This increased

physiological fidelity is likely to explain the observed sensitivity to cisplatin in our study,

facilitating the replication of cellular senescence and other key responses. Importantly, tubuloids

may capture biological reactions that are difficult to detect in conventional two-dimensional cell

cultures.

The primary focus of this study was cisplatin-induced nephropathy. However, our

overarching goal goes beyond this single study to develop a strong pathophysiological model for

CKD using human kidney-derived tubuloids. The precise mechanisms underlying CKD

following cisplatin chemotherapy remain largely unknown. However, even low-dose cisplatin

has been shown to cause long-term renal pathologies similar to CKD [38]. Cisplatin-induced

nephropathy is caused by the uptake of cisplatin into proximal tubular epithelial cells via OCT2,

resulting in direct damage to nuclear DNA. This DNA damage initiates a cascade of DNA

damage response (DDR) processes: some cells die via programmed cell death, such as apoptosis,

while others activate the cell cycle in an attempt to compensate for the loss of damaged cells.

When DNA damage reaches a critical threshold, it causes cell cycle arrest, particularly in the

G2/M phase[2], preventing further division, a sign of cellular senescence.

Senescent cells have been shown to activate the SASP, resulting in the secretion of a

variety of inflammatory and fibrotic cytokines. This cascade causes inflammation and fibrosis in

the surrounding tissue while remaining resistant to cell death. This mechanism, which is driven

by cellular senescence and SASP induced by DNA damage, is a significant contributor to renal

fibrosis. It is an important part of the transition from AKI to CKD, as well as the broader

pathophysiology of CKD, which goes beyond cisplatin-induced nephropathy.

In our tubuloid model, we found not only a highly differentiated three-dimensional

structure, but also pathophysiological responses to cisplatin, such as DDR, structural collapse,

dedifferentiation, cell death, epithelial-mesenchymal transition (EMT), cellular senescence,

SASP activation, and fibrotic changes. Our findings suggest that tubuloids have the potential to

serve as a precise disease model for cisplatin-induced nephropathy, faithfully replicating the

majority of pathological mechanisms. This underscores their commitment to developing a

comprehensive CKD model.

Moving forward, we plan to improve the sophistication of our disease models by

incorporating coculture systems and microfluidic technologies. Specifically, we intend to

coculture tubuloids with other relevant cell types, including vascular endothelial cells and

fibroblasts. Additionally, we plan to simulate physiological urine flow using microfluidic devices

to better mimic the *in vivo* renal microenvironment [39].

The successful development of a highly precise disease model for cisplatin-induced

nephropathy using human kidney-derived tubuloids represents a significant advance in kidney

disease research. Using tubuloids as a reliable kidney disease model has great potential for

improving our understanding of various kidney diseases. With their ability to faithfully replicate

cellular senescence, SASP, and fibrosis, tubuloids appear to be a promising tool for modeling

CKD, a condition defined by fibrosis as a final common pathological pathway.

In conclusion, our development of tubuloids as a precise disease model for cisplatin-induced

nephropathy is a significant step forward in kidney disease research, especially as a presice

pathophyisiological model based on patient-derived materials, not animal models or iPSC kidney

organoids. Through continued refinement and innovation, we anticipate that tubuloids will play

an important role in sensing drug toxicity in a semi-personalized fashion and advancing our

understanding of CKD.

Disclosure Statement

The authors declare that they have no conflicts of interest.

419
Acknowledgement

We would like to thank the study participants who kindly allowed to give us the pieces of their

resected kidneys.

Funding

This work was supported by Leading Initiative for Excellent Young Researchers (LEADER)

from Ministry of Education, Culture, Sports, Science and Technology (to Y.M.), Grant-in-Aid

for Research Activity Start-up (22K20881) and Grand-in-Aid for Scientific Research (B)

(24K03249) from Japan Society for the Promotion of Science (to Y.M.), Innovation Idea Contest

from Tokyo Medical and Dental University (TMDU) (in 2022 to Y.M. and in 2023 to Y.N.),

Next Generation Researcher Training Unit from TMDU (to Y.M.) and Priority Research Areas

Grant from TMDU (to Y.M.), Research Grant from Uehara Memorial Foundation (to Y.M.),

Research Grant (Lifestyle-related diseases) from MSD Life Science Foundation (to Y.M.),

Medical Research Grant from Takeda Science Foundation (to Y.M.), and Academic Support

from Bayer Yakuhin, Ltd. (to Y.M.).

Data availability statements

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Yutaro Mori (y-mori.kid@tmd.ac.jp).

Author contributions

Y.N., M.M., Y.S., I.O., R.S. and Y.M. performed the experiments, collected and analyzed data,

and wrote the manuscript. Y.M., M.M., Y.S., I.O., and R.S. established the hRPTECs and S.M.

helped the procedure. T.F., H.K., F.A., K.S., T.M., E.S., and S.U. supported the data analysis.

Y.W., S.Y. and Y.F. resected the patients' kidneys as standard treatment for malignant diseases.

Y.M. developed experimental strategy, supervised the project, and edited the manuscript. All

authors discussed the results and implications and commented on the manuscript.

Disclosure

The authors declare that they have no conflicts of interest.

Figure Legends

Figure 1. Formation of highly polarized tubuloid structures from primary human renal

proximal tubular epithelial cells (hRPTECs).

(A) Schematic overview of the establishment of tubuloids from hRPTECs isolated from adult

human kidney.

(B) Image of the human resected kidney utilized during the establishment of hRPTECs.

(C) Representative image showing hRPTECs. Scale bars: 50 µm.

(D) Two-week protocol for generating tubuloids from 2D-cultured hRPTECs.

(E) Representative image demonstrating the seeding of hRPTECs on an ultralow attachment

plate. Scale bars: 25 µm.

(F) Representative images of tubuloids at different magnifications ($\times$10, $\times$40). Scale bars: 100

µm, 25 µm.

(G) Representative image of tubuloids at $\times$40 magnification, showing multiple layers along the Z

axis. Scale bars: 25 µm.

(H) Immunostaining of makers for renal proximal tubular epithelial cells (LRP2/Megalin and

LTL). Scale bars: 25 µm.

Figure 2. Effects of cisplatin treatment on tubuloid structure, DNA damage response,

differentiation loss, cell death, and epithelial-mesenchymal transition.

(A) Schematic representation of the pathway by which cisplatin induces DNA damage in

proximal tubular epithelial cells.

(B) Representative images of tubuloids exposed to cisplatin (0.2, 2.0, 20.0 µg/mL) or control.

(C) Immunostaining of marker for DNA damage (γH2AX) in tuboloids exposed to cisplatin (0.2,

2.0, 20.0 µg/mL) or control. Scale bars: 100 µm.

(D) Western blotting of γH2AX in tuboloids exposed to cisplatin (0.2, 2.0, 20.0 µg/mL) or

control.

(E) Statistical analysis of γH2AX expression (n = 4).

(F) Immunostaining of a differentiation marker for epithelial cells, LRP2/Megalin in tuboloids

exposed to cisplatin (20.0 µg/mL) or control. Scale bars: 25 µm.

(G) Representative plots of flowcytometry measuring LRP2/Megalin expression in tuboloids

exposed to cisplatin (20.0 µg/mL) or control.

(H) Statistical analysis of LRP2/Megalin expression (n = 3).

(I) Immunostaining of a differentiation marker for epithelial cells, Na^+/K^+-ATPase in tuboloids

exposed to cisplatin (20.0 µg/mL) or control. Scale bars: 25 µm**.**

(J) Western blotting of Na^+/K^+-ATPase in tuboloids exposed to cisplatin (20.0 µg/mL) or control.

(K) Statistical analysis of LRP2/Megalin expression (n = 4).

(L) Immunostaining of an epithelial injury marker, Kidney Injury Molecule-1 (KIM-1) in

tuboloids exposed to cisplatin (20.0 µg/mL) or control. Scale bars: 25 µm.

(M) Western blotting of KIM-1 in tuboloids exposed to cisplatin (20.0 µg/mL) or control.

(N) Statistical analysis of KIM-1 expression (n = 4).

(O) Immunostaining of an apoptosis marker, Cleaved Caspase-3, in tuboloids exposed to

cisplatin (20.0 µg/mL) or control. Scale bars: 25 µm.

(P) Representative plots of flowcytometry measuring Cleaved Caspase-3 expression in tuboloids

exposed to cisplatin (20.0 µg/mL) or control.

(Q) Statistical analysis of Cleaved Caspase-3 expression (n = 4).

(R) Immunostaining of intermediate filament, Vimentin, in tubuloids exposed to cisplatin (20.0

µg/mL) or control. Scale bars: 100 µm.

(S) Western blotting of Vimentin in tubuloids exposed to cisplatin (20 µg/mL) or control.

(T) Statistical analysis of Vimentin expression (n = 4).

All data are presented as mean ± standard error (SE). Significance is presented by *p < 0.05, **p

< 0.01, ***p < 0.001.

Figure.3 Tubuloids exhibit cellular senescence and release inflammatory cytokines

following repeated cisplatin treatment.

(A) Experimental design for repeated cisplatin injury.

(B) Representative images of SA-β-GAL staining of tubuloids exposed to cisplatin (0.2 µg/mL)

5 times repeatedly or control. Scale bars: 100 µm.

(C) Western blotting of p21 in tubuloids exposed to cisplatin (0.2 µg/mL) 5 times repeatedly or

control.

(D) Statistical analysis of p21 expression (n = 4).

(E) Western blotting of p16 in tubuloids exposed to cisplatin (0.2, 2.0 µg/mL) 5 times repeatedly

or control.

(F) Statistical analysis of p16 expression (n = 4).

(G) Immunostaining of p16 in tubuloids exposed to cisplatin (2.0 µg/mL) 5 times repeatedly or

control. Scale bars: 100 µm.

(H) Immunostaining of an inflammatory cytokine, IL-1β, in tubuloids exposed to cisplatin (2.0

µg/mL) 5 times repeatedly or control. Scale bars: 100 µm.

(I, J) Quantitative PCR of IL-1β **(I)** and IL-6 **(J)** in tubuloids exposed to cisplatin (0.2, 2.0

µg/mL) 5 times repeatedly or control (n = 3). Statical analysis was applied to delta-delta Ct value,

not fold increase.

All data are presented as mean ± standard error (SE). Significance is presented by *p < 0.05, **p

< 0.01.

Figure 4. Secretion released from cisplatin-treated tubuloids strongly promotes fibrosis *in*

***vitro*.**

(A) Experimental design for fibrosis bioassay.

(B) A representative image of Mouse Primary Kidney Fibroblasts (MPKFs). Scale bar: 25 µm.

(C) Representative images of fibrosis bioassay stained with anti-α-SMA antibody on MPKFs

cultured with Conditioned Media (CM) harvested from tubuloids exposed to cisplatin (0.2, 2.0

µg/mL) or control, or directly with cisplatin-containing media. Scale bars: 100 µm.

(D) Quantification of αSMA-positive area in MPKFs cultured with CM or directly with cisplatin-

containing media measured by Image J (*n* = 8-10 in each condition).

(E) Representative images of fibrosis bioassay stained with anti-Collagen type I antibody on

MPKFs cultured with Conditioned Media (CM) harvested from tubuloids exposed to cisplatin

(0.2, 2.0 µg/mL) or control, or directly with cisplatin-containing media. Scale bars: 100 µm.

(F) Quantification of Collagen type I-positive area in MPKFs cultured with CM or directly with

cisplatin-containing media measured by Image J (*n* = 8-10 in each condition).

All data are presented as mean ± standard error (SE). Significance is presented by *p < 0.05, **p

< 0.01, ***p < 0.001, ****p < 0.0001.

References

1. Humphreys, B.D., *Mechanisms of Renal Fibrosis.* Annu Rev Physiol, 2018. **80**: p. 309-326.
2. Yang, L., et al., *Epithelial cell cycle arrest in G2/M mediates kidney fibrosis after injury.* Nat Med, 2010. **16**(5): p. 535-43, 1p following 143.
3. Sturmlechner, I., et al., *Cellular senescence in renal ageing and disease.* Nat Rev Nephrol, 2017. **13**(2): p. 77-89.
4. Docherty, M.H., et al., *Cellular Senescence in the Kidney.* J Am Soc Nephrol, 2019. **30**(5): p. 726-736.
5. Mylonas, K.J., et al., *Cellular senescence inhibits renal regeneration after injury in mice, with senolytic treatment promoting repair.* Sci Transl Med, 2021. **13**(594).
6. Takasato, M., et al., *Kidney organoids from human iPS cells contain multiple lineages and model human nephrogenesis.* Nature, 2015. **526**(7574): p. 564-8.
7. Liu, M., et al., *Studying Kidney Diseases Using Organoid Models.* Front Cell Dev Biol, 2022. **10**: p. 845401.
8. Schutgens, F., et al., *Tubuloids derived from human adult kidney and urine for personalized disease modeling.* Nat Biotechnol, 2019. **37**(3): p. 303-313.
9. Xu, Y., et al., *Adult human kidney organoids originate from CD24.* Nat Genet, 2022. **54**(11): p. 1690-1701.
10. Mori, Y., et al., *KIM-1 mediates fatty acid uptake by renal tubular cells to promote progressive diabetic kidney disease.* Cell Metab, 2021. **33**(5): p. 1042-1061 e7.
11. Latcha, S., et al., *Long-Term Renal Outcomes after Cisplatin Treatment.* Clin J Am Soc Nephrol, 2016. **11**(7): p. 1173-1179.
12. McSweeney, K.R., et al., *Mechanisms of Cisplatin-Induced Acute Kidney Injury: Pathological Mechanisms, Pharmacological Interventions, and Genetic Mitigations.* Cancers (Basel), 2021. **13**(7).
13. Miller, R.P., et al., *Mechanisms of Cisplatin nephrotoxicity.* Toxins (Basel), 2010. **2**(11): p. 2490-518.
14. Sato, Y., M. Takahashi, and M. Yanagita, *Pathophysiology of AKI to CKD progression.* Semin Nephrol, 2020. **40**(2): p. 206-215.
15. Ferenbach, D.A. and J.V. Bonventre, *Mechanisms of maladaptive repair after AKI leading to accelerated kidney ageing and CKD.* Nat Rev Nephrol, 2015. **11**(5): p. 264-76.
16. Ichimura, T., et al., *Kidney injury molecule-1 is a phosphatidylserine receptor that confers a phagocytic phenotype on epithelial cells.* J Clin Invest, 2008. **118**(5): p. 1657-68.
17. Cong, W., et al., *Implications of the Wnt5a/CaMKII pathway in retinoic acid-induced myogenic tongue abnormalities of developing mice.* Sci Rep, 2014. **4**: p. 6082.
18. Takai, J., et al., *heterozygous mutant mice exhibit reduced inflammatory responses and impaired bacterial clearance.* iScience, 2021. **24**(8): p. 102836.
19. Nielsen, R., E.I. Christensen, and H. Birn, *Megalin and cubilin in proximal tubule protein reabsorption: from experimental models to human disease.* Kidney Int, 2016. **89**(1): p. 58-67.
20. Mody, H., et al., *Pharmacodynamic Modeling to Evaluate the Impact of Cimetidine, an OCT2 Inhibitor, on the Anticancer Effects of Cisplatin.* Cells, 2022. **12**(1).

21. Kuo, L.J. and L.X. Yang, *Gamma-H2AX - a novel biomarker for DNA double-strand*
breaks. In Vivo, 2008. **22**(3): p. 305-9.
22. Lovisa, S., et al., *Epithelial-to-mesenchymal transition induces cell cycle arrest and*
parenchymal damage in renal fibrosis. Nat Med, 2015. **21**(9): p. 998-1009.
23. Grande, M.T., et al., *Snail1-induced partial epithelial-to-mesenchymal transition drives*
renal fibrosis in mice and can be targeted to reverse established disease. Nat Med, 2015.
21(9): p. 989-97.
24. Patteson, A.E., et al., *Vimentin protects cells against nuclear rupture and DNA damage*
during migration. J Cell Biol, 2019. **218**(12): p. 4079-4092.
25. Li, S., et al., *Tubular cell senescence promotes maladaptive kidney repair and chronic*
kidney disease after cisplatin nephrotoxicity. JCI Insight, 2023. **8**(8).
26. Mohamad Kamal, N.S., et al., *Aging of the cells: Insight into cellular senescence and*
detection Methods. Eur J Cell Biol, 2020. **99**(6): p. 151108.
27. Yan, J., et al., *The role of p21 in cellular senescence and aging-related diseases.* Mol
Cells, 2024. **47**(11): p. 100113.
28. Liu, B.C., et al., *Renal tubule injury: a driving force toward chronic kidney disease.*
Kidney Int, 2018. **93**(3): p. 568-579.
29. LeBleu, V.S., et al., *Origin and function of myofibroblasts in kidney fibrosis.* Nat Med,
2013. **19**(8): p. 1047-53.
30. Gupta, N., et al., *Modeling injury and repair in kidney organoids reveals that*
homologous recombination governs tubular intrinsic repair. Sci Transl Med, 2022.
14(634): p. eabj4772.
31. Hiratsuka, K., et al., *Organoid-on-a-chip model of human ARPKD reveals*
mechanosensing pathomechanisms for drug discovery. Sci Adv, 2022. **8**(38): p.
eabq0866.
32. Tabibzadeh, N., et al., *Navigating the kidney organoid: insights into assessment and*
enhancement of nephron function. Am J Physiol Renal Physiol, 2023. **325**(6): p. F695-
F706.
33. Yousef Yengej, F.A., et al., *Kidney Organoids and Tubuloids.* Cells, 2020. **9**(6).
34. Yousef Yengej, F.A., et al., *Tubuloid differentiation to model the human distal nephron*
and collecting duct in health and disease. Cell Rep, 2023. **43**(1): p. 113614.
35. Wiraja, C., et al., *Nephrotoxicity Assessment with Human Kidney Tubuloids using*
Spherical Nucleic Acid-Based mRNA Nanoflares. Nano Lett, 2021. **21**(13): p. 5850-5858.
36. Vidal Yucha, S.E., et al., *"3D, human renal proximal tubule (RPTEC-TERT1) organoids*
'tubuloids' for translatable evaluation of nephrotoxins in high-throughput". PLoS One,
2022. **17**(11): p. e0277937.
37. Yousef Yengej, F.A., et al., *Tubuloid differentiation to model the human distal nephron*
and collecting duct in health and disease. Cell Rep, 2024. **43**(1): p. 113614.
38. Fu, Y., et al., *Chronic effects of repeated low-dose cisplatin treatment in mouse kidneys*
and renal tubular cells. Am J Physiol Renal Physiol, 2019. **317**(6): p. F1582-F1592.
39. Gijzen, L., et al., *Culture and analysis of kidney tubuloids and perfused tubuloid cells-on-*
a-chip. Nat Protoc, 2021. **16**(4): p. 2023-2050.
628
629

Figure.1

Figure.2

Figure.3

Figure.4

Figure.5

DDR: DNA Damage Response
SASP: Senescence Associated Secretary Phenotype
EMT: Epithelial-Mesenchymal Transition

Figure.1

A

| Human Kidney | hRPTECs (human Renal Proximal Tubular Epithelial Cells) | Tubuloids |

1 week
Culture in serum-free media containing EGF

1~2 week expansion
Culture in media containing Matrigel with growth factors

B Human Kidney

C hRPTECs (×40)

D

Day 1 — hRPTECs on ultra-low attachment plate
Day 2 — + Matrigel
Day 3 — + Tubuloid Media
Change Culture Medium × 2~3
Day 14 — Forming Tubuloids

E hRPTECs on Ultra-low attachment Plate (×40)

F Tubuloids (×10) Tubuloid (×40)

G Tubuloid (×40)

Move in Z-axis direction

H Merge LRP2/Megalin LTL DAPI

Figure.2

Figure.3

Figure.4

Figure.5

DDR: DNA Damage Response

SASP: Senescence Associated Secretary Phenotype

EMT: Epithelial-Mesenchymal Transition