The impact of SARS-CoV-2 on fatigue and quality of life in the early phase of the pandemic: prevalence of post COVID-19 condition among unvaccinated individuals in a Dutch population-based serosurveillance cohort.

Cheyenne C.E. van Hagen[1]*, Elizabeth N. Mutubuki[1,2]*, Siméon de Bruijn[1], Eric R.A. Vos[1], Gerco den Hartog[1,3], Fiona R.M. van der Klis[1], Cees C. van den Wijngaard[1], Hester E. de Melker[1], Albert Jan van Hoek[1]

[1]National Institute for Public Health and the Environment (RIVM), Center for Infectious Disease Control, Bilthoven, the Netherlands

[2]National Institute for Public Health and the Environment (RIVM), Center for Health and Society, Bilthoven, the Netherlands

[3]Laboratory of Medical Immunology, Radboud UMC, Nijmegen, The Netherlands

* These authors contributed equally to this work.

Running title: Fatigue and quality of life after SARS-CoV-2 infection

Corresponding author

Cheyenne C.E. van Hagen, Centre for Infectious Disease Control Netherlands, National Institute for Public Health and the Environment (RIVM), Antonie van Leeuwenhoeklaan 9, 3721 MA Bilthoven, the Netherlands

E-mail: cheyenne.van.hagen@rivm.nl

Abstract

Objectives: We studied post-COVID-19 condition by investigating health-related quality of life and
fatigue in the general Dutch population in the early phase of the pandemic, including symptomatic
and asymptomatic infections among unvaccinated individuals.

Methods: (Still) unvaccinated participants aged ≥15 years were selected from the February 2021
round of the nationwide seroepidemiological PIENTER Corona cohort study. We assessed associations
between the time since serologically-identified SARS-CoV-2 infection and four outcome measures:
health utility (Short-Form 6 Dimensions), mental health and physical health (Short Form Health Survey
12) and fatigue (Checklist Individual Strength subscale fatigue). Per outcome, cutoff points were
selected at each 5% increment (5-75%) along the cumulative distribution of those uninfected. At each
cutoff, multivariable logistic regression models (score below cutoff yes/no) were fitted adjusted for
infection history, age, sex, education level, comorbidities, and restriction intensity.

Results: At the cutoff of the lowest 15th percentile among uninfected, significant differences between
uninfected (n=4,569) and infected ≤4 months ago (n=351) were observed for health utility (OR
[95%CI]: 1.6 [1.2-2.2]), physical health (1.9 [1.5-2.5]) and fatigue (1.6 [1.3-2.1]), but not for mental
health (1.2 [0.9-1.6]). There were no significant differences between uninfected and infected >4
months ago (n=327) for all outcomes at any cutoff of the cumulative distribution, with post-hoc
analysis showing a power to detect prevalence differences as low as 7%.

Conclusions: In the first year of the pandemic, data from this Dutch population-based
seroepidemiological cohort showed that unvaccinated individuals with a SARS-CoV-2 infection ≤4
months ago reported poorer health utility and physical health, and more severe fatigue compared to
those uninfected. Interestingly, for those infected >4 months ago differences remained below the
detection limit, suggesting a lower population prevalence of post-COVID-19 condition than currently
found in literature for this period.

Keywords: Health-related quality of life, mental health, physical health, post-covid-19 condition,
fatigue, SARS-CoV-2 infection, seroepidemiology

Background

Identifying and measuring the overall impact of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) on health-related quality of life (HRQoL) is crucial for informing public health strategies. During the pandemic, HRQoL of the general population was negatively affected as a result of both physical (e.g. infection or disease) and psychological (e.g. anxiety or social isolation) factors (1-3). With some groups more at risk of (severe) SARS-CoV-2 infection, different groups were disproportionately impacted (4). Clinical presentations of SARS-CoV-2 can range from no or mild symptoms to critical illness and death. In a proportion of patients, symptoms develop or remain after the acute phase resulting in long-term complaints, a condition referred to as Long COVID or Post-COVID-19 Condition (PCC) (5-8). PCC is defined by the World Health Organization as "the continuation or development of new symptoms 3 months after the initial SARS-CoV-2 infection, with these symptoms lasting for at least 2 months with no other explanation" (9). Symptoms of PCC include shortness of breath and cognitive dysfunction, with the most common symptom being fatigue (9, 10). Due to the long duration and/or detrimental impact of these symptoms, PCC may form a significant proportion of the overall disease burden of SARS-CoV-2 (11). Therefore, the impact of PCC on the population in terms of HRQoL is important to investigate.

The burden of PCC in the population depends on SARS-CoV-2 infection rates and the proportion of individuals developing PCC after infection. Although the prevalence of PCC has been studied in prospective cohorts and in cross-sectional population samples, where patients are identified by a positive SARS-CoV-2 PCR or lateral flow test (12, 13), discrepancies exist in PCC prevalence estimates (ranging from 9% to 81%) (10). These discrepencies predominantly arise due to variations in definitions of PCC, the population under investigation, the identification of (re-)infection/COVID-19, or other factors such as SARS-CoV-2 variant, time since infection or COVID-19 vaccination. Vaccination and previous infection contribute to immunological heterogeneity in current populations, making it difficult to establish PCC prevalence after more recent SARS-CoV-2 variants (14). To avoid vaccination effects on PCC development, studying early-pandemic SARS-CoV-2 infections, before the rollout of vaccination programs, is insightful and could function as a benchmark for more recent SARS-CoV-2 variants amidst mostly vaccinated populations. In addition, studies that rely on positive SARS-CoV-2 testing are prone to selection bias due to testing behavior and asymptomatic infections. Therefore, to establish the true prevalence of PCC and its impact on HRQoL on a population level, population-based serological studies are required because such studies provide a more comprehensive and unbiased assessment of the infection rates in the general population (15).

Although a population-based and/or serological approach was previously used to assess PCC in some

studies (16-19), these often involved a more heterogenous immunological landscape, included only

children or focused on a variety of PCC-related symptoms rather than looking at the HRQoL. To our

knowledge, there are no population-based serological studies investigating HRQoL after SARS-CoV-2

infection in the pre-vaccination era. Hence, using Dutch population-based serosurveillance data from

the first year of the COVID-19 pandemic, the aim of this study was to assess the relationship between

the time since wild-type SARS-CoV-2 infection and HRQoL and fatigue as endpoints linked to PCC

among unvaccinated individuals.

Materials and methods

Study design and population

The PIENTER Corona (PICO) study is a prospective population-based cohort study aimed at monitoring humoral immunity against SARS-CoV-2 in the Dutch population. The PICO study started in April 2020 (PICO1) when 3,244 participants aged 2-92 years were enrolled (20). These participants were sampled from the PIENTER3 serosurvey, an existing cohort established in 2016/2017 (21), for which participants had been randomly selected (using age strata) from the Dutch population registry. In the second round, in June 2020 (PICO2), the study sample was supplemented with an additional sample of 4,606 randomly-selected participants from the Dutch population registry in order to enhance national coverage and overall power (combined response rate, 21.4%) (22). Subsequent rounds were conducted in September 2020 (PICO3) and February 2021 (PICO4) using the same study sample as in PICO2 with exception to dropouts and non-responders (n=1,320) (Figure 1).

In every study round, participants were asked to return a finger-prick blood sample in a microtainer by mail and complete a questionnaire (online or on paper). The questionnaire collected information on demographics, experienced COVID-19 like symptoms (i.e., respiratory, gastro-intestinal, systemic, etc.), SARS-CoV-2 testing, COVID-19 vaccination, clinical risk factors for disease, contact patterns, HRQoL and fatigue (only from PICO4 onwards for those 15 years and older). Blood samples were tested for immunoglobulin G (IgG) antibodies against SARS-CoV-2 Spike-S1 and Nucleoprotein (N) proteins using a previously described validated fluorescent bead-based multiplex-immunoassay (23). Briefly, serum samples were centrifuged and stored at -20 degrees Celsius before analysis. The analysis process included dilution and incubation of samples with S1- (Sino Biological, 40591-V08H) and N- (Sino Biological, 40588-V08B) coupled beads in SM01-buffer (Surmodics, USA) supplemented with 2% FCS and shaking of samples for 45 minutes in the dark at room temperature. Subsequently, plates underwent three washes with PBS and were incubated for 30 minutes with PE-conjugated goat anti-human IgG (Jackson ImmunoResearch, 109-116-098) and washed again. Finally, samples were acquired on a Luminex LX200 or FlexMap3D. A 5-parameter logistic fit was used to interpolate antibody concentrations from pooled sera calibrated against the World Health Organization standard (NIBSC, 20/136). The cut-off for seropositivity was set at a Spike-S1-specific IgG antibody concentration of 10.08 binding antibody units per milliliter (BAU/mL) and a N-specific IgG antibody concentration of 14.3 BAU/mL, in accordance with international standards (22-25). This yielded a specificity and a sensitivity of 99.9% and 94.3% respectively for anti-S1 and 98.0% and 86.0% for anti-N (22, 24-26).

In this study we included all unvaccinated PICO4 participants with questionnaire data on HRQoL and

fatigue and at least one blood sample in PICO1-4. COVID-19 vaccination status was based on the Dutch

COVID-19 vaccination registry (CIMS) (27) and complemented with self-reported questionnaire data

if CIMS was unavailable. The PICO study protocol was approved by the Medical Ethics Committee MEC-

U, the Netherlands (Clinical Trial Registration NTR8473). All participants provided written informed

consent.

Time since SARS-CoV-2 infection

To assess SARS-CoV-2 infection, we distinguished between participants with serological proof of

previous SARS-CoV-2 infection (seropositives/cases) and those without (seronegatives/uninfected).

Proof of infection was defined as the presence of serum immunoglobulin G (IgG) antibodies against

the SARS-CoV-2 Spike-S1 antigen in rounds PICO1-4. Participants who were Spike-S1 seropositive at

least once were considered a case, irrespective of negative PCR/rapid antigen tests or serostatus in

subsequent rounds. To assess time since SARS-CoV-2 infection, cases were further classified according

to time since seroconversion; we distinguished participants that seroconverted in PICO1-3, i.e., before

or in September 2020 (cases >4 months) and participants that seroconverted in PICO4, i.e., after

September 2020 (cases ≤4 months). The uninfected group consisted of all participants who had not

been Spike-S1 seropositive through PICO1-4. Uninfected participants who missed serological data in

PICO4 or self-reported a positive PCR or rapid antigen test in one of the study rounds were excluded.

Thus, three SARS-CoV-2 serostatus groups were ultimately identified: uninfected, cases >4 month and

cases ≤4 months.

Outcome measures

HRQoL: Short Form Health Survey 12 (SF-12) and Short-Form Six Dimensions (SF-6D)

HRQoL was assessed in terms of mental health, physical health and health utility. To measure these,

the Dutch translation of the SF-12 version 1 was used. The SF-12 consists of 12 questions from eight

health dimensions that can be summarized into a physical health score (physical component summary;

PCS) and a mental health score (mental component summary; MCS). The PCS comprises the following

health dimensions; physical functioning, role limitations due to physical health problems, bodily pain

and general health, whereas the MCS consists of vitality (energy/fatigue), social functioning, role

limitations due to emotional problems and mental health (psychological distress and psychological

well-being). The summarized scores were weighted using orthogonal regression coefficients from the

general Dutch population (28). PCS and MCS scores range from 0 (lowest health) to 100 (highest

health) (29). To convert results of the SF-12 into health utility scores the SF-6D was used (30). The SF-

6D uses seven items from the SF-12 and consists of six dimensions: physical functioning, role

limitations, social functioning, pain, mental health and vitality (30). SF-6D scores range from 0.3 to 1,

with higher scores indicating better health (30, 31). In the online questionnaire, participants received

the SF-12 version 1 UK, while the paper-based questionnaire contained the SF-12 version 1 US. In the

SF-12 version 1 UK, item 12 (social functioning) has a six-scale answer instead of a five-scale answer

that is used in the SF-12 version 1 US. Participants who selected the additional answer option (i.e.

option 3 of the six-scale answer) in the SF-12 version 1 UK were equally randomized to a category

below and a category above in order to analyze all questionnaires in the SF-12 version 1 US format.

Fatigue: Checklist Individual Strength (CIS)

Fatigue severity was assessed using the fatigue subscale of the Checklist Individual Strength (CIS-

fatigue). The CIS-fatigue is an 8-item questionnaire validated to assess chronic fatigue (32, 33). Each

item is scored on a 7-point Likert scale. Scores can range from 8 to 56, with fatigue severity increasing

with higher scores (34).

Statistical analysis

Participant demographic (age, sex and educational level) and health characteristics (number of

comorbidities, health utility, physical health, mental health and fatigue) were presented using

proportions (%) and frequencies (n). Information collected in PICO4 were used, except for educational

level and number of comorbidities which were determined at baseline.

In the analyses we looked at four different outcomes: health utility (SF-6D), mental health (MCS),

physical health (PCS), and fatigue (CIS). To enable comparison between the four outcome variables

with different scales and score ranges, we opted for a uniform approach. In this approach, we

separately plotted the cumulative distribution of each outcome for all three SARS-CoV-2 serostatus

groups to visualize differences between these groups over the entire range of the outcome. Since

higher CIS scores indicate worse outcomes, the cumulative distribution of the CIS scale was inverted

for comparability. The cumulative distribution of individuals in the uninfected group guided the

selection of cut-off points. Cut-off points were determined at each 5% increment along the cumulative

distribution of the uninfected group (5% to 75%), and at each cut-off individuals were assigned 0 (score

above the cut-off) or 1 (score on or below the cut-off). For each of the cut-off points multivariable

logistic regression models were fitted separately, per outcome. The following confounders were

added to the multivariable logistic regression models: age group (18-35/36-65/66+ years), sex

(male/female), educational level (low/intermediate/high), number of comorbidities (none/one/two

or more) and Stringency Index (75/79). The Stringency Index measures the intensity of restrictions in

the Netherlands on a scale of 0 to 100 (strictest) as quantified by the COVID-19 Stringency Index of

the Oxford Coronavirus Government Response Tracker (OxCGRT) (35, 36). The Stringency Index was

dichotomized into score 75 and 79 because the scores remained steady during the inclusion period

(78.7 from 11 February to 2 March 2021 and 75.0 from 3 March to 6 April 2021). Complete case data

on all variables were used for the analyses. Adjusted odds ratios (OR) and 95% confidence intervals

(CI) were provided and a p-value of $p<0.05$ was considered statistically significant.

Post-hoc analyses were conducted to explore the power to detect a minimal difference in prevalence

between cases >4 months and those uninfected given the observed cases in our study. With those

uninfected as reference group, an iterative process was performed in which per outcome, the

prevalence of cases >4 months meeting the cut-off (assigned a 1) was increased until the CI of the OR

was above 1. In this process the cases >4 months were included at random. To account for

stochasticity, an average of ten independent samples was taken for each prevalence point, for each

cut-off point.

Data was cleaned in SAS (94 M7 English) and analyzed in R (version 4.1.0).

Results

Demographic and clinical characteristics of the participants

From a total of 6,530 participants, 1,155 participants did not meet inclusion criteria of which 294 (4.5%) were excluded because of being vaccinated or missing vaccination data (Figure 1). Of 5,375 eligible participants (supplementary file – Table 1), complete case analysis was conducted with 5,247 participants of which 4,569 were uninfected, 327 cases >4 months and 351 cases ≤4 months (Table 1), thus excluding 2% of participants (missings). All groups had a higher proportion of females compared to males (55-58%). The majority of the participants in all groups were aged between 36-65 years (48-52%) and had a high or intermediate educational level (79-82%). Among cases ≤4 months there were fewer participants with comorbidities (27%) compared to those uninfected and cases >4 months (respectively 34% and 35%). In all groups, mean SF-6D scores were 0.8. Mean MCS (46-48) and PCS scores (53-55) were also similar between groups. Mean CIS scores of those uninfected and cases >4 months were comparable (22), but those of cases ≤4 months deviated (25).

Cumulative distribution

For health utility, the cumulative distribution of SF-6D scores showed that cases ≤4 months had a higher proportion of individuals with a lower score, compared to those uninfected and cases >4 months (Figure 2, upper panel); among those uninfected and cases >4 months respectively 16% and 17% had an SF-6D score below 0.66, whereas for cases ≤4 months this was 22% (supplementary file – Table 2). Looking at mental health, cases ≤4 months had a higher proportion of individuals with lower MCS scores compared to those uninfected, which was also observed for cases >4 months across a smaller part of the cumulative distribution (Figure 3, upper panel); 15% of those uninfected scored below MCS 37, whereas for cases >4 months and cases ≤4 months this was respectively 18% and 19% (supplementary file – Table 3). For physical health, cases ≤4 months also had a higher proportion of individuals with lower PCS scores compared to those uninfected and cases >4 months (Figure 4, upper panel); among cases ≤4 months 22% had a PCS score below 48, whereas for those uninfected and cases >4 months this was 15% and 13% respectively (supplementary file – Table 4). For fatigue, cases ≤4 months had a higher proportion of individuals with higher CIS scores throughout most of the cumulative distribution (Figure 5, upper panel); 15% of those uninfected and cases >4 months scored above 35, whereas for cases ≤4 months this was 23% (supplementary file – Table 5).

Multivariable logistic regression

The results of the logistic regression analyses showed that statistically significant differences between the groups differed based on the cut-off point of the outcome used. The 15% cut-off point corresponds to an SF-6D score of 0.66, an MCS score of 36.77, a PCS score of 48.36 and a CIS score of 35

(supplementary file – Table 2-5). This cut-off represents clinically relevant severity for the fatigue (CIS)
scale (34), hence the regression results for the 15% cut-off point are presented (to note: the regression
results for the other cut-off points can be found in the supplementary files (supplementary file – Table
6-19)). At the 15% cut-off, prevalence of cases ≤4 months was 5.9, 4.3, 6.9 and 7.9 percentage point
higher compared to those uninfected (prevalence cases ≤4 months minus prevalence uninfected) for
SF-6D, MCS, PCS and CIS respectively (supplementary file – Table 2-5). In the multivariable logistic
regression, significant differences between those uninfected and cases ≤4 months were observed for
health utility (OR [CI]: 1.60 [1.15-2.18]), physical health (OR [CI]: 1.92 [1.45-2.53]) and fatigue (OR [CI]:
1.63 [1.25-2.12]), but not for mental health (OR [CI]: 1.19 [0.88-1.57]) (Table 2). For all outcomes, there
was no significant difference observed between those uninfected and cases >4 months at the 15% cut-
off point (or any of the other cut-off points).

At the 15% cut-off point, multivariable logistic regressions (Table 2) showed that age groups 35-65 and
66+ years were significantly associated with poorer physical health, but with better health utility,
mental health, and fatigue (p<0.001) compared to age group 18-35 years. Female sex (p<0.001) and
having comorbidities (p<0.001 to p=0.001) were significantly associated with poorer outcomes on all
four scales, except for mental health and having one comorbidity (p=0.001). Low educational level was
significantly associated with better mental health (p=0.014), but poorer health utility (p=0.039) and
physical health (p<0.001). A higher stringency index was significantly associated with better health
utility and fatigue (p=0.010).

Post-hoc analysis

In the post-hoc analysis we investigated the power to detect a minimal difference in prevalence
between cases >4 months and those uninfected required to detect a significant difference (with those
uninfected as reference group). Given the 327 cases >4 months, a difference in prevalence at the 15%
cut-off would only become significant when the prevalence in this group was at least 4.1% points
higher than that of the uninfected for the SF-6D, 5.5% for the MCS, 3.3% for the PCS and 5.2% for the
CIS (supplementary file – Figure 1-4). Using the given cut-offs, the largest possible difference while
having a non-significant outcome of the logistic regression would be 7.3%, 7.4%, 4.7% or 7.0%
respectively.

Discussion

In this nationwide Dutch cohort study, we assessed the relationship between time since serologically identified SARS-CoV-2 infection and HRQoL (health utility, physical health, mental health) and fatigue among unvaccinated individuals in the first year of the COVID-19 pandemic. Our findings show that recently infected participants (≤4 months ago) more often had poor outcomes in health utility, physical health and fatigue compared to those uninfected (at the 15% cut-off). Almost no significant differences were observed for mental health. For participants infected more than 4 months ago, no statistically significant differences were found compared to those uninfected, though post-hoc analyses showed a power to detect prevalence differences as low as 7%.

These observations are interesting, as they suggest that at the population level (including those symptomatic and asymptomatic) the impact of long-term sequelae after a SARS-CoV-2 infection was lower in our sample compared to the lowest estimated PCC prevalence as currently found in literature (9%) (10). The limited impact that we measured may be explained by a combination of factors. Firstly, in this study we focused on predominantly wild-type SARS-CoV-2 infected individuals, since the Alpha variant of concern started circulating in the end of 2020 and was not dominant until week 7 of 2021 (15-21 February 2021). With the onset of the Alpha variant coinciding with the beginning of the Dutch vaccination campaign for susceptible populations, disentangling the effects of the variant, age and vaccination on PCC endpoints becomes difficult. As a result, there is no evident consensus on the between-variant differences for PCC prior to Omicron as well as the influence of vaccination on PCC development and progression. The definition used, population studied, and variation in vaccination status resulted in large heterogeneity (14). Some studies report that the wild-type SARS-CoV-2 variant is associated with similar odds of long-term symptoms as the Alpha or Delta variants, after adjusting for vaccination status (37, 38). On the other hand, other vaccination-controlled evidence suggests that early pandemic variants held higher risk of PCC (14, 39). A large U.S. cohort study revealed a significant decrease in PCC incidence from pre-Delta to Omicron, primarily attributable to vaccination, with minor contributions from time-related factors such as variants (40). Secondly, our approach of serological SARS-CoV-2 detection is superior to methodologies used in many other studies (e.g. diagnostic SARS-CoV-2 testing) in terms of sensitivity and specificity, given the high test specifics of our assay (23, 41). This enabled us to include mild and asymptomatically infected individuals, who had milder acute COVID-19 and subsequently may have had lower risk for PCC (42). Thirdly, the individuals at risk for severe COVID-19, such as those with underlying conditions, may have also been at risk for long-term health impacts. However, they were less likely to be infected due to adapted behaviors (26, 43, 44). Since our study was randomly sampled from the Dutch population, the impact of several risk factors such as hospitalization, admission to intensive care unit, having comorbidities, smoking and high body

mass index (45, 46) are less visible from our sample. This is because the majority of our participants
experienced mild infections, and risk groups were not overrepresented in our population-based
sample. Fourthly, because of the sampling, there is variability in time since infection for those infected
more than 4 months ago. The majority of these cases are likely to have occurred in March and April
2020, given the epidemic curve in the Netherlands before September 2020. This equates to 10 to 11
months since infection, which is long and can contribute to the lower observed prevalence. Further
subdivision of our cases >4 months into exact round of seroconversion was not possible due to the
small numbers (PICO1: N=64; PICO2: N=40; PICO3: N=20).

We observed differences in health utility between those uninfected and cases ≤4 months, which is in
line with the study by Poudel (2021). Poudel (2021) reported that quality of life was impacted more
during the acute phase of SARS-CoV-2 infection (47). Although HRQoL can also be impacted among
PCC patients, we did not observe any significant differences in health utility between those uninfected
and cases >4 months. There were no differences in mental health observed in our study between
those uninfected and cases >4 months or ≤4 months. This could be because the severe restrictions
that were in place in February 2021 (schools were closed, people had to work from home and an
evening curfew was in place from January 20th until April 28th) had an impact on the entire population,
irrespective of infection status. Literature also shows that although society as a whole was affected by
the pandemic, some groups in society were disproportionally affected such as females and younger
age groups (48). On the other hand, in a study examining HRQoL over the pandemic years, we did not
observe a relationship between quality of life and the stringency of non-pharmaceutical interventions,
nor did we find an indication that HRQoL was very different in February 2021 compared to the period
after (49).

There are some limitations to consider in the interpretation of our results. Firstly, because we use
serological data, acute infections with SARS-CoV-2 just before blood sampling might not have been
detected in our analysis as it can take two to three weeks to seroconvert after primary infection (50).
Similarly, some individuals may have been misclassified in the cases or uninfected groups due to false
negative or false positive serological testing. Few individuals do not seroconvert after infection (51-
56). To minimize the effect of misclassification on our estimation, we excluded participants that
reported a positive PCR or rapid antigen test from the uninfected group. However, since widescale
PCR testing was not available in the Netherlands until June 2020, there is still a possibility of
misclassification. Given that the validation studies of the assay used showed a high sensitivity and
specificity (23, 41), we believe these consist of a small minority, and hence the overall impact on our
results is limited. Moreover, the use of serological data to identify cases enabled identification of
infections, independent of testing accessibility, policies and adherence, which is a major strength of

this study. Secondly, there might be some heterogeneity in SARS-CoV-2 infection variants in our cases

≤4 months. Given that the Alpha variant of concern started circulating in the end of 2020, but was not

dominant until week 7 of 2021 (15-21 February 2021), we expect this to only be a minority of infections

and thus only marginally impact our results (57). Thirdly, reinfections were not taken into account in

our study. However, as very few reinfections occurred early in the pandemic (58), this most likely did

not affect our results (59, 60). Finally, in this study we included a study population that was randomly

selected from the Dutch population in order to ascertain representativeness of the Dutch population.

However, despite these efforts, participants of Dutch descent and potentially more health-oriented

persons were overrepresented, a phenomenon commonly seen in other studies as well. In addition,

since the vaccination rollout in the Netherlands initially prioritized vulnerable groups and healthcare

workers, the exclusion of vaccinated individuals in our study may have affected the representativeness

of our results.

Focusing on early-pandemic serologically identified infections among unvaccinated individuals

provides valuable insights into the overall temporal health dynamics following SARS-CoV-2 infection

in a largely naïve population. With waning vaccine-induced immunity and discontinued testing policy,

these results may reflect current PCC susceptibility among symptomatic and asymptomatic SARS-CoV-

2 infections. In addition, they may serve as a benchmark for PCC estimates following infections with

later SARS-CoV-2 variants in more immunologically heterogeneous populations. Our results could also

shed light into suspectibility to other respiratory infections for which the population is largely naïve

(37, 61, 62).

Conclusion

In a Dutch randomly-selected population-based seroepidemiological cohort of SARS-CoV-2 infected

unvaccinated individuals, those infected in the previous four months more often reported poor health

utility and physical health, and were more often severely fatigued compared to those uninfected.

HRQoL and fatigue remained below the detection limit for those infected more than four months ago,

suggesting a relatively low prevalence of PCC.

Acknowledgements

We would like to thank our colleagues from the National Institute of Public Health and Environment (RIVM), Centre of Infectious Disease Control for their contributions regarding logistics, laboratory analyses, methodological insights and manuscript reviewing.

This work was supported by the Ministry of Health, Welfare and Sports (VWS), the Netherlands. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. There was no additional external funding received for this study.

Data availability

The data that support the findings of this study can be requested via the PICO study website (https://www.rivm.nl/en/pienter-corona-study/information-for-researchers). Restrictions may apply to the availability of these data.

Conflicts of interest

The authors report no conflicts of interest.

References

1.   Bäuerle A, Steinbach J, Schweda A, Beckord J, Hetkamp M, Weismüller B, et al. Mental Health Burden of the COVID-19 Outbreak in Germany: Predictors of Mental Health Impairment. J Prim Care Community Health. 2020;11:2150132720953682.

2.   Hay JW, Gong CL, Jiao X, Zawadzki NK, Zawadzki RS, Pickard AS, et al. A US Population Health Survey on the Impact of COVID-19 Using the EQ-5D-5L. J Gen Intern Med. 2021;36(5):1292-301.

3.   Lim SL, Woo KL, Lim E, Ng F, Chan MY, Gandhi M. Impact of COVID-19 on health-related quality of life in patients with cardiovascular disease: a multi-ethnic Asian study. Health Qual Life Outcomes. 2020;18(1):387.

4.   RIVM. Risk groups and COVID-19: RIVM - National Institute for Public Health and the Environment; 2023 [Available from: https://www.rivm.nl/en/coronavirus-covid-19/risk-groups.

5.   Cirulli ET, Schiabor Barrett KM, Riffle S, Bolze A, Neveux I, Dabe S, et al. Long-term COVID-19 symptoms in a large unselected population. medRxiv. 2020:2020.10.07.20208702.

6.   Petersen MS, Kristiansen MF, Hanusson KD, Danielsen ME, B ÁS, Gaini S, et al. Long COVID in the Faroe Islands: A Longitudinal Study Among Nonhospitalized Patients. Clin Infect Dis. 2021;73(11):e4058-e63.

7.   Nalbandian A, Sehgal K, Gupta A, Madhavan MV, McGroder C, Stevens JS, et al. Post-acute COVID-19 syndrome. Nat Med. 2021;27(4):601-15.

8.   Ziauddeen N, Gurdasani D, O'Hara ME, Hastie C, Roderick P, Yao G, et al. Characteristics and impact of Long Covid: Findings from an online survey. PLOS ONE. 2022;17(3):e0264331.

9.   Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-19 condition by a Delphi consensus. Lancet Infect Dis. 2022;22(4):e102-e7.

10.   Chen C, Haupert SR, Zimmermann L, Shi X, Fritsche LG, Mukherjee B. Global Prevalence of Post-Coronavirus Disease 2019 (COVID-19) Condition or Long COVID: A Meta-Analysis and Systematic Review. J Infect Dis. 2022;226(9):1593-607.

11.   Costantino V, Grafton Q, Kompas T, Chu L, Honeyman D, Notaras A, et al. The public health and economic burden of long COVID in Australia, 2022–24: a modelling study. Medical Journal of Australia. 2024;221(4):217-23.

12.   Ballering AV, van Zon SKR, Olde Hartman TC, Rosmalen JGM. Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study. Lancet. 2022;400(10350):452-61.

13.   van der Maaden T, Mutubuki EN, de Bruijn S, Leung KY, Knoop H, Slootweg J, et al. Prevalence and Severity of Symptoms 3 Months After Infection With SARS-CoV-2 Compared to Test-Negative and Population Controls in the Netherlands. J Infect Dis. 2023;227(9):1059-67.

14.   Fernández-de-Las-Peñas C, Notarte KI, Peligro PJ, Velasco JV, Ocampo MJ, Henry BM, et al. Long-COVID Symptoms in Individuals Infected with Different SARS-CoV-2 Variants of Concern: A Systematic Review of the Literature. Viruses. 2022;14(12).

15.   Kritsotakis EI. On the importance of population-based serological surveys of SARS-CoV-2 without overlooking their inherent uncertainties. Public Health in Practice. 2020;1:100013.

16.   Dumont R, Richard V, Lorthe E, Loizeau A, Pennacchio F, Zaballa M-E, et al. A population-based serological study of post-COVID syndrome prevalence and risk factors in children and adolescents. Nature Communications. 2022;13(1):7086.

17.   Blankenburg J, Wekenborg MK, Reichert J, Kirsten C, Kahre E, Haag L, et al. Comparison of mental health outcomes in seropositive and seronegative adolescents during the COVID19 pandemic. Scientific Reports. 2022;12(1):2246.

18.   Matta J, Wiernik E, Robineau O, Carrat F, Touvier M, Severi G, et al. Association of Self-reported COVID-19 Infection and SARS-CoV-2 Serology Test Results With Persistent Physical Symptoms Among French Adults During the COVID-19 Pandemic. JAMA Intern Med. 2022;182(1):19-25.

19.   Robineau O, Wiernik E, Lemogne C, de Lamballerie X, Ninove L, Blanché H, et al. Persistent symptoms after the first wave of COVID-19 in relation to SARS-CoV-2 serology and experience of

425 acute symptoms: A nested survey in a population-based cohort. Lancet Reg Health Eur.
426 2022;17:100363.

20. Vos ERA, den Hartog G, Schepp RM, Kaaijk P, van Vliet J, Helm K, et al. Nationwide
428 seroprevalence of SARS-CoV-2 and identification of risk factors in the general population of the
429 Netherlands during the first epidemic wave. J Epidemiol Community Health. 2020;75(6):489-95.

21. Verberk JDM, Vos RA, Mollema L, van Vliet J, van Weert JWM, de Melker HE, et al. Third
431 national biobank for population-based seroprevalence studies in the Netherlands, including the
432 Caribbean Netherlands. BMC Infect Dis. 2019;19(1):470.

22. Vos ERA, van Boven M, den Hartog G, Backer JA, Klinkenberg D, van Hagen CCE, et al.
434 Associations Between Measures of Social Distancing and Severe Acute Respiratory Syndrome
435 Coronavirus 2 Seropositivity: A Nationwide Population-based Study in the Netherlands. Clin Infect
436 Dis. 2021;73(12):2318-21.

23. den Hartog G, Schepp RM, Kuijer M, GeurtsvanKessel C, van Beek J, Rots N, et al. SARS-CoV-
438 2-Specific Antibody Detection for Seroepidemiology: A Multiplex Analysis Approach Accounting for
439 Accurate Seroprevalence. J Infect Dis. 2020;222(9):1452-61.

24. Geers D, Shamier MC, Bogers S, den Hartog G, Gommers L, Nieuwkoop NN, et al. SARS-CoV-2
441 variants of concern partially escape humoral but not T-cell responses in COVID-19 convalescent
442 donors and vaccinees. Sci Immunol. 2021;6(59).

25. van den Hoogen LL, Smits G, van Hagen CCE, Wong D, Vos ERA, van Boven M, et al.
444 Seropositivity to Nucleoprotein to detect mild and asymptomatic SARS-CoV-2 infections: A
445 complementary tool to detect breakthrough infections after COVID-19 vaccination? Vaccine.
446 2022;40(15):2251-7.

26. Vos ERA, van Hagen CCE, Wong D, Smits G, Kuijer M, Wijmenga-Monsuur AJ, et al. SARS-CoV-
448 2 Seroprevalence Trends in the Netherlands in the Variant of Concern Era: Input for Future
449 Response. Influenza Other Respir Viruses. 2024;18(6):e13312.

27. CBS. Cims: COVID-vaccinatie Informatie- en Monitoringssysteem: Centraal Bureau voor de
451 Statistiek; [Available from: https://www.cbs.nl/nl-nl/onze-diensten/maatwerk-en-
452 microdata/microdata-zelf-onderzoek-doen/microdatabestanden/cims-covid-vaccinatie-informatie-
453 en-monitoringssysteem.

28. Mols F, Pelle AJ, Kupper N. Normative data of the SF-12 health survey with validation using
455 postmyocardial infarction patients in the Dutch population. Qual Life Res. 2009;18(4):403-14.

29. Ware JE, Kosinski M, Keller SD. SF-12: How to score the SF-12 physical and mental health
457 summary scales. Boston, MA: The Health Institute, New England Medical Center; 1995.

30. Brazier JE, Roberts J. The estimation of a preference-based measure of health from the SF-
459 12. Med Care. 2004;42(9):851-9.

31. van Stel HF, Buskens E. Comparison of the SF-6D and the EQ-5D in patients with coronary
461 heart disease. Health Qual Life Outcomes. 2006;4:20.

32. Vercoulen JH, Swanink CM, Fennis JF, Galama JM, van der Meer JW, Bleijenberg G.
463 Dimensional assessment of chronic fatigue syndrome. J Psychosom Res. 1994;38(5):383-92.

33. Worm-Smeitink M, Gielissen M, Bloot L, van Laarhoven HWM, van Engelen BGM, van Riel P,
465 et al. The assessment of fatigue: Psychometric qualities and norms for the Checklist individual
466 strength. J Psychosom Res. 2017;98:40-6.

34. van Engelen E. Uitgebreide toelichting van het meetinstrument - Checklist Individuele
468 Spankracht (CIS): Meetinstrumenten in de zorg; 2019 [Available from:
469 https://meetinstrumentenzorg.nl/wp-content/uploads/instrumenten/CIS-form.pdf.

35. Hale T, Angrist N, Goldszmidt R, Kira B, Petherick A, Phillips T, et al. A global panel database
471 of pandemic policies (Oxford COVID-19 Government Response Tracker). Nat Hum Behav.
472 2021;5(4):529-38.

36. Roser M. What is the COVID-19 Stringency Index? : Our World in Data; 2021 [Available from:
https://ourworldindata.org/metrics-explained-covid19-stringency-index.

37.   Beale S, Yavlinsky A, Fong WLE, Nguyen VG, Kovar J, Vos T, et al. Long-term outcomes of
SARS-CoV-2 variants and other respiratory infections: evidence from the Virus Watch prospective
cohort in England. Epidemiol Infect. 2024;152:e77.
38.   Azzolini E, Levi R, Sarti R, Pozzi C, Mollura M, Mantovani A, et al. Association Between
BNT162b2 Vaccination and Long COVID After Infections Not Requiring Hospitalization in Health Care
Workers. Jama. 2022;328(7):676-8.
39.   Fernández-de-Las-Peñas C, Cancela-Cilleruelo I, Rodríguez-Jiménez J, Gómez-Mayordomo V,
Pellicer-Valero OJ, Martín-Guerrero JD, et al. Associated-Onset Symptoms and Post-COVID-19
Symptoms in Hospitalized COVID-19 Survivors Infected with Wuhan, Alpha or Delta SARS-CoV-2
Variant. Pathogens. 2022;11(7).
40.   Xie Y, Choi T, Al-Aly Z. Postacute Sequelae of SARS-CoV-2 Infection in the Pre-Delta, Delta,
and Omicron Eras. N Engl J Med. 2024;391(6):515-25.
41.   den Hartog G, Vos ERA, van den Hoogen LL, van Boven M, Schepp RM, Smits G, et al.
Persistence of Antibodies to Severe Acute Respiratory Syndrome Coronavirus 2 in Relation to
Symptoms in a Nationwide Prospective Study. Clin Infect Dis. 2021;73(12):2155-62.
42.   Tsampasian V, Elghazaly H, Chattopadhyay R, Debski M, Naing TKP, Garg P, et al. Risk Factors
Associated With Post-COVID-19 Condition: A Systematic Review and Meta-analysis. JAMA Intern
Med. 2023;183(6):566-80.
43.   RIVM. Naleving van en draagvlak voor de basis gedragsregels: RIVM - National Institute for
Public Health and the Environment; 2024 [Available from:
https://www.rivm.nl/gedragsonderzoek/trendonderzoek.
44.   RIVM. Ondanks versoepelingen nog (deels) geïsoleerd leven: RIVM - National Institute for
Public Health and the Environment; 2022 [Available from:
https://www.rivm.nl/gedragsonderzoek/ondanks-versoepelingen-nog-deels-geisoleerd-leven.
45.   Subramanian A, Nirantharakumar K, Hughes S, Myles P, Williams T, Gokhale KM, et al.
Symptoms and risk factors for long COVID in non-hospitalized adults. Nat Med. 2022;28(8):1706-14.
46.   Tziolos NR, Ioannou P, Baliou S, Kofteridis DP. Long COVID-19 Pathophysiology: What Do We
Know So Far? Microorganisms. 2023;11(10).
47.   Poudel AN, Zhu S, Cooper N, Roderick P, Alwan N, Tarrant C, et al. Impact of Covid-19 on
health-related quality of life of patients: A structured review. PLoS One. 2021;16(10):e0259164.
48.   RIVM. More than 1 in 5 young people had suicidal thoughts during last lockdown: RIVM -
National Institute for Public Health and the Environment; 2022 [Available from:
https://www.rivm.nl/en/news/more-than-1-in-5-young-people-had-suicidal-thoughts-during-last-
lockdown.
49.   van Hagen CCE, Huiberts AJ, Mutubuki EN, de Melker HE, Vos ERA, van de Wijgert JHHM, et
al. Health-related quality of life during the COVID-19 pandemic: The impact of restrictive measures
using data from two Dutch population-based cohort studies. PLOS ONE. 2024;19(3):e0300324.
50.   Guo L, Ren L, Yang S, Xiao M, Chang D, Yang F, et al. Profiling Early Humoral Response to
Diagnose Novel Coronavirus Disease (COVID-19). Clin Infect Dis. 2020;71(15):778-85.
51.   Oved K, Olmer L, Shemer-Avni Y, Wolf T, Supino-Rosin L, Prajgrod G, et al. Multi-center
nationwide comparison of seven serology assays reveals a SARS-CoV-2 non-responding seronegative
subpopulation. EClinicalMedicine. 2020;29:100651.
52.   Masiá M, Telenti G, Fernández M, García JA, Agulló V, Padilla S, et al. SARS-CoV-2
Seroconversion and Viral Clearance in Patients Hospitalized With COVID-19: Viral Load Predicts
Antibody Response. Open Forum Infect Dis. 2021;8(2):ofab005.
53.   Wellinghausen N, Plonné D, Voss M, Ivanova R, Frodl R, Deininger S. SARS-CoV-2-IgG
response is different in COVID-19 outpatients and asymptomatic contact persons. J Clin Virol.
2020;130:104542.
54.   Thiruvengadam R, Chattopadhyay S, Mehdi F, Desiraju BK, Chaudhuri S, Singh S, et al.
Longitudinal Serology of SARS-CoV-2-Infected Individuals in India: A Prospective Cohort Study. Am J
Trop Med Hyg. 2021;105(1):66-72.

55. Pathela P, Crawley A, Weiss D, Maldin B, Cornell J, Purdin J, et al. Seroprevalence of Severe
Acute Respiratory Syndrome Coronavirus 2 Following the Largest Initial Epidemic Wave in the United
States: Findings From New York City, 13 May to 21 July 2020. J Infect Dis. 2021;224(2):196-206.
56. Liu W, Russell RM, Bibollet-Ruche F, Skelly AN, Sherrill-Mix S, Freeman DA, et al. Predictors
of Nonseroconversion after SARS-CoV-2 Infection. Emerg Infect Dis. 2021;27(9):2454-8.
57. RIVM. Variants of the coronavirus SARS-CoV-2: RIVM - National Institute for Public Health
and the Environment; 2024 [Available from: https://www.rivm.nl/en/coronavirus-covid-
19/current/variants.
58. de Boer PT, van de Kassteele J, Vos ERA, van Asten L, Dongelmans DA, van Gageldonk-
Lafeber AB, et al. Age-specific severity of severe acute respiratory syndrome coronavirus 2 in
February 2020 to June 2021 in the Netherlands. Influenza and Other Respiratory Viruses.
2023;17(8):e13174.
59. Medić S, Anastassopoulou C, Lozanov-Crvenković Z, Vuković V, Dragnić N, Petrović V, et al.
Risk and severity of SARS-CoV-2 reinfections during 2020-2022 in Vojvodina, Serbia: A population-
level observational study. Lancet Reg Health Eur. 2022;20:100453.
60. O Murchu E, Byrne P, Carty PG, De Gascun C, Keogan M, O'Neill M, et al. Quantifying the risk
of SARS-CoV-2 reinfection over time. Rev Med Virol. 2022;32(1):e2260.
61. Brown M, Gerrard J, McKinlay L, Marquess J, Sparrow T, Andrews R. Ongoing symptoms and
functional impairment 12 weeks after testing positive for SARS-CoV-2 or influenza in Australia: an
observational cohort study. BMJ Public Health. 2023;1(1):e000060.
62. Xie Y, Choi T, Al-Aly Z. Long-term outcomes following hospital admission for COVID-19 versus
seasonal influenza: a cohort study. Lancet Infect Dis. 2024;24(3):239-55.

Figure 1. Flowchart of participant recruitment in PICO1-4

Table 1: Participant demographic and health characteristics*

	UNINFECTED	CASES >4 MONTHS	CASES ≤4 MONTHS
N	4,569	327	351
SEX = FEMALE *% (N)*	55.5 (2,534)	55.4 (181)	57.8 (203)
AGE GROUPS *% (N)*			
15-35	21.7 (990)	28.7 (94)	32.2 (113)
36-65	52.0 (2,375)	47.7 (156)	50.4 (177)
66+	26.4 (1,204)	23.5 (77)	17.4 (61)
EDUCATIONAL LEVEL[A] *% (N)*			
HIGH	47.7 (2,178)	43.7 (143)	38.7 (136)
INTERMEDIATE	31.8 (1,455)	37.9 (124)	39.9 (140)
LOW	20.5 (936)	18.3 (60)	21.4 (75)
NUMBER OF COMORBIDITIES[B] *% (N)*			
NONE	65.9 (3,010)	64.8 (212)	73.2 (257)
ONE	24.6 (1,126)	27.2 (89)	19.9 (70)
TWO OR MORE	9.5 (433)	8.0 (26)	6.8 (24)
HEALTH UTILITY (SF-6D) *MEAN (SD)*	0.82 (0.11)	0.81 (0.11)	0.79 (0.13)
MENTAL HEALTH (MCS) *MEAN (SD)*	47.83 (10.26)	47.16 (11.13)	46.36 (10.87)
PHYSICAL HEALTH (PCS) *MEAN (SD)*	54.53 (7.54)	54.93 (6.86)	53.18 (8.74)
FATIGUE (CIS) *MEAN (SD)*	21.67 (11.67)	21.73 (11.44)	24.77 (12.96)

* All characteristics were determined in PICO4, except for educational level and comorbidities (determined at baseline).

[A] Educational level was classified as low (no education or primary education), intermediate (secondary school or vocational training), or high (bachelor's degree, university).

[B] Included comorbidities are asthma, cancer, cardiovascular disease, diabetes (type 1 & 2), immunodeficiency, pulmonary disease and renal disease.

Figure 2. Health utility – cumulative distribution and logistic regression analyses*

* Each 5% increment (5-75%) along the cumulative distribution of the control group is marked with a grey bar. The 15% point of the distribution is marked with a dark grey bar.

Figure 3. Mental health – cumulative distribution and logistic regression analyses*

* Each 5% increment (5-75%) along the cumulative distribution of the control group is marked with a grey bar. The 15% point of the distribution is marked with a dark grey bar.

Figure 4. Physical health – cumulative distribution and logistic regression analyses*

* Each 5% increment (5-75%) along the cumulative distribution of the control group is marked with a grey bar. The 15% point of the distribution is marked with a dark grey bar.

Figure 5. Fatigue – cumulative distribution and logistic regression analyses*

* Each 5% increment (5-75%) along the cumulative distribution of the control group is marked with a grey bar. The 15% point of the distribution is marked with a dark grey bar.

Table 2. Multivariable logistic regression results for SF-6D, MCS, PCS and CIS at the 15% cut-off.

		SF-6D		MCS		PCS		CIS	
		OR [95%CI]	p-value	OR [95%CI]	p-value	OR [95%CI]	p-value	OR [95%CI]	p-value
	Intercept	5106.63	0.045	1.39	0.933	81.09	0.290	1784.32	0.038
SEROLOGICAL STATUS	Uninfected	1		1		1		1	
	Cases >4 months	1.15 [0.78-1.64]	0.467	1.15 [0.84-1.54]	0.377	0.87 [0.61-1.22]	0.436	1.03 [0.75-1.39]	0.845
	Cases ≤4 months	1.60 [1.15-2.18]	0.004	1.19 [0.88-1.57]	0.242	1.92 [1.45-2.53]	<0.001	1.63 [1.25-2.12]	<0.001
AGE GROUP	18-35 years	1		1		1		1	
	36-65 years	0.58 [0.46-0.72]	<0.001	0.41 [0.34-0.48]	<0.001	1.79 [1.40-2.30]	<0.001	0.58 [0.48-0.69]	<0.001
	65+ years	0.40 [0.30-0.54]	<0.001	0.15 [0.11-0.20]	<0.001	2.53 [1.93-3.34]	<0.001	0.27 [0.21-0.35]	<0.001
EDUCATIONAL LEVEL	High	1		1		1		1	
	Intermediate	1.26 [1.02-1.56]	0.032	0.85 [0.72-1.01]	0.065	1.45 [1.20-1.74]	<0.001	1.14 [0.96-1.35]	0.135
	Low	1.31 [1.01-1.70]	0.039	0.74 [0.59-0.94]	0.014	1.66 [1.36-2.03]	<0.001	1.07 [0.87-1.33]	0.511
SEX	Male	1		1		1		1	
	Female	1.62 [1.33-1.98]	<0.001	1.49 [1.26-1.75]	<0.001	1.59 [1.35-1.88]	<0.001	1.87 [1.59-2.20]	<0.001
COMORBIDITIES	No comorbidities	1		1		1		1	
	1 comorbidity	1.52 [1.21-1.89]	<0.001	1.20 [0.99-1.45]	0.056	2.08 [1.74-2.49]	<0.001	1.77 [1.48-2.12]	<0.001
	2+ comorbidities	2.46 [1.81-3.31]	<0.001	1.65 [1.22-2.20]	0.001	4.29 [3.41-5.37]	<0.001	3.21 [2.49-4.11]	<0.001
STRINGENCY INDEX	75	1		1		1		1	
	79	0.87 [0.78-0.97]	0.010	0.98 [0.89-1.08]	0.707	0.91 [0.82-1.01]	0.065	0.89 [0.81-0.97]	0.010