1
2　　　　　A brain vital sign study of subconcussion in female and male high school student athletes
3
4

5　　　Ryan C.N. D'Arcy [1,2,3]*, David McCarthy[4], Derek Harrison[4], Zander Levenberg[4], Julian Wan[4], Aidan
6　Hepburn[4], Eric D. Kirby [1,5], Tanja Yardley[6], Nikita Yamada-Bagg[2], Shaun D. Fickling[1,2], Thayne A. Munce[7],
7　　　　　　　David W. Dodick[8,9], Christopher Ahmad[10,11], & Ken Shubin Stein[9,12]
8
9
10
11
1)　BrainNET, Health and Technology District, Surrey, BC, Canada;
2)　Centre for Neurology Studies, HealthTech Connex, Surrey, BC, Canada;
3)　Faculty of Engineering Science, Simon Fraser University, Burnaby, BC, Canada;
4)　Brentwood College School, BRAIN team, Mill Bay, BC, Canada;
5)　Faculty of Individualized Interdisciplinary Studies, Simon Fraser University, Burnaby, BC,
Canada;
6)　Surrey Neuroplasticity Clinic, HealthTech Connex, Surrey, BC, Canada;
7)　Environmental Influences on Health & Disease Group, Sanford Research, Sioux Falls, SD,
USA;
8)　Mayo Clinic College of Medicine, Rochester, MN, USA;
9)　Atria Academy of Science and Medicine, New York City, NY, USA;
10)　New York Presbyterian Hospital, New York City, NY, USA;
11)　Columbia University Irving Medical Center, New York City, NY, USA;
12)　Hospital for Special Surgery, New York City, NY, USA.
26
27
28　*Corresponding author: Ryan C.N. D'Arcy, PhD, PlEng, Centre for Neurology Studies, HealthTech Connex,
29　Surrey, BC Canada ryan@healthtechconnex.com

Abstract

Emerging evidence from neurophysiological brain vital sign studies show repeatable sensitivity
to cumulative subconcussive impairments over a season of contact sports. The current study addressed
the need for a large prospective study comparing a low-contact control group to high-contact group.
Importantly, the study also expanded the scope of neurophysiological changes related to repetitive head
impacts to include female athletes in addition to male athletes. In total, 89 high school student athletes
underwent 231 brain vital sign scans over a full calendar year. The results replicated prior subconcussive
cognitive impairments (N400 delays) and sensory impairments (N100 amplitude reductions) in male
athletes and demonstrated similar subconcussive impairments for the first time in female athletes.
While there was no significant subconcussive difference between female and male athletes, female
athletes show overall larger responses in general. The findings demonstrated that subconcussive
impairments are detectable in a controlled experimental comparison for both female and male high
school athletes. The study highlights the opportunity to monitor subconcussive changes in cognitive
processing for both female and male athletes to help advance prevention, mitigation and management
efforts aimed at reducing athletes' risk of potential long-term negative health outcomes related to
cumulative exposure to repetitive head impacts.
51

Introduction

Background

Head impact exposure from participating in contact or collision sports is commonly associated
with concussion; however, subconcussion is a growing concern for athletes as well as a developing
research area in parallel (Smith et al., 2018; 2019). Subconcussion has been increasingly implicated with
potential short-term consequences on the developing brains and longer-term concerns related to chronic
traumatic encephalopathy (McKee et al. 2009; Guskiewicz et al. 2005; Dioso et al. 2022). Concussion
research has also historically been primarily focused on males (Valera et al. 2021; Snedaker et al. 2022).
Therefore, emerging subconcussive research must be inclusive of both female and male athletes. This is a
timely priority as new intervention options begin to emerge from the clinical trials literature (Breuer et al.
2023).
Operationally defined, a subconcussive impact is a mechanical force transmitted to the brain
below the threshold for a diagnosis of an acute concussive injury. The effects of these repetitive head
impacts (RHI) may neither be detectable to players nor to observers; however, it has been shown that RHI
over time can result in both microstructural alterations and functional brain impairment (Mainwaring et
al. 2018). The extent of impairment is highly correlated with the frequency of exposure to these impacts
(Fickling, Smith, et al. 2021; Fickling, Poel, et al. 2021; Hirad et al. 2019; Saghafi et al. 2018; Fickling et al.
2019).
Recent research has increasingly focused on the role of cumulative RHI exposure rather than a
concussion as a singular, acute traumatic event. Evidence suggests that even a single practice session
involving head contact, such as heading a ball in soccer, can result in impairment (Nowak et al. 2020).
Magnetic resonance imaging (MRI) studies have demonstrated that players exposed to RHI over the

course of a season of sport show significant structural abnormalities compared to controls over the same
time period (Hirad et al. 2019; Saghafi et al. 2018; Veksler et al. 2020). Both Hirad et al. (2019) and Saghafi
et al. (2018) used diffusion-tensor imaging, an MRI-derived technique, to show abnormalities in the white
matter of a group of football athletes were related to different measures of head impact exposure.
Interestingly, while detecting differences in permeability and white matter microstructure in the brains of
contact sport athletes, Veksler et al. (2020) did not find any statistical differences in blood-brain barrier
permeability between non-contact athletes and non-athlete healthy controls, leading the authors to
merge the groups into a single control group.
A major challenge for current medical imaging approaches, including MRI, relates to the need for
accessible, objective, and sensitive measures of subconcussive impacts at points-of-care (PoC). Portable
electroencephalography (EEG) provides the evaluation of individual neurophysiological event-related
potentials (ERPs). These have recently been translated into a brain vital signs framework, facilitating
access to sensitive measures for various PoC applications (Ghosh Hajra et al. 2016). Three well-established
ERP responses are extracted within the brain vital signs framework: 1) the N100 for auditory sensation
(Davis 1939); 2) the P300 for basic attention (Sutton et al. 1967); and 3) the N400 for cognitive processing
(Kutas and Federmeier 2011). Brain vital signs have been successfully implemented in the evaluation of
both concussion and subconcussion (Fickling et al. 2019; Fickling, Smith, et al. 2021; Fickling, Poel, et al.
2021).
Identification of subconcussive cognitive impairments over a season of contact sports was initially
reported in male Junior-A ice hockey players (Fickling et al. 2019, N=23). Subsequent studies (Fickling,
Smith, et al. 2021, N=23a; Fickling, Poel, et al. 2021, N=15b; Breuer et al. 2023, N=30) replicated and
confirmed identification of accumulated subconcussive exposure in male athletes for different age ranges
(12-14, 16-21, & 21-28, respectively) and different contact sports, including ice hockey, tackle football,
and mixed martial arts. Importantly, this work demonstrated strong, significant, and linearly predictive
relationships between brain vital sign changes and exposure to subconcussive impacts (Fickling, Smith, et
al. 2021; Fickling, Poel, et al. 2021). These relationships were demonstrated both when subconcussive
impacts were quantified directly by impact sensors or indirectly by the number of games and practices
played. A pattern of subconcussive changes have emerged, identified by changes in the N400 (cognitive
processing) and N100 (sensory processing). The N400 and N100 components have been implicated in the
rate of information processing and synchronous pyramidal neural activation (Luck 2014): the two are
interdependent factors reflecting an association with cognitive processing (Kutas and Federmeier 2011)
and auditory sensation (Davis 1939). While the replications of N400 and N100 subconcussive changes in
male athletes are noteworthy, a single prospective study with a large sample size including female athletes
is critical.

Objectives & Hypothesis

The study was conducted at Brentwood College School (a Grade 8-12 high school in Mill Bay,
British Columbia, Canada) by student researchers from the Brentwood Research, Action, and Innovation
in Neuroscience (BRAIN) team. Brain vital signs were monitored in 89 female and male high school
students (15-17 years of age) participating in high-contact (e.g. rugby, ice hockey, and soccer) and low-
contact sports (e.g. rowing, climbing, and tennis) over three academic terms across the school year (Figure

1). With respect to matching participants in high-contact versus low-contact sports, Brentwood is a private
boarding school, in which all students follow a structured program with commonly matched ages, daily
schedules, diets, academics, athletics, and sleep routines. The study used a mixed-model, repeated
measures design evaluated across 3 terms before and after each of the three school year terms as well as
after the summer break, resulting in 7 possible scan timepoints. The objectives were to replicate prior
subconcussion findings in male athletes, as well as in female athletes, with the primary hypothesis
predicting that brain vital sign subconcussive differences would be detectable in the comparisons for high-
contact sports relative to low-contact sports. Subsequent analyses were conducted to explore specific
differences between female and male subconcussive and overall neurophysiological results.
125
126

128   **Figure 1:** Overview of the experimental design and study timeline (N: Number of participants; n: brain
129   vital signs scans completed).
130
131

Results

In the current study, 89 female and male student athletes were scanned a total of 231 times
before and after varying terms of contact or non-contact sports. All three brain vital signs, the N100,
P300, and N400, were verified at both the group- and individual- levels. Peak amplitude and latency
values were examined in the multivariate analysis of variance (MANOVA), which were first examined
with univariate tests of normality and Levene's test of equality of error variances. N100 latency and
N400 amplitude failed tests of normality ($p < 0.05$), while all amplitude and latency values passed
Levene's test ($p > 0.05$). Therefore, non-parametric Kruskal-Wallis test results are also included in
Supplemental Data 1 to demonstrate the robustness of the following MANOVA results.
142

<u>High-contact versus low-contact effects:</u>

Figure 2a shows brain vital sign differences in the N400 latency and N100 amplitude, while

figure 2b provides the underlying waveform differences. The findings confirmed the consistent

replication of subconcussive changes in the N400 and N100 at the overall group level, with a significant

delay in N400 peak latency and a significant increase in N100 peak amplitude. Specifically, the MANOVA

showed a significant main effect of condition between high- versus low-contact groups [$F(6,104) = 7.95$,

$p < 0.001$, $\eta_p^2 = 0.314$] (Table 1). Separate univariate between-subjects tests on the outcome variables

revealed significant effects of group on the N400 latency, [$F(1) = 35.79$, $p < 0.001$, $\eta_p^2 = 0.247$], and N100

amplitude, [$F(1) = 7.88$, $p = 0.006$, $\eta_p^2 = 0.067$] (Figure 2, Table 2). Figure 3 provides the individual scatter

plot data for both the N400 latency and N100 amplitude effects, including means and standard

deviations. The current results addressed the planned high- versus low-contact group comparison to

build on prior pre- versus post- season differences in male athletes. Specific pre- versus post-

comparisons are provided as Supplemental data for comparison (Supplemental Figure 1).

Figure 2: (A): Overall group (female and male combined) brain vital sign radar plot comparing high-

versus low-contact groups. **(B)** Group-level grand average ERP waveforms from tones (top) and word

pairs (bottom) for overall group comparing high- versus low-contact group differences. Deviant tones

and incongruent word pair waveforms are plotted, with standard error shaded. N100, P300, and N400

brain vital sign components of the high-contact group are highlighted with labels and arrows. Asterisks

denote significant effects in both the brain vital sign and waveform results.

Table 1: MANOVA multivariate effect results for high- and low-contact scans.

MANOVA	DF	PILLAI	*F*-VALUE	*P*-VALUE
GROUP	6,104	0.314	**7.947**	**<0.001***
SEX	6, 104	0.148	**3.020**	**0.009***
GROUP X SEX	6, 104	0.003	0.047	1.000

168 **Table 2:** MANOVA results for the group effect on each brain vital sign component.

Brain vital sign	High-contact mean (standard deviation)	Low-contact mean (standard deviation)	*F*-value	*p*-value
N100 Amplitude	-4.60 (2.82)	-3.15 (2.11)	**7.883**	**0.006***
N100 Latency	100.78 (16.58)	100.78 (25.64)	0.750	0.388
P300 Amplitude	4.52 (2.84)	3.86 (3.26)	0.508	0.477
P300 Latency	274.77 (40.03)	270.14 (35.22)	0.304	0.583
N400 Amplitude	-3.49 (2.85)	-3.35 (2.26)	0.001	0.977
N400 Latency	433.50 (70.42)	359.16 (57.86)	**35.786**	**<0.001***

171
Figure 3: Individual scatter plot data, including group means and standard deviations (middle) showing a
significant N400 latency and N100 amplitude differences for high-contact (red) versus low-contact
(blue).

<u>Female compared to male subconcussive effects:</u>
There was no significant sex by contact group interaction present [$F(6,104) = 0.047$, $p = 1.00$, η_p^2
$= 0.003$] (Table 1), indicating female and male athletes experienced similar subconcussive effects. Figure
4a shows brain vital sign radar plots for female and male athletes for high- versus low-contact and figure
4b provides the corresponding waveform results. The MANOVA confirmed a significant main effect for
female athletes compared to male athletes [$F(6, 104) = 3.02$, $p = 0.009$, $\eta_p^2 = 0.148$] (Table 1) across
these scans. Separate univariate between-subjects tests revealed significant effects for the P300
amplitude [$F(1) = 7.45$, $p = 0.007$, $\eta_p^2 = 0.064$] and N400 amplitude [$F(1) = 4.71$, $p = 0.032$, $\eta_p^2 = 0.041$]
(Table 3). Note in Figure 4b that the vertical y-axis (Amplitude) for female athletes and male athletes is

fixed, highlighting the overall increased response amplitude in female responses relative to male
responses across all peaks.
187

188
Figure 4: (A) Radar plots comparing individual-level brain vital signs of the high-contact and low-contact
groups for both female and male athletes. **(B)** Group-level grand average ERP waveforms from tones
(top) and word pairs (bottom) for female athletes (left, dark shading) and male athletes (right, light
shading) comparing high- versus low-contact group differences. Deviant tones and incongruent word
pair waveforms are plotted (with standard error shading). N100, P300, and N400 brain vital sign
components of the high-contact group are highlighted with labels and arrows. Note in Figure 4b that the
vertical y-axis (Amplitude) for female athletes and male athletes is fixed, highlighting the overall
increased response amplitude in female responses relative to male responses across all peaks.
197
198
199 **Table 3:** MANOVA results for the main effect of sex on each brain vital sign component (high- and low-
200 contact scans).

Brain vital sign component	Female athlete mean (standard deviation)	Male athlete mean (standard deviation)	*F*-value	*p*-value
N100 Amplitude	-4.29 (2.47)	-3.36 (2.47)	2.427	0.122
N100 Latency	94.28 (15.32)	101.86 (25.74)	2.597	0.110
P300 Amplitude	5.18 (3.24)	3.47 (2.85)	**7.447**	**0.007***
P300 Latency	273.21 (37.50)	271.17 (37.04)	0.025	0.874
N400 Amplitude	-4.07 (2.96)	-3.00 (2.08)	**4.707**	**0.032***
N400 Latency	391.67 (78.34)	385.91 (69.232)	0.053	0.819

<u>Female compared to male effects:</u>

To further characterize brain vital sign profiles, the multivariate main effect of sex, and the

univariate results above (Table 3) for female and male athletes, contact condition was collapsed across

high- and low-contact scans to create Figure 5. P300 and N400 amplitude differences accounts for a

significant main effect with a near significant effect in the N100 amplitude. The trends driving the main

effect can be seen in Figure 6, which showed the individual scatter plots across female athletes and male

athletes, including means and standard deviations. To further characterize female and male effects with

a larger and closely weighted group number, Figure 7 provides radar plot and waveform results from

female and male athletes for all collected scans (N=202 scans). The purpose of this comparison was to

include all scans as an evaluation of robust differences between female and male athletes.

Corresponding statistics are in Supplemental Data 3.

Figure 5: Comparison of all female and male high- and low-contact scans showing amplitude differences.
(A) Radar plot showing overall female and male brain vital sign difference. **(B)** Group-level waveforms
from deviant tones (top) and incongruent word pairs (bottom) are plotted (standard error shading) for
female athletes (dark blue) and male athletes (light blue). N100, P300, and N400 components are
highlighted with arrows.
221
222

223
224 **Figure 6:** Scatter plots of individual-level data points for all scans of the amplitudes (top) and latencies
225 (bottom) for all brain vital signs components in female athletes (left) and male athletes (right).
226 Significant differences are highlighted by the asterisk, all p-values are reported as well (Bonferroni
227 adjusted).
228
229

230
Figure 7: Comparison of all female and male scans. **(A)** Radar plot showing individual-level brain vital
sign differences across all six components. **(B)** Group-level waveforms from deviant tones (top) and
incongruent word pairs (bottom) are plotted (standard error shading) confirming lower amplitudes in
male athletes (highlighted with arrows).

The current analyses examined high- versus low-contact, using equivalent exposure durations to
prior studies. For additional analyses (N=202 scans), contact exposure levels were compared between
low-contact exposure and both a single term and two or more terms of high-contact exposure. These
results were closely consistent with both subconcussive and sex effects above and are provided as
Supplemental Data 4.

Discussion

The current findings replicated prior studies of subconcussive changes in brain vital signs. The
study used a large, prospective, repeated measures design to compare high- versus low-contact exposure
across both female and male athletes. The findings supported the primary hypothesis that prior
subconcussive differences in the N400 and N100 would be detectable in a high- versus low- contact
comparison (Figure 2), not only for male athletes but also in female athletes (Figure 4). Additional
comparisons of female and male results showed significantly larger P300 and N400 responses for female
athletes (Figure 6).

Subconcussive analyses

Subconcussive N400 changes replicate four previous studies on subconcussion (Fickling et al.
2019; Fickling, Smith, et al. 2021; Fickling, Poel, et al. 2021; Breuer et al. 2023). The N400 is commonly
associated with high-order cognitive semantic processing (D'Arcy et al. 2004; Kutas and Federmeier 2011)
and measures changes in semantic processing, suggesting the initial demonstration of cognitive changes
due to contact (Connolly and D'Arcy 2000; D'Arcy et al. 2003; Gawryluk et al. 2010; Ghosh Hajra et al.
2018). It is noteworthy that these replications have spanned across different sports and age ranges in
male athletes. However, the specific N400 changes have varied with respect to latency delays and/or
amplitude reductions, with the former directly measuring the rate of information processing and the latter
related to synchronous pyramidal neural activation (Luck 2014). The two are interdependent factors
reflecting a relative impairment in cognitive processing. While the initial results for female athletes
showed comparable N400 latency delays, a specific pattern of change in N400 latency versus amplitude
remains to be characterized.

Similarly, subconcussive changes in the N100 have been replicated across a number of prior
studies. While the studies have reported only N100 amplitude changes, the direction of the effects has
varied (i.e., decreases, increases, and correlated variability (Fickling et al. 2019; Fickling, Smith, et al. 2021;
Fickling, Poel, et al. 2021). In prior linear regression analyses of the relationship between brain vital signs
and helmet mounted accelerometers, inclusion of the N100 with the N400 provided additional
explanatory variance (Fickling, Poel, et al. 2021). Consequently, N100 changes suggest that subconcussive
impairments can span from lower-level sensory processing to higher level cognitive processing. As with
the N400 above, the common preliminary results showing N100 changes in both female and male athletes
do not rule out the possibility of specific differences. Indeed, examination of the female and male
waveforms shows early sensory and attention differences in other ERP components (see below).

Of note, the current findings are the first to compare brain vital signs between high- and low-
contact groups using a between-subjects design. To anchor to prior studies, a direct evaluation of the
within-subjects pre- versus post- design has also demonstrated a significant delay in the N400 latency
(Supplemental Data 2). The contribution of a control group is an important step, with the interpretation
further underpinned by the nature of the closely matched comparison (i.e., related to questions about
whether the effects are indeed attributable to subconcussive impact exposure). The student athletes in
this study attended a private school with full time residence. Accordingly, all students follow a highly
structured daily and weekly schedule, matched across programs, activities, diet, sleep, physical fitness,
rest, and related factors.

Another novel aspect of the current study was the ability to begin exploring differences between
a single term versus two or more terms of exposure to subconcussive impacts (Supplemental Data 4).
While preliminary, the findings suggested that there may be a graded effect of exposure on the N400 peak
latency (Supplemental Data 4). These results highlighted important questions related to tracking the
extent of exposure with increased temporal resolution to characterize the changes over time. As reflected
in the study design (Figure 1), a large cohort of these students were also re-evaluated after the summer
break to determine whether this duration of rest resulted in a return to pre- baseline levels following post-
subconcussive changes. Evaluation of these results are currently being analyzed as a follow-up study.
While the emerging subconcussive findings are concerning, the demonstrated utility of using brain vital
signs to objectively identify acute cognitive changes associated with repeated impact exposure has

enabled clinical trials on potential interventions. A recent study on subconcussion focused on a
prospective randomized placebo-controlled trial evaluating the effect of a dietary supplement effects
created to reduce neuroinflammation and support energy production, on brain structure and function in
Junior-A ice hockey players (Breuer et al. 2023). Breuer et al. (2023) randomized players into two groups:
a group that took a supplement containing a combination of non-pharmacological ingredients shown to
support brain health (Synaquell™ - Thorne Research Inc., NY, USA) daily over the season and a placebo
control group who followed the same protocol without supplementation. Multivariate analyses showed
significant neurocognitive pre- to post- season changes in only the control group, with significant negative
impacts on the N100 and N400 along with reduced saccadic eye movement (King-Devick Test) and
neurofilament light chain measures.
304
Female and male comparison
Robust P300 and N400 differences between female and male athletes were shown with and
without subconcussive contact exposure as a factor. While the significant P300 and N400 differences
demonstrated larger amplitudes for female athletes, examination of the overall waveforms showed this
finding generally across all ERP responses. There are at least two potential factors related to interpretation
of the difference between female and male athletes. One factor may relate to general attention difference
during the approximately 6-minute test time, whereby female athletes showed increased sustained
attention on average relative to male athletes. In this case, improved sustained attention to the task
would result in a higher signal-to-noise ratio in the signal averaged ERPs and consequently higher response
amplitudes. Alternatively, another factor may relate to a fundamental difference in ERP response size
between female and male individuals. While these two factors are not mutually exclusive, the notable
differences in baseline variance suggested sustained attention during the test time is likely a factor (Figure
5). Differences in attention between female and male participants has been noted in prior literature, but
these results are still mixed (Davidson, Cave, and Sellner 2000; Solianik, Brazaitis, and Skurvydas 2016;
Riley et al. 2016).
As mentioned above, despite the subconcussive similarities between female and male athletes, it
is too early to exclude differences, as an "absence of evidence is not evidence of absence." For instance,
examination of the ERP waveforms (Figure 4) shows noteworthy subconcussive differences in responses
outside of the N100, P300, and N400. In particular, there appears to be a subconcussive reduction in the
N200-P200 difference in female athletes, which is not present in male athletes. The N200-P200 is
associated with early perceptual-attentional processing (Portella et al. 2012; D'Arcy, Connolly, and
Crocker 2000; Patel and Azzam 2005), and further research should explore this in the context of sex-
specific changes in subconcussion and head-impact exposure. With respect to this, the clinical importance
of even small attentional impairments can consequently have a larger impact on high-level processing for
individuals in attention demanding professions. For example, we identified early on in clinical concussion
evaluation that these changes can have a compounding effect on high demand jobs such as air traffic
controllers relative to less attentionally-demanding positions (Mateer and D'Arcy 2000). A comparable
impact may be a consideration for enhanced sustained attention processing in female athletes. Given
recent advances in understanding functional neuroanatomical differences (Andrushko et al. 2023), further
studies needed to better characterize female- versus male- specific subconcussive profiles.

Limitations and Future Directions

The current study included data across a large sample made up of smaller subsets across different
sports, with differing contact levels. While this was beneficial to create a large sample size for this study,
larger subsets of groups pertaining to each specific sport would help elucidate sport-specific differences.
Similarly, history of high-contact sport participation and concussion are important variables that may
affect the current state of participants. Nonetheless, the current study is the first in a continuing
investigation that will subsequently focus on specific sports and recovery times following subconcussive
impairment in contact sport. Future studies will expand to focus on recovery effects and differences
following summer break. Also, the current study combined subjects in grades 10-12 in a highly structured
school environment; however, previous work has shown differing neurological developmental rates
between female and male adolescents (Paus 2010; Kaczkurkin, Raznahan, and Satterthwaite 2019; Gur
and Gur 2016), which may have been a factor for female and male-specific profiles. Future work will
include larger age subset sizes and longitudinal analyses, as well as controlling for age as a co-variate in
future models, to have a better understanding of developmental differences between female and male
individuals. Finally, the current study did not track important biological factors that differ between female
and male athletes. These factors include but are not limited to menstrual cycle phase at time of exposure
or in the case of multiple exposures, when the majority of exposures occurred (menstrual phase has been
a predictor of mild TBI outcome (Wunderle et al. 2014) and progesterone shows neuroprotective as a TBI
treatment (Wright et al. 2007)); body fat percentages given recent understanding of fat being a major
endocrine organ (Musi and Guardado-Mendoza 2014); and diet given the understanding of diet impact
on inflammation and brain function (Patel et al. 2023; Markovic et al. 2021). With clear neurophysiological
differences identified between females and males, future work must include expanded characterization
of these key biological factors to understand change that may affect results at the time of evaluation.
Lastly, to better understand the resulting effect of these subconcussive neurophysiological changes and
differences, additional study variables should include heart rate variability data to understand clinical and
sub-clinical autonomic dysfunction and time to recovery, as well as changes in sleep and endocrine levels
due to RHI, as TBI may lead to endocrine changes, specifically in developing adolescents (Richmond and
Rogol 2014).

Conclusion

With increasing attention on subconcussive impairments and potential interventions to limit
future risks (e.g., long-term neuropsychiatric impairment, chronic traumatic encephalopathy – CTE), the
current findings provide a large sample demonstration of subconcussive changes between closely
matched high- versus low-contact student athletes that replicate previous investigations. The findings
provide strong support for continued focus on the role of subconcussive injuries in sport-related brain
health research. Importantly, as this field of study has historically focused on male athletes, inclusion of
female athletes in the current findings represents a critical milestone towards understanding specific
neurophysiological differences before and after exposure to subconcussive impacts. Initial results strongly
support continued investigations that span beyond male athletes in helmeted contact sports.

Methods

A total of 89 high school students were scanned 231 times over the year (34 females and 55
males). In total, 8 scans were removed due to attrition, 12 scans had a data collection error, failing to
produce scan data, 5 due to dropping all epochs for a condition (i.e. deviant tones), and 4 due to excess
noise (N=202 scans). For analysis and results we ended with N=81, n=202 (33 females and 48 males). The
current study focused on a scan-based between contact groups analysis, instead of a pre-post subject
based within-subjects analysis. I.e. results from a subject scan after a season of low contact sport would
be characterized as a low contact data point (not a benchmark or pre data point), while if that same
subject had a separate scan later in the school year after 2 terms of high contact, that scan would be
characterized as a high contact data point. This was done to maximize the total data point number for
analyses and provide more equal weighting across groups, as opposed to a repeated measure subject-
based analysis including only subjects that had not had any contact immediately prior to the study.
All healthy participants were in grades 10 through 12 (age range: 15-17), fluent in English, no
history of concussion in the past 6-months, no self-reported problems with brain function, and had no
hearing problems. All participants were recruited from Brentwood College School athletic programs,
which involve either high-contact (e.g., rugby, ice-hockey, soccer, field hockey, jujitsu, and basketball) or
no-contact/low-contact sports (e.g., rowing, tennis, and climbing) over three distinct three-month terms
in the school year (Fig. 1). As described above, Brentwood is a private boarding school in which all students
follow a structured daily schedule and a similar diet, exercise, and sleep schedule throughout the
academic year. Accordingly, participants were closely matched across a wide number of typical lifestyle
factors. The study had Advarra institutional research ethics board approval and all participants along with
their guardians provided informed consent.

Data Collection

The brain vital sign framework (Ghosh Hajra et al. 2016) was employed with the NeuroCatch®
Platform (Version 1.1), which uses EEG to rapidly extract the N100, P300, and N400 ERPs. All three ERPs
were stimulated using a compressed and standardized auditory stimulus sequence (approximately 6
minutes) consisting of sets of tones and spoken word pairs. The brain vital signs framework utilizes an
oddball paradigm, involving frequent standard tones with infrequent oddball tones, which elicit the N100
and P300 components. The tone stimuli precede word pairs, which are either matched or mismatched
semantically-related word-pair primes (e.g., match: bread-butter; mismatch: bread-window) to elicit the
N400 component. Participants passively listened to the stimuli and decided if the word pairs matched or
mismatched.
Participants were fitted with an elasticized cloth cap containing an 8-channel EEG amplifier with
standardized electrode locations (Fz, Cz, & Pz). Three additional electrodes were placed on the
participant's forehead to record the ground (GND) and electrooculogram (vEOG, hEOG) signals. A
reference electrode was clipped to the participant's right earlobe. Standard EEG skin preparation (70%
isopropyl alcohol wipes) and conductive gel products (SignaGel) were used to ensure appropriate contact
for all EEG sensors. Distractions were mitigated by performing the scans in a quiet and closed classroom.

The same environment was used for all scans. Participants were asked to listen attentively to the stimuli,
but no active response was required. To reduce motor and ocular artifacts, participants were instructed
to sit motionlessly, and maintain visual fixation on a cross positioned at eye level 2 m away.
419
420

Data Processing

The NeuroCatch® Platform was calibrated for any trigger latency delays. The most common delay
was identified and applied to shift all trigger data. EEG data were processed using a fourth order
Butterworth filter (0.5-10 Hz) and a Notch filter (60 Hz). Adaptive filtering using electrooculogram
channels was used to correct for ocular artifacts. EEG was epoched based on stimulus events using an
epoch from -100ms pre-stimulus to 1000ms post-stimulus. All epochs were linearly detrended and any
epochs with noise exceeding ±75μV were rejected. N100, P300, and N400 peaks were calculated using
the maximal peak within a standard temporal window and manually verified. Amplitude and latency
metrics from these peaks were then linearly transformed into standardized scores on a scale from 0 to
100, derived from entire group means (Fickling et al. 2019). All preprocessing was completed in Python
using the Scipy and MNE libraries.
432
433

Statistical Analysis

All individual-level statistical analyses of peak amplitude and latencies were performed using
SPSS (Version 29.0.0, IBM, NY, USA). In the current study, individual-level can be defined as amplitudes
and latencies from each individual scan data, not derived from the group-level average waveforms. To
assess differences in brain vital signs between contact groups and sex, brain vital sign scores were
compared using a multivariate analysis of variance (MANOVA) (**Between:** Group: low-contact and high-
contact; Sex: Female and Male) on N100, P300, and N400 amplitudes and latencies for only scans
characterized as low-contact or high-contact for two or more terms. To further assess differences in
brain vital signs between sex in a larger number of scans, brain vital sign scores were compared using a
multivariate analysis of variance (MANOVA) (**Between:** Sex: Female and Male) on N100, P300, and N400
amplitudes and latencies for all 202 scans collected. All univariate results are adjusted for multiple
comparisons using Bonferroni adjustment. All statistics were also rerun only with subjects with less than
50% of epochs dropped (12 of 202 scans dropped) due to the 75μV thresholding measures mentioned
above. To keep our participant number large and because the analysis resulted in very similar outcomes,
the current manuscript reports the statistics from the full group data, including subjects with more than
50% of epochs dropped.
450
451

Acknowledgements

454   The authors would like to acknowledge the important leadership of Dr. Aynsley Smith (Mayo
455   Clinic, retired), whose passionate leadership for research advances in sports-related concussion and
456   ever-lasting priority to translate research advances to young females and males served as an important
457   inspiration for this study. The authors would also like to acknowledge the technical assistance by Joshua
458   Ighalo and the assistance in data collection by Laila Reed.
459

Funding

461   This work has been supported in part by funding from the Natural Sciences and Engineering
462   Research Council of Canada (Discovery Grant, PI R.C.N. D'Arcy; CGS M, E.D. Kirby).
463

Competing interests

Some of the authors (EDK, TY, NYB, SDF, RCND) have financial and/or business interests in
HealthTech Connex Inc., which may be affected by the research reported in this paper. Over the past 36
months, DWD reports the following competing interests: Consulting: Amgen, Atria, CapiThera Ltd.,
Cerecin, Ceruvia Lifesciences LLC, CoolTech, Ctrl M, Allergan, AbbVie, Biohaven, GlaxoSmithKline,
Lundbeck, Eli Lilly, Novartis, Impel, Satsuma, Theranica, WL Gore, Genentech, Nocira, Perfood, Praxis,
AYYA Biosciences, Revance, Pfizer. Honoraria: American Academy of Neurology, Headache Cooperative
of the Pacific, Canadian Headache Society, MF Med Ed Research, Biopharm Communications, CEA Group
Holding Company (Clinical Education Alliance LLC), Teva (speaking), Amgen (speaking), Eli Lilly
(speaking), Lundbeck (speaking), Pfizer (speaking), Vector Psychometric Group, Clinical Care Solutions,
CME Outfitters, Curry Rockefeller Group, DeepBench, Global Access Meetings, KLJ Associates, Academy
for Continued Healthcare Learning, Majallin LLC, Medlogix Communications, Medica Communications
LLC, MJH Lifesciences, Miller Medical Communications, WebMD Health/Medscape, Wolters Kluwer,
Oxford University Press, Cambridge University Press. Non-profit board membership: American Brain
Foundation, American Migraine Foundation, ONE Neurology, Precon Health Foundation, International
Headache Society Global Patient Advocacy Coalition, Atria Health Collaborative, Arizona Brain Injury
Alliance, Domestic Violence HOPE Foundation/Panfila. Research support: Department of Defense,
National Institutes of Health, Henry Jackson Foundation, Sperling Foundation, American Migraine
Foundation, Henry Jackson Foundation, Patient Centered Outcomes Research Institute (PCORI). Stock
options/shareholder/patents/board of directors: Aural analytics (options), Axon Therapeutics, ExSano
(options), Palion (options), Man and Science, Healint (options), Theranica (options), Second
Opinion/Mobile Health (options), Epien (options/board), Nocira (options), Matterhorn (shares/board),
Ontologics (shares/board), King-Devick Technologies (options/board), Precon Health (options/board),
AYYA Biosciences (options), Axon Therapeutics (options/board), Cephalgia Group (options/board), Atria
Health (options/employee). Patent 17189376.1-1466:vTitle: Onabotulinum Toxin Dosage Regimen for
Chronic Migraine Prophylaxis (Non-royalty bearing). Patent application submitted: Synaquell® (Precon
Health).
491

Author Contributions

RCND assisted with project study design, manuscript drafting, and acted as lab PI. DM

contributed with data collection and project study design. DH contributed with data collection, data

processing, and manuscript editing. ZL contributed with data collection and manuscript editing. JW

contributed with data collection and manuscript editing. AH contributed with data collection, statistical

analysis, and manuscript editing. EDK contributed with data processing, statistical analysis, and

manuscript drafting. TY, NYB, SDF, TAM, DD, CA, and KSS contributed with manuscript editing.

501 **References**

Andrushko, Justin W., Shie Rinat, Eric D. Kirby, Julia Dahlby, Chelsea Ekstrand, and Lara A.
Boyd. 2023. "Females Exhibit Smaller Volumes of Brain Activation and Lower Inter-
Subject Variability during Motor Tasks." *Scientific Reports* 13 (1): 17698.
https://doi.org/10.1038/s41598-023-44871-4.
Breuer, Logan T., Shaun D. Fickling, David A. Krause, Ryan C. N. D'Arcy, Tory O. Frizzell,
Tania Gendron, and Michael J. Stuart. 2023. "The Effects of a Dietary Supplement on
Brain Function and Structure in Junior A Ice Hockey Players: A Prospective Randomized
Trial." medRxiv. https://doi.org/10.1101/2023.04.03.23288079.
Connolly, John F, and Ryan C. N D'Arcy. 2000. "Innovations in Neuropsychological Assessment
Using Event-Related Brain Potentials." *International Journal of Psychophysiology* 37 (1):
31–47. https://doi.org/10.1016/S0167-8760(00)00093-3.
D'Arcy, Ryan C. N, John F Connolly, and Susan F Crocker. 2000. "Latency Shifts in the N2b
Component Track Phonological Deviations in Spoken Words." *Clinical Neurophysiology*
111 (1): 40–44. https://doi.org/10.1016/S1388-2457(99)00210-2.
D'Arcy, Ryan C. N, Yannick Marchand, Gail A Eskes, Edmund R Harrison, Stephen J Phillips,
Alma Major, and John F Connolly. 2003. "Electrophysiological Assessment of Language
Function Following Stroke." *Clinical Neurophysiology* 114 (4): 662–72.
https://doi.org/10.1016/S1388-2457(03)00007-5.
D'Arcy, Ryan C.N., John F. Connolly, Elisabet Service, Colin S. Hawco, and Michael E.
Houlihan. 2004. "Separating Phonological and Semantic Processing in Auditory
Sentence Processing: A High-Resolution Event-Related Brain Potential Study." *Human*
Brain Mapping. https://doi.org/10.1002/hbm.20008.
Davidson, Heather, Kyle R Cave, and Daniela Sellner. 2000. "Differences in Visual Attention
and Task Interference between Males and Females Reflect Differences in Brain
Laterality." *Neuropsychologia* 38 (4): 508–19. https://doi.org/10.1016/S0028-
3932(99)00084-6.
Davis, P. A. 1939. "Effects of Acoustic Stimuli on the Waking Human Brain." *Journal of*
Neurophysiology 2 (6): 494–99. https://doi.org/10.1152/jn.1939.2.6.494.
Dioso, Emma, John Cerillo, Mohammed Azab, Devon Foster, Isaac Smith, Owen Leary, Michael
Goutnik, and Brandon Lucke-Wold. 2022. "Subconcussion, Concussion, and Cognitive
Decline: The Impact of Sports Related Collisions." *Journal of Medical Research and*
Surgery 3 (4): 54–63. https://doi.org/10.52916/jmrs224081.
Fickling, Shaun D, Daniel N Poel, Jason C Dorman, Ryan C N D'Arcy, and Thayne A Munce.
2021. "Subconcussive Changes in Youth Football Players: Objective Evidence Using
Brain Vital Signs and Instrumented Accelerometers." *Brain Communications* 4 (2):
fcab286. https://doi.org/10.1093/braincomms/fcab286.
Fickling, Shaun D., Aynsley M. Smith, Gabriela Pawlowski, Sujoy Ghosh Hajra, Careesa C. Liu,
Kyle Farrell, Janelle Jorgensen, Xiaowei Song, Michael J. Stuart, and Ryan C.N. D'Arcy.

541   2019. "Brain Vital Signs Detect Concussion-Related Neurophysiological Impairments in
542   Ice Hockey." *Brain* 142 (2): 255–62. https://doi.org/10.1093/brain/awy317.
543 Fickling, Shaun D, Aynsley M Smith, Michael J Stuart, David W Dodick, Kyle Farrell, Sara C
544   Pender, and Ryan C N D'Arcy. 2021. "Subconcussive Brain Vital Signs Changes Predict
545   Head-Impact Exposure in Ice Hockey Players." *Brain Communications* 3 (2): fcab019.
546   https://doi.org/10.1093/braincomms/fcab019.
547 Gawryluk, Jodie R., Ryan C.N. D'Arcy, John F. Connolly, and Donald F. Weaver. 2010.
548   "Improving the Clinical Assessment of Consciousness with Advances in
549   Electrophysiological and Neuroimaging Techniques." *BMC Neurology*.
550   https://doi.org/10.1186/1471-2377-10-11.
551 Ghosh Hajra, Sujoy, Careesa C. Liu, Xiaowei Song, Shaun D. Fickling, Teresa P. L. Cheung,
552   and Ryan C. N. D'Arcy. 2018. "Multimodal Characterization of the Semantic N400
553   Response within a Rapid Evaluation Brain Vital Sign Framework." *Journal of*
554   *Translational Medicine* 16 (June): 151. https://doi.org/10.1186/s12967-018-1527-2.
555 Ghosh Hajra, Sujoy, Careesa C. Liu, Xiaowei Song, Shaun Fickling, Luke E. Liu, Gabriela
556   Pawlowski, Janelle K. Jorgensen, et al. 2016. "Developing Brain Vital Signs: Initial
557   Framework for Monitoring Brain Function Changes Over Time." *Frontiers in*
558   *Neuroscience* 10. https://www.frontiersin.org/article/10.3389/fnins.2016.00211.
559 Gur, Raquel E., and Ruben C. Gur. 2016. "Sex Differences in Brain and Behavior in
560   Adolescence: Findings from the Philadelphia Neurodevelopmental Cohort."
561   *Neuroscience & Biobehavioral Reviews*, The Adolescent Brain, 70 (November): 159–70.
562   https://doi.org/10.1016/j.neubiorev.2016.07.035.
563 Guskiewicz, Kevin M., Stephen W. Marshall, Julian Bailes, Michael McCrea, Robert C. Cantu,
564   Christopher Randolph, and Barry D. Jordan. 2005. "Association between Recurrent
565   Concussion and Late-Life Cognitive Impairment in Retired Professional Football
566   Players." *Neurosurgery* 57 (4): 719.
567   https://doi.org/10.1227/01.NEU.0000175725.75780.DD.
568 Hirad, Adnan A., Jeffrey J. Bazarian, Kian Merchant-Borna, Frank E. Garcea, Sarah
569   Heilbronner, David Paul, Eric B. Hintz, et al. 2019. "A Common Neural Signature of Brain
570   Injury in Concussion and Subconcussion." *Science Advances* 5 (8): eaau3460.
571   https://doi.org/10.1126/sciadv.aau3460.
572 Kaczkurkin, Antonia N., Armin Raznahan, and Theodore D. Satterthwaite. 2019. "Sex
573   Differences in the Developing Brain: Insights from Multimodal Neuroimaging."
574   *Neuropsychopharmacology* 44 (1): 71–85. https://doi.org/10.1038/s41386-018-0111-z.
575 Kutas, Marta, and Kara D. Federmeier. 2011. "Thirty Years and Counting: Finding Meaning in
576   the N400 Component of the Event-Related Brain Potential (ERP)." *Annual Review of*
577   *Psychology*. https://doi.org/10.1146/annurev.psych.093008.131123.
578 Luck, Steven J. 2014. *An Introduction to the Event-Related Potential Technique, Second*
579   *Edition*. MIT Press.
580 Mainwaring, Lynda, Kaleigh M. Ferdinand Pennock, Sandhya Mylabathula, and Benjamin Z.
581   Alavie. 2018. "Subconcussive Head Impacts in Sport: A Systematic Review of the
582   Evidence." *International Journal of Psychophysiology*, Long-term outcomes of sport-
583   related brain injuries: A psychophysiological perspective, 132 (October): 39–54.
584   https://doi.org/10.1016/j.ijpsycho.2018.01.007.
585 Markovic, Shaun J., Melinda Fitzgerald, Jeremiah J. Peiffer, Brendan R. Scott, Stephanie R.
586   Rainey-Smith, Hamid R. Sohrabi, and Belinda M. Brown. 2021. "The Impact of Exercise,
587   Sleep, and Diet on Neurocognitive Recovery from Mild Traumatic Brain Injury in Older
588   Adults: A Narrative Review." *Ageing Research Reviews* 68 (July): 101322.
589   https://doi.org/10.1016/j.arr.2021.101322.

Mateer, Catherine A., and Ryan C. N. D'Arcy. 2000. "Current Concepts in Assessment and
Management." In *Neuropsychological Management of Mild Traumatic Brain Injury*, edited
by Sarah A. Raskin and Catherine A. Mateer. New York, NY, USA: Oxford University
Press.
McKee, Ann C., Robert C. Cantu, Christopher J. Nowinski, E. Tessa Hedley-Whyte, Brandon E.
Gavett, Andrew E. Budson, Veronica E. Santini, Hyo-Soon Lee, Caroline A. Kubilus, and
Robert A. Stern. 2009. "Chronic Traumatic Encephalopathy in Athletes: Progressive
Tauopathy After Repetitive Head Injury." *Journal of Neuropathology & Experimental
Neurology* 68 (7): 709–35. https://doi.org/10.1097/NEN.0b013e3181a9d503.
Miller, Jeff, Rolf Ulrich, and Wolfgang Schwarz. 2009. "Why Jackknifing Yields Good Latency
Estimates." *Psychophysiology* 46 (2): 300–312. https://doi.org/10.1111/j.1469-
8986.2008.00761.x.
Musi, Nicolas, and Rodolfo Guardado-Mendoza. 2014. "Chapter 14 - Adipose Tissue as an
Endocrine Organ." In *Cellular Endocrinology in Health and Disease*, edited by Alfredo
Ulloa-Aguirre and P. Michael Conn, 229–37. Boston: Academic Press.
https://doi.org/10.1016/B978-0-12-408134-5.00014-7.
Nowak, Madeleine K., Zachary W. Bevilacqua, Keisuke Ejima, Megan E. Huibregtse, Zhongxue
Chen, Timothy D. Mickleborough, Sharlene D. Newman, and Keisuke Kawata. 2020.
"Neuro-Ophthalmologic Response to Repetitive Subconcussive Head Impacts: A
Randomized Clinical Trial." *JAMA Ophthalmology* 138 (4): 350–57.
https://doi.org/10.1001/jamaophthalmol.2019.6128.
Patel, Palak R., Patrick Armistead-Jehle, Nicholas R. Eltman, Kelly M. Heath, David X. Cifu, and
Randel L. Swanson. 2023. "Brain Injury: How Dietary Patterns Impact Long-Term
Outcomes." *Current Physical Medicine and Rehabilitation Reports* 11 (3): 367–76.
https://doi.org/10.1007/s40141-023-00413-7.
Patel, Salil H., and Pierre N. Azzam. 2005. "Characterization of N200 and P300: Selected
Studies of the Event-Related Potential." *International Journal of Medical Sciences* 2 (4):
147–54.
Paus, Tomáš. 2010. "Chapter 2 - Sex Differences in the Human Brain: A Developmental
Perspective." In *Progress in Brain Research*, edited by Ivanka Savic, 186:13–28. Sex
Differences in the Human Brain, Their Underpinnings and Implications. Elsevier.
https://doi.org/10.1016/B978-0-444-53630-3.00002-6.
Portella, Claudio, Sergio Machado, Oscar Arias-Carrión, Alexander T. Sack, Julio Guilherme
Silva, Marco Orsini, Marco Antonio Araujo Leite, et al. 2012. "Relationship Between
Early and Late Stages of Information Processing: An Event-Related Potential Study."
Neurology International 4 (3): e16. https://doi.org/10.4081/ni.2012.e16.
Richmond, Erick, and Alan D. Rogol. 2014. "Traumatic Brain Injury: Endocrine Consequences in
Children and Adults." *Endocrine* 45 (1): 3–8. https://doi.org/10.1007/s12020-013-0049-1.
Riley, Elizabeth, Hidefusa Okabe, Laura Germine, Jeremy Wilmer, Michael Esterman, and
Joseph DeGutis. 2016. "Gender Differences in Sustained Attentional Control Relate to
Gender Inequality across Countries." *PLoS ONE* 11 (11): e0165100.
https://doi.org/10.1371/journal.pone.0165100.
Saghafi, Behrouz, Gowtham Murugesan, Elizabeth Davenport, Ben Wagner, Jillian Urban,
Mireille Kelley, Derek Jones, et al. 2018. "Quantifying the Association between White
Matter Integrity Changes and Subconcussive Head Impact Exposure from a Single
Season of Youth and High School Football Using 3D Convolutional Neural Networks."
Proceedings of SPIE--the International Society for Optical Engineering 10575
(February): 105750E. https://doi.org/10.1117/12.2293023.
Smulders, Fren T. Y. 2010. "Simplifying Jackknifing of ERPs and Getting More out of It:
Retrieving Estimates of Participants' Latencies." *Psychophysiology* 47 (2): 387–92.

https://doi.org/10.1111/j.1469-8986.2009.00934.x.
Snedaker, Katherine Price, Jennifer P. Lundine, Angela H. Ciccia, Mohammad Nadir Haider,
and Katy H. O'Brien. 2022. "Gaps in Concussion Management across School-Aged
Children." *Brain Injury* 36 (6): 714–21. https://doi.org/10.1080/02699052.2022.2034954.
Solianik, Rima, Marius Brazaitis, and Albertas Skurvydas. 2016. "Sex-Related Differences in
Attention and Memory." *Medicina* 52 (6): 372–77.
https://doi.org/10.1016/j.medici.2016.11.007.
Sutton, Samuel, Patricia Tueting, Joseph Zubin, and E. R. John. 1967. "Information Delivery
and the Sensory Evoked Potential." *Science* 155 (3768): 1436–39.
https://doi.org/10.1126/science.155.3768.1436.
Ulrich, Rolf, and Jeff Miller. 2001. "Using the Jackknife-Based Scoring Method for Measuring
LRP Onset Effects in Factorial Designs." *Psychophysiology* 38 (5): 816–27.
https://doi.org/10.1111/1469-8986.3850816.
Valera, Eve M., Annie-Lori C. Joseph, Katherine Snedaker, Matthew J. Breiding, Courtney L.
Robertson, Angela Colantonio, Harvey Levin, et al. 2021. "Understanding Traumatic
Brain Injury in Females: A State-of-the-Art Summary and Future Directions." *The Journal*
of Head Trauma Rehabilitation 36 (1): E1–17.
https://doi.org/10.1097/HTR.0000000000000652.
Veksler, Ronel, Udi Vazana, Yonatan Serlin, Ofer Prager, Jonathan Ofer, Nofar Shemen,
Andrew M Fisher, et al. 2020. "Slow Blood-to-Brain Transport Underlies Enduring Barrier
Dysfunction in American Football Players." *Brain* 143 (6): 1826–42.
https://doi.org/10.1093/brain/awaa140.Markovic, Shaun J., Melinda Fitzgerald, Jeremiah
J. Peiffer, Brendan R. Scott, Stephanie R. Rainey-Smith, Hamid R. Sohrabi, and Belinda
M. Brown. 2021. "The Impact of Exercise, Sleep, and Diet on Neurocognitive Recovery
from Mild Traumatic Brain Injury in Older Adults: A Narrative Review." *Ageing Research*
Reviews 68 (July): 101322. https://doi.org/10.1016/j.arr.2021.101322.
Wright, David W., Arthur L. Kellermann, Vicki S. Hertzberg, Pamela L. Clark, Michael Frankel,
Felicia C. Goldstein, Jeffrey P. Salomone, et al. 2007. "ProTECT: A Randomized Clinical
Trial of Progesterone for Acute Traumatic Brain Injury." *Annals of Emergency Medicine*
49 (4): 391-402.e2. https://doi.org/10.1016/j.annemergmed.2006.07.932.
Wunderle, Kathryn, Kathleen M. Hoeger, Erin Wasserman, and Jeffrey J. Bazarian. 2014.
"Menstrual Phase as Predictor of Outcome After Mild Traumatic Brain Injury in Women."
The Journal of Head Trauma Rehabilitation 29 (5): E1.
https://doi.org/10.1097/HTR.0000000000000006.
674