Machine Learning Model Utilization for Mortality Prediction in Mechanically Ventilated ICU Patients

Yiming Liu[a], Xin Xu[a], Yingqi Wang[a], Zhiyuan Zhao[a] and Maryam Pishgar[a,*]

[a]*Andrew and Erna Viterbi School of Engineering, University of Southern California (USC), 3650 McClintock Ave, Los Angeles, CA 90089, United States of America*

ARTICLE INFO

Keywords:
Mechanical ventilation
ICU mortality
Deep learning
Machine learning

ABSTRACT

Background: The requirement of mechanical ventilation has increased in recent years. Patients in the intensive care unit (ICU) who undergo mechanical ventilation often experience serious illness and critical physiological states, contributing to a high risk of mortality. Prediction of mortality for mechanically ventilated ICU patients offers an early alert mechanism to help physicians implement more targeted treatments to mitigate the risk of mortality.

Methods: We extracted medical information of patients with invasive mechanical ventilation in ICU admission from the Medical Information Mart for Intensive Care III (MIMIC III) dataset. This includes demographics, disease severity, diagnosis, and laboratory test results. Patients who met inclusion criteria were randomly divided into the train set (n=11,549, 70%), the test set (n=2,475, 15%), and the validation set (n=2,475, 15%). The oversampling method was utilized to resolve the imbalance dataset. After literature research, clinical expertise and ablation study, we selected 12 variables for model establishment, which were less than the 66 features in the best existing literature. We proposed a deep learning neural network model to predict the ICU mortality of mechanically ventilated patients, and established 7 baseline machine learning (ML) models for comparison, including k-nearest neighbors (KNN), logistic regression, decision tree, random forest, bagging, XGBoost, and support vector machine (SVM). Area under the Receiver Operating Characteristic Curve (AUROC) was used as an evaluation metric for model performance.

Results: By using 16,499 mechanically ventilated patients in the MIMIC III database, the deep learning model demonstrated the best performance, showing a 7.06% improvement over the best existing literature. The neural network achieved an AUROC score of 0.879 with 95% Confidence Interval (CI) [0.861-0.896] and an accuracy score of 0.859 on the test set. These results far exceeded those of the best existing literature.

Conclusions: The proposed model demonstrated an exceptional ability to predict ICU mortality among mechanically ventilated patients. Through SHAP analysis, we found that respiratory failure is a significant indicator of the mortality prediction compared to other related respiratory dysfunction diseases. We also incorporated mechanical ventilation duration as a variable for the first time in our prediction model. We observed that patients with higher mortality rates tended to have longer mechanical ventilation times. This highlights the model's potential in guiding clinical decisions by indicating that longer mechanical ventilation may not necessarily enhance patient survival chances.

1. BACKGROUND

An intensive care unit (ICU) is designated for individuals facing severe illnesses or injuries. Most of these patients require assistance from medical equipment, such as mechanical ventilation, to sustain regular bodily functions, and they need to be monitored continuously and intensively [1, 2, 3].

In the ICU, more than 25% of patients require mechanical ventilation [4]. The use of invasive mechanical ventilation is associated with a high risk of mortality and various complications [5]. Mechanical ventilation is a key method for supporting organ function in ICU patients [6]. In the United States, approximately 40% of ICU patients receive support for invasive mechanical ventilation at any given time, contributing to 12% of overall hospital expenditures [7]. Mortality rates are notably high among those requiring mechanical ventilation [8].

With the growing life expectancies and the survival time extension of individuals with chronic conditions, the utilization of mechanical ventilation for artificial support is anticipated to increase [9, 10]. Mechanically ventilated patients usually experience acute respiratory failure or reduced lung function due to an underlying condition, such as pneumonia, sepsis, or heart disease [11, 12, 13]. Alternatively, the need for respiratory assistance may arise from neurological disability, disorder consciousness, or fatigue after significant surgical procedures [14].

In recent years, machine learning algorithms have been widely employed to predict diverse critical health outcomes [15, 16], more specially associated with mechanical ventilation [17, 18]. Developing a mortality prediction model for patients with mechanical ventilation may offer valuable support to ICU physicians for timely alerts and informed clinical judgment [19].

Neural network modeling has gained widespread recognition for its effectiveness and it has become powerful tools for sophisticated modeling in various domains [20]. Neural networks employ a multi-layered structure to autonomously generate distinctive features, and convert input features

---

*Corresponding author.
E-mail address:* pishgar@usc.edu (Maryam Pishgar).

through activation functions. As a result, neural networks demonstrate a distinct advantage over foundational machine learning models in capturing nonlinearities, particularly when an abundance of data is accessible [21, 22].

The primary aim of this research was to establish a deep learning model designed for forecasting the mortality of ICU patients undergoing mechanical ventilation, utilizing comprehensive patient medical history data. We used fewer predictive variables yet more accurate results, which achieved a higher evaluation performance than the best existing literature. The inclusion of additional variables CHF and respiratory failure significantly enhanced the results of our proposed model. The predictive model was achieved based on guidelines of the Transparent Reporting of Individual Prognostic or Diagnostic Multivariate Predictive Model (TRIPOD) initiative.

2. METHODOLOGY

2.1. Data Source and Study Design

The Medical Information Mart for Intensive Care (MIMIC-III) database, which is a medical information sharing initiative containing rich clinical patient data, was used in our study. Specific data, including patients' clinical physiological parameters and disease diagnosis reports, were extracted from the database to cover specific patient cohorts. The reason for selecting the MIMIC-III database was its provision of a substantial amount of real-world patient information that contributes to a more comprehensive understanding of the research questions and hypotheses. Following the completion of data extraction, necessary data preprocessing was conducted to ensure data quality and alignment with the requirements for model training. The data from the MIMIC-III database provided a crucial foundation for our study, offering robust support for in-depth analysis and model construction, benefiting medical institutions and researchers.

2.2. Patient extraction

Our study focused on adult patients who underwent invasive mechanical ventilation during their ICU stay. Figure 1 shows the patient extraction process. First, we selected 61,532 patients with ICU stays records and extracted 28,861 patients whose records indicated a ventilation duration greater than 0. Among those patients, we excluded patients under 18 or over 90 years of age upon ICU admission, as well as patients with missing records of relevant physiological indicators. Then we cleaned the dataset by removing the missing values in the dataset. In the end, we extracted a total 16,499 patients who met the established inclusion criteria for the final analyses.

2.3. Statistical analysis between cohorts

The train and validation cohorts were compared using Chi-Square tests and two-sided t-tests with a significance level of $P < 0.05$ to help determine whether there are significant differences between the train set and the validation set. Chi-Square tests were utilized for comparing categorical variables, while t-tests were employed for continuous variables.

Figure 1: Flow diagram of the selection process of patients.

2.4. Feature selection

We started with 65 variables based on literature research and expert opinion. 14 key predictors were selected for subsequent analysis. The subject IDs and ICU stay IDs serve as the unique identifiers for patients and records of ICU admission, respectively. All physiological test indicators and disease diagnoses were referred to ICD-9 codes. Table 1 illustrates the proposed 14 predictors, including: (i) age: patients' age when entering the ICU; (ii) respiratory dysfunction: all diseases related to 'respiratory' in the diagnostic table; (iii) SAPS II: Simplified Acute Physiology Score II; (iv) maximum hemoglobin: patients' maximum value of blood hemoglobin in the lab events records; (v) minimum lactate: patients' minimum levels of lactate in the lab events records; (vi) respiratory failure: patients diagnosed with respiratory failure; (vii) minimum BUN: patients' minimum levels of blood urea nitrogen in the lab events records; (viii) CHF: patients diagnosed with chronic heart failure; (ix) diabetes: patients diagnosed with diabetes; (x) Malignancy: patients diagnosed with malignancy; (xi) maximum $PaCO_2$: patients' maximum levels of partial pressure of carbon dioxide in the arterial blood; (xii) minimum $PaCO_2$: patients' minimum levels of partial pressure of carbon dioxide in the arterial blood; (xiii) vent duration: the duration of invasive mechanical ventilation; (xiv) maximum PaO_2: patients' maximum levels of partial pressure of oxygen in the arterial blood.

First, we applied the XGBoost model to get the feature importance of these variables and we selected the top 5 important features: age, respiratory dysfunction, SAPS II score, maximum hemoglobin, and minimum lactate. The XGBoost is a powerful ML model which has been used widely for feature selection [19, 23]. This model includes regularization parameters such as 'gamma', 'alpha', and 'lambda' that help prevent overfitting, ensuring that the

Table 1

Features category table. Demographic: Age. Disease severity: SAPS II. Diagnosis: Respiratory dysfunction, Respiratory failure, CHF, Diabetes, Malignancy. Laboratory results: Maximum hemoglobin (g/dl), Minimum lactate (mmol/L), Minimum BUN (mg/dl), Minimum PaO2(mmHg), Maximum PaCO2(mmHg), Minimum PaCO2(mmHg). Others: Vent Duration (Hour).

Category	Features	Category	Features
Demographic	Age (years)	Laboratory results	Maximum hemoglobin (g/dl)
Disease severity	SAPS II		Minimum lactate (mmol/L)
Diagnosis	Respiratory dysfunction		Minimum BUN (mg/dl)
	Respiratory failure		Minimum PaO2(mmHg)
	CHF		Maximum PaCO2(mmHg)
	Diabetes		Minimum PaCO2(mmHg)
	Malignancy	Others	Vent Duration (Hour)

selected features are not chosen purely based on noise in the data.

After conducting related research and considering expert opinion, we also added other possible influencing factors: malignancy, BUN (Blood urea nitrogen), CHF (Congestive heart failure), diabetes, vent duration, respiratory failure, maximum PaCO2, minimum PaCO2, and maximum PaO2 as our variables. We decided to include these factors because their impact on respiratory health is well-documented and have the potential to significantly impact outcomes for patients requiring mechanical ventilation.

Malignancies can directly affect the respiratory system, such as lung cancer or metastases to the lungs, leading to compromised lung function. Especially in advanced stages, it can be a crucial factor leading to respiratory failure and the need for mechanical ventilation [24]. BUN is a medical test that measures the amount of urea nitrogen found in blood. High BUN levels can be associated with conditions that may lead to respiratory failure, such as severe infections, sepsis, or organ dysfunction. The decision to use mechanical ventilation is based on a combination of factors, including the underlying condition causing the high BUN level [25]. CHF can cause pulmonary effusion (pulmonary edema), leading to severe breathing difficulties and respiratory failure, thus requiring the use of mechanical ventilation to support breathing [23]. Diabetes can contribute to conditions such as respiratory infections, acute respiratory distress syndrome (ARDS), or other respiratory complications that may lead to the need for mechanical ventilation. Ventilator duration is a critical factor in the management of patients requiring mechanical ventilation, particularly in ICU. The use of a ventilator can prolong the lifespan of patients, and the duration of ventilator use may also be a factor that can affect the outcome. During mechanical ventilation, the goal is to maintain adequate gas exchange and ensure that the patient's blood is fully oxygenated, which includes removing carbon dioxide (CO2) from the body. If the PaCO2 levels are too high (hypercapnia) or too low (hypocapnia), it indicates that the patient is not effectively ventilating, which can lead to respiratory acidosis and potentially life-threatening

complications. Monitoring the maximum PaO2 level helps doctors evaluate the oxygenation of blood in a patient's lungs and whether adjustments to the ventilator settings are necessary to maintain optimal oxygen levels [26]. We also considered patients with various respiratory system diseases specifically in respiratory dysfunction from the MIMIC III database.

2.5. Ablation process

We planned to determine if the currently selected 14 features would negatively impact the model's performance. We decided to progressively eliminate variables that had a negative effect on the model's performance, assessing the model's performance on the validation set by calculating the 95% CI for AUROC. We sequentially removed one variable at a time and assessed the resulting degradation in model performance. The variable that causes the most significant deterioration in model performance is identified and removed in each round. This process was repeated until further removal of variables does not result in a noticeable improvement in model performance. This approach allows us to filter out variables that do not contribute significantly to the predictive power of the model. After applying this iterative feature selection process, we retained 12 out of the 14 initially selected variables. We excluded the variables malignancy and respiratory dysfunction from the final set of features, as they were found to have a negative impact on the model's performance.

2.6. Modeling

The dataset was imbalanced between the number of survivors and non-survivors, with 14,441 numbers of survivors and 2,058 non-survivors. The Synthetic Minority Oversampling Technique (SMOTE) method was used to address the data imbalance issue. Moreover, the train_test_split method was utilized for hierarchical stratified sampling. The dataset was split into three groups: train set, test set and validation set. We proposed a novel deep learning neural network to predict the mortality of ICU patients with mechanical ventilation. Seven baseline ML models were established for result comparison, including KNN, logistic

regression, decision tree, random forest, bagging, XGBoost, SVM[27, 28, 29, 30, 31, 32, 33, 34].

The proposed model is a fully connected neural network comprising an input layer with a dimensionality of 12, followed by a batch normalization (BN) layer for input normalization to improve the stability of the model [35]. Subsequently, three hidden layers are incorporated, each utilizing the rectified linear unit (ReLU) activation function. Between these hidden layers, dropout (DP) layers are employed to randomly discard 50% of the neurons, mitigating overfitting [36]. The first hidden layer consists of 100 neurons, the second hidden layer consists of 50 neurons, and the third hidden layer consists of 25 neurons. The model concludes with an output layer containing a single neuron, utilizing the sigmoid activation function for binary classification with an output probabilities between 0 and 1. This architecture is designed to capture complex patterns in the data while addressing potential overfitting through the strategic use of dropout layers. Figure 2 shows the architecture of our NN model. The model is trained with the Adam optimizer, using binary cross-entropy as the loss function and the AUROC as the evaluation criterion. The training process is trained for 100 epochs with a batch size of 256. The model iteratively refines its parameters to minimize the loss function and enhance AUROC performance, aiming to improve its ability to discriminate between positive and negative instances.

Figure 2: Neural network architecture op-level. This figure shows the details of the neural network architecture.

For the KNN model, we performed Grid Search CV to find the optimal n_neighbors parameter within the range of 1 to 20. The cross-validation (cv) was set to 5, with AUROC employed as the evaluation metric. Regarding the Logistic Regression model, the maximum iteration was set to 1000, and 'liblinear' was chosen as the solver. We utilized Grid Search CV across all baseline ML models to identify the best hyperparameter values, optimizing the performance of each prediction model. This process automated the optimization of hyperparameters.

The best model was chosen based on its performance in AUROC on the validation set. Also, we calculated accuracy for evaluating our models' performance.

In our project, we extracted the dataset using BigQuery, performed data cleaning and conducted model training using Python 3.9.17. The models were derived from the Python libraries scikit-learn 1.2.2 and TensorFlow 2.14.0.

3. RESULTS

3.1. *Cohort Comparison*

We obtained 16,499 patients from the MIMIC-III database for model establishment in the patient extraction part. The cohort was then randomly split into a 70% train set, 15% test set, and 15% validation set, respectively, allocating 11,549 patients to the train set, 2,475 patients to the test set, and 2,475 patients to the validation set. The train and validation cohorts were used to train the models. The model with the highest AUROC value was chosen as the best prediction model which was utilized for further assessment on the test set. Table 2 illustrates the comparison of the train set and validation set. In terms of respiratory failure, the proportion of patients (36%) was slightly greater than the validation cohort (35%). The train set (34%) also had slightly higher the share of patients with diabetes than the validation set (33%). However, the p-values of respiratory failure and diabetes were 0.380 and 0.248, respectively, showing no significant difference between cohorts. Moreover, there was no significant difference among all variables between cohorts based on their p-values.

In addition, a detailed comparison between the survivors group and the non-survivors group was presented in Table 3. Diagnosis and target variables were displayed on the number of diagnosed patients or non-survivors with their proportion, other variables were calculated on median with standard deviation. The p-values between two subgroups were calculated using the t-test, with the significant level set as $P < 0.05$. All variables had significant differences between the two groups, indicating a higher association with mortality, except malignancy.
'

3.2. *Ablation Study on Variable*

In our validation set, we evaluated the impact of each variable on the model performance by dropping one at a time, measured by AUROC. Firstly, we collected 65 variables from related literature research and applied feature importance to get 14 variables. After training the model, we found that the model with 14 features yields an AUROC of 0.862. After excluding the 'malignancy' variable, the model achieved a slightly increase in AUROC to 0.864. This suggested that 'malignancy' may not significantly contribute to the outcome. After removing the 'respiratory dysfunction' variable, the AUROC improved to 0.866. This iterative process indicated a potential improvement in model performance. Respiratory failure is a type of disease within the category of 'respiratory dysfunction'. Retaining respiratory failure while removing respiratory dysfunction indicates that the confirmed diagnosis of respiratory failure has a higher impact on the patient's survival rate compared to other respiratory dysfunctions. In the end, we obtained the 12 most important variables for

Table 2

Characteristics between train cohort (N=11,549) and validation cohort (N=2,475) with P value. Demographic: Age. Disease severity: SAPS II. Diagnosis: Respiratory dysfunction, Respiratory failure, CHF, Diabetes, Malignancy. Laboratory results: Maximum hemoglobin (g/dl), Minimum lactate (mmol/L), Minimum BUN (mg/dl), Minimum PaO2(mmHg), Maximum PaCO2(mmHg), Minimum PaCO2(mmHg). Others: Vent Duration (Hour).

	Train cohort (N=11,549)	Validation cohort (N=2,475)	P
Demographic			
Age (years)	63.6(22.0)	63.6(22.0)	0.985
Disease severity			
SAPS II	40.4(19.0)	40.2(18.0)	0.515
Diagnosis			
Respiratory dysfunction	4,273(0.37)	892(0.36)	0.382
Respiratory failure	4,114(0.36)	858(0.35)	0.380
CHF	4,109(0.36)	888(0.36)	0.795
Diabetes	3,908(0.34)	807(0.33)	0.248
Malignancy	1,468(0.13)	297(0.12)	0.350
Laboratory results			
Maximum hemoglobin (g/dl)	12.2(2.6)	12.2(2.4)	0.414
Minimum lactate (mmol/L)	1.3(0.6)	1.3(0.6)	0.612
Minimum BUN (mg/dl)	13.3(8.0)	13.4(8.0)	0.653
Minimum PaO2(mmHg)	67.7(45.0)	68.3(46.0)	0.531
Maximum PaCO2(mmHg)	58.9(17.0)	58.0(16.0)	0.056
Minimum PaCO2(mmHg)	31.1(8.0)	31.2(8.0)	0.728
Treatment			
Vent Duration (Hour)	98.2(102.2)	97.7(95.1)	0.900
Target			
Dead in ICU or not	1,441(0.12)	307(0.12)	0.947

Table 3

Characteristics between survivors (N=14,441) and non-survivors (N=2,058) with P value. Demographic: Age. Disease severity: SAPS II. Diagnosis: Respiratory dysfunction, Respiratory failure, CHF, Diabetes, Malignancy. Laboratory results: Maximum hemoglobin (g/dl), Minimum lactate (mmol/L), Minimum BUN (mg/dl), Minimum PaO2(mmHg), Maximum PaCO2(mmHg), Minimum PaCO2(mmHg). Others: Vent Duration (Hour).

	Survivors (N=14,441)	Non-survivors (N=2,058)	P
Demographic			
Age (years)	63.2(21.0)	66.6(22.0)	<0.0001
Disease severity			
SAPS II	38.6(17.0)	52.4(23.0)	<0.0001
Diagnosis			
Respiratory dysfunction	4,918(0.34)	1,112(0.54)	<0.0001
Respiratory failure	4,721(0.33))	1,090(0.53)	<0.0001
CHF	5,085(0.35)	789(0.38)	0.0060
Diabetes	4,956(0.34)	600(0.29)	<0.0001
Malignancy	1,867(0.13)	251 (0.12)	0.3715
Laboratory results			
Maximum hemoglobin (g/dl)	12.3(2.5)	11.8(2.8)	<0.0001
Minimum lactate (mmol/L)	1.2(0.5)	1.9(1.0)	<0.0001
Minimum BUN (mg/dl)	12.3(8.0)	19.7(13.8)	<0.0001
Minimum PaO2(mmHg)	69.1(48.0)	56.6(32.0)	<0.0001
Maximum PaCO2(mmHg)	58.1(15.0)	62.7(23.0)	<0.0001
Minimum PaCO2(mmHg)	31.4(7.0)	28.7(8.0)	<0.0001
Treatment			
Vent Duration (Hour)	89.4(81.7)	167.2(194.0)	<0.0001

model establishment. Figure 3, Figure 4 and Figure 5 showed the AUROC improvement process of deleting insignificant features one by one.

Figure 3: AUROCs boxplots of neural network models with 14 features. Each boxplot displays the AUROC with 95% CI after deleting the corresponding variables. The column baseline shows the result that keeps all the variables.

3.3. *Evaluation results*

Table 4 and Table 5 show the detailed results summary of our proposed model and baseline ML models. Our proposed neural network model resulted in test set, validation set, and train set, AUROC=0.879, 95% CI = [0.860-0.896], AUROC=0.866, 95% CI = [0.846-0.886], and AUROC=0.958, 95% CI = [0.955-0.960]. The baseline models encompassed KNN, logistic regression, decision tree, random forest, bagging, XGBoost, and SVM algorithms, yielded the following scores: 0.605, 95% CI [0.578-0.634], 0.851, 95% CI = [0.829-0.871], 0.623, 95% CI = [0.595-0.652]], 0.809, 95% CI = [0.784-0.833], 0.765, 95% CI = [0.734-0.794], 0.854, 95% CI = [0.832-0.872], and 0.851, 95% CI = [0.828-0.874], respectively. We also calculated the accuracy score of KNN 0.809, logistic regression 0.783, decision tree 0.809, random forest 0.860, neural network 0.859, bagging 0.845, XGBoost 0.878, and SVM 0.881. These scores provided insights into how well the models would perform on new data.

Figure 6 and Figure 7 display Receiver Operating Characteristic (ROC) curves of our proposed model neural network and the seven baseline models, KNN, Logistic Regression, Decision Tree, Random Forest, Bagging, XGBoost, and SVM on test set and validation set. Noticed that all models except KNN and logistic regression had smooth ROC curves and higher AUROC values.

Figure 4: AUROCs boxplots of neural network models with 13 features. Each boxplot displays the AUROC with 95% CI after deleting the corresponding variables. The column baseline shows the result that keeps all the variables. Malignancy has been deleted.

Figure 5: AUROCs boxplots of neural network models with 12 features. Each boxplot displays the AUROC with 95% CI after deleting the corresponding variables. The column baseline shows the result that keeps all the variables. Respiratory dysfunction has been deleted.

Figure 8 displays the AUROC boxplots of our proposed model and baseline models. We observed that the AUROCs

Table 4

Evaluation results and confidence interval for proposed model. Trained the model on different sets: test set, validation set and train set. The evaluation metrics included AUROC, AUROC 95% CI, precision, recall value, accuracy score, and F1 score.

Models	AUROC	AUROC 95% CI	Accuracy
Proposed model performance on test set	0.879	[0.860-0.896]	0.859
Proposed model performance on validation set	0.866	[0.846-0.886]	0.855
Proposed model performance on train set	0.958	[0.955-0.960]	0.881

Table 5

Evaluation metrics and confidence interval for seven models. We used a total of seven different models, KNN, Logistic Regression, Decision Tree, Random Forest, Bagging, XGBoost, and SVM. The evaluation metrics included AUROC, AUROC 95% CI and accuracy score.

Models	AUROC	AUROC 95% CI	Accuracy
KNN	0.605	[0.578-0.634]	0.809
Logistic Regression	0.851	[0.829-0.871]	0.783
Decision Tree	0.623	[0.595-0.652]	0.809
RF	0.809	[0.784-0.833]	0.860
Bagging	0.765	[0.734-0.794]	0.845
XGBoost	0.854	[0.832-0.872]	0.876
SVM	0.851	[0.828-0.874]	0.881

of LR, neural network, XGBoost, and SVM exceeding 0.8, showing the strong predictive ability. Among these boxplots, the mean AUROC of our proposed model exceeds the maximum value of the AUROCs of other baseline models.

Figure 7: ROC curves of the eight models for the validation set. KNN, logistic regression, decision tree, random forest, neural network, bagging, XGBoost, SVM.

Figure 6: ROC curves of the eight models for the test set. KNN, logistic regression, decision tree, random forest, neural network, bagging, XGBoost, SVM.

3.4. SHAP analysis

SHAP (SHapley Additive exPlanations) is a method used in ML to understand the impact of individual variables on model predictions. It provides a way to interpret the output of any ML model by quantifying the contribution of each feature to the predicted outcome [37]. Figure 9, along with Table 6, displays the SHAP (SHapley Additive exPlanations) values

for the test set, providing a detailed analysis of how each variable affects the model's prediction. The SHAP analysis identifies 'respiratory failure' as the most significant predictor, followed by 'minimum Lactate', 'saps ii', 'vent duration', 'minimum BUN', 'diabetes', 'maximum PaCO2', 'CHF', 'minimum PaCO2', 'age', 'minimum PaO2' and 'maximum hemoglobin'. Here, the 'respiratory failure' shows a notable positive impact on the model's predictions. The 'minimum PaO2' and 'maximum hemoglobin', in comparison, demonstrate more moderate effects. The shift in the order of feature importance in the training set highlights the importance

Figure 8: AUROCs boxplots of the eight models, KNN, Logistic Regression, Decision Tree, Random Forest, neural network, Bagging, XGBoost, and SVM. The upper line of the single boxplot represents the maximum value of AUROC, the lower line represents the minimum value of AUROC, and the middle line represents the mean value of AUROC.

Figure 9: SHAP value based on neural network model for the test set. Predictors: Respiratory failure, diabetes, age, SAPS II Score, maximum hemoglobin, minimum lactate, CHF, vent duration, minimum bun, minimum PaCO2, maximum PaCO2, minimum PaO2.

Figure 10: Feature importance is based on Neural Network model for the test set. Predictors: Respiratory failure, diabetes, age, SAPS II Score, maximum hemoglobin, minimum lactate, CHF, vent duration, minimum bun, minimum PaCO2, maximum PaCO2, minimum PaO2.

of considering diverse metrics for a comprehensive model evaluation.

The difference between the feature importance ranking in Figure 10 and the average SHAP value in Table 6 could be attributed to the different methodologies underlying these two approaches. It was also worth mentioning that while 'CHF' and 'diabetes' held moderate importance in the SHAP ranking, they appeared as top contributors in the feature importance. This divergence underscores the complexity of variable interactions within the model and highlights the necessity of employing multiple interpretability methods to fully understand the model's behavior.

Furthermore, the minimal impact of 'Minimum PaCO2' and 'Maximum Hemoglobin' on the model output, as indicated by their low feature importance and SHAP values, suggests that these factors are less discriminative for the predictive task at hand. The comprehensive analysis of these indicators provides valuable insights into the model's decision-making process, guiding practitioners in refining the model and focusing on the most pertinent predictors for outcome prediction.

4. DISCUSSION

4.1. *Existing model compilation summary*

In our study, we proposed a NN model to predict the ICU mortality of patients undergoing invasive mechanical ventilation. The result of our mortality prediction model was better than the best existing literature by Y.Zhu et al [19]. The

effectiveness of our model demonstrated 7.06% improvement in AUROC.

Although the existing literature result had effectively anticipated mortality rates among ICU patients, it exhibited certain limitations. They used a total of 66 variables to predict the model outcome. This approach may raise concerns related to model complexity and overfitting. Additionally, the outcome of their research was considered to be unsatisfactory and inadequate for practical use in clinical exercising.

Table 6

Average SHAP value for 12 features in the test set. The 12 predictors: Respiratory failure, diabetes, age, SAPS II Score, maximum hemoglobin, minimum lactate, CHF, vent duration, minimum bun, minimum PaCO2, maximum PaCO2, minimum PaO2.

Features	Average SHAP Value (Test Set)
Respiratory Failure	0.096
Minimum Lactate	0.080
SAPS II	0.075
Vent Duration	0.075
Minimum Bun	0.052
Diabetes	0.043
Maximum PaCO2	0.040
CHF	0.034
Minimum PaCO2	0.025
Age	0.019
Minimum PaO2	0.010
Maximum Hemoglobin	0.008

For our research, we used advanced feature selection techniques to select only 12 variables as our features. The results of our model had a significant improvement, which provided a more reliable result for clinical use and highlights the efficiency and effectiveness of our model in delivering superior predictive performance with a more concise feature set. Moreover, we used the train set, validation set and test set for evaluation. The best existing model only used train and test sets. Using different sets for hyperparameter tuning and model evaluation could avoid information leakage and enhance the model's generalization assessment capabilities. Furthermore, our proposed neural network model was easy to be set up and replicated, had fewer layers which helps prevent overfitting and trains faster than other deep neural networks. These features made our model efficient and practical for real-world applications.

We found that respiratory failure had a higher association with patients mortality than respiratory dysfunction from SHAP analysis and feature importance, which means the condition of respiratory failure was much more important than the general comprehensive disease of respiratory dysfunction. This suggested that clinicians should pay more attention to patients with a history of respiratory failure. The vent duration was also a clinically meaningful variable that directly reflects the severity and duration of respiratory failure. It also provided insight into the patient's respiratory status and the level of support needed. From the SHAP value analysis (Figure 7), we found that the longer the ventilation use time, the higher the patient's mortality rate, proving that prolonged use of a ventilator does not improve patient survival.

Additionally, the SVM and XGBoost baseline models had AUROCs of 0.841 and 0.825, respectively, which were similar to the performance of our proposed NN model. We observed that an overlap of AUROC boxplots of these two models in Figure 8, which illustrates the performances of these two

models were also powerful in predicting mortality since NN had multiple layers and needed more time for training.

4.2. *Study limitations*

In our model development, we used train and validation datasets to construct the model. The test dataset was for evaluating the performance of the model. The train, validation and test sets were all from the MIMIC-III database. However, using independent datasets from different systems will be beneficial for testing the performance of the model. The MIMIC III database is a large but outdated database, which contains the dataset of related ICU patients only between 2001 and 2012. Exploring newer datasets could enhance the predictive capabilities of our model. Additionally, integrating other types of data, such as images and text, could further improve the accuracy and utility of our results.

5. CONCLUSION

The goal of our paper is to build a novel neural network model to predict the mortality of ICU patients undergoing mechanical ventilation. Compared with the results of baseline ML models and existing literature, deep learning methods for modeling ICU patient data in the MIMIC II database to predict mechanical ventilation mortality have provided significant improvements in predicting observed outcomes. This improvement may be due to the efficiency of variables, such as the time series variables we selected for predicting the model. Our framework provides valuable support for clinicians to identify patients at high risk of death in the ICU. This predictive tool is particularly beneficial for patients and clinicians, as it can assess the time a patient leaves the ICU and guide clinicians in arranging patient treatment plans.

Future research could focus on validating our methods with datasets from different healthcare systems or exploring their applicability to various diseases and outcomes. Exploring the applicability of our methods to various diseases and outcomes can uncover new insights and potential applications. Additionally, other researchers could explore different types of data based on our dataset and can lead to innovative research directions. For example, integrating image data or text data with our existing dataset could provide a more comprehensive understanding of patient health and improve the accuracy of our predictions.

Acknowledgment

The authors extend their gratitude to the creators of MIMIC-III for furnishing a thorough and inclusive public electronic health record (EHR) dataset.

References

[1] H. Xia, B. J. Daley, A. Petrie, and X. Zhao, "A neural network model for mortality prediction in ICU," IEEE Xplore, Sep. 01, 2012. URL: https://ieeexplore.ieee.org/abstract/document/6420380.

[2] G. Thiéry et al., "From Mechanical Ventilation to Intensive Care Medicine: A Challenge for Bosnia and Herzegovina," Bosnian Journal

of Basic Medical Sciences, vol. 9, no. 1, pp. S69–S76, Oct. 2009, URL: https://doi.org/10.17305/bjbms.2009.2766.

[3] T. Amador, S. Saturnino, A. Veloso, and N. Ziviani, "Early identification of ICU patients at risk of complications: Regularization based on robustness and stability of explanations," Artificial Intelligence in Medicine, p. 102283, Mar. 2022, URL: https://doi.org/10.1016/j.artmed.2022.102283.

[4] H. Wunsch, J. Wagner, M. Herlim, D. H. Chong, A. A. Kramer, and S. D. Halpern, "ICU Occupancy and Mechanical Ventilator Use in the United States*," Critical Care Medicine, vol. 41, no. 12, pp. 2712–2719, Dec. 2013, URL: https://doi.org/10.1097/ccm.0b013e318298a139.

[5] Y.-S. Hung et al., "Clinical characteristics and survival outcomes of terminally ill patients undergoing withdrawal of mechanical ventilation," Journal of the Formosan Medical Association, vol. 117, no. 9, pp. 798–805, Sep. 2018, URL: https://doi.org/10.1016/j.jfma.2017.09.014.

[6] M. Alemayehu, A. Azazh, H. Hussien, and A. Baru, "Characteristics and Outcomes of Mechanically Ventilated Patients at Adult ICU of Selected Public Hospitals in Addis Ababa, Ethiopia," Open Access Emergency Medicine, vol. Volume 14, pp. 395–404, Aug. 2022, URL: https://doi.org/10.2147/oaem.s369752.

[7] A. ESTEBAN et al., "How Is Mechanical Ventilation Employed in the Intensive Care Unit?," American Journal of Respiratory and Critical Care Medicine, vol. 161, no. 5, pp. 1450–1458, May 2000, URL: https://doi.org/10.1164/ajrccm.161.5.9902018.

[8] H. Wunsch, W. T. Linde-Zwirble, D. C. Angus, M. E. Hartman, E. B. Milbrandt, and J. M. Kahn, "The epidemiology of mechanical ventilation use in the United States*," Critical Care Medicine, vol. 38, no. 10, pp. 1947–1953, Oct. 2010, URL: https://doi.org/10.1097/ccm.0b013e3181ef4460.

[9] D. Pertab, "Principles of mechanical ventilation – a critical review," British Journal of Nursing, vol. 18, no. 15, pp. 915–918, Aug. 2009, URL: https://doi.org/10.12968/bjon.2009.18.15.43560.

[10] S. S. Carson, C. E. Cox, G. M. Holmes, A. Howard, and T. S. Carey, "The Changing Epidemiology of Mechanical Ventilation: A Population-Based Study," Journal of Intensive Care Medicine, vol. 21, no. 3, pp. 173–182, May 2006, URL: https://doi.org/10.1177/0885066605282784.

[11] L. Brochard, A. Slutsky, and A. Pesenti, "Mechanical Ventilation to Minimize Progression of Lung Injury in Acute Respiratory Failure," American Journal of Respiratory and Critical Care Medicine, vol. 195, no. 4, pp. 438–442, Feb. 2017, URL: https://doi.org/10.1164/rccm.201605-1081cp.

[12] R. G. Brower and H. E. Fessler, "MECHANICAL VENTILATION IN ACUTE LUNG INJURY AND ACUTE RESPIRATORY DISTRESS SYNDROME," Clinics in Chest Medicine, vol. 21, no. 3, pp. 491–510, Sep. 2000, URL: https://doi.org/10.1016/s0272-5231(05)70161-1.

[13] Y. Jia, C. Kaul, T. Lawton, R. Murray-Smith, and I. Habli, "Prediction of weaning from mechanical ventilation using Convolutional Neural Networks," Artificial Intelligence in Medicine, vol. 117, p. 102087, Jul. 2021. URL: https://doi.org/10.1016/j.artmed.2021.102087.

[14] N. Prasad, L.-F. Cheng, C. Chivers, M. Draugelis, and B. E. Engelhardt, "A Reinforcement Learning Approach to Weaning of Mechanical Ventilation in Intensive Care Units," arXiv:1704.06300 [cs], Apr. 2017, URL: https://arxiv.org/abs/1704.06300.

[15] M. Pishgar, M. Razo, J. Theis and H. Darabi, "Process Mining Model to Predict Mortality in Paralytic Ileus Patients," 2021 International Conference on Cyber-Physical Social Intelligence (ICCSI), Beijing, China, 2021, pp. 1-6. URL: https://arxiv.org/abs/2108.01267.

[16] J. Gao, Y. Lu, Ian Raymond Domingo, Kamiar Alaei, and M. Pishgar, "Predicting Sepsis Mortality Using Machine Learning Methods," medRxiv (Cold Spring Harbor Laboratory), Mar. 2024, URL: https://www.medrxiv.org/content/10.1101/2024.03.14.24304184v1.

[17] J. H. Kim, Y. S. Kwon, and M. S. Baek, "Machine Learning Models to Predict 30-Day Mortality in Mechanically Ventilated Patients," Journal of Clinical Medicine, vol. 10, no. 10, p. 2172, May 2021. URL: https://doi.org/10.3390/jcm10102172.

[18] A. Abujaber, A. Fadlalla, D. Gammoh, H. Abdelrahman, M. Mollazehi, and A. El-Menyar, "Prediction of in-hospital mortality in patients on

mechanical ventilation post traumatic brain injury: machine learning approach," BMC Medical Informatics and Decision Making, vol. 20, no. 1, Dec. 2020, URL: https://doi.org/10.1186/s12911-020-01363-z.

[19] Y. Zhu et al., "Machine Learning Prediction Models for Mechanically Ventilated Patients: Analyses of the MIMIC-III Database," Frontiers in Medicine, vol. 8, Jul. 2021, URL: https://doi.org/10.3389/fmed.2021.662340.

[20] R. Féraud and F. Clérot, "A methodology to explain neural network classification," Neural Networks, vol. 15, no. 2, pp. 237–246, Mar. 2002, URL: https://doi.org/10.1016/s0893-6080(01)00127-7.

[21] M. Pishgar, J. Theis, M. Del Rios, A. Ardati, H. Anahideh, and H. Darabi, "Prediction of unplanned 30-day readmission for ICU patients with heart failure," BMC Medical Informatics and Decision Making, vol. 22, no. 1, May 2022, URL: https://doi.org/10.1186/s12911-022-01857-y.

[22] Á. Silva, P. Cortez, M. F. Santos, L. Gomes, and J. Neves, "Mortality assessment in intensive care units via adverse events using artificial neural networks," Artificial Intelligence in Medicine, vol. 36, no. 3, pp. 223–234, Mar. 2006, URL: https://doi.org/10.1016/j.artmed.2005.07.006.

[23] J. H. Kim, Y. S. Kwon, and M. S. Baek, "Machine Learning Models to Predict 30-Day Mortality in Mechanically Ventilated Patients," Journal of Clinical Medicine, vol. 10, no. 10, p. 2172, May 2021, URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8157228/.

[24] Al-Dorzi, Hasan M et al. "Characteristics and predictors of mortality of patients with hematologic malignancies requiring invasive mechanical ventilation." Annals of thoracic medicine vol. 12,4, 2017, URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5656944/.

[25] Giri, Mohan et al. "Blood Urea Nitrogen Is Associated with In-Hospital Mortality in Critically Ill Patients with Acute Exacerbation of Chronic Obstructive Pulmonary Disease: A Propensity Score Matching Analysis." Journal of clinical medicine vol. 11,22 6709. 13 Nov. 2022, URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9699438/.

[26] Prediletto, Irene et al. "Standardizing PaO2 for PaCO2 in P/F ratio predicts in-hospital mortality in acute respiratory failure due to Covid-19: A pilot prospective study." European journal of internal medicine vol. 92 (2021): 48-54. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8222796/.

[27] K. Taunk, S. De, S. Verma and A. Swetapadma, "A Brief Review of Nearest Neighbor Algorithm for Learning and Classification," 2019 International Conference on Intelligent Computing and Control Systems (ICCS), Madurai, India, 2019, pp. 1255-1260, URL: https://ieeexplore.ieee.org/document/9065747.

[28] Ranganathan P, Pramesh CS, Aggarwal R. Common pitfalls in statistical analysis: Logistic regression. Perspect Clin Res. 2017 Jul-Sep;8(3):148-151, URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5543767/.

[29] Song YY, Lu Y. Decision tree methods: applications for classification and prediction. Shanghai Arch Psychiatry. 2015 Apr 25;27(2):130-5. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4466856/.

[30] Breiman, L. Random Forests. Machine Learning 45, 5–32 (2001), URL: https://doi.org/10.1023/A:1010933404324.

[31] G. Singh and M. Sachan, "Multi-layer perceptron (MLP) neural network technique for offline handwritten Gurmukhi character recognition," 2014 IEEE International Conference on Computational Intelligence and Computing Research, Coimbatore, India, 2014, pp. 1-5, URL: https://ieeexplore.ieee.org/document/7238334.

[32] Breiman, L. Bagging predictors. Mach Learn 24, 123–140 (1996). URL: https://doi.org/10.1007/BF00058655.

[33] Chen, Tianqi & Guestrin, Carlos. (2016). XGBoost: A Scalable Tree Boosting System. 785-794. 10.1145/2939672.2939785, URL: https://arxiv.org/abs/1603.02754.

[34] M. A. Hearst, S. T. Dumais, E. Osuna, J. Platt and B. Scholkopf, "Support vector machines," in IEEE Intelligent Systems and their Applications, vol. 13, no. 4, pp. 18-28, July-Aug. 1998, URL: https://ieeexplore.ieee.org/document/708428.

[35] S. Santurkar, D. Tsipras, A. Ilyas, and A. Madry, "How Does Batch Normalization Help Optimization?,"

Neural Information Processing Systems, 2018. URL: https://proceedings.neurips.cc/paper/2018/hash/905056c1ac1dad141560467e0a99e1cf-Abstract.html.

[36] N. Srivastava, G. Hinton, A. Krizhevsky, and R. Salakhutdinov, "Dropout: A Simple Way to Prevent Neural Networks from Overfitting," Journal of Machine Learning Research, vol. 15, pp. 1929–1958, 2014, URL: https://jmlr.org/papers/v15/srivastava14a.html.

[37] R. I. Hamilton and P. N. Papadopoulos, "Using SHAP Values and Machine Learning to Understand Trends in the Transient Stability Limit," vol. 36, no. 5, pp. 4000-4011, Sept. 2021, URL: https://arxiv.org/abs/2302.06274.