1 Prospective study of *Candida auris* **nucleic-acids in wastewater solids in 190 wastewater**
2 **treatment plants in the United States suggests widespread occurrence**
3

4 **Alessandro Zulli[1], Elana M. G. Chan[1], Bridgette Shelden[2], Dorothea Duong[2], Xiang-Ru S.**
5 **Xu[2], Bradley J. White[2], Marlene K. Wolfe[3], Alexandria B. Boehm[1*]**
6
7 1. Department of Civil and Environmental Engineering, Stanford University, 473 Via Ortega,
8 Stanford, California, 94305
9 2. Verily Life Sciences LLC, South San Francisco, CA, USA
10 3. Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory
11 University, 1518 Clifton Rd, Atlanta, GA, USA, 30322
12
13 * Corresponding author: aboehm@stanford.edu
14
15

16
Abstract

Candida auris is an emerging, multidrug-resistant fungal pathogen that poses a significant

public health threat in healthcare settings. Despite yearly clinical cases rapidly increasing from

77 to 8,131 in the last decade, surveillance data on its distribution and prevalence remains

limited. We implemented a novel assay for *C. auris* detection on a nationwide scale

prospectively from September 2023 to March 2024, analyzing a total of 13,842 samples from

190 wastewater treatment plants across 41 U.S. states. Assays were extensively validated

through comparison to other known assays and internal controls. Of these 190 wastewater

treatment plants, *C. auris* was detected in the wastewater solids of 65 of them (34.2%) with

1.45% of all samples having detectable levels of *C. auris*. Detections varied seasonally, with

2.00% of samples positive in autumn versus 1.01% in winter (p<0.0001). The frequency of

detection in wastewater was significantly associated with states having older populations

(p<0.001), sewersheds containing more hospitals (p<0.0001), and sewersheds containing more

nursing homes (p<0.001). These associations are in agreement with known *C. auris*

epidemiology. This nationwide study demonstrates the viability of wastewater surveillance for *C.

auris* surveillance, and further highlights the value of wastewater surveillance when clinical

testing is constrained.

36
Introduction

Healthcare associated infections are rapidly emerging as a global public health threat due to

increasing levels of antimicrobial resistance and an aging worldwide population (1–3). Among

these infections, *Candida auris,* first identified in 2009 in Japan, is an opportunistic, drug-

resistant fungal pathogen with significant levels of associated morbidity and mortality (3–6). It

has now spread to over 40 countries and has been designated as a priority pathogen by both

international organizations, such as the World Health Organization (WHO), and national

organizations, such as the Center for Disease Control and Prevention (CDC) in the United

States, where it is now a nationally notifiable condition (4, 6, 7). The estimated crude mortality

rate within the United States for *C. auris* infections was 34% between 2017-2022 (5).

C. auris prevention and control efforts are exacerbated by the fungus' hardiness on fomites and high levels of transmissibility (6–8). The fungus has been found to persist for at least 7 days on both moist and dry surfaces, increasing the frequency and chance of transmission events in healthcare settings (9, 10). Of particular concern are high-touch plastic surfaces within healthcare facilities where *C. auris* was found to be viable for up to 14 days (10). In addition, *C. auris* is resistant to many common disinfectants so proper decontamination protocols are difficult and time intensive in already strained healthcare facilities (11). Further complicating containment efforts are the different presentations in colonized individuals compared to those with acute infections, with most colonized individuals presenting as asymptomatic (4). A lack of screening in high-risk patients and misidentification of specimens as other *Candida* subspecies has also contributed to the rapid spread of *C. auris* (5, 12). In the United States, clinical cases rose from 77 between 2013-2016 to 8,131 in 2022 and are projected to continue rising (5, 13).

Despite this tremendous increase in cases and the accompanying screening efforts, clinically available data are still sparse, with many institutions not speciating *Candida* cases resulting in underreporting cases in long-term care facilities and nursing homes (13, 14). Many of these facilities do not have the necessary equipment or human capital to implement speciation efforts and screening, which has been shown to be a necessary part of successful containment efforts (12). Alternative approaches to clinical surveillance are therefore necessary to better track both the spread and severity of outbreaks. Wastewater represents a naturally composite sample which is particularly useful for disease surveillance as it contains urine, feces, vomit, saliva, sloughed skin cells, sputum, and other excretions. Wastewater surveillance has been successfully used to track a wide variety of human pathogens including enterovirus*,* SARS-CoV-2, influenza A virus*,* and flavivirus (15–17)*.* Like these viruses, *C. auris* has been detected in urine and stool samples from colonized or infected individuals, though it is typically diagnosed

through presence in the blood or dermis of infected patients (4, 10, 18). Furthermore, recent

studies have demonstrated the persistence and subsequent detection of *C. auris* in wastewater

samples in Nevada and Florida (19). Early detection of this emerging pathogen through

environmental surveillance may complement clinical surveillance efforts and inform containment

and screening procedures (20).

In this paper, we present the results of a United States-wide wastewater monitoring effort for *C.*

auris and compare the data to publicly available, recent, clinical case data. We also compare

results to available demographic indicators and healthcare facility locations. Detections of *C.*

auris in wastewater demonstrate wide geographical occurrence , and we demonstrate that these

wastewater detections appear to be associated with age and presence of hospitals and nursing

homes in adjacent communities. These results demonstrate the advantages of using

wastewater surveillance for the detection of emerging pathogens and for supplementing clinical

testing efforts.

Methods

RT-PCR assays. We designed a novel probe for *Candida auris* to use in conjunction with

previously published specific and sensitive forward and reverse primers (21). The assay targets

the region of the genome encoding the 5.8S ribosomal RNA, all of ITS2, and a fragment of 28S

ribosomal RNA. To design the probe, *C. auris* genome sequences were downloaded from the

National Center for Biotechnology Information (NCBI) and aligned to identify conserved regions

located between the primers. The probe was designed *in silico* using Primer3Plus

(https://primer3plus.com/). Parameters used in the design process (e.g., sequence length, GC

content, and melt temperatures) are provided in Table S1. The primers and probe were then

screened for specificity *in silico* by blasting the sequences in NCBI and *in vitro* against fungal

panels, or cDNA sequences. The *in silico* analysis was performed by excluding all *C. auris*

genomes from a BLAST search, thereby only giving off-target matches (22).

The primers and probe were tested *in vitro* for specificity and sensitivity using a fungal panel

(NATCTVPOS-BD, Zeptomatrix Buffalo, NY) and *Candida auris* Satoh et Makimura (ATCC

MYA-5001) purchased from American Type Culture Collection (ATCC, Manassas, VA). The

fungal panel NATCTVPOS-BD includes chemically inactivated *Candida albicans*, *C. krusei*, *C.*

glabrata, and *Trichomoniasis vaginalis*. Nucleic acids were extracted from fungi using Chemagic

Viral DNA/RNA 300 Kit H96 for Chemagic 360 (PerkinElmer, Waltham, MA).

Nucleic acids were used undiluted as template in digital droplet RT-PCR (ddRT-PCR) singleton

assays for sensitivity and specificity testing in single wells. The concentration of targets used in

the *in vitro* specificity testing was between 10^3 and 10^4 copies per well. Negative RT-PCR

controls were included on each plate.

We used an RT-PCR assay for the in vitro sensitivity and specificity testing as the goal was to

ultimately multiplex the *C. auris* assay in a panel that includes RNA-genome viruses. The

implications of the use of RT-PCR, and a comparison to results obtained using PCR are further

investigated, as described below. Regardless, this means the assay detects both DNA and RNA

targets.

Clinical surveillance data. The National Notifiable Disease Surveillance System (NNDSS) is a

nationwide collaboration to which all levels of healthcare providers share health information

about nationally notifiable infectious and noninfectious diseases (23). Case reports are compiled

on a weekly basis using uniform case definitions from 50 states, the District of Columbia, New

York City, and 5 territories (American Samoa, Guam, Northern Mariana Islands, Puerto Rico,

U.S. Virgin Islands) (23). For this study, we used publicly available information on NNDSS for

Candida auris from 11 September 2023 to 1 March 2024 (https://data.cdc.gov/api/views/x9gk-

5huc/rows.csv?accessType=DOWNLOAD). Dates of case determination varied but included the

following in order of preference: date of disease start, date of diagnosis, date of laboratory

result, date of first report to public health system, or date of state report (24). States which did

not report clinical case data were omitted from analyses. It should be noted that this database is

known to be incomplete as many local jurisdictions do not submit data; additional data on C.

auris cases or Candida cases may be available from other local or regional sources that are not

publicly available or currently updated.

Wastewater data: **sample collection.** Wastewater measurements were made prospectively as

part of a wastewater surveillance program. Between 11 September 2023 and 1 March 2024,

wastewater samples (either 24-hour composited influent or grab samples from the primary

clarifier, Table S1) were collected by wastewater treatment plant (WWTP) staff using sterile

containers. Samples were typically obtained approximately three times per week and shipped

overnight to the laboratory at 4°C where they were processed immediately with no storage.

Samples were collected from 190 distinct WWTPs across a total of 41 states over the course of

the study period (Figure 1). A total of 13,842 samples were collected and analyzed as part of

this study.

Wastewater data: **pre-analytical processing.** *C. auris* is expected to be associated with the

solid phase of wastewater based on previous empirical measurements across liquid and solid

phases in wastewater and its size (diameter = 2.5-5 µm), so in this study we made

measurements in the solid phase of wastewater (25, 26). Details of the isolation of solids from

the samples are provided in other peer-reviewed publications (27). In short, samples were

centrifuged to dewater the solids. One aliquot of dewatered solids was used for nucleic-acid

extractions and another was used to determine the dry weight of the solids using an oven (27).

Approximately 75 mg of dewatered solids was added per milliliter of DNA/RNA shield (Zymo,

Irvine, CA) spiked with bovine coronavirus vaccine as a positive extraction control. This mixture

of solids and buffer was previously shown to have minimal inhibition.(28) The mixture was

homogenized after the addition of grinding balls, and centrifuged; the supernatant was used

immediately for nucleic-acid extractions. Nucleic-acids were extracted from 6-10 replicate

aliquots of the supernatant using the Chemagic Viral DNA/RNA 300 kit H96 for the Perkin Elmer

Chemagic 360 (Perkin Elmer, Waltham, MA) followed by PCR inhibitor removal with the Zymo

OneStep-96 PCR Inhibitor Removal kit (Zymo Research, Irvine, CA). Whether 6 or 10 replicates

were used is specified in Table S1. The suspension volume entered into the nucleic-acid

extraction process was 300 μl and 50 μl of nucleic-acids were retrieved after the inhibitor

removal kit. Negative extraction controls consisted of BCoV-vaccine spiked DNA/RNA Shield.

Nucleic-acids were used immediately as template in PCR as described next, with no storage.

Wastewater data: **analytical processing.** We used droplet digital reverse transcription

polymerase chain reaction (ddRT-PCR) to measure nucleic-acids in wastewater. We measured

pepper mild mottle virus (PMMoV) as an endogenous positive control as concentrations tend to

be extremely elevated in wastewater samples; we measured BCoV as an endogenous, spiked-

in control (28–30). Methods applied to wastewater solids to measure PMMoV and BCoV in a

duplex reaction are provided in detail elsewhere (27). The *C. auris* assay (Table 1) was run in

multiplex using a probe-mixing approach. The exact assays that *C. auris* was multiplexed with

varied slightly over the duration of the project for some of the plants (information is provided in

Table S2). Multiplexing *C. auris* with these targets did not affect *C. auris* quantification (see SI

Figure S1). The results from those other assays are not provided herein. The PMMoV/BCoV

duplex and *C. auris* multiplex assays were run on 96-well plates. Each 96-well PCR plate of

wastewater samples included PCR positive controls for each target assayed on the plate in 1

well, PCR negative no template controls in two wells, and extraction negative controls in two

wells. PCR positive controls consisted of *C. auris* DNA. Each of the 6 or 10 replicate nucleic-

acid extracts were run in their own wells to measure *C. auris*. Two randomly selected or 10

replicate nucleic-acid extracts were run in their own wells to measure PMMoV and BCoV (Table

S1).

Table 1. Primers and probes used in this study for detection of *C. auris* nucleic acids. Primers
and probes were purchased from Integrated DNA Technologies (Coralville, IA, USA). The
probes contained fluorescent molecule HEX and quenchers (5′ HEX/ZEN/3′ IBFQ); HEX,
hexachloro-fluorescein; ZEN, a proprietary internal quencher from Integrated DNA Technologies
(Coralville, IA, USA); and IBFQ, Iowa Black FQ. Amplicon size is 163 basepairs.

Forward	CGCACATTGCGCCTTGGGGTA
Reverse	GTAGTCCTACCTGATTTGAGGCGAC
Probe	CTTCTCACCAATCTTCGCGGT

189
190
ddRT-PCR was performed on 20-µl samples from a 22 µl-reaction volume, prepared using 5.5

µl of template mixed with 5.5 µl of One-Step RT-ddPCR Advanced Kit for Probes (Bio-Rad

1863021), 2.2 µl of 200 U/µl Reverse Transcriptase, 1.1 µl of 300 mM dithiothreitol (DTT), and

primers and probes mixtures at a final concentration of 900 nM and 250 nM, respectively.

Primer and probes for assays were purchased from Integrated DNA Technologies (IDT, San

Diego, CA) (Table 1). *C. auris* was measured in reactions with undiluted template whereas

PMMoV and BCoV were run on template diluted 1:100 in molecular grade water. It is important

to note that an RT step was used because during the prospective study, we were measuring a

large number of RNA genome viruses. Therefore, the *C. auris* target quantified should be

interpreted as copies of RNA plus DNA.

Droplets were generated using the AutoDG Automated Droplet Generator (Bio-Rad, Hercules,

CA). PCR was performed using Mastercycler Pro (Eppendforf, Enfield, CT) with with the

following cycling conditions: reverse transcription at 50°C for 60 minutes, enzyme activation at

95°C for 5 minutes, 40 cycles of denaturation at 95°C for 30 seconds and annealing and

extension at 59°C (for *C. auris*) or 56°C (for PMMoV and BCoV) for 30 seconds, enzyme

deactivation at 98°C for 10 minutes then an indefinite hold at 4°C. The ramp rate for

temperature changes were set to 2°C/second and the final hold at 4°C was performed for a

minimum of 30 minutes to allow the droplets to stabilize. Droplets were analyzed using the

QX200 or the QX600 Droplet Reader (Bio-Rad). A well had to have over 10,000 droplets for

inclusion in the analysis. All liquid transfers were performed using the Agilent Bravo (Agilent

Technologies, Santa Clara, CA).

Thresholding was done using QuantaSoft™ Analysis Pro Software (Bio-Rad, version 1.0.596)

and QX Manager Software (Bio-Rad, version 2.0). Replicate wells were merged for analysis of

each sample. In order for a sample to be recorded as positive, it had to have at least 3 positive

droplets. Concentrations of RNA targets were converted to concentrations in units of copies

(cp)/g dry weight using dimensional analysis. The total error is reported as standard deviations.

Three positive droplets across 6 merged wells corresponds to a concentration between ~1000

cp/g; the range in values is a result of the range in the equivalent mass of dry solids added to

the wells. Data collected as part of the study are available from the Stanford Digital Repository

(https://purl.stanford.edu/hv291tt0888).

Wastewater: Confirmatory analyses. We selected seven samples (Table S3) with relatively

high concentrations of *C. auris,* as measured above, to test another published *C. auris* assay by

Barber et al. (25) for detecting *C. auris* in wastewater. The assay used by Barber et al. targets

the same region of the *C. auris* genome as the assay in Table 1 but uses different primers and

probes (Table S4) (25). Archived nucleic acid extracts, stored at -80°C, from those samples

were thawed and then run as template using the Barber et al. assay as well as the assay

described herein (Table 1). The samples were run in duplex with an assay that targets the N

gene of SARS-CoV-2 (28). *C. auris* was run using a probe labeled with fluorescein amidite

(FAM). The methods described above for the analytical procedures (number of replicate wells,

negative and positive controls) were used.

These same seven samples were assayed with and without the reverse transcription step.

Nucleic-acid extracts were thawed and run exactly as described for the prospective study

except that the RT and DTT were not added to the mastermix.

We selected a subset of five of the seven samples (Table S3) for Sanger sequencing of the

PCR amplicon. One µl of archived (at -80°C) nucleic acid extracts was used as the template in a

50-µl PCR reaction prepared according to manufacturer's protocol with Q5® High-Fidelity 2X

Master Mix (New England Biolabs, Ipswich, MA) using the primers in Table 2. The thermocycler

program was run at an annealing temperature of 67°C for 30 seconds and extension at 72°C for

30 seconds. *C. auris* DNA was used as a positive control. PCR products were run on a 2%

agarose gel stained with SYBR Safe gel stain (Thermofisher, Fremont, CA) and amplicons of

the expected size (259 bp) were excised and gel purified using Zymo gel DNA recovery kit

(Irvine, CA). The purified DNA was then sent to Molecular Cloning Laboratories (South San

Francisco, CA) for sanger sequencing. Sequences were downloaded and aligned with *C. auris*

genomes using Snapgene software.

250
251
252

Figure 1. Location of wastewater treatment plants (WWTPs) with *Candida auris* DNA data included in the paper. Each blue dot represents the location of a WWTP.

Demographic data. *C. auris* hospitalizations are most common among older adults (5). We used 5-year estimates from the 2022 American Community Survey (ACS) at the state level to identify the median age of states with participating WWTPs (n = 41 states) (31).

Hospitals and nursing homes data. *C. auris* is particularly a concern in hospitals and nursing homes (5, 14, 20). We obtained the locations of hospitals and nursing homes in the United States from the US Department of Homeland Security's Homeland Infrastructure Foundation-Level Data (HIFLD) database (https://gii.dhs.gov/HIFLD) (32, 33). We selected all types of hospitals and nursing homes with an "open" status and then determined the number of hospitals and nursing homes in each sewershed using the Tabulate Intersection geoprocessing tool in ArcGIS Pro (version 3.1.1). Sewershed boundaries were provided by most WWTPs (n = 115).

For WWTPs that did not provide a sewershed (n = 75), we approximated sewershed boundaries

based on the zip codes serviced by the WWTP and USA ZIP Code Boundaries geospatial data

set published by Esri (Source: TomTom, US Postal Service, Esri) (34).

Data analysis. We made comparisons on the basis of percent of positive detections in

wastewater. Percentage of positive detections of *C. auris* was calculated by counting all positive

observations for a single WWTP and dividing by the total number of observations. Wastewater

measurements in each state were aggregated by Morbidity and Mortality Weekly Report

(MMWR) week, and we calculated the percentage of wastewater measurements that were

positive for *C. auris* each week.

Spearman's rank correlation coefficients were used to assess the association between weekly

(MMWR weeks) publicly available NNDSS clinical data and wastewater percent positive

detections (hereafter "detections") of *C. auris* for each state (n=18 states with wastewater and

clinical data). Case data from NNDSS, available on a weekly basis for each state, was used

after adjusting for population (cases per million people, Fig 2a).

We tested whether concentrations measured by the assay presented in Table 1 were different

from those measured by the Barber et al. assay using RT-ddPCR on 9 representative samples

(Table S3, Figure S2) (25). We ran the same samples using RT-ddPCR and ddPCR with no RT

step (Table S3, Figure S2). Differences between these controls were assessed using a paired t-

test.

We also assessed the relationship between *C. auris* percent positive detections and several

other contextual and demographic factors. We explored seasonality by assessing *C. auris*

detections in wastewater samples collected in fall (defined as Sept 1 to Nov 30) and winter

(defined as Dec 1 to Feb 29). We then assessed the association between *C. auris* detections

and age at the state level. We grouped WWTPs by those in states with a median age less than

versus greater than or equal to the national median age (39 years) (31). Lastly, we assessed

the association between *C. auris* detections and hospitals and nursing homes at the sewershed

level. We grouped WWTPs by those with a sewershed intersecting (1) less than versus greater

than or equal to the median number of hospitals among all sewersheds (median: 1 hospital) and

(2) less than versus greater than or equal to the median number of nursing homes among all

sewersheds (median: 7 nursing homes). For seasonality, we compared the percentage of

wastewater samples positive for *C. auris* in each season. For age, hospitals, and nursing

homes, we compared the percentage of wastewater samples that were positive for *C. auris*

across WWTPs in the less than median group versus the greater than or equal to median group.

The null hypothesis tested for each of these variables was that there were no differences

between the two groups.

Statistical significance and 95% confidence intervals between the groups described above were

assessed through bootstrapping (35, 36). For each variable and group, a distribution of values

was generated through random sampling with replacement 10,000 times. Significance was then

calculated using the proportion of values below our calculated statistic (i.e. whether the

calculated statistics were above the 95% confidence intervals). This same method was used

when assessing differences between assays (Fig S2) and seasonal differences in detection. In

those cases, grouping was done based on assay and season, respectively.

315
Results

QA/QC. The *C. auris* assay was found to be sensitive and specific. The *in silico* analysis

indicated no primer or probe sequence was more than an 85% match to off-target matches, and

there was no overlap found between combinations of forward primer off-targets, reverse primer

off-targets, and probe off-targets. These factors indicate a high level of specificity in the primer-

probe combination used. The in vitro testing confirmed sensitivity and specificity of the *C. auris*

assay as *C. auris* nucleic-acids were positive and the remaining non-target nucleic-acids were

negative.

Results are reported following the Environmental Microbiology Minimal Information (EMMI)

guidelines (see SI and Figure S3).(37) All positive and negative controls performed as expected,

indicating acceptable assay performance. Median (IQR) BCoV recoveries across all wastewater

samples were 1.08 (0.82, 1.43) indicating good recovery across all samples. Recoveries

exceeding 1 are the result of uncertainties in the measurement of BCoV added to the buffer

matrix. PMMoV levels were elevated in all samples indicating lack of gross extraction failures

(median = 4.8×10^8 cp/g, min = 5.0×10^5 cp/g, max = 2.02×10^{11} cp/g).

As a further validation step, *C. auris* was quantified in a subset of samples also using the Barber

et al. assay (Figure S2) (25). Concentrations measured by the two assays closely matched, with

no statistically significant differences (p>0.05). Comparisons between concentrations measured

using RT-ddPCR and ddPCR showed similar results (Figure S2); no significant differences were

found (p>0.05). Additionally, the PCR amplicons obtained using the forward and reverse

primers were sequenced and the sequences matched the *C. auris* genome segment targeted

(Figure S4).

National Overview. The study period spanned 11 September 2023 to 1 March 2024.

Wastewater data were available from 190 distinct WWTPs in 41 states (Table S1). The number

of WWTPs in these states ranged between 1–57 WWTPs (median: 2 WWTPs), with population

coverage of the sewersheds ranging from 0.13% to 59.5% of the population of each individual

state (median: 5.75%). The number of wastewater samples collected and analyzed from

individual WWTPs ranged between 23–173 samples (median: 72 samples) during the study

period. In total, 13,842 wastewater samples were collected and analyzed across all WWTPs

during the study period. *C. auris* concentrations in these samples ranged from below the limit of

detection to 1,456,751 cp/g, with 1.45% (n = 200 samples) of samples above the lowest

detectable concentration. Sixty five (34.21%) WWTPs across 26 states had at least one

detectable *C. auris* concentration during the study period, and the percentage of positive

samples from these WWTPs ranged between 0.58%–46.48% (median: 1.49%). Time series

heatmaps of *C. auris* detections at each of the 190 WWTPs are provided in Figure S5.

Seasonally, 2.00% (n=6754) of observations were positive during autumn (defined as Sep 1 to

Nov 30), and 1.01%(n=7245) of observations were positive during winter (defined as Dec 1 to

Feb 29). This difference was statistically significant when assessed by a bootstrapping

approach (p<0.0001).

Clinical data from the National Notifiable Disease Surveillance System (NNDSS) were

downloaded for comparison to wastewater data, and no significant correlations were found

between wastewater detections and population adjusted clinical case rates (Spearman's rho

between -0.10 and 0.23 for the 18 states, all p>0.05). Several states had obvious divergence

between wastewater positivity rates and population adjusted NNDSS clinical case rates (**Figure

2**). For example, in Maine, 13.3% of wastewater samples were positive for *C. auris* despite not

reporting any clinical cases during the study period to NNDSS (**Figure 2**). Twelve states which

did not report *C. auris* cases to the NNDSS were found to have positive detections of *C. auris*.

Figure 3 shows percentage positive detections as a function of population normalized clinical

cases and highlights the limited clinical data available.

a. Cases

b. Wastewater

Figure 2: Map showing percentage of positive wastewater samples and population normalized case numbers from NNDSS across the United States. **Figure 2a** shows the normalized number of cases per million in each state as reported to NNDSS during the study period. **Figure 2b** shows the percentage of positive wastewater samples in each of the states surveilled during the study period. Yellow represents low normalized values while black represents high normalized values. Gray states represent areas without available data.

380
381 **Figure 3.** Comparison of wastewater percent positive detections and total clinical cases as
382 reported by NNDSS. Only states with clinical data reported to NNDSS are included. Each
383 labeled point represents the percentage of positive wastewater samples in that state and the
384 number of cases reported to NNDSS. The state abbreviation is above each data point.
385
Population demographics and healthcare facility locations and *Candida auris.*

WWTPs were grouped using three separate statistics: median age of the state, number of

hospitals in the sewershed, and number of nursing homes in the sewershed (Fig 4). The median

age in the United States is 39 years as of the latest American Community Survey (ACS) data

(31). Seventy-eight WWTPs were located in a state with a median age at or above the national

median age; 112 WWTPs were located in a state with a median age below the national median

age. As shown in **Figure 4a**, among WWTPs in states with a median age at or above the

national median age, 2.03% (95% CI: [0.80-3.50]) of all wastewater samples were positive for *C.*

auris. Among WWTPs in states with a median age below the national median age, 0.83% (95%

CI: [0.34-1.38]) of all wastewater samples were positive for *C. auris*. This difference was

statistically significant when assessed by bootstrapping (p < 0.001).

The median number of hospitals in the sewersheds was 1 hospital (range: 0–74 hospitals).

WWTPs were grouped by whether they contained 0 hospitals (fewer than the median, n = 54) or

1 or more hospitals in their sewersheds (equal to or greater than the median, n = 136). We

found that 1.78% (95% CI: [0.92-2.90], n = 10,019) of samples from WWTPs with at least 1

hospital in their sewersheds were positive for *C. auris* whereas 0.62% (95% CI: [0.34-1.11], n =

3,819) of samples from WWTPs with no hospital in their sewershed were positive for *C. auris*

(**Figure 4b**). This difference was significant when assessed by bootstrapping (p < 0.0001).

The median number of nursing homes in the 190 sewersheds was 7 (range: 0–420 nursing

homes). WWTPs were grouped by whether they contained 0–6 nursing homes (fewer than the

median, n = 91) or 7 or more nursing homes in their sewersheds (equal to or greater than the

median, n = 99). We found that 1.92% (95% CI: [0.89–3.46], n = 7,544) of samples from

WWTPs with 7 or more nursing homes in their sewersheds were positive for *C. auris* whereas

0.87% (95% CI: [0.47–1.47], n = 6,294) of samples from WWTPs with 0–7 nursing homes in

their sewersheds were positive for *C. auris* (**Figure 4c**). This difference was significant when

assessed by bootstrapping (p < 0.001).

415
Figure 4. Comparison of wastewater detections across different groups. *** indicates p<0.001,
**** indicates p<0.0001. **Figure 4a** shows the percentage of positive samples for states at or
above the median age compared to those below. **Figure 4b** shows the percentage of positive
samples when grouped by number of hospitals at the sewershed level. **Figure 4c** shows the
same statistic when grouped by number of nursing homes, again at the sewershed level. Error
bars represent 95% confidence intervals derived from the bootstrapping approach.
422

Discussion

Previous studies at the sewershed scale have shown that detection of *C. auris* in wastewater is

indicative of ongoing outbreaks in the contributing population (19, 25). To our knowledge, this

study represents the first nationwide wastewater monitoring effort of *C. auris* and comparison to

national clinical case data (23). We first demonstrate the sensitivity and specificity of the assay

used in this study through comparisons to published assays, sequencing (Figure S2, S4), and in

silico analyses. The widespread detection of *C. auris* in wastewater suggests a significant gap in

clinical case data reported to the NNDSS. Indeed, it is known that many local jurisdictions do

not provide data for inclusion in NNDSS. We found that *C. auris* detections are more frequent in

states with older populations and in sewersheds with higher numbers of nursing homes and

hospitals. These findings are consistent with *C. auris* infections being more common in older,

more susceptible populations and further support the assertion that clinical case data may be

incomplete.

C. auris has gone from initial detection in the United States in 2013 to being declared an

"Urgent" and notifiable pathogen in 2019 by the United States (CDC) (14, 20, 38). The rate of

testing volume and clinical positives has rapidly increased, with 17 states identifying their first *C.*

auris cases between 2019 to 2021 (14). Despite this, clinical testing for *C. auris* is far from

ubiquitous and not conducted uniformly across the United States (14, 39). On-site PCR testing

of clinical specimens is rare, and samples often have to be sent to public health or reference

laboratories, further increasing the cost of testing and the turnaround time before the data

reaches public health officials (39). Detections of *C. auris* nucleic-acids in wastewater collected

throughout the United States demonstrate a need for increased clinical testing and reporting

capacity, and the capability of wastewater testing to serve as a sentinel surveillance mechanism

for emerging pathogens.

C. auris is an opportunistic pathogen that often infects the elderly and immunocompromised

within long-term healthcare facilities (10, 20, 40). Following our detection of *C. auris*, we

investigated potential demographics and healthcare facilities related to its pathology using data

from the ACS and US Department of Homeland Security's HIFLD database (31, 41). We

analyzed age, number of hospitals, and number of nursing homes as plausible factors impacting

the presence of *C. auris* within a sewershed or state. We found significant differences between

the percentage of positive detections of *C. auris* in states with a median age less than versus

greater than or equal to the national median age. Similarly, we found significant differences

between the percentage of positive detections of *C. auris* in sewersheds with less than versus

greater than or equal to the median number of hospitals and nursing homes. States with a

population above the national median age were more than three times as likely to detect *C.*

auris in a given wastewater sample. At the sewershed level, locations with 1 or more hospitals

had three times the amount of *C. auris* detections than locations with no hospitals. Similarly,

sewersheds with greater than or equal to the median number of nursing homes had over twice

as many detections of *C. auris* as those with fewer than the median number of nursing homes.

These findings are consistent with the known epidemiology of *C. auris*, as infected patients'

median age is 68 and outbreaks are prevalent in long-term healthcare facilities and nursing

homes (5, 14)*.* The presence of an at-risk population appears to be strongly associated with *C.*

auris wastewater detections, suggesting public health efforts are appropriately focused on these

at-risk populations.

Significant steps were taken to ensure the validity of detection results. A similar assay was used

by Barber et al. for the detection of *C. auris* in southern Nevada during an outbreak (25). We

tested nine samples from three different states using both the assay presented in this study and

the one used by Barber et al. (25). No statistically significant difference was found between the

two assays, which both targeted the same region of an internally transcribed spacer region. Our

assay was further validated using a subset of 5 of these 9 samples through sequencing. The

amplicons generated by our forward and reverse primers independently and accurately matched

the reference amplicon, indicating specific amplification (Figure S4). Together, these results

demonstrate that the assay used in this study is highly specific. Finally, we used an RT-PCR

assay to detect *C. auris* which means the assay should detect both RNA and DNA. We

measured some samples using both RT-PCR and PCR and the results indicated that most of

the target nucleic acids in those samples were DNA. This was somewhat surprising since we

expect the DNA target (located in the region of the genome that encodes ribosomal RNA)

should be transcribed into RNA by *C. auris*. Future work will need to be done to further

investigate the relative proportion of target RNA versus DNA *C. auris* targets in wastewater.

Regardless, the use of an RT-PCR assay, allowing detection of both RNA and DNA, should

increase the sensitivity of the assay (by detecting both nucleic-acids) yet perhaps complicate

interpretation of measured concentrations. For this reason, we only use the data in

presence/absence format.

Overall, our results demonstrate that *C. auris* wastewater surveillance can provide timely

information on geographical distribution and help identify at-risk populations; wastewater data

can be available within 24-48 hours of sample collection. As an emerging disease, clinical

testing for *C. auris* largely depends on shipping samples to public health laboratories (7, 13, 14).

Wastewater surveillance can help fill this gap for municipalities and states without the capital to

fund testing systems, providing community level information as to the prevalence, and more

importantly, the spread of *C. auris*. These data are critical to public health responses,

particularly when dealing with pathogens that do not or did not have robust clinical testing

systems such as Dengue virus, SARS-CoV-2, and *C. auris* (14, 16, 17, 39). *C. auris* wastewater

surveillance data can then be used to implement stricter screening protocols, cleaning

protocols, and other non-pharmaceutical interventions that can prevent the spread of illness

amongst vulnerable populations (9, 10, 20).

There are limitations associated with this study. Importantly, the availability of clinical data was

limited due to the recent classification in 2018 of *C. auris* as a notifiable disease, making in-

depth analyses of associations between wastewater detections and clinical data on infections

unfeasible. In addition, it is well understood that the NNDSS database may not contain all the

information on C. auris positive cases in the United States; local jurisdictions may have local

data to inform public health response to infections and those data were not available for this

work. Local jurisdictions are encouraged to use these wastewater data to compare to their local

case rates. Further, wastewater sampling was not uniform across the United States or within

individual states, which could introduce biases into the analyses. Within the demographic data

available at the sewershed level, information was limited as to the size of nursing homes and

hospitals, which might be important factors controlling associations with wastewater data on *C.*

auris. Lastly, we were unable to link specific wastewater concentrations to population-level

incidence. Further experiments are necessary to understand the shedding patterns of *C. auris* in

human excretions as to provide this direct link to disease occurrence in the contributing

population. These experiments are particularly important to the determination of the significance

of wastewater detections, as *C. auris* contributions to wastewater could potentially be from

colonized individuals or individuals with acute infection. Finally, an additional limitation is that we

cannot rule out the potential for *C. auris* detections in wastewater to be from zoonotic or

environmental sources. Detections have been shown to match underlying expected

demographic indicators such as age and presence of nursing homes, further lending credence

to the source of detections being human. While there is no evidence to support the possibility of

zoonotic or environmental contributions of *C. auris* nucleic-acids to wastewater, further research

is needed to see if such evidence might exist. There have been case reports of *C. auris*

infections in canines, and *Candida spp*. have been detected using culture and other molecular

methods from biofilms in sewer pipes (42–45). It should be noted that wastewater detection of

Candida auris collected at the wastewater treatment plant scale does not imply community-

spread of the pathogen, but is consistent with its presence in one or more individuals within the

sewershed.

Acknowledgements. We acknowledge all the wastewater treatment plant staff who provided

samples for this project.

References

541 1. Ageing and health. https://www.who.int/news-room/fact-sheets/detail/ageing-and-health.

542   Retrieved 17 January 2024.

543 2. Hospital-Acquired Infections - ClinicalKey.

544   https://www.clinicalkey.com/#!/content/playContent/1-s2.0-S0039610911001514. Retrieved

545   16 January 2024.

546 3. Serra-Burriel M, Keys M, Campillo-Artero C, Agodi A, Barchitta M, Gikas A, Palos C,

547   López-Casasnovas G. 2020. Impact of multi-drug resistant bacteria on economic and

548   clinical outcomes of healthcare-associated infections in adults: Systematic review and

549   meta-analysis. PLOS ONE 15:e0227139.

550 4. Ahmad S, Alfouzan W. 2021. Candida auris: Epidemiology, Diagnosis, Pathogenesis,

551   Antifungal Susceptibility, and Infection Control Measures to Combat the Spread of

552   Infections in Healthcare Facilities. Microorganisms 9:807.

553 5. Benedict K, Forsberg K, Gold JAW, Baggs J, Lyman M. Candida auris–Associated

554   Hospitalizations, United States, 2017–2022 - Volume 29, Number 7—July 2023 - Emerging

555   Infectious Diseases journal - CDC https://doi.org/10.3201/eid2907.230540.

556 6. Lone SA, Ahmad A. 2019. Candida auris—the growing menace to global health. Mycoses

557   62:620–637.

558 7. 2023. Surveillance for Candida auris | Candida auris | Fungal Diseases | CDC.

559   https://www.cdc.gov/fungal/candida-auris/c-auris-surveillance.html. Retrieved 17 January

560   2024.

561 8. Sengupta S, Marimuthu K, Stewardson A, Harbarth S, Durante A, Singh S. 2020.

562   Challenges in Identification of Candida auris in Hospital Laboratories: Comparison

563   Between HIC and LMIC. Infect Control Hosp Epidemiol 41:s158–s158.

9.   Piedrahita CT, Cadnum JL, Jencson AL, Shaikh AA, Ghannoum MA, Donskey CJ. 2017.

Environmental Surfaces in Healthcare Facilities are a Potential Source for Transmission of

Candida auris and Other Candida Species. Infect Control Hosp Epidemiol 38:1107–1109.

10.   Welsh RM, Bentz ML, Shams A, Houston H, Lyons A, Rose LJ, Litvintseva AP. 2017.

Survival, Persistence, and Isolation of the Emerging Multidrug-Resistant Pathogenic Yeast

Candida auris on a Plastic Health Care Surface. J Clin Microbiol 55:2996–3005.

11.   US EPA O. 2020. List P: Antimicrobial Products Registered with EPA for Claims Against

Candida Auris. Overviews and Factsheets. https://www.epa.gov/pesticide-registration/list-

p-antimicrobial-products-registered-epa-claims-against-candida-auris. Retrieved 25

January 2024.

12.   Karmarkar EN, O'Donnell K, Prestel C, Forsberg K, Gade L, Jain S, Schan D, Chow N,

McDermott D, Rossow J, Toda M, Ruiz R, Hun S, Dale JL, Gross A, Maruca T, Glowicz J,

Brooks R, Bagheri H, Nelson T, Gualandi N, Khwaja Z, Horwich-Scholefield S, Jacobs J,

Cheung M, Walters M, Jacobs-Slifka K, Stone ND, Mikhail L, Chaturvedi S, Klein L,

Vagnone PS, Schneider E, Berkow EL, Jackson BR, Vallabhaneni S, Zahn M, Epson E.

2021. Rapid Assessment and Containment of Candida auris Transmission in Postacute

Care Settings—Orange County, California, 2019. Ann Intern Med 174:1554–1562.

13.   2023. Tracking Candida auris | Candida auris | Fungal Diseases | CDC.

https://www.cdc.gov/fungal/candida-auris/tracking-c-auris.html. Retrieved 17 January

2024.

14.   Worsening Spread of Candida auris in the United States, 2019 to 2021 | Annals of Internal

Medicine. https://www.acpjournals.org/doi/full/10.7326/M22-3469. Retrieved 17 January

2024.

15.   Boehm AB, Hughes B, Doung D, Chan-Herur V, Buchman A, Wolfe MK, White BJ. 2022.

Wastewater surveillance of human influenza, metapneumovirus, parainfluenza, respiratory

syncytial virus (RSV), rhinovirus, and seasonal coronaviruses during the COVID-19

590   pandemic. preprint. Infectious Diseases (except HIV/AIDS).

591 16. Peccia J, Zulli A, Brackney DE, Grubaugh ND, Kaplan EH, Casanovas-Massana A, Ko AI,

592   Malik AA, Wang D, Wang M, Warren JL, Weinberger DM, Arnold W, Omer SB. 2020.

593   Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics.

594   Nat Biotechnol 38:1164–1167.

595 17. Wolfe MK, Paulos AH, Zulli A, Duong D, Shelden B, White BJ, Boehm AB. 2024.

596   Wastewater Detection of Emerging Arbovirus Infections: Case Study of Dengue in the

597   United States. Environ Sci Technol Lett 11:9–15.

598 18. Pacilli M, Kerins JL, Clegg WJ, Walblay KA, Adil H, Kemble SK, Xydis S, McPherson TD,

599   Lin MY, Hayden MK, Froilan MC, Soda E, Tang AS, Valley A, Forsberg K, Gable P,

600   Moulton-Meissner H, Sexton DJ, Jacobs Slifka KM, Vallabhaneni S, Walters MS, Black

601   SR. 2020. Regional Emergence of Candida auris in Chicago and Lessons Learned From

602   Intensive Follow-up at 1 Ventilator-Capable Skilled Nursing Facility. Clin Infect Dis

603   71:e718–e725.

604 19. Babler K, Sharkey M, Arenas S, Amirali A, Beaver C, Comerford S, Goodman K, Grills G,

605   Holung M, Kobetz E, Laine J, Lamar W, Mason C, Pronty D, Reding B, Schürer S,

606   Schaefer Solle N, Stevenson M, Vidović D, Solo-Gabriele H, Shukla B. 2023. Detection of

607   the clinically persistent, pathogenic yeast spp. Candida auris from hospital and municipal

608   wastewater in Miami-Dade County, Florida. Sci Total Environ 898:165459.

609 20. 2023. Infection Prevention and Control for Candida auris | Candida auris | Fungal

610   Diseases | CDC. https://www.cdc.gov/fungal/candida-auris/c-auris-infection-control.html.

611   Retrieved 25 January 2024.

612 21. Kordalewska M, Zhao Y, Lockhart SR, Chowdhary A, Berrio I, Perlin DS. 2017. Rapid and

613   Accurate Molecular Identification of the Emerging Multidrug-Resistant Pathogen Candida

614   auris. J Clin Microbiol 55:2445–2452.

615 22. BLAST: Basic Local Alignment Search Tool. https://blast.ncbi.nlm.nih.gov/Blast.cgi.

Retrieved 29 February 2024.

23.   National Notifiable Diseases Surveillance System (NNDSS) - Healthy People 2030 |

health.gov. https://health.gov/healthypeople/objectives-and-data/data-sources-and-

methods/data-sources/national-notifiable-diseases-surveillance-system-nndss. Retrieved

26 January 2024.

24.   Weston E (CDC/OID/NCIRD). Guidance on classifying STD case reports into MMWR

week.

25.   Barber C, Crank K, Papp K, Innes GK, Schmitz BW, Chavez J, Rossi A, Gerrity D. 2023.

Community-Scale Wastewater Surveillance of Candida auris during an Ongoing Outbreak

in Southern Nevada. Environ Sci Technol 57:1755–1763.

26.   Roldan-Hernandez L, Boehm AB. 2023. Adsorption of Respiratory Syncytial Virus,

Rhinovirus, SARS-CoV-2, and F+ Bacteriophage MS2 RNA onto Wastewater Solids from

Raw Wastewater. Environ Sci Technol 57:13346–13355.

27.   Boehm A, Wolfe M, Wigginton K, Bidwell A, White B, Duong D, Hughes B, Chan-Herur V,

Naughton C, Bischel H. 2023. Human viral nucleic acids concentrations in wastewater

solids from Central and Coastal California, USA. Boehm Res Group Stanf.

28.   Human viral nucleic acids concentrations in wastewater solids from Central and Coastal

California USA | Scientific Data. https://www-nature-

com.stanford.idm.oclc.org/articles/s41597-023-02297-7. Retrieved 14 October 2023.

29.   McClary-Gutierrez JS, Aanderud ZT, Al-faliti M, Duvallet C, Gonzalez R, Guzman J, Holm

RH, Jahne MA, Kantor RS, Katsivelis P, Kuhn KG, Langan LM, Mansfeldt C, McLellan SL,

Grijalva LMM, Murnane KS, Naughton CC, Packman AI, Paraskevopoulos S, Radniecki

TS, Roman FA, Shrestha A, Stadler LB, Steele JA, Swalla BM, Vikesland P, Wartell B,

Wilusz CJ, Wong JCC, Boehm AB, Halden RU, Bibby K, Vela JD. 2021. Standardizing

data reporting in the research community to enhance the utility of open data for SARS-

CoV-2 wastewater surveillance. Environ Sci Water Res Technol 7:1545–1551.

30. Rosario K, Symonds EM, Sinigalliano C, Stewart J, Breitbart M. 2009. Pepper mild mottle

643   virus as an indicator of fecal pollution. Appl Environ Microbiol 75:7261–7267.

644   31. U.S. Census Bureau. S0101: Age and Sex - Census Bureau Table. Am Community Surv

645   ACS 1-Year Estim Subj Tables Table S0101.

646   https://data.census.gov/table/ACSST1Y2022.S0101?q=median%20age%20by%20state&g

647   =010XX00US. Retrieved 12 February 2024.

648   32.   Nursing Homes. https://hifld-geoplatform.hub.arcgis.com/datasets/geoplatform::nursing-

649   homes/about. Retrieved 13 March 2024.

650   33.   Hospitals. https://hifld-geoplatform.hub.arcgis.com/datasets/geoplatform::hospitals/about.

651   Retrieved 13 March 2024.

652   34.   USA ZIP Code Boundaries.

653   https://hub.arcgis.com/datasets/d6f7ee6129e241cc9b6f75978e47128b_0/about. Retrieved

654   6 March 2024.

655   35. Efron B. 1979. Bootstrap Methods: Another Look at the Jackknife. Ann Stat 7:1–26.

656   36. DICICCIO T, EFRON B. 1992. More accurate confidence intervals in exponential families.

657   Biometrika 79:231–245.

658   37. Borchardt MA, Boehm AB, Salit M, Spencer SK, Wigginton KR, Noble RT. 2021. The

659   Environmental Microbiology Minimum Information (EMMI) Guidelines: qPCR and dPCR

660   Quality and Reporting for Environmental Microbiology. Environ Sci Technol 55:10210–

661   10223.

662   38. Centers for Disease Control and Prevention (U.S.). 2019. Antibiotic resistance threats in

663   the United States, 2019. Centers for Disease Control and Prevention (U.S.).

664   39. Arenas S, Patel S, Seely SO, Pagan PP, Warde PR, Tamrakar LJ, Parekh DJ, Ferreira T,

665   Zhou Y, Gershengorn HB, Shukla BS. 2023. Operational impact of decreased turnaround

666   times for Candida auris screening tests in a tertiary academic medical center. Antimicrob

667   Steward Healthc Epidemiol ASHE 3:e176.

668   40.   Schwartz IS, Dingle TC. 2019. Candida auris. CMAJ Can Med Assoc J 191:E865.

669   41.   HIFLD Open. https://hifld-geoplatform.hub.arcgis.com/pages/hifld-open. Retrieved 19

670   March 2024.

671   42.   White TC, Esquivel BD, Rouse Salcido EM, Schweiker AM, dos Santos AR, Gade L, Petro

672   E, KuKanich B, KuKanich KS. 2024. Candida auris detected in the oral cavity of a dog in

673   Kansas. mBio 15:e03080-23.

674   43.   Yadav A, Wang Y, Jain K, Panwar VAR, Kaur H, Kasana V, Xu J, Chowdhary A. 2023.

675   Candida auris in Dog Ears. J Fungi 9:720.

676   44.   Cavalheiro M, Teixeira MC. 2018. Candida Biofilms: Threats, Challenges, and Promising

677   Strategies. Front Med 5:28.

678   45.   Nobile CJ, Johnson AD. 2015. Candida albicans Biofilms and Human Disease. Annu Rev

679   Microbiol 69:71–92.