iGTP: Learning interpretable cellular embedding for inferring biological mechanisms

underlying single-cell transcriptomics

Kang-Lin Hsieh[1,†], Kai Zhang[2,†], Yan Chu[3,4,†], Lishan Yu[2], Xiaoyang Li[5,6], Nuo Hu[5,7], Isha

Kawosa[5,8], Patrick G. Pilié[1], Pratip K. Bhattacharya[9], Degui Zhi[4], Xiaoqian Jiang[2], Zhongming

Zhao[5,10,*], Yulin Dai[5,*]

[1]Department of Genitourinary Medical Oncology, Division of Cancer Medicine, UT MD Anderson

Cancer Center, Houston, TX 77030, USA

[2]Center for Secure Artificial Intelligence for Healthcare, School of Biomedical Informatics, The

University of Texas Health Science Center at Houston, Houston, TX 77030, USA

[3]Department of Radiation Physics, Division of Radiation Oncology, UT MD Anderson Cancer

Center, Houston, TX 77030, USA

[4]Center for Artificial Intelligence and Genome Informatics, School of Biomedical Informatics, The

University of Texas Health Science Center at Houston, Houston, TX 77030, USA

[5]Center for Precision Health, McWilliams School of Biomedical Informatics, The University of

Texas Health Science Center at Houston, Houston, TX 77030, USA

[6]Department of Biostatistics and Data Science, School of Public Health, The University of Texas

Health Science Center at Houston, Houston, TX 77030, USA

[7]Department of Statistics, Rice University, Houston, TX 77005, USA

[8]Department of Computer Science and Engineering, Texas A&M University, College Station, TX

77843, USA

[9] Department of Cancer Systems Imaging, The University of Texas MD Anderson Cancer

Center, Houston, TX 77030, USA

[10]Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN

37203, USA

†These three authors contributed equally to this work.

*To whom correspondence should be addressed:

Yulin Dai, Ph.D.

Center for Precision Health

McWilliams School of Biomedical Informatics

The University of Texas Health Science Center at Houston

7000 Fannin St. Suite E760A Houston, TX 77030

Phone: 713-500-3462

Email: Yulin.Dai@uth.tmc.edu

Zhongming Zhao, Ph.D.

Center for Precision Health

McWilliams School of Biomedical Informatics

The University of Texas Health Science Center at Houston

7000 Fannin St., Suite 600, Houston, TX 77030

Phone: 713-500-3631

Email: Zhongming.Zhao@uth.tmc.edu

Abstract

Deep-learning models like Variational AutoEncoder have enabled low dimensional cellular embedding representation for large-scale single-cell transcriptomes and shown great flexibility in downstream tasks. However, biologically meaningful latent space is usually missing if no specific structure is designed. Here, we engineered a novel interpretable generative transcriptional program (iGTP) framework that could model the importance of transcriptional program (TP) space and protein-protein interactions (PPI) between different biological states. We demonstrated the performance of iGTP in a diverse biological context using gene ontology, canonical pathway, and different PPI curation. iGTP not only elucidated the ground truth of cellular responses but also surpassed other deep learning models and traditional bioinformatics methods in functional enrichment tasks. By integrating the latent layer with a graph neural network framework, iGTP could effectively infer cellular responses to perturbations. Lastly, we applied iGTP TP embeddings with a latent diffusion model to accurately generate cell embeddings for specific cell types and states. We anticipate that iGTP will offer insights at both PPI and TP levels and holds promise for predicting responses to novel perturbations.

Keywords: single-cell transcriptomics, interpretable deep learning, transcriptional program, Variational AutoEncoder, Graph Neural Network.

Introduction

With the rapid advent of single-cell technology, millions of cellular transcriptomes have been generated by large consortia [e.g. Human Cell Atlas[1], Tabula Sapiens[2]] and various individual studies to advance our understanding of the human body, developmental biology, tumor heterogeneity, and immune system dynamics, leading to more precise diagnostic and therapeutic strategies[3]. The massive increase in cellular transcriptomes also provides opportunities to use deep-learning frameworks to learn and represent heterogeneous biological functions.

The deep-learning framework Variational AutoEncoder (VAE)[4], has been widely adapted for dimension reduction[5] and a variety of other applications[6,7,8] in high-dimensional single-cell RNA-seq data analysis. By leveraging its probabilistic nature, VAE effectively addresses challenges such as batch effect correction[7], imputation[9], and perturbation response prediction[8,10], which shows significant advantages over traditional statistical methods in terms of scalability and the ability to detect nonlinear patterns[11,12]. Deep generative models excel in their specific modeling tasks but often lack interpretability and struggle to provide biologically relevant latent representations of protein-protein interactions (PPI) and biological pathways. Without implementing specific model structures, these models cannot directly capture variations in biological processes[8]. Elmarakeby et al. (P-NET)[13] introduced a new fashion of using fully connected layers to encode various levels of biological information, including genes, pathways, and biological processes, for outcome prediction, providing a biologically meaningful interpretation via the weight between layers. However, P-NET is primarily designed for prediction purposes and is not a generative model. More recently, methods such as single-cell embedded topic model (scETM)[14] engineered the Decoder layer to ensure the embedding layers accurately represent gene and topic embeddings, as well as prior gene modules. However, the topics from the scETM cannot be directly linked to biological meaning. The VAE

Enhanced by Gene Annotations (VEGA)[15] designed a VAE model to learn gene module

variables (GMV) space via employing a sparse linear Decoder with masking to preserve a

connection from a GMV to an output gene exclusively if the gene is recognized as part of the

specific gene module. However, this one-layer Decoder simplified the linkage between genes

and GMV, neglecting to incorporate prior knowledge like PPIs.

To advance beyond these constraints, we propose iGTP (interpretable generative transcriptional

program), a VAE with a masked multi-layer linear Decoder informed by biological

pathways/transcriptional programs (TPs) and protein-protein interactions. iGTP offers the

following functions: 1) direct interpretable latent space for TPs related to biological state

alteration and use these high dimensions to represent cellular embeddings; 2) uncover

previously unknown associations between PPIs/TPs and cellular state alteration; 3) infer TP

activity in unseen data using pretrained model; 4) infer gene expression alteration for unseen

perturbation; 5) assess the relative importance for TPs and PPIs, providing new insights for the

underlying biological mechanism; 6) demonstrate the generative capability of TP embeddings in

creating new cells corresponding to specified cell types and condition.

Results

iGTP framework

We developed a customized VAE model to capture the biologically meaningful embedding in

latent spaces for scRNA-seq data, enabling the interpretation of transcriptional shifts and

leveraging Graph Neural Networks (GNN) for propagating gene alteration inferences through

the PPI network (Fig. 1). To ensure a biologically interpretable latent layer, we engineered a

sparse multilayer Decoder with prior biological knowledge of transcription programs and PPIs,

which explicitly guides the model to learn the differential TP and PPI responsible for cellular

states shift. Similar to the previous graph-enhanced gene activation and repression simulator

(GEARS) model[16], we further incorporate the latent TP embeddings with gene-perturbation PPI

embedding to predict the perturbation response of TPs and genes on unseen scenarios of

certain genes.

Distilling biological insights with interpretable latent space framework

We first evaluate the capacity of the iGTP model to unveil the ground-truth biological processes

in interferon-β-treated Peripheral Blood Mononuclear Cells (PBMC) dataset[17]. We input the

single-cell dataset and enforce iGTP to learn the gene ontology (GO) as the TP and PC

commons as the PPIs. After training, we reparametrized the latent space Z generated from

iGTP on 2D t-distributed stochastic neighbor embedding (TSNE). As shown in Fig. 2a & 2b,

iGTP differentiated both cell types and stimulation states. In Fig. 2c, we found that the TP

"GO:0035456: response to interferon-β" simulation had overall higher activities in stimulated

cells, confirming the ability of iGTP to capture pathway activity in its latent space. Comparable

cellular "GO:0035455: response to interferon-α" and "GO:0035456: response to interferon-β"

were observed in immune cells[18]. The iGTP framework captured the higher PPI activity between

ISH15 and IFNB1 (Interferon Beta 1) within the BP TP of "GO:0035456: response to interferon-β"

in the stimulated cells (Fig. 2d). We also benchmarked against the conventional single sample

Gene Set Enrichment Analysis (ssGSEA) method[19,20] to calculate the activity of the same TPs in

each cell. As shown in Fig. 2e, the TP activity of response to interferon-β is much lower (or

close to resting status) in stimulated adaptive immune cells (CD8+ T cells, CD4+ T cells, and B

cells) than in other stimulated innate immune cells (NK cells, Dendritic cells, FCGR3A+

monocytes, and CD14+ monocytes). Compared to Fig. 2c & 2e, both algorithms exhibited

similar patterns, but iGTP demonstrated a stronger change in biological activity. The

phenomenon was further validated by a recently published paper[18], suggesting ssGSEA and

iGTP could capture the response to interferon-β activity. As shown in Fig. 2f, the volcano plot for

differential TPs across all cell types between stimulated vs control with plaid-based Bayers

factor as the fold changes magnitude (Table S1). The top stimulated TPs were BP of

"GO:0009615: response to virus", BP of "GO:0034340: response to type I interferon", and other

interferon-related TPs. We also benchmarked the conventional gene-set enrichment analysis

(GSEA) method to measure the activity of the same TPs in each cell type with iGTP results. In

Fig. 2g, we showed the TPs in both Bayers factor from iGTP and normalized enrichment score

from GSEA. We pinpointed that BP of "GO:0009615: response to virus" and other interferon-

related TPs were in the top-right corners of CD8+ T cells (Table S2) and CD14+ monocytes

(Table S3). Interestingly, the BP TP of "GO:0045088: regulation of innate immune response"

was predicted to be much lower in CD8+ T cells than CD14+ monocytes by iGTP. On the other

hand, GSEA predicted it among the top TPs in both cell types, which is contradicted by the fact

that TP "regulation of innate immune response" is mainly regulated in innate immune cells

instead of CD8+ T cells[21]. Collectively, these findings imply that iGTP TP accurately represents

the anticipated primary biological pathways in PBMC, thereby potentially serving as a valuable

tool for projecting cells from other datasets into an interpretable framework. This approach

enables the detailed exploration of cell-type-specific patterns at the level of cellular processes.

Lastly, we leveraged this PBMC dataset to comprehensively compare our iGTP model and

another deep learning model VEGA, with different knowledge of PPI and TP databases. Then,

we evaluated the model performance using NMI, ARI, silhouette metrics from scvi-tools[22], and

graph connectivity for evaluation. Overall, iGTP (GO TP+PPI compiled) outperforms ssGSEA

and VEGA, as evidenced by the average of all evaluation metrics (Fig. 2h).

iGTP model captures both known and unknown TP activities related to brain cellular

function and Alzheimer's disease.

To explore the generalizability of our iGTP model, we analyzed brain cells from Alzheimer's

disease (AD) and cognitively normal (CN) individuals who share similar AD risk backgrounds.

We adapted AD and brain function-related TP curation from our previous work[23]. The

normalized mean TP activities by each cell type and disease condition were visualized in Fig. 3a.

We identified the TP activities that were enriched with corresponding cell types, such as BP of

"adaptive immune response in Microglia", cellular component (CC) of Synapse in both inhibitory

neurons and excitatory neurons, suggesting TP embedding from our iGTP model could capture

meaningful biological activities. Moreover, we also identified a few well-known AD-related TP

with more activated states, such as molecular function (MF) of "GO:0048156: Tau protein

binding" in AD excitatory neurons[24,25] and BP of "GO:1990000: Amyloid fibril formation" in AD

excitatory and inhibitory neurons[26]. The embeddings of these TPs were further visualized in 2D

(Fig. 3b) for astrocyte and microglia. The dot shifts between AD and CN indicate our TP

embeddings indicate biological differences in AD and CN. More interestingly, we delineated PPI

connections, PTPRC-CD4[27] and SORL1-APOE, linked to BP "GO:0002250: Adaptive immune

response" and BP of "GO:1990000: Amyloid fibril formation" respectively, across cell types. Fig.

3c illustrates the cell-type-specific PPI activities in excitatory neurons and microglia between AD

and CN cells. To identify potential unknown GO TPs that predominantly distinguish AD from CN

by iGTP, we highlighted the 15 most variable TPs (Fig. 3d), including a few neurotransmitter

receptors (such as MF of "GO:0008066: glutamate receptor activity", MF of "GO:0016917:

GABA receptor activity", and MF of "GO:0005539: glycosaminoglycan binding") have higher

activities in excitatory neurons, inhibitory neurons, and OPCs. MHC protein complex was mainly

expressed in Microglia, OPCs, and Oligodendrocytes. The corresponding TP activities from

ssGSEA (Fig. S1) exhibit a median Spearman correlation coefficient ρ of 0.74 across cell types

for these TPs. As shown in Fig. 3e, the difference in cellular embedding distribution between

AD and CN indicates our TP embeddings in excitatory neurons and microglia in synapse-related

functions (BP of "GO:0050808: synapse organization" and CC of "GO:0005874: microtubule").

In our analysis of these TPs, we specifically identified PPIs with significant differences between

AD and CN cells. For example, the NTRK2-DYNLL1 link in the BP of synapse organization

showed a notable disparity, where a variant in NTRK2 has been previously associated with AD

in earlier research[28]. As Fig. 3f illustrates, the PPI between MTUS1-CLTC in the BP of

microtubule formation exhibits distinct cellular activities in AD compared to CN individuals. This

observation is consistent with previous studies showing elevated MTUS1 expression in specific

brain regions of AD patients relative to CN controls[29].

Generalizability to predict out-of-sample single-cell PBMC data

Next, we examined if iGPT could generalize to accurately infer an interpretable latent

representation of data not seen during training (out-of-sample data). We evaluated iGTP in two

settings for this purpose. In the first case, we assessed the biological generalization of iGTP's

inference by excluding some pairs from the same dataset during training. Specifically, we

explored whether the inferred iGTP activities for the omitted cells conveyed meaningful

biological information while not in training. For this, we excluded cells infected by influenza from

the COVID-19 dataset during training. Then, we inferred the PBMC "GO:0009615: response to

virus"-related TP activities on the removed samples (influenza) using the model trained by the

remaining cells. As shown in Fig. 4a, CD14+ monocytes and dendritic cells from severe

influenza patients tend to have higher activities than cells from COVID-19 Low-Dose Naltrexone

(LDN) treatment or healthy controls in BP TP of "GO:0034340: response to type 1 interferon"

and TP "GO:0009615: response to virus". We selected the 6 interferon-related TPs (Table S4)

and 4806 related protein-protein interactions (PPIs) which represent the biological activities of

virus invasion across various cell types. We tested the Pearson correlation coefficient r of the

Bayes factor in pretrained samples and predicted samples. All cell types, except for B cells,

have an intra-sample correlation of $r > 0.4$ (Fig. 4b & 4c). Next, we explored whether the iGTP

model could predict this unseen similar cellular condition population from a new batch study.

Here, we trained our model using the COVID-19 dataset and predicted the TP activities in the

Kang et al. dataset. As shown in Fig. 4d, CD14+ monocytes and dendritic cells from the Kang et

al. dataset tend to have higher activities than in interferon-β stimulated cells and lower activities

in resting cells from COVID-19 dataset in BP TP of "GO:0034340: response to type 1 interferon"

and TP "GO:0009615: response to virus". We further tested the coefficient r of the Bayes factor

between the COVID-19 dataset and predicted the Kang et al. dataset by each cell type. The

overall prediction performance in Fig. 4e & 4f failed to reach the performance in Fig. 4b & 4c,

suggesting the virus infection conditions on heterogeneous individuals have a more complex

impact than the cellular response to stimulation of interferon-β only[30,31].

Inference unseen perturbation on genes using pretrained TP embeddings

Next, we investigated if our distinct TP embedding and Decoder architecture could effectively

capture the impact of perturbations on gene expression levels, thereby offering *in silico*

assessment of the perturbation response. Consequently, we have integrated our TP embedding

with a perturbation relationship graph, which is an adaptation from GEARS, a deep learning

framework based on Graph Neural Network (GNN) architecture[16]. The Norman et al. dataset[32]

contained 105 single CRISPRa perturbations on K562 chronic myelogenous leukemia cell line.

As shown in Fig. 5a, we input the gene matrix of unperturbed K562 cells in the same study to

obtain the unperturbed TP embeddings. We combined TP embedding with knockdown gene

embedding generated from the perturbation graph, where edges connecting genes are

extracted from GO (Fig. 1). After two layers of GNN training, gene embedding representing its

spatial pattern inherited in the perturbation graph further concatenates TP embedding as the

input of perturbation prediction layers. As shown in Fig. 5a, we predicted the mean difference

after perturbation of MAPK1 and compared the top 20 most differentially expressed genes with

the real perturbation distribution. We benchmarked our iGTP framework with GEARS and

identified a similar overlap (10 out of 20 genes) and trend of the top 20 most differentially

expressed genes (Fig. 5b). Lastly, we observed a much larger perturbation in genes within

perturbed TP than in genes out of perturbed TP with t-test p-value 1.97^{-72} and Mann-Whitney U

test p-value 1.24^{-76} (Fig. 5c).

SHAP importance of TPs in predicting PBMC states and cell-type-specific PPIs

We examined the performance of iGTP and the relationship between the layers of the Decoder.

Initially, we evaluated if the iGTP embedding can differentiate the stimulation state in the PBMC

dataset from Kang et al (Fig. 6a). We used the embeddings of iGTP TP, iGTP PPI, and VEGA,

as well as conventional ssGSEA and raw gene expression features to measure the AUC for

PBMC labels. iGTP TP embedding had the highest AUC (0.9904) in our comparison. We then

utilized SHapley Additive exPlanations (SHAP) values to validate the contributions of TP

embeddings in differentiating stimulation state. Fig. 6b presents the SHAP values for the top 10

TPs and the aggregate value for the remaining TPs. Consistent with expectations, many of the

highest-ranking TPs closely corresponded with those exhibiting the highest Bayes factors (Fig.

2f). Furthermore, TPs associated with shared functions, such as BP "GO:0009615: response to

virus" and BP "GO:0043331: response to dsRNA", exhibited similar SHAP value distributions. In

Fig. 6c & 6d, we further dissected the PPI embeddings in these two TPs among different cell

types. Specifically, we selected the top 10 PPIs with the largest MAD differences between

stimulation and control statuses from those shared by CD8+ T cells and CD14+ monocytes.

Interestingly, we found that the BST2-TRIM25 PPI within the "GO:0035455: response to

interferon-α" TP had the highest difference. The interferon-inducible protein, BST-2 (or, tetherin),

plays an important role in adaptive immune response[33] and TRIM25, also known as tripartite

motif-containing 25, can regulate the immune response of viral infection[34]. Absolute MAD has a

higher value of this PPI in CD8+ T cells than CD14+ monocytes. On the other hand, the

RSAD2-APOC1 PPI within the "response to type one interferon" TP was the second most

differentiable between CD8+ T cells and CD14 + monocytes (Fig. 6d). Recent research

indicates that RSAD2 is up-regulated when exposed to type one interferon at the monocyte's

single-cell level[18]. These findings suggest that our iGTP model could capture the top TPs and

fine-grained cell-type-specific responses PPIs response to interferon-β stimulation. To delineate

the relationship between layers of the iGTP Decoder, Sankey Diagram (Fig. 6e) shows the

relative weight flow, in between TP "GO:0035455: response to interferon-α" to IFITM3 and

GATA3 genes within the TP. Intuitively, the thickness of the link between TP and PPIs reflected

the normalized importance of PPIs to the TP. The PPI IFITM3-STAT3 was found to have the

largest contribution to this TP, with the IFITM3/STAT3 axis playing a key role in regulating

immune responses through interferon signaling. IFITM3 modulates the phosphorylation and

activation of STAT3, thereby influencing gene expression related to inflammation and

immunity[35]. Similarly, the thickness between the PPIs and genes suggested the relative

importance of gene to this PPI. The final layer indicated the relative importance of genes within

this TP, with gene IFITM3 making the largest contribution.

Conditional generation of new cells using diffusion model and TP embeddings

The denoising diffusion probabilistic models (DDPM)[36], widely used in image generation, starts

with random noise and iteratively refines it into a coherent image by reversing a diffusion

process through a series of conditionally learned denoising steps. To further explore the

generative application of our iGTP model, we revised and integrate one recent single-cell latent

diffusion model scDiffusion[37] into our iGTP framework and use iGTP TP embedding as the

latent input X_0. To develop a robust pretrained model, we utilized the previous Ahern et al.[38]

COVID-19 PBMC dataset, which contains over 600,000 cells, to pretrain the iGTP model with

non-redundant GO TPs. Then, we input Kang et al.[17] dataset through this pretrained encoder to

obtain the TP embeddings (X_0) of ~13,000 PBMC cells with interferon-beta stimulation and

control. As shown in Fig. 7a&7b, most diffusion-generated control CD4+ T cells were located in

the overlapping region of control state cells and CD4+ T cells, suggesting the potential of

conditionally generating data on specific cell types and states. We designed a classifier to

distinguish control CD4+ T cells (including both diffusion-generated and original) from other cell

types, using TP embeddings as features. This approach achieved an accuracy of 0.98 on the

test set (Fig. 7c) and an area under the precision-recall curve (AUPRC) of 0.98 ± 0.00 in five-

fold cross-validation (Fig. 7d).

Discussion

The deep-learning framework VAE has gained more attention for the scalability and flexibility to

handle the growing single-cell RNA-seq data. In this work, we developed an interpretable

generative transcriptional program (iGTP) VAE-based framework that quantifies the biological

networks composed of TPs and PPIs in driving the cellular state alteration. We used examples

in the PMBC dataset with interferon-β treatment dataset, COVID-19 PMBC dataset, and AD

dataset, suggesting our iGTP model could accurately capture the known biological process and

discover new TP that was not well-characterized before. Moreover, we tested our model

robustness within intra-sample and inter-sample tasks, showing the VAE has the potential

transferability. Lastly, we integrated one GNN-based module to provide *in silico* perturbation

estimation for the known perturbation effect. The integration of this feature substantially

augments our model's proficiency in accurately interpolating responses to perturbations at each

TP. This advancement demonstrates substantial potential in facilitating extensive *in silico*

predictions, which could be pivotal for strategizing and optimizing real-world experimental

approaches. Finally, we demonstrated an application of our iGTP TP embeddings by integrating

iGTP with a latent diffusion model, successfully generating accurate new cell embeddings for

specified cell types and states.

In this study, we used gene expression to approximate protein, PPI, and TP level activities. We

innovatively designed a biologically meaningful Decoder that integrates prior knowledge,

enabling the VAE model to capture the hierarchical structures of biological processes among

TPs, PPIs, and genes. In the Kang et al PBMC dataset, we successfully replicated the BP of

"GO:0009615: response to virus", BP of "GO:0034340: response to type I interferon" as the

biological processes corresponding to interferon-β treatment (Fig. 2f & 6b). In addition, the

model not only differentiates the cell state (w/wo interferon-β stimulation) but also captures the

activity difference between monocytes and T cells (Fig. 2c). Moreover, we identified several key

PPIs with cell-type-specific importance, such as BST2-TRIM25 and RSAD2-APOC1, which

exhibited the highest MAD differences between stimulated and control conditions for CD4+ T

cells and CD14+ monocytes (Fig. 6c & 6d). This indicates that these PPIs within the TPs

contribute significantly to the alteration of cellular states. In the AD dataset, we confirmed iGTP

could capture the cell-type-specific TP activities and pinpointed a few novel TPs might be

responsible for the molecular mechanism for AD pathogenesis. Even though we obtained a

Spearman correlation coefficient ρ of 0.74, some top differential TP was predicted to flip in cells

from AD and CN individuals between the results of iGTP and ssGSEA. For example, iGTP

identified the higher activity of CC "GO:0005874: microtubule" in synapse, oligodendrocytes,

and astrocytes for cells from AD individuals, while the activity signals were predominant in

synapse only in the result of ssGSEA (Fig. S1). In oligodendrocytes, microtubules play a crucial

role in extending processes to contact axons and elongating the myelin sheath. In astrocytes,

neighboring cells and the extracellular environment can influence changes in cell shape and

branching[39]. The BP "GO:0050808: synapse organization" is essential for neuron

communication. In Alzheimer's, amyloid plaques and tau tangles disrupt this, leading to

cognitive decline. OPCs play a role in signal transmission by interacting with neurons and

forming neuronal-OPC synapses, which exhibit synaptic plasticity like that observed in neuron-

neuron synapses[40]. However, while iGTP identifies signals in OPCs, ssGSEA fails to detect

much activity (see Fig. S1). This discrepancy might be raised by the methodology difference. As

ssGSEA[19,20] calculates each cellular TP score by comparing the distribution of gene expression

ranks inside and outside the TP, it may fail to capture the complex, non-linear relationships that

iGTP can. Therefore, further analysis incorporating ground truth knowledge about TP activities

is warranted.

We evaluated the transferability of iGTP on PBMC datasets subjected to different immune

stimulations: one from a virus (COVID-19 and influenza) and the other from interferon-β. Viral

invasion can stimulate interferon-β production, boosting the immune response in a similar way

to interferon-β stimulation. The model was trained on COVID-19 patients in various conditions,

including critical, severe, mild, receiving COVID-19 treatment, and normal. It can predict the

biological activity in two TPs: the BP of "GO:0009615: response to virus" and the BP of

"GO:0034340: response to type I interferon" in previously unseen patients with severe influenza

infection. The model also shows the difference in biological activity linked to the conditions (Fig.

4a). When the entire COVID-19 dataset was used to train iGTP and predict the Kang et al.

PBMC dataset, the model could differentiate the cell state with and without interferon-β

stimulation (Fig. 4d). Both results suggest that iGTP can transfer knowledge from one dataset to

another while maintaining some common biological rationale. However, overall prediction

performance in the same dataset is better than in different datasets (Fig. 4b, 4c, 4e & 4f). This

suggests that virus infection conditions in diverse individuals have a more complex impact than

the cellular response to stimulation of interferon-β alone.

As shown in Fig.6a, our iGTP model obtained a very close AUROC among conventional

bioinformatics approaches and other deep-learning-based methods in predicting PBMC data

with stimulation or not. Compared to traditional bioinformatics tools, deep learning-based

methods demonstrate significant advantages in learning feature representation within high-

dimensional spaces. These vectorized spaces could be further integrated with other frameworks

(such as GEARS) to handle a more complex task, such as response to perturbation prediction.

Although the reconstruction rate of iGTP is lower compared to the VEGA model, particularly

against designs using fully connected layers or a single sparse masker layer (as employed by

VEGA), this trade-off is intentional to enhance the interpretability of biological mechanisms in

our approach. We envision GEAR in iGTP, which could also be useful to prioritize drugs based

on pathway expression in cancer, as studying the response of specific cell populations may

inform drug sensitivity and resistance. Integrating drug response prediction models with such

explanatory models could benefit designing novel therapeutic strategies.

Methods and Materials

iGTP framework

iGTP is a deep learning framework based on the Variational AutoEncoder (VAE) structure and

utilizes an Encoder with fully connected layers to process input gene expression data (X). We

transformed gene expression data into a biologically interpretable latent space to approximate

the activities of protein-protein interaction (PPI), and biological pathway or transcriptional

program (TP). From the TP space, a vector Z undergoes decoding through a specifically

designed network to reconstruct the gene features $\hat{X}$.

iGTP differs from the traditional VAE model in that the Decoder network is designed to

incorporate prior knowledge of the relationship between TP and genes. Specifically, we employ

a sequence of masked fully connected layers as the Decoder network to reconstruct the gene

features $\hat{X}$.

The vector of gene features $G = \{g_1, g_2, , ..., g_{N_{Gene}}\}$ were encoded to derive latent variables

$Z = \{z_1, ..., z_{N_{TP}}\}$. Each latent variable of Z corresponds to a single transcriptome program. To

ensure that each dimension of Z is a TP (scalar), we enforce the Decoder network to embed

only (masked) linear operations, i.e. masked fully connected layers.

To reconstruct the gene features, Z is transformed by a Decoder network with two masked fully

connected layers. First, Z is transferred into gene by going through the first masked fully

connected layer, whose weights are a binary matrix $W^{TP-Gene}$. The weights of $W^{TP-Gene}$ are

predefined using prior knowledge of TP-gene relations: for each entry $W^{TP-gene}(i, j)$, its value is

one if the i-th gene is a member of j-th TP, and zero otherwise. Second, the output matrix,

denoted as O, will go through another masked fully connected layer, $W^{Gene-Gene}$, to reconstruct

the gene features G. Each entry of $W^{Gene-Gene}(i, j)$ denotes whether the i-th gene and j-th

gene has PPI relations: one if the relation exist and zero otherwise. The weight matrix

$W^{Gene-Gene}$ is, therefore, a symmetric matrix. The Decoder network reconstructs gene feature

vector $\widehat{G} = \{\hat{g}_1, \hat{g}_2, , ..., \hat{g}_{N_{Gene}}\}$ as following:

$$O = Relu(W^{TP-Gene}Z)$$

$$\widehat{G} = W^{Gene-Gene}O$$

392   Like the traditional VAE, the objective of training iGTP is to reconstruct gene expression X from

393   the TP space Z by maximizing the reconstruction likelihood,

$$p(X|\theta) = \int p(X|Z, \theta)p(Z|\theta)dZ$$

where θ is the parameters of iGTP, X is the input gene expression data, and Z is the embedding

in the TP space.

Model training

The raw scRNA-seq read count data X is fed into iGTP, which is a matrix of C rows (cells) and

N_{gene} columns (genes). The training of iGTP follows the training of traditional VAEs by forming a

data compression and reconstruction task. The Encoder network to maps the X to embedding

vector Z, and the Decoder neural networks reconstruct $\tilde{X}$ to approximate the input. iGTP is

trained by optimizing the reconstruction loss as the KL divergence loss as below,

$$loss = \sum_{i=1}^{C} \sum_{j=1}^{N_{gene}} \text{MSE}(X, \tilde{X}) + \beta \sum KL\big(q(Z|X)\big|p(Z)\big)$$

Measuring Differential TP activity via Plaid-based Bayes Factor

Different treatment conditions often elicit varying activities and TPs. To identify these differential

activity genes and/or TPs, we utilize the Bayesian differential gene expression method inspired

by Lopez et al.[7].

The method assumes cell independence and utilizes a Monte Carlo sampling approach by

randomly sampling cells from the two treatment groups, denoted as g_a, g_b respectively. For

each TP q, the method constructs two hypotheses for any pair of cells $x_a \in g_a$ and $x_b \in g_b$ to

test if the TP q has different activations in the two cells from different conditions.

$$\mathcal{H}_0^q := E_g[Z_a^q] > E_g[Z_b^q]; \; \mathcal{H}_1^q := E_g[Z_a^q] \leq E_g[Z_b^q]$$

410   The log-Bayes factor K is defined as

$$K := \log_e \frac{p(\mathcal{H}_0^q | x_a, x_b)}{p(\mathcal{H}_1^q | x_a, x_b)}$$

, where the sign and magnitude of K denote likelihood of each hypothesis and the significance

level, respectively.

To achieve robust results, the method aggregates the Bayes factors of a group of cells and

computes their average. In iGTP, we use a threshold of 5 (K larger than 5) to denote significant

differentiation of TP activity. The process of randomly sampling cells from different groups can

introduce bias, especially when dealing with the presence of various cell types, some of which

may have very small numbers. To address this challenge, we draw inspiration from the

Geometric Sketching method proposed by Hie et al.[41] Geometric sketching is a sampling

method to summarize the single-cell transcriptomic landscape. The method highlights rare cell

states compared to traditional uniform sampling methods. This approach allows us to

subsample large single-cell RNA sequencing (scRNA-seq) datasets while preserving the

representation of rare cell states. Geometric Sketching samples cells from cell types according

to each cell type's area of geometric plaid covering in the latent space. The sampling approach

is less prone to bias by more evenly sampling according to cell types of landscape, rather than

uniform sampling which may underrepresent the rare cell types.

iGTP perturbation prediction function iGEARS and perturbation embedding integration

We further integrated our TP embedding with a perturbation relationship graph adapted from

GEARS[16] and named iGEARS, to predict the *in silico* impact of gene perturbation on TPs and

genes. iGEARS leverages prior knowledge of gene-gene relationships using a GO-derived

knowledge graph from GEARS. Intuitively, genes that are involved in similar pathways should

impact the expression of similar genes after perturbation. iGEARS functionalizes this graph-

based inductive bias using a GNN architecture. Specifically, for cells from perturbation dataset

such as Norman et al.[32], $D = \left\{ \left(g_1, \dots, g_{N_{Gene}} \right)_c, P(c) \mid c = 1, \dots, C \right\}$, where $\left(g_1, \dots, g_{N_{Gene}} \right)_c$ is the

expression of cell c and $P(c)$ stands for the perturbation of cell c ,from perturbation types

$\boldsymbol{P} = \{P_1, \dots, P_M\}$. We initially applied the unperturbed cells into iGTP Encoder $f_{iGTP} : \mathbb{R}^{N_{Gene}} \to$

$\mathbb{R}^{N_{TP}}$ that maps each cell into TPs of unperturbed cell in N_{TP}-dimensional embedding. Then, a

GNN-based Encoder $f_{(pert_{graph})} : \boldsymbol{Z} \to \mathbb{R}^d$ is adopted to map each perturbation P to a d -

dimensional perturbation embedding (emb$_p$). Lastly, we concatenated the emb$_p$ with each TP

embedding of unperturbed cells, resulting in an integrated (d+1)-dimensional embedding. This

integrated embedding was used as input for the cross-TP MLP Decoder $f_{(pert_{dec})} : \mathbb{R}^{N_{TP}} \to$

$\mathbb{R}^{N_{Gene}}$ to predict gene expression alteration to unperturbed cells in response to unseen

perturbation type P (Fig. 1 step 4). We evaluated the prediction performance using top the 20

mean gene expression alteration in perturbated cell $P(c)$ (Fig. 1 step 5).

Latent diffusion model for conditionally generation of cell types and phenotypes using iGTP and TP embeddings

To further evaluate the generative capabilities of our iGTP model, we integrated its TP

embeddings into the latent diffusion model, scDiffusion[37], which is a generative framework

consisting of autoencoder, diffusion process, denoising network, and a conditional classifier.

This integration enables the generation of new cells corresponding to specific cell types and

phenotypes. Specifically, the pre-trained autoencoder was adapted from iGTP Ahern et al.[38]

PBMC dataset and an single-cell dataset Kang et al. [17] was processed through this autoencoder

to generate TP embeddings, which were subsequently L2-normalized as latent representations.

The diffusion process further produces a series of progressively noisy embeddings, which serve

as the training data for the denoising network. In scDiffusion, the denoising network is trained to

model the reverse diffusion process, while the classifier is designed to guide the generation

toward specific cell types and states. Notably, the classifier is trained on intermediate states

derived from the diffusion process, with a particular focus on states at step 0 and step T/2,

ensuring robust and accurate conditional generation.

Our experiment consisted of two primary components: training the denoising network and the

classifier model. Both models were adapted from the foundational architectures of scDiffusion,

with modifications to accommodate the updated embedding and layer dimensions. The

denoising network, implemented as a skip-connected multilayer perceptron with an input

dimension of 1303 (GO TPs) and hidden layer dimensions of [4096, 2048, 2048], was trained to

model the reversed diffusion process. This network was trained for 800,000 steps with a

minibatch size of 12 and a learning rate of 1^{-4}. The classifier, a feed-forward neural network with

an input dimension of 1303 and appropriately sized hidden layers, was trained on noised

samples at steps 0 and T/2. Training was conducted with a minibatch size of 256, a learning

rate of 5^{-5}, and a total of 10,000 steps. After training the scDiffusion model, we generated

hundreds of synthetic samples corresponding to specific selected cell types. For additional

details, please refer to the scDiffusion[37].

Datasets and preprocessing

Kang et al. PBMC dataset

The Kang et al. dataset[17], (accessed by 10/23/2023 from GSE96583), comprises two states of

Peripheral Blood Mononuclear Cells (PBMC): a control state and another stimulated with

interferon-β. Data was processed by Seurat[42] and filtered with the following criteria "percent.mt

<= 50 & nCount_RNA > 677 & nFeature_RNA > 500 & multiplets == 'singlet'". In total, we

obtained a gene matrix of 13,672 cells by 35,635 genes. Cell type annotation and states were

directly adapted from the original MetaData file in the GSE96583 repository. Megakaryocytes

were excluded for their small population. We then converted the Seurat object to Scanpy

AnnData object[43] for further analysis.

Ahern et al. COVID-19 PBMC dataset

The Ahern et al.[38] dataset was downloaded from the Chan Zuckerberg (CZ) CellxGene portal[44]

(accessed by 11/10/2023). The dataset provided an extensive scRNA-seq analysis to construct

a blood atlas of patients across a spectrum of COVID-19 severity levels, comparing it with cases

of influenza, sepsis, and healthy volunteers. We conducted the standard quality control process

using Scanpy[43]. Specifically, we filtered cells with 200 expressed genes and genes expressed in

less than three cells. Then, we kept the cell with the following criteria [n_genes_by_counts <

10000; pct_counts_mt < 20; total_counts < 50000]. Cell type annotation was adapted from the

original dataset and cell types were renamed with the Kang et al dataset, leading to 599,044

cells in total, including the following sample sources: COVID-19 in-patient critical (COVID_CRIT),

COVID-19 in-patient severe (COVID_SEV); COVID-19 in-patient mild (COVID_MILD);

COVID_LDN (COVID-19 in-patient with low-dose Naltrexone treatment), Influenza and Normal

(healthy volunteer). Detailed description for each phenotype can be found in Ahern et al.[38].

Alzheimer's Disease single-cell RNA-seq and polygenic risk score process

We adapted snRNA-seq data from Synapse portal (syn2580853, accessed by 4/15/2023),

which contains 454 individuals from Religious Orders Study/Memory and Aging Project

(ROS/MAP).   We obtained the matched whole-genome-sequence (WGS) data from the

Synapse portal (syn11724057, accessed by 11/10/2022). LDPred2[45] was used to estimate the

individual's risk by adapting the variant effect size from Wightman et al. GWAS summary

statistics[46]. In total, we obtained 407 individuals with matched snRNA-seq and WGS data. To

obtain individuals with similar high-risk PRS backgrounds, we selected the top 20 quantiles of

PRS. This subset included 53 high-risk Alzheimer's disease (AD) cases and 15 high-risk

cognitively normal (CN) individuals for comparison. Data was processed by Seurat[42] and filtered

with the following criteria "percent.mt <= 50 & nFeature_RNA > 200". We adapted the cell type

annotation from the original study.

Norman et al. CRISPR Perturbation dataset

507 Norman et al.[32] raw data was downloaded from scPerturb[47] processed data (accessed by

508 11/17/2023), which contained 105 single CRISPR activation (CRISPRa) perturbations with at

509 least 20 cells. We used the same QC process as the *Ahern et al. dataset.* After conducting

510 pretraining on the Norman et al. dataset.

Transcriptional program curation

We incorporated three TP resources. The GO curation (Biological Process: BP, Molecular

Function: MF, and Cellular Component: CC) comprises 1303 non-redundant GO terms from

WebGestalt[48,49] (accessed on 5/7/2021). Lastly, we curated 2082 canonical pathways from three

major resources: Kyoto Encyclopedia of Genes and Genomes (KEGG), Reactome, and

BioCarta pathways from the Molecular Signatures Database[50] (MSigDB 7.4 C2 category,

accessed on 5/10/2021). The brain and AD-related curation is adapted from our previous work,

containing 167 TPs[23].

Human protein-protein interaction (PPI) curation

Our human PPI curation included one public database and one in-house curation.

Pathway Commons curation

Pathway Commons is an integrated resource of publicly available biological pathways
information, biochemical reactions, biomolecular complex assembly, transport, catalysis events,
and physical interactions involving proteins, DNA, RNA, and small molecules. We downloaded
the Pathway Commons V11 "All.hgnc.txt", which contains the PPI curations from 18 databases[51].
We extracted the file annotation with human source only and used the annotation of
"INTERACTION_TYPE = interaction only", leading to 485,383 human protein-protein
interactions and 17,733 unique proteins.

PPI compiled curation

PPI compiled was curated from 6 different resources: BioGRID[48,49,52], ESCAPE[48], HINT[53],
IRefIndex[54], ReactomeFI[55,56], and STRING[57]. In short, we extracted all the human PPI with
physical interaction evidence from each of these resources. Specifically, BioGRID is version
"BIOGRID-ALL-3.5.171" was downloaded 4/15/2019. ESCAPE is a database for integrating high
content published data collected from human and mouse embryonic stem cells
(http://www.maayanlab.net/ESCAPE/). All human interactions were accessed by 4/8/2019.
High-quality INTeractomes (HINT) database was curated for PPI with experiment validation and
was accessed by 4/6/2019. iRefIndex 15.0 for human "9606.mitab.22012018" was used
(accessed by 4/10/2019). ReactomeFI (version 2017) for protein complex was downloaded
(accessed by 4/11/2019). The STRING version was downloaded. All the PPIs from six
resources were merged, including 419,971 protein-protein interactions and 21,216 unique
proteins[55].

Benchmark with VEGA and ssGSEA

We followed the procedure of Kang et al. for the PBMC dataset from VEGA[15]. Specifically, we

selected the top 7,000 highly variable genes and used GO terms containing at least 10% of

these genes to construct the hidden layer. This hidden layer was then used to assess clustering

performance and calculate the AUC for 'stimulated' state label prediction.

We used default setting of "runEscape" function in R package "escape"[58] to conduct the single

sample Gene Set Enrichment Analysis (ssGSEA) analyses[19,20] for each cell in the single-cell

dataset. The resulting cell-by-TP matrix was subsequently used for differential analysis and to

calculate the AUC for 'stimulated' state label prediction.

DATA AVAILABILITY

All the data generated or analyzed in this study is available from the authors upon reasonable

request. The overall framework can be downloaded from https://github.com/davidroad/iGTP.

ACKNOWLEDGEMENTS

We would like to thank Brisa S. Fernandes from UTHealth at Houston for constructive

discussion. The authors acknowledge the High-Performance Computing for Research facility at

the Center for Secure Artificial Intelligence for Healthcare, McWilliams School of Biomedical

Informatics, University of Texas Health Science Center at Houston, and MD Anderson Cancer

Center for providing the computational resources that supported the research presented in this

paper.

FUNDING

This research was partially supported by National Institutes of Health grants awarded to Y.D.,

L.Y., and Z.Z. (R21AG087299), and to Z.Z (U01AG079847, R03AG077191, R01LM012806,

R01DE030122, and R01DE029818). K.H was fully supported by Mulva Family Fund (CFS:

600655-80-120663-19). P.P was partially supported by National Cancer Institutes grant

(NCT04947254, NCT03748641**).** We thanked the resource support from the Cancer Prevention

and Research Institute of Texas (CPRIT RP240610). I.K. is a CPRIT summer intern in the

Biomedical Informatics, Genomics, and Translational Cancer Research Training Program (BIG-

TCR) funded by the Cancer Prevention & Research Institute of Texas (CPRIT RP210045).

References

1. Regev, A. *et al.* The Human Cell Atlas. *Elife* **6**, (2017).

2. Tabula Sapiens Consortium* *et al.* The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. *Science* **376**, eabl4896 (2022).

3. Jovic, D. *et al.* Single-cell RNA sequencing technologies and applications: A brief overview. *Clin. Transl. Med.* **12**, e694 (2022).

4. Kingma, D. P. & Welling, M. Auto-Encoding Variational Bayes. (2013).

5. Wang, D. & Gu, J. VASC: Dimension Reduction and Visualization of Single-cell RNA-seq Data by Deep Variational Autoencoder. *Genomics Proteomics Bioinformatics* **16**, 320–331 (2018).

6. Way, G. P. & Greene, C. S. Extracting a biologically relevant latent space from cancer transcriptomes with variational autoencoders. *Pac. Symp. Biocomput.* **23**, 80–91 (2018).

7. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for single-cell transcriptomics. *Nat. Methods* **15**, 1053–1058 (2018).

8. Lotfollahi, M., Wolf, F. A. & Theis, F. J. scGen predicts single-cell perturbation responses. *Nat. Methods* **16**, 715–721 (2019).

9. Grønbech, C. H. *et al.* scVAE: variational auto-encoders for single-cell gene expression data. *Bioinformatics* **36**, 4415–4422 (2020).

10. Jia, P. *et al.* Deep generative neural network for accurate drug response imputation. *Nat. Commun.* **12**, 1740 (2021).

11. Tran, D. *et al.* Fast and precise single-cell data analysis using a hierarchical autoencoder. *Nat. Commun.* **12**, 1029 (2021).

12. Eraslan, G., Simon, L. M., Mircea, M., Mueller, N. S. & Theis, F. J. Single-cell RNA-seq denoising using a deep count autoencoder. *Nat. Commun.* **10**, 390 (2019).

13. Elmarakeby, H. A. *et al.* Biologically informed deep neural network for prostate cancer discovery. *Nature* **598**, 348–352 (2021).

14. Zhao, Y., Cai, H., Zhang, Z., Tang, J. & Li, Y. Learning interpretable cellular and gene

signature embeddings from single-cell transcriptomic data. *Nat. Commun.* **12**, 5261 (2021).

15. Seninge, L., Anastopoulos, I., Ding, H. & Stuart, J. VEGA is an interpretable generative

model for inferring biological network activity in single-cell transcriptomics. *Nat. Commun.*

12, 5684 (2021).

16. Roohani, Y., Huang, K. & Leskovec, J. Predicting transcriptional outcomes of novel

multigene perturbations with GEARS. *Nat. Biotechnol.* (2023) doi:10.1038/s41587-023-

01905-6.

17. Kang, H. M. *et al.* Multiplexed droplet single-cell RNA-sequencing using natural genetic

variation. *Nat. Biotechnol.* **36**, 89–94 (2018).

18. Cui, A. *et al.* Dictionary of immune responses to cytokines at single-cell resolution. *Nature*

625, 377–384 (2024).

19. Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven

cancers require TBK1. *Nature* **462**, 108–112 (2009).

20. Abazeed, M. E. *et al.* Integrative radiogenomic profiling of squamous cell lung cancer.

Cancer Res. **73**, 6289–6298 (2013).

21. Marshall, J. S., Warrington, R., Watson, W. & Kim, H. L. An introduction to immunology and

immunopathology. *Allergy Asthma Clin. Immunol.* **14**, 49 (2018).

22. Gayoso, A. *et al.* A Python library for probabilistic analysis of single-cell omics data. *Nat.*

Biotechnol. **40**, 163–166 (2022).

23. Li, X. *et al.* Genetically-regulated pathway-polygenic risk score (GRPa-PRS): A risk

stratification method to identify genetically regulated pathways in polygenic diseases.

medRxiv (2023) doi:10.1101/2023.06.19.23291621.

24. St George-Hyslop, P. H. Piecing together Alzheimer's. *Sci. Am.* **283**, 76–83 (2000).

25. Perl, D. P. Neuropathology of Alzheimer's disease. *Mt. Sinai J. Med.* **77**, 32–42 (2010).

26. Hampel, H. *et al.* The Amyloid-β Pathway in Alzheimer's Disease. *Mol. Psychiatry* **26**,

5481–5503 (2021).

27.   Arcuri, C., Mecca, C., Bianchi, R., Giambanco, I. & Donato, R. The Pathophysiological Role

of Microglia in Dynamic Surveillance, Phagocytosis and Structural Remodeling of the

Developing CNS. *Front. Mol. Neurosci.* **10**, 191 (2017).

28.   Chen, Z. *et al.* Genetic association of neurotrophic tyrosine kinase receptor type 2 (NTRK2)

With Alzheimer's disease. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **147**, 363–369

(2008).

29.   Chung, J. *et al.* Genome-wide association study of Alzheimer's disease endophenotypes at

prediagnosis stages. *Alzheimers. Dement.* **14**, 623–633 (2018).

30.   Zhu, L. *et al.* Single-Cell Sequencing of Peripheral Mononuclear Cells Reveals Distinct

Immune Response Landscapes of COVID-19 and Influenza Patients. *Immunity* **53**, 685-

696.e3 (2020).

31.   Zhang, J.-Y. *et al.* Single-cell landscape of immunological responses in patients with

COVID-19. *Nat. Immunol.* **21**, 1107–1118 (2020).

32.   Norman, T. M. *et al.* Exploring genetic interaction manifolds constructed from rich single-

cell phenotypes. *Science* **365**, 786–793 (2019).

33.   Urata, S. *et al.* BST-2 controls T cell proliferation and exhaustion by shaping the early

distribution of a persistent viral infection. *PLoS Pathog.* **14**, e1007172 (2018).

34.   Gack, M. U. *et al.* TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated

antiviral activity. *Nature* **446**, 916–920 (2007).

35.   Prelli Bozzo, C. *et al.* IFITM proteins promote SARS-CoV-2 infection and are targets for

virus inhibition in vitro. *Nat. Commun.* **12**, 1–13 (2021).

36.   Jonathan, H., Ajay, J. & Pieter, A. Denoising Diffusion Probabilistic Models. *arXiv [cs.LG]*

(2020).

37.   Luo, E., Hao, M., Wei, L. & Zhang, X. scDiffusion: conditional generation of high-quality

single-cell data using diffusion model. *Bioinformatics* **40**, btae518 (2024).

38. COvid-19 Multi-omics Blood ATlas (COMBAT) Consortium. Electronic address:

julian.knight@well.ox.ac.uk & COvid-19 Multi-omics Blood ATlas (COMBAT) Consortium. A

blood atlas of COVID-19 defines hallmarks of disease severity and specificity. *Cell* **185**,

916-938.e58 (2022).

39. Weigel, M., Wang, L. & Fu, M.-M. Microtubule organization and dynamics in

oligodendrocytes, astrocytes, and microglia. *Dev. Neurobiol.* **81**, 310–320 (2021).

40. Zou, P., Wu, C., Liu, T. C.-Y., Duan, R. & Yang, L. Oligodendrocyte progenitor cells in

Alzheimer's disease: from physiology to pathology. *Transl. Neurodegener.* **12**, 52 (2023).

41. Hie, B., Cho, H., DeMeo, B., Bryson, B. & Berger, B. Geometric sketching compactly

summarizes the single-cell transcriptomic landscape. *Cell Syst.* **8**, 483-493.e7 (2019).

42. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

43. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* **19**, 15 (2018).

44. CZI Single-Cell Biology Program *et al.* CZ CELLxGENE Discover: A single-cell data

platform for scalable exploration, analysis and modeling of aggregated data. *bioRxiv*

2023.10.30.563174 (2023) doi:10.1101/2023.10.30.563174.

45. Privé, F., Arbel, J. & Vilhjálmsson, B. J. LDpred2: better, faster, stronger. *Bioinformatics* **36**,

5424–5431 (2021).

46. Wightman, D. P. *et al.* A genome-wide association study with 1,126,563 individuals

identifies new risk loci for Alzheimer's disease. *Nat. Genet.* **53**, 1276–1282 (2021).

47. Peidli, S. *et al.* scPerturb: harmonized single-cell perturbation data. *Nat. Methods* 1–10

(2024).

48. Xu, H. *et al.* ESCAPE: database for integrating high-content published data collected from

human and mouse embryonic stem cells. *Database* **2013**, bat045 (2013).

49. Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set analysis

toolkit with revamped UIs and APIs. *Nucleic Acids Res.* **47**, W199–W205 (2019).

50. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set

collection. *Cell Syst.* **1**, 417–425 (2015).

51. Rodchenkov, I. *et al.* Pathway Commons 2019 Update: integration, analysis and

exploration of pathway data. *Nucleic Acids Res.* **48**, D489–D497 (2020).

52. Oughtred, R. *et al.* The BioGRID interaction database: 2019 update. *Nucleic Acids Res.* **47**,

D529–D541 (2019).

53. Das, J. & Yu, H. HINT: High-quality protein interactomes and their applications in

understanding human disease. *BMC Syst. Biol.* **6**, 92 (2012).

54. Razick, S., Magklaras, G. & Donaldson, I. M. iRefIndex: a consolidated protein interaction

database with provenance. *BMC Bioinformatics* **9**, 405 (2008).

55. Patil, A. & Nakamura, H. HINT: a database of annotated protein-protein interactions and

their homologs. *Biophysics* **1**, 21–24 (2005).

56. Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res.* **50**,

D687–D692 (2022).

57. Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased

coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*

Acids Res. **47**, D607–D613 (2019).

58. Borcherding, N. *et al.* Mapping the immune environment in clear cell renal carcinoma by

single-cell genomics. *Commun. Biol.* **4**, 122 (2021).

697
698

     Figures and Figure Legends

Fig.1. Overview of iGTP framework: The single-cell data is processed by the iGTP non-linear

Encoder, generating the transcriptional program (TP) embedding (z) using reparameterization

for input TPs. The Decoder consists of two layers: 1) The TP-Gene layer, 2) The Gene-Gene

layer. Both layers apply a sparse matrix to the gradient to eliminate non-existent connections,

leaving only the true connections. For instance, in the TP-Gene layer, the value is one if a gene

is present in a given TP; otherwise, it's zero. During backpropagation, the weight of the true

connection is retained, while all others are forced to zero. The two layers are used to

reconstruct the gene expression within each cell, allowing each TP to be learned by TP-related

genes and the gene-gene interaction within those genes. The TP embedding Z and gene layer

is then extracted for visualization and differential TP analysis (step 1). Additionally, TP

embedding Z (step 2) can be concatenated with gene perturbation embeddings (emb_p) derived

from one PPI-guided gene relationship graph (step 3). This graph is generated using genetic

perturbation data and graph propagation of the genetic perturbation impact (P on gene G_p),

constructed from gene-gene relationship derived from gene ontology (GO). The concatenated

embeddings are then processed through a two-layer Multilayer Perceptron (MLP), designed to

associate TPs with genes. The results in the model outputting altered gene expressions (step 4)

represented the predicted overall impact of perturbations on other genes. The top altered gene

expressions were ordered and highlighted using a boxplot (step 5). Finally, the TP embedding Z

could be used for latent diffusion model to generate cells for given cell types and condition (step

6).

Fig.2. Benchmarking the model performance of iGTP on stimulated and resting PBMC

dataset. First, we evaluate the ability of iGTP to distinguish between different cell types and

stimulation states. Panel a. labeled the cells on the t-SNE plot based on their cell types, while

panel b. labeled them based on the stimulation condition. c. The score indicated the activity of

interferon beta in each cell. Higher activity (red) was observed in interferon beta-stimulated cells

compared to the unstimulated cells (blue). NK cells, monocytes, and dendritic cells with higher

biological activity than other cells. d. We labeled the PPI score (ISH15 and IFNAR1) on the

same map and find that stimulated T cells (CD4+ and CD8+) have lower activity than stimulated

monocytes. Panel e indicates that the conventional ssGESA results are similar to the iGTP

results for the TP (panel c). f, The featured biological pathways ($|log_e(Bayes factor)| > 3$ &

mean absolute difference (MAD) > 6) were highlighted in the volcano plot. Additionally, a

positive ln (Bayes factor) indicates a Transcriptional program (TP) with higher activity in

stimulated cells than unstimulated cells. A higher MAD indicates a more significant difference

within the latent space between stimulated and unstimulated cells. g. The correlation between

the Bayes factor and NES (GSEA) was compared in two different cell types, CD8+ T cells and

CD14+ monocytes, focusing on highly differentiated TPs based on the Bayes factor results. h.

Several state-of-the-art methods, including ssGESA, VEGA, and iGTP, were benchmarked

using performance metrics from scvi-tools.

Fig.3. TP embeddings characterize biological activities of brain-related and AD-related functions. a. Heatmap for visualizing the mean TP embedding values in each cell type and condition for 14 representative TPs for brain-related and AD-related functions. The mean TP embedding values were conducted z-score transformation within each TP. b. Dot plot visualizing the cellular relationship in TP embeddings between two TPs across two cell types, stratified by AD in red and CN groups in blue. c. Dot plot visualizing the cellular relationship in TP embeddings between two PPI pairs across two cell types, stratified by AD (red) and CN (blue) groups. d. Heatmap for visualizing the mean TP embedding values in each cell type and

condition for the top 15 most differential TPs in AD and CN from GO TP sets. The mean TP

embedding values underwent a z-score transformation within each TP. e. Dot plot visualizing

the cellular relationship in TP embedding values between two top TPs across two associated

cell types, stratified by AD (red) and CN (blue) groups. f. Dot plot visualizing the cellular

relationship in TP embeddings between two TP PPI pairs across two associated cell types,

stratified by AD (red) and CN (blue) groups.

Fig4. Prediction performance of intra- and inter- sample for representative TPs and PPIs.

a. Distribution of TP activities in COVID-19 PBMC training samples, alongside predictions of

intra-sample activities in the testing dataset. Specifically, the sample source labeled with an

asterisk (*) represents the unseen intra-sample (testing sample) from the same COVID-19

PBMC dataset. For this testing sample, TP activity predictions were made using a pretrained

model, which was trained on all other cell types. Two TPs were used to demonstrate the related

biological mechanism in CD14+ monocytes and dendritic cells, respectively. b. Bar plot for each

cell type shows the distribution of Pearson correlation coefficient *r* between the Bayes factor (BF)

of selected TPs from GO in pretrained samples and the corresponding predicted TP BF values

for the unseen intra-samples (testing sample) on the same COVID-19 PBMC dataset. c. Bar plot

for each cell type shows the distribution of Pearson correlation coefficient *r* between the PPIs of

the selected 6 interferon-related TPs from GO in pretrained samples and the corresponding

predicted PPI BF values for the unseen intra-sample (testing sample) on the same COVID-19

PBMC dataset. d. Distribution of TP activities in COVID-19 PBMC training samples, alongside

predictions of inter-sample activities in the Kang et al. PBMC dataset. Specifically, the sample

source labeled with an asterisk (*) represents the unseen intra-sample (testing sample) from

Kang et al. PBMC dataset. For this testing sample, TP activity predictions were made using a

pretrained model, which was trained on all cell types from COVID-19 PBMC dataset. Two TPs

were used to demonstrate the related biological mechanism in CD14+ monocytes and dendritic

cells, respectively. e. Bar plot for each cell type shows the distribution of Pearson correlation

coefficient *r* between the BF of selected TPs from GO in pretrained samples and the

corresponding predicted TP BF values for the unseen inter-sample (testing sample) of the Kang

et al. PBMC dataset. f. Bar plot for each cell type shows the distribution of Pearson correlation

coefficient *r* between the PPIs of the selected 6 interferon-related TPs from GO in pretrained

sample and the corresponding predicted PPI BF values for the unseen intra-sample (testing

sample) of the Kang et al. PBMC dataset.

Fig5. The Comparison between GEARS and iGEARS on the perturbation prediction

iGTP perturbation (iGEARs) assesses the change in gene expression and benchmark with

GEARs. a&b, Changes in gene expression in iGEARs and GEARs. Violin plot and Boxplot

showed experimentally measured gene expression difference after perturbing the corresponding

gene, where top 20 post-perturbation genes in response to perturbation. Genes were ordered

by change magnitude. The red dots show the mean change in gene expression predicted by our

model. The green dotted line represents unperturbed control gene expression. a&b showed top

20 variated genes after perturbation on gene MAPK1 in iGTP. c. We investigated the change in

gene expression over MAPK1 perturbation. We tested whether genes in the TP group

containing MAPK1 or genes in the TP group excluding MAPK1 differed significantly, using t-

tests and Mann-Whitney U-tests to compare their mean differences.

Fig6. Model performance and interpretability for TP, PPI, and gene on Kang et al. PBMC

dataset. a. Benchmark prediction as stimulated cell label on Kang et al. PBMC dataset. b.

SHAP value derived from XGBoost, utilizing all TP embedding z values as the features (X) and

the cellular state (either 'stimulated' or 'control') as the label (Y). Beeswarm plot for the top 10

TPs with the biggest absolute SHAP values and the sum of all other features. c & d, Barplot for

top 10 MAD difference of PPIs derived from stim vs ctrl states in CD14+ monocyte and CD8+ T

cell for "GO:0035455: response to interferon-α" and "GO:0034340: response to type I interferon",

respectively. e. Sankey diagram illustrated relative weight flow in Decoder of iGTP framework

for TP "GO:0035455: response to interferon-α" and their related genes within TP, showing how

TP embeddings towards reconstruction of gene expression levels through PPIs and relative

gene importance from certain TP.

Fig7. Latent Diffusion Model to generate given cell type and condition using TP

embeddings on Kang et al. PBMC dataset. a. t-SNE embedding for PMBC datasets and

diffusion generated control CD4+ T cells by conditions. b. t-SNE embedding for PMBC datasets

and diffusion generated control CD4+ T cells by cell types. c. Confusion matrix illustrating the

prediction performance on the test set of PBMC datasets. In this analysis, diffusion-generated

control CD4+ T cells and actual control CD4+ T cells were labeled as 1, while resting cells were

labeled as 0. d. The AUPRC plot evaluating the performance of predicting control CD4+ T cells

using 5-fold cross-validation.