Optimised methods for the targeted surveillance of extended-spectrum beta-lactamase producing *Escherichia coli* in human stool

Sarah Gallichan[1], Sally Forrest[1], Esther Picton-Barlow[1], Claudia McKeown[1], Maria Moore[1], Eva Heinz[1,2,] Nicholas A Feasey[1, 4], Joseph M Lewis[1,3], Fabrice E Graf[1, *]

[1]Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK

[2]Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, UK

[3]Department of Clinical Infection, Microbiology & Immunology, University of Liverpool, Liverpool, UK

[4]The School of Medicine, University of St Andrews, St Andrews, UK

Keywords

Molecular epidemiology; Antimicrobial resistance; Transmission; Surveillance; Microbiological methods; DNA extraction

*Correspondence:

Fabrice Graf

Liverpool School of Tropical Medicine

Pembroke Place

Liverpool, UK

Abstract

Understanding transmission pathways of important opportunistic, drug resistant

pathogens, such as extended-spectrum beta-lactamase (ESBL) producing

Escherichia coli, is essential to implementing targeted prevention strategies to

interrupt transmission and reduce the number of infections. To link transmission of

ESBL-producing *E. coli* (ESBL-EC) between two sources, single nucleotide resolution

of *E. coli* strains as well as *E. coli* diversity within and between samples is required.

However, the microbiological methods to best track these pathogens are unclear. Here

we compared different steps in the microbiological workflow to determine the impact

different pre-enrichment broths, pre-enrichment incubation times, selection in pre-

enrichment, selective plating, and DNA extraction methods had on recovering ESBL-

EC from human stool samples, with the aim to acquire high quality DNA for sequencing

and genomic epidemiology. We demonstrate that using a 4-hour pre-enrichment in

Buffered Peptone Water, plating on cefotaxime supplemented MacConkey agar and

extracting DNA using Lucigen MasterPure DNA Purification kit improves the recovery

of ESBL-EC from human stool and produced high-quality DNA for whole genome

sequencing. We conclude that our optimised workflow can be applied for single

nucleotide variant analysis of an ESBL-EC from stool.

2

Introduction

The spread of antimicrobial resistance (AMR) in bacterial pathogens is a global threat to public health. Of particular concern are infections caused by extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli*, which is on the World Health Organisations (WHO) priority AMR pathogen list[1]. ESBL-producing *E. coli* (ESBL-EC) are resistant to aminopenicillins and 3rd-generation cephalosporin (3GC) antibiotics, which are commonly used to treat infections caused by Gram-negative bacteria. ESBL-EC infections can result in a higher mortality and longer hospital stays and may require access to reserve antibiotics such as carbapenems[2–4].

Infection prevention is preferable to cure and will concurrently reduce antibiotic use. To prevent the acquisition of enteric pathogens in care facilities, it is critical to understand their transmission pathways. This is challenging for near ubiquitous organisms such as *E. coli*, which are frequently part of a healthy gut flora and can also be opportunistic pathogens[5]. Typically, human faecal samples have been shown to contain one to three different *E. coli* genotypes at any one time[6] although as many as nine or more different *E. coli* genotypes per stool sample have been described[7]. The number of multidrug resistant *E. coli* strains detected has also been shown to increase with host age[8]. Therefore, differentiating intraspecies diversity of different *E. coli* strains requires comparisons at single nucleotide variation (SNV) resolution[9]. To track transmission events of ESBL-EC in care settings, microbiological methods need to capture the within-host diversity of ESBL-EC within and between different samples accurately and cost-effectively, even when present at low levels. Consensus on which microbiological methods are optimal and a comprehensive comparison of the impact of different microbiological methods are currently lacking.

Currently, the most common healthcare diagnostic method of ESBL-EC isolation from

human stool is direct plating on cefotaxime-supplemented MacConkey or

chromogenic agar and confirming the production of ESBL using the double disk

diffusion method[10,11]. This is often followed by molecular investigation of ESBL

resistance genes via multiplex PCR[12,13]. While these methods are effective in

identifying the presence of an ESBL-EC to inform treatment options, they lack the

resolution to infer transmission because they do not allow us to clearly distinguish

between closely related *E. coli*[14]. The distinction between two different clades within

the same ST of *E. coli* can be as little as 70 nucleotides, therefore cannot be accurately

distinguished using PCR methods[15].

Surveillance studies focused on describing transmission pathways pre-enrich samples

with low microbial load, i.e., from rectal swabs, to increase detection of present ESBL-

EC[16,17]. Combined with whole-genome sequencing (WGS) this allows for accurate

clustering of isolates at SNV resolution. However, there is little consensus between

studies, or data on controlled comparisons, on the impact of pre-enrichment broth,

pre-enrichment duration, selective agar, and method of DNA extraction has on sample

diversity. Here, we describe an optimized approach for targeted surveillance of ESBL-

EC from human stool samples at SNV resolution to inform transmission modelling with

the potential to capture within-host diversity.

Methods

Pre-Enrich → **Isolate** → **Extract** → **Sequence**

What is the **optimal pre-enrichment broth** and **incubation time** for the recovery of ESBL-producing *E. coli* from stool?

What is the **optimal selective agar** for the isolation of ESBL-producing *E. coli*?

What is the **optimal method for extracting DNA** from ESBL-producing *E. coli* for whole-genome sequencing?

Can **ESBL-producing *E. coli* SNV** be accurately captured using this laboratory method?

112   *Figure 1 Graphical summary of the main research questions and comparisons preformed for each step of the microbiological*

113   *processing of ESBL-EC from stool*

ESBL-EC growth dynamics in different pre-enrichment broths

Pre-enrichment broths (Brain-Heart Infusion, Tryptic Soy, Buffered Peptone Water and Davis Minimal) commonly used in the recovery of ESBL-EC from clinical sources were selected for comparison in this study. These pre-enrichment broths were prepared according to the manufacturer's recommendations **(Supplementary Table 1)**. We used three reference strains encoding ESBLs: NCTC 13341 ($bla_{CTX-M-15}$); NCTC 13476 (bla_{IMP}); NCTC 13846 ($bla_{CTX-M-27}$), with the latter also encoding plasmid-derived colistin resistance **(Supplementary Table 2)**. We further included a clinical isolate, CAB17W, encoding $bla_{CTX-M-15}$ **(Supplementary Table 2)**. The growth of single ESBL-EC isolates in different pre-enrichment broths was determined to establish growth dynamics. Growth of the reference ESBL-EC (NCTC 13441) in four pre-enrichment broths (BPW, DM, BHI, and TS) was monitored every 10 minutes through regular optical density readings at 600 nm wavelength in the CLARIOstar plate reader (BMG Labtech, Ottenberg, Germany) for 24 h. A population logistic model was fitted to the resulting data with the Growthcurver package[18]. The recovery of *E. coli* from different pre-enrichment broths with varying incubation times was tested. *E. coli* (NCTC 13441) was inoculated (approximately 2×10^7 CFU/mL) into 5 mL of four pre-enrichment broths (BPW, DM, BHI, and TS) and grown shaking at 37 °C and 220 rpm. After 2, 4, 6 and 8 hours, 1:10 serial dilutions of each pre-enrichment culture were performed and dilutions 10^{-4} to 10^{-12} were spotted onto LB agar and incubated at 37°C overnight. The following day, the number of colony forming units (CFUs) recovered from each broth was calculated.

Stool was collected from three healthy volunteers, defined as individuals not using antibiotics or experiencing any gastrointestinal problems. The stool was resuspended in a stool diluent solution (1:5 w/v) prepared according to the maltodextrin-trehalose

method with the addition of 10% glycerol [19]. The stool slurry was mixed using a

magnetic stirrer for 15 minutes and allowed to sediment for 5 minutes. Supernatants

from each of the three stool slurries were mixed in equal proportions and stored in

cryotubes at - 80 °C. Before use in experiments, the stool slurry was rapidly thawed in

a water bath at 37°C for 10 minutes and vortexed thoroughly. To determine the optimal

pre-enrichment broth and incubation time for the recovery of ESBL-EC from stool, a

spiked stool model was developed using 1 mL of the pooled stool slurry and a 1 μL

spike of the NCTC 13441 or CAB17W (approximately 1 x 10^5 CFU/mL). The ESBL-

non-producing *E. coli* strain (NCTC 12241) was used as a negative control to

determine if the antibiotic supplement sufficiently inhibited growth. The spiked stool

models were pre-enriched using TS and BPW with and without the addition of

selection (1 μg/ mL cefotaxime) and incubated for 4 and 18 hours, shaking at 37 °C

and 220 rpm. Following incubation, 1:10 serial dilutions of each pre-enrichment culture

were made and dilutions (10^{-4} to 10^{-7} for the 4-hour incubation and 10^{-5} to 10^{-8} for the

18-hour incubation) were plated onto LB agar and CHROMagar in 10 μL spots and

incubated at 37 °C overnight. The number of CFUs recovered from each broth was

calculated.

Recovery of ESBL-EC using different selective agars

Selective agar (MacConkey, Membrane Lactose Glucoronide Agar (MLGA) and

CHROMagar) commonly used in the recovery of ESBL-EC from environmental, clinical

and food sources were selected for comparison in this study. These selective agars

were prepared according to the manufacturer's recommendations **(Supplementary**

Table 1). MacConkey and MLGA agars were supplemented to a final concentration of

1 µg/mL cefotaxime (Melford, Suffolk, UK). To assess the recovery of ESBL-EC from

stool using selective agar, the stool spike model described above was used. The

spiked stool was pre-enriched in BPW for 4 hours, shaking at 37°C and 220 rpm.

Following incubation, 1:10 serial dilutions of each spiked stool sample were made, and

10^{-4} to 10^{-7} dilutions were plated onto MacConkey agar, CHROMagar and MLGA

plates in 10 μL spots and incubated at 37 °C overnight. The number of CFUs

recovered from each selective agar was calculated.

Limit of ESBL-EC detection with pre-enrichment compared with direct plating

A series of 100 μL of CAB17W ranging from 1 CFU/mL to 1×10^8 CFU/mL was spiked

into 100 μL of stool slurry. The spiked stools were serially diluted (1:10) and 10 μL of

each dilution was spotted onto cefotaxime supplemented MacConkey agar and

incubated at 37 °C overnight. A 96 deep well plate was then inoculated with 10 μL of

the remaining stool spikes in 1 mL of BPW, this was done in triplicate for each spike

input concentration. The deep well plate was then sealed with a breathable sealing

membrane and placed in an Innova 42R 19 mm Orbit shaking incubator set at 37 °C

and 220 rpm. After 4 hours the plate was removed from the incubator, each well was

again diluted in series 1:10 and spots of 10μL were plated onto cefotaxime

supplemented MacConkey and incubated at 37°C overnight. The number of CFUs

recovered were then calculated.

DNA extraction kit comparison

A single colony of each of four *E. coli* strains (**Supplementary Table 2:** NCTC 13441,

NCTC 13476, NCTC 13846, CAB17W) were inoculated in 5 mL of LB with and grown

overnight. Then, cultures were diluted to an OD600 of 2 (~ 2×10^9 CFU/mL). The cells

were washed three times in phosphate-buffered saline (PBS). The resulting pellets

were resuspended in 1 mL of PBS and split into three tubes with 250 µL and used as

the input for each extraction method. To assess the yield, quality, efficiency, and cost

of five commercially available kits **(Supplementary Table 1)** and a simple boiling

method[20], we extracted DNA from each of the four *E. coli* isolates in technical

triplicates (n=12 for each extraction method), with three experimental replicates (n=36

extractions per method) on different days. The DNA yield was then measured using

the Qubit 4 Fluorometer (Agilent Technologies, California, USA) and the Agilent

TapeStation System 4150 (Agilent Technologies, California, USA) was used to

determine the DNA integrity. The DNA quality was assessed by measuring the

A260/230 and A260/280 absorbance ratios using the NanoPhotometer (Implen,

California, United States). Scores **(Supplementary Table 3)** were allocated according

to the criteria in **Supplementary Table 4**. Each DNA extraction method was then

ranked according to the score given, from highest (1) to lowest (6) preforming in each

category: DNA yield, DNA quality, cost, hands-on time, ease of protocol and total

protocol time.

Single nucleotide variant analysis

To determine whether our protocol would provide sufficient resolution for the detection

of single nucleotide variants in a stool sample, we processed a spiked stool sample

and an ESBL-EC positive rectal swab. For the spiked stool sample, we inoculated 10

µL of a 100-fold diluted overnight CAB17W culture (~ 8 x 10^7 CFU/mL) into 1 mL of

the pooled stool slurry, vortexed the mixture thoroughly and added 100 µL to 5 mL of

BPW in a 15 mL Falcon tube. The ESBL-EC positive rectal swab was obtained with

permission from a participant in an observational cohort study of faecal ESBL carriage

in hospital patients and care home residents across Liverpool, UK. The rectal swab

10

was placed directly in 5 mL of BPW in a 15 mL Falcon tube. Falcon tubes were

incubated for 4 hours at 37 °C, shaking at 220 rpm. After 4 hours, 100 μL of the culture

was spread on cefotaxime supplemented (1 μg/mL) MacConkey agar and incubated

at 37 °C overnight. The following day, we picked 7 single colonies, restreaked each

colony on another cefotaxime supplemented MacConkey agar and incubated at 37 °C

overnight. DNA was then extracted from each of the purified colonies using the

MasterPure Complete DNA and RNA Purification Kit (Lucigen, Wisconsin, USA)

according to the manufacturer's recommendations. The DNA concentrations were

then measured using the Qubit 4 Fluorometer (Agilent Technologies, California, USA)

and the DNA sent for whole-genome sequencing using the NovaSeq platform (Illumina

Inc, California, USA) at Azenta Life Sciences (Frankfurt, Germany). Sequence data

(reads) were obtained as fastq files. Sequencing adaptors were removed from the

reads using Trimmomatic version 0.39[21] and the quality of the trimmed reads was

assessed using the quality metrics provided by FastQC version 0.11.9[22]. ARIBA[23] was

used to call the AMR genes using the CARD database[24] and to determine ESBL-EC

multilocus sequence type as defined by the seven-gene Achtman scheme[25] hosted at

pubMLST (https://pubmlst.org/). Single nucleotide variants (SNVs) were called

between the relevant reference genome (ST167: GCF_025398915.1; ST131:

GCF_004358405.1; ST1193: GCF_003344465.1) and the single pick sequences

using snippy version 4.3.6 (https://github.com/tseemann/snippy). We then generated

a core genome alignment of all the single pick sequences using snippy-core (with

default settings). All non-nucleotide characters and potentially recombinant sequence

regions were removed from the multisequence alignment using the snippy-

clean_full_aln and Gubbins version 2.3.4[26] respectively. We then extracted the SNVs

from the cleaned multisequence alignment using snp-sites (using the -c option) and

11

created a pairwise SNV distance comparison between each single pick sequence

using snp-dists version 0.8.2 (https://github.com/tseemann/snp-dists).

Statistical analysis

Measurements were summarised as mean and standard deviation or 95% confidence

interval, calculated as one standard deviation above and below the mean value. Box-

and-whisker plots, where presented, show medians as a horizontal line, 25th and 75th

centiles as boxes, and at most 1.5 times interquartile range as whiskers. Analysis of

variance (ANOVA) tests were used to determine the significance of any observed

differences between groups. $P < 0.05$ was considered statistically significant and,

where appropriate, P values were adjusted for multiple comparisons using the

Bonferroni correction.   A post-hoc Tukey test was used for pairwise sample

comparisons when ANOVA tests were significant. All statistical analysis was

performed using R version 4.1.3[27]. All R code is accessible from our Github repository

(https://github.com/SarahGallichan/ESBL-EC_methods.git)

Results

ESBL-EC recovery using different pre-enrichment conditions and selective

agars

We assessed the growth dynamics of two ESBL-EC strains (NCTC 13441 and

CAB17W) in different pre-enrichment broths to determine the optimal incubation time

for recovery of bacteria in log growth phase **(Supplementary Figure 1).** To summarise

the dynamics of the resulting growth curves, the data was fitted to a population logistic

model **(Supplementary Table 5)**. ESBL-EC reached midpoint of log-phase after

approximately 4 hours in BHI, TS, and DM; and after approximately 7 hours in BPW.

Therefore, we chose incubation times of between 2 and 8 hours to quantify ESBL-EC recovery from different broths and assessed bacterial count in CFU from each broth after 4 incubation time points (2, 4, 6 and 8 hours). Across all 4 time points, BHI resulted in the highest and DM the lowest *E. coli* recovery **(Supplementary Figure 2)**. We performed pair-wise broth comparisons at each time point. There was no significant difference between *E. coli* growth in BHI, BPW and TS across all 4 time points **(Supplementary Table 6)**. However, *E. coli* counts in DM was significantly lower than BHI and BPW after 2 hours; BHI, BPW and TS after 4 hours; and BHI and TS after 8 hours of incubation. The largest increase in amount of *E. coli* CFU recovered between the two time points for each broth was from 2 to 4 hours of incubation **(Supplementary Table 6)**, which is within the log growth phase of *E. coli* in each broth. Therefore, we selected 4 hours of incubation to investigate the growth of ESBL-EC from spiked stool.

We selected TS (a high nutrient broth with multiple amino acid sources) and BPW (a low nutrient broth with one amino acid source) for the spike experiments, adding defined amounts of ESBL-EC (NCTC 13441 and CAB17W) to stool samples. Donated stool was confirmed to have no growth of cefotaxime (a 3GC) resistant bacteria by plating on 1µg/mL cefotaxime supplemented MacConkey. We assessed the recovery of these spiked strains, testing whether the addition of selection with cefotaxime and the length of incubation time influenced the concentration of spiked ESBL-EC strains we were able to recover **(Figure 2)**. After 4 hours of incubation, there was no significant difference in CAB17W growth (i.e. CFU concentration) between TS and BPW and whether antibiotic was supplemented or not. However, at 4 hours the growth of NCTC 13441 in the TS without antibiotic supplement was 2.75 CFU/mL (CI: 2.28 –

13

3.33, p = 0.0004) more than in BPW without supplement, while there was no significant

difference between the growth in either broth at 4 hours when supplemented with

cefotaxime.

At 18 hours the effects of the addition of cefotaxime and broth on growth were different

for each strain. Antibiotic supplement significantly increased the recovery of NCTC

13441 after 18 hours compared to no supplementation in TS, by 100.78 CFU/mL (CI:

39.81 – 251.19, p = 0.01). There was no significant difference in growth of NCTC

13441 at 18 hours between TS and BPW. However, recovery of CAB17W was

significantly higher in BPW after 18 hours of incubation compared to TS, by a

difference of 68 CFU/mL (CI: 51.11 – 91.26, p = 0.0001) with antibiotic supplement

and 41.16 CFU/mL (CI: 33.15 – 51.10, p < 0.0001) without an antibiotic supplement.

While an 18-hour pre-enrichment in antibiotic supplemented broth best recovered the

highest CFU/mL for both NCTC 13441 and CAB17W, there were inconsistencies

between broths depending on the strain. Therefore, we concluded that a 4-hour pre-

enrichment in BPW was better for a more consistent increase in both ESBL-EC strains

without requiring an antibiotic supplement.

Next, we then compared ESBL-EC recovery on three different ESBL-EC selective

agars (CHROMagar, MLGA and MacConkey) from our stool spike models. ESBL-EC

was recovered from all agars, with no significant differences in CFU/mL

(Supplementary Table 8). However, there was a clear difference in the cost per

sample of each selective agar, with MacConkey being the most cost-effective and

CHROMagar being the most expensive **(Supplementary Table 1)**.

The limit of ESBL-EC detection from CAB17W spiked stool was then determined for

4-hour pre-enrichment in BPW and compared with directly plating on cefotaxime

supplemented MacConkey. Overall, the mean difference in ESBL-EC recovery

between 4-hour pre-enrichment in BPW and direct plating was 47 CFU/mL (CI: 9.65 –

233.08, $p < 0.001$) **(Supplementary Figure 3)**. The limit of detection using direct

plating was 1.91 CFU/mL and with a 4-hour pre-enrichment, the limit of detection

improved to 0.05 CFU/mL.

Large variability in yield and quality of DNA using different extraction methods

DNA from the same input concentration of four different *E. coli* strains

(Supplementary Table 2) was extracted using five commercially available kits **(Table**

3) and a simple boiling method. The DNA yield was measured using the Qubit

fluorometer **(Figure 3)**. The boiling method resulted in the highest mean DNA yield

overall for the Qubit (16.66 ng/mL, 95% CI: 16.16 – 17.16) while the DNeasy kit

resulted in the lowest (3.28 ng/mL; 95% CI: 4.28 – 2.28) **(Figure 3)**.

Figure 3 *DNA yield from the six extraction methods measured using the Qubit. Technical (each sample extracted in triplicates) and experimental replicates (each extraction method performed three separate times) are plotted for each extraction kit (n = 36). The outliers are shown as points outside of the boxplot whiskers.*

The DNA quality was assessed using the A260/280 absorbance ratio. Downstream

applications requiring high quality DNA, such as short-read and especially long-read

sequencing platforms, require that the A260/A280 absorbance ratio is in the range

between 1.8 to 2.1. The Promega Wizard kit and the DNeasy kit had the largest

proportion of technical and experimental replicates (67%) within this range while the

Zymo Miniprep kit had the lowest proportion (11%) **(Figure 4A).** Additionally, the DNA

integrity was assessed with the DNA Integrity Number (DIN), which is a numerical

assessment (1 to 10) of the DNA integrity determined by the TapeStation system

algorithm. A DIN score of 1 indicated very degraded DNA and DIN 10 indicated highly

intact DNA. The Lucigen MasterPure and the Zymo miniprep kits had the highest DIN

followed closely by the Promega Wizard **(Table 1).**

Table 1 *Mean of measurements (technical and experimental replicates averaged*
across three independent experiments) for each DNA extraction method, including
the 95% confidence intervals, for the tested E. coli and proportion (percentage of
technical replicates) within defined absorbance ratio range.

	Boiling	DNeasy	Lucigen MasterPure	Promega Wizard	NEB Monarch	Zymo Miniprep
DNA yield (ng/ µL)	16.66 ± 4.66	3.28 ± 1.00	13.37 ± 0.58	7.37 ± 1.62	7.35 ± 0.97	4.85 ± 0.48
Absorbance ratio (A260/A280)	1.74 ± 0.06	1.99 ± 0.15	2.1 ± 0.03	2.06 ± 0.13	1.74 ± 0.15	1.66 ± 0.17
Proportion within range (1.8 – 2.1)	14%	67%	50%	67%	42%	11%
Absorbance ratio (A260/A230)	0.70 ± 0.04	3.00 ± 0.73	2.89 ± 0.60	4.44 ± 1.27	1.2 ± 0.63	1.74 ± 0.41
Proportion within range (2.0 – 3.0)	0%	50%	59%	0%	8%	0%
DNA Integrity Number	1.97 ± 1.56	5.40 ± 0.50	7.40 ± 0.35	7.18 ± 0.45	6.08 ± 0.45	7.40 ± 0.25

The purity of extracted DNA was assessed using the 260/230 absorbance ratio. DNA

with a 260/230 absorbance ratio of 2.0 – 3.0 are considered pure and mostly free of

contaminants (such as phenols) [28]. Lucigen Masterpure had the largest proportion of

replicates (59%) within the suggested A260/A230 absorbance range, while the simple

boiling method, the Zymo miniprep kit and the Promega Wizard kit failed to yield DNA

within this purity range **(Figure 4B).**

Figure 4 *Absorbance ratios **(A)** A260/280 and **(B)** A260/230 indicating the presence of contamination and quality of DNA, respectively, of the extracted DNA extracted using six extraction methods. Technical (each sample extracted in*

triplicates) and experimental replicates (each extraction method preformed three separate times) are plotted for each extraction kit (n = 36). The outliers are shown as points outside of the boxplot whiskers.

Using all the measurement data, cost, ease of protocol (technical difficulty and

equipment requirements), and time aspects (hands-on and total-protocol time) from

the comparison of the five commercially available DNA extraction kits, a ranking matrix

(Table 2) was produced based on the scores for each extraction method

(Supplementary Table 3). Simple boiling scored highly for yield, cost, and time but

low for the purity scores (A260/230, A260/280 and DIN). Commercial kits performed

better in this regard, with the Promega HMW and Qiagen DNeasy ranking highest for

the A260/280 and the MasterPure ranking highest for the A260/230 absorbance ratio

and the DIN score. All other rankings were variable between kits. The Lucigen

MasterPure Complete DNA and RNA Purification Kit performed best for the extraction

of high yield and quality DNA for sequencing to investigate single nucleotide variants

of ESBL-EC in stool.

Table 2 *Ranking of different DNA extraction methods, highest (1) to lowest (6).*

DNA extraction kit	Cost	Time		Mean DNA yield (ng/ µL)	DNA quality	Difficulty
		Hands-on	Total			
Qiagen DNeasy	5	2	3	2	5	3
Lucigen Masterpure	2	4	5	1	1	6
Promega HMW	5	4	5	2	1	5
NEB Monarch	3	1	2	2	4	2
Zymo Miniprep	4	2	3	2	1	3
Boiling	1	1	1	1	6	1

Validation of workflow

To investigate whether our workflow of a 4-hour pre-enrichment in BPW, plating on cefotaxime supplemented MacConkey and extraction using the MasterPure kit was sufficient for investigation of SNVs in closely related ESBL-EC strains, we performed both a control experiment where we processed an ESBL-negative stool spiked with a single clinical ESBL-EC strain (CAB17W) and assessed an ESBL-EC positive patient rectal swab using our workflow. Seven colonies from each of the cefotaxime supplemented MacConkey agar plates were then picked and sequenced. SNV analysis of the spiked stool confirmed that there were no SNVs between any of the single colony pick genomes as expected for a clonal spike **(Supplementary Table 10)**. Analysis of the ESBL-EC positive rectal swab revealed that the patient had two different ESBL-EC sequence types ST131 that had bla$_{CTX-M-27}$ and ST1193 that had bla$_{CTX-M-15}$. SNV analysis of each ESBL-EC sequence type showed that there were 5 SNV differences between the ST131 single picks **(Supplementary Table 11)** and 1 SNV difference between one of the ST1193 picks and the other four single picks **(Supplementary Table 12)**. We therefore concluded that our workflow was appropriate for SNV analysis of ESBL-EC from stool.

Discussion

Although *E. coli* is the dominating species of Enterobacteriaceae in the gut of healthy human adults, *E. coli* only constitute around 0.5% - 5% of the gut microbial community [29]. This makes it difficult for methods such as shotgun metagenomics to disentangle the diversity of pathogenic and non-pathogenic *E. coli* within a highly complex human gut microbiome without using costly very deep sequencing [6,30]. On the other end of the spectrum, the analysis of single colony picks from stool samples directly plated on

antibiotic supplemented agar, as is practice in many routine diagnostic laboratories, is

insufficient to capture the intraspecies diversity of ESBL-EC, which can be substantial,

as each colony would need to be analysed which would be extremely time-, cost- and

labour-intensive[31,32]. Both methods are thus not feasible to implement as routine tool

to monitor transmission of ESBL-EC in healthcare settings. Therefore, we investigated

the impact of different microbiological approaches to optimise the recovery of ESBL-

EC from human stool. We compared different methods at each stage of laboratory

processing; pre-enrichment, isolation on selective agar, and DNA extraction, and were

able to optimise a rapid and cost-effective approach for the recovery of ESBL-EC for

targeted surveillance of ESBL-EC colonisation with the potential for sub-species

resolution.

Numerous transmission studies have highlighted the challenge of capturing ESBL-EC

variants present in small numbers in stool[33–35]. *E. coli* are only a small part of the

diversity present within complex matrices and clinical samples (e.g., stool), increasing

the likelihood of a test resulting in false negatives when stool is directly plated on

supplemented agar[36,37]. To improve ESBL-EC detection for targeted surveillance,

previous studies used pre-enrichment broth before plating on supplemented agar and

were successful in recovering multiple ESBL-EC genotypes from human stool[17].

Indeed, we also showed that pre-enrichment is an effective tool to increase sensitivity

for ESBL-EC detection from stool, thus improving the amount of ESBL-EC recovered

on supplemented agar.

The standard practice of pre-enriching a sample for 18 to 24 hours is not conducive to

rapid diagnosis and patient intervention[38,39]. A further concern with longer pre-

22

enrichment is the increased risk of creating competition with other lineages or species

and skewing the relative abundance. Further, the potential of movement of mobile

genetic elements (e.g. plasmids) can bias the recovery of the original strain[40]. We

found that growth of ESBL-EC was variable and strain-dependent after an 18-hour

pre-enrichment with and without the addition of an antibiotic supplement, suggesting

that there is indeed potential for competition bias with longer incubation times.

Therefore, we investigated a short pre-enrichment and found that a 4-hour pre-

enrichment yielded sufficient growth for reliable downstream identification of ESBL-

EC. Hence, we concluded that 4-hour pre-enrichment was not only preferable for

consistent ESBL-EC recovery but also did not necessitate the addition of an antibiotic

supplement and reduced the time and cost.

There are a variety of recommended supplemented agars available for ESBL-EC

detection. Cefotaxime supplemented MacConkey agar is commonly used for

surveillance studies and clinical microbiology laboratories since it is readily available

and the cheapest of the three agars compared here[10,11,41]. However, chromogenic

agar (CHROMagar ESBL[42]) is an increasingly popular alternative to MacConkey since

it is effective at distinguishing between *E. coli* and other ESBL-producing

Enterobacteriaceae using colour[38,43,44]. In addition to these two agars, we also tested

ESBL-EC recovery on cefotaxime supplemented Membrane Lactose Glucuronide

Agar (MLGA), which is commonly used for detecting faecal contamination in water

sources[45]. We found that all three agars were effective in recovering from ESBL-EC

from pre-enriched stool, allowing for flexibility with selective agar choice; we hence

selected cefotaxime supplemented MacConkey as the most cost-effective in our

setting.

We tested a wide variety of DNA extraction methods including commercial DNA

purification kits and the boiling method as the simplest and cheapest method of DNA

extraction from *E. coli*[46]. Since transmission studies require high resolution typing to

clearly distinguish between two *E. coli* strains, high quality DNA is required.

Commercially available kits provide standardised reagents, consumables and

validated DNA extraction and purification methods. Depending on the extraction

protocol, specialised equipment e.g. refrigerated centrifuges, skilled personnel, and

an increased amount of time and cost are needed to extract DNA[47]. Taking this into

consideration, along with the cost-per-sample, quality, and yield of DNA we compared

six DNA extraction methods. While the boiling method was the simplest, cheapest,

least time-consuming DNA extraction method with the highest yield and thus arguably

of high value for highly time-sensitive applications or in cost-restricted setting, we show

that the Lucigen MasterPure produced comparable DNA yield with much higher quality

scores that were consistently reproducible.

Finally, we validated our optimised ESBL-EC recovery method using a spiked stool

sample and a patient rectal swab. For the spiked stool sample, all sequenced 7

colonies were identical and clonality was confirmed at single nucleotide resolution.

With the patient rectal swab, we demonstrated the value of multiple colony picks

through the recovery of two different ESBL-EC sequence types and the analysis of

variation within each sequence type at single nucleotide resolution. The analysis of

both samples at suggests that our method is suitable for studies aiming to infer

transmission using SNV. While the methods we show here have focused on the

recovery of ESBL-EC from stool, we suggest that these methods are also applicable

24

or easily adaptable for the recovery other drug resistant Enterobacteriaceae. ESBL-

producing *Klebsiella* and *Enterobacter* species occupy a similar niche to ESBL-EC and

are all recovered from stool through direct plating on supplemented agar[10]. With the

increasing prevalence of drug resistant infections worldwide, it has become crucial to

understand transmission pathways to put targeted preventative measures in place to

interrupt transmission.

Our study demonstrates that a 4-hour pre-enrichment in either BPW or TS is effective

for the recovery of ESBL-EC from stool without the addition of a third-generation

cephalosporin and cefotaxime supplemented MacConkey, chromogenic agar and

MLGA are all viable options for the recovery of pre-enriched ESBL-EC. The quality

and quantity of DNA using different extraction kits is variable and not all yield high-

quality DNA required for whole genome sequencing. The MasterPure Complete DNA

and RNA Purification Kit produced a DNA yield that was comparable to the simple

boiling method, but with higher quality scores that were consistently reproducible.

Altogether this optimised method was sufficient for SNV analysis of an ESBL-EC

spiked stool sample.

Ethics

The healthy volunteers gave confidential consent for the use and storage of stool

samples (LSTM Research Tissue Bank RTB/2022/007). The observational cohort

study of hospital patients and care home residents in facilities in Liverpool was

approved by the National Research Ethics Service Greater Manchester South ethics

committee (ref: 22/NW/0343). Written informed consent was obtained from

participants or consultees, as appropriate.

Supplementary materials and data availability

Supplementary Tables 1 – 12 and Supplementary Figures 1 - 3 are accessible in the supplementary material of this manuscript. The R scripts used to generate the figures, supplementary figures and data analyses in this manuscript are available from the GitHub repository https://github.com/SarahGallichan/ESBL-EC_methods.git. Protocols are available on Protocols.io (dx.doi.org/10.17504/protocols.io.kxygxyk3dl8j/v). Reads from sequenced isolates in this study are accessible in the Sequence Read Archive (SRA) using BioProject ID: PRJNA1095376.

Funding

This work was supported by iiCON (infection innovation consortium) via UK Research and Innovation (107136) and Unilever (MA-2021-00523N).

Author contributions

Conceptualisation and method development was done by S.G., S.F., M.M., E.H., J.M.L., N.A.F., and F.E.G. Investigation was undertaken by S.G., S.F., E.P.-B, and C.M. Data analysis was done by S.G., J.M.L. and F.E.G. The original draft was prepared by S.G., E.H., J.M.L., N.A.F., and F.E.G and then reviewed and edited by all authors. Supervision was provided by E.H., J.M.L., N.A.F., and F.E.G.

Acknowledgements

We would like to thank Ross Gray for the illustrations used in Figure 1 of this manuscript, Finlay Hitchman for his assistance with the pre-enrichment broth

520 comparison work, and all the clinical staff and study participants that facilitated this

521 study.

References

1. WHO publishes list of bacteria for which new antibiotics are urgently needed.

https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-

which-new-antibiotics-are-urgently-needed.

2. Schwaber, M. J. & Carmeli, Y. Mortality and delay in effective therapy associated

with extended-spectrum beta-lactamase production in Enterobacteriaceae

bacteraemia: a systematic review and meta-analysis. *J Antimicrob Chemother*

60, 913–920 (2007).

3. Melzer, M. & Petersen, I. Mortality following bacteraemic infection caused by

extended spectrum beta-lactamase (ESBL) producing E. coli compared to non-

ESBL producing E. coli. *Journal of Infection* **55**, 254–259 (2007).

4. Blom, A., Ahl, J., Månsson, F., Resman, F. & Tham, J. The prevalence of ESBL-

producing Enterobacteriaceae in a nursing home setting compared with elderly

living at home: a cross-sectional comparison. (2016) doi:10.1186/s12879-016-

1430-5.

5. Bettelheim, K. A. Escherichia coli in the normal flora of humans and animals. in

Escherichia coli: mechanisms of virulence (ed. Sussman, M.) 85–109

(Cambridge University Press, Cambridge, 1997).

6. Foster-Nyarko, E. & Pallen, M. J. The microbial ecology of Escherichia coli in

the vertebrate gut. *FEMS Microbiol Rev* **46**, (2022).

7. Alm, E. A. , W. S. T. , and G. D. M. The niche of Escherichia coli. in *In Population*

Genetics of Bacteria: A Tribute to Thomas S. Whittam (eds. Walk, S. T. & Feng,

P. C. H.) 68–89 (ASM Press, Washington, DC, USA, 2011).

8. Vollmerhausen, T. L. *et al.* Population structure and uropathogenic virulence

associated genes of faecal Escherichia coli from healthy young and elderly

adults. *J Med Microbiol* **60**, 574–581 (2011).

9. Balloux, F. *et al.* From Theory to Practice: Translating Whole-Genome

Sequencing (WGS) into the Clinic. *Trends Microbiol* **26**, 1035–1048 (2018).

10. Public Health England. *UK Standards for Microbiology Investigations Detection*

of Enterobacteriaceae Producing Extended Spectrum β-Lactamases-

Standards-for-Microbiology-Investigations-Smi-Quality-and-Consistency-in-

Clinical-Laboratories. (2016).

11. World Health Organization & Advisory Group on Integrated Surveillance on

AMR. *WHO Integrated Global Surveillance on ESBL-Producing E. Coli Using a*

"One Health" Approach: Implementation and Opportunities.

https://www.who.int/publications/i/item/9789240021402 (2021).

12. Bradford, P. A. Extended-spectrum beta-lactamases in the 21st century:

characterization, epidemiology, and detection of this important resistance threat.

Clin Microbiol Rev **14**, 933–951 (2001).

13. Naas, T., Cotellon, G., Ergani, A. & Nordmann, P. Real-time PCR for detection

of blaOXA-48 genes from stools. *J Antimicrob Chemother* **68**, 101–104 (2013).

14. Wu, G. *et al.* Comparative Analysis of ESBL-Positive Escherichia coli Isolates

from Animals and Humans from the UK, The Netherlands and Germany. *PLoS*

One **8**, e75392 (2013).

15. Petty, N. K. *et al.* Global dissemination of a multidrug resistant Escherichia coli

clone. *Proc Natl Acad Sci U S A* **111**, 5694–5699 (2014).

29

16.   Toombs-Ruane, L. J. *et al.* Carriage of Extended-Spectrum-Beta-Lactamase-

and AmpC Beta-Lactamase-Producing Escherichia coli Strains from Humans

and Pets in the Same Households. *Appl Environ Microbiol* **86**, (2020).

17.   Ludden, C. *et al.* Defining nosocomial transmission of Escherichia coli and

antimicrobial resistance genes: a genomic surveillance study. *Lancet Microbe*

2, e472–e480 (2021).

18.   Sprouffske, K. & Wagner, A. Growthcurver: An R package for obtaining

interpretable metrics from microbial growth curves. *BMC Bioinformatics* **17**, 1–4

(2016).

19.   Burz, S. D. *et al.* A Guide for Ex Vivo Handling and Storage of Stool Samples

Intended for Fecal Microbiota Transplantation. *Sci Rep* **9**, (2019).

20.   Dashti, A. A., Jadaon, M. M., Abdulsamad, A. M. & Dashti, H. M. Heat Treatment

of Bacteria: A Simple Method of DNA Extraction for Molecular Techniques.

Kuwait Medical Journal **41**, 117–122 (2009).

21.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

22.   Andrews, S. FastQC: A Quality Control tool for High Throughput Sequence

Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (2010).

23.   Hunt, M. *et al.* ARIBA: rapid antimicrobial resistance genotyping directly from

sequencing reads. *Microb Genom* **3**, (2017).

24.   Alcock, B. P. *et al.* CARD 2023: expanded curation, support for machine

learning, and resistome prediction at the Comprehensive Antibiotic Resistance

Database. *Nucleic Acids Res* **51**, D690–D699 (2023).

25.   Wirth, T. *et al.* Sex and virulence in Escherichia coli: an evolutionary perspective.

Mol Microbiol **60**, 1136–1151 (2006).

26.   Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of
recombinant bacterial whole genome sequences using Gubbins. *Nucleic Acids*
Res **43**, e15–e15 (2015).

27.   R Core Team. _R: A Language and Environment for Statistical Computing_. *R*
Foundation for Statistical Computing, Vienna, Austria. <https://www.R-
project.org/> (2023).

28.   DNA   Purification | DNA   Extraction   Methods | Promega.
https://www.promega.co.uk/resources/guides/nucleic-acid-analysis/dna-
purification/.

29.   Tenaillon, O., Skurnik, D., Picard, B. & Denamur, E. The population genetics of
commensal Escherichia coli. *Nat Rev Microbiol* **8**, 207–217 (2010).

30.   Huttenhower, C. *et al.* Structure, function and diversity of the healthy human
microbiome. *Nature 2012 486:7402* **486**, 207–214 (2012).

31.   Stoesser, N. *et al.* Extensive Within-Host Diversity in Fecally Carried Extended-
Spectrum-Beta-Lactamase-Producing Escherichia coli Isolates: Implications for
Transmission Analyses. *J Clin Microbiol* **53**, 2122–2131 (2015).

32.   Pallen, M. J. *et al.* Genomic diversity of Escherichia coli from healthy children in
rural Gambia. doi:10.7717/peerj.10572.

33.   Aguilar-Bultet, L. *et al.* Within-host genetic diversity of extended-spectrum beta-
lactamase-producing Enterobacterales in long-term colonized patients. *Nature*
Communications 2023 14:1 **14**, 1–14 (2023).

34.   Adler, A. *et al.* Transmission dynamics of ESBL-producing Escherichia coli
clones in rehabilitation wards at a tertiary care centre. *Clinical Microbiology and*
Infection **18**, E497–E505 (2012).

35.   Duval, A. *et al.* Close proximity interactions support transmission of ESBL-K.

pneumoniae but not ESBL-E. coli in healthcare settings. *PLoS Comput Biol* **15**,

e1006496 (2019).

36.   Peto, L. *et al.* Selective culture enrichment and sequencing of feces to enhance

detection of antimicrobial resistance genes in third-generation cephalosporin

resistant Enterobacteriaceae. *PLoS One* **14**, (2019).

37.   Huttenhower, C. *et al.* Structure, function and diversity of the healthy human

microbiome. *Nature* **486**, 207–214 (2012).

38.   Rondinaud, E., Ruppé, E., Matheron, S., Lucet, J. C. & Armand-Lefevre, L.

Screening   methods   for   intestinal   carriage   of   multidrug-resistant

Enterobacterales: interest of enrichment broth. *Diagn Microbiol Infect Dis* **97**,

115079 (2020).

39.   Jazmati, T., Hamprecht, A. & Jazmati, N. Comparison of stool samples and

rectal swabs with and without pre-enrichment for the detection of third-

generation   cephalosporin-resistant   Enterobacterales   (3GCREB).

doi:10.1007/s10096-021-04250-1/Published.

40.   Conrad, C. C., Stanford, K., McAllister, T. A., Thomas, J. & Reuter, T.

Competition during enrichment of pathogenic Escherichia coli may result in

culture bias. *https://doi.org/10.1139/facets-2016-0007* **1**, 114–126 (2016).

41.   Jacob, M. E., Keelara, S., Aidara-Kane, A., Matheu Alvarez, J. R. & Fedorka-

Cray, P. J. Optimizing a Screening Protocol for Potential Extended-Spectrum β-

Lactamase Escherichia coli on MacConkey Agar for Use in a Global

Surveillance Program. *J Clin Microbiol* **58**, (2020).

42.   CHROMagar™   ESBL   -   Chromagar.

https://www.chromagar.com/en/product/chromagar-esbl/.

43.   El-Jade, M. R. *et al.* ESBL Detection: Comparison of a Commercially Available

Chromogenic Test for Third Generation Cephalosporine Resistance and

Automated Susceptibility Testing in Enterobacericeae. *PLoS One* **11**,

e0160203 (2016).

44.   Réglier-Poupet, H. *et al.* Performance of chromID ESBL, a chromogenic medium

for detection of Enterobacteriaceae producing extended-spectrum β-

lactamases. *J Med Microbiol* **57**, 310–315 (2008).

45.   Sartory, D. P. & Howard, L. A medium detecting β-glucuronidase for the

simultaneous membrane filtration enumeration of Escherichia coli and coliforms

from drinking water. *Lett Appl Microbiol* **15**, 273–276 (1992).

46.   Shah DH, S. S. B. T. C. D. *Molecular Detection of Food Borne Pathogens*.

(Taylor & Francis Group, CRC Press, USA, 2010).

47.   Barbosa, C., Nogueira, S., Gadanho, M. & Chaves, S. DNA extraction: finding

the most suitable method. *Molecular Microbial Diagnostic Methods: Pathways*

to Implementation for the Food and Water Industries 135–154 (2016)

doi:10.1016/B978-0-12-416999-9.00007-1.