1 **Title:**
2 # Vaccination mitigates climate-driven disruptions to malaria control
3

4 **Authors**
5 Benjamin L. Rice[1]*, Estelle Raobson[2,3], Sylviane Miharisoa[4], Mahery Rebaliha[5,6], Joseph Lewinski[6],
6 Hanitriniaina Raharinirina[2], Bryan Greenhouse[7], Christopher D. Golden[5,8], Gabriel A. Vecchi[9,10], Amy
7 Wesolowski[11], Bryan Grenfell[1,12], C. Jessica E. Metcalf[1,12]
8
Affiliations
1. Department of Ecology and Evolutionary Biology, Princeton University; Princeton, NJ, USA.
2. Mention Zoologie et Biodiversité Animale, University of Antananarivo; Antananarivo,
Madagascar.
3. Association Vahatra; Antananarivo, Madagascar.
4. Institut Pasteur de Madagascar; Antananarivo, Madagascar.
5. Madagascar Health and Environmental Research (MAHERY); Maroantsetra, Madagascar.
6. Multisectoral Malaria Project, Catholic Relief Services; Baltimore, MD, USA.
7. EPPIcenter Program, Division of HIV, Infectious Diseases, and Global Medicine, Department of
Medicine, University of California San Francisco; San Francisco, CA, USA.
8. Department of Nutrition, Harvard TH Chan School of Public Health; Boston, MA, USA.
9. Department of Geosciences, Princeton University; NJ, USA.
10. High Meadows Environmental Institute, Princeton University; NJ, USA.
11. Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health; Baltimore,
MD, USA.
12. Princeton School of Public and International Affairs, Princeton University; Princeton, NJ, USA.
25
26 *Corresponding author. Email: b.rice@princeton.edu
27
28

Abstract

Increasingly extreme weather events in high malaria burden areas threaten progress to disease
control targets. Yet, data on the impact of these events on control programs remain rare. Using
unique data from Madagascar, we estimate high rates of infection in the wake of two major
tropical cyclones. Evidence that infection rebounds rapidly during gaps in interventions indicates
maintaining continuity of coverage is crucial to limiting burden. Relative to other interventions,
recently available malaria vaccines have a longer duration of protection, with the potential to
address interruptions in prevention deployment. Evaluating this use case, we quantify the
reduction in symptomatic infections expected for a range of vaccination scenarios. We find long-
lasting interventions such as vaccination are a key mitigation measure against climatic
disruptions to disease control.

Main Text

Malaria remains a major global health challenge (*1, 2*). Much work has investigated how the coincident challenge of climate change threatens progress to malaria control (*1*). Rising temperatures modify mosquito vector dynamics (*3–5*), altering infection risk (*6, 7*), but these impacts are often focused in low burden areas at the altitudinal and latitudinal margins of malaria transmission (*8–10*). By contrast, extreme weather events, with some types likely to increase in frequency or severity due to climate change (*11–14*), disrupt public health efforts (*15–19*), including in areas where the burden of malaria is concentrated. Data on the impact of climate-related disasters on malaria infection remain rare (*18, 20, 21*), despite high rates of exposure to such events in malaria endemic settings and the clear importance of disruptions amidst the changing landscape of malaria control activities.

In response to stalled progress in malaria control in high burden countries and the development of new treatment and prevention tools (*1*), the World Health Organization (WHO) and national malaria control programs are preparing the large-scale deployment of new interventions. These include scaling the mass distribution of antimalarial chemoprophylaxis and treatment (e.g., seasonal or perennial malaria chemoprophylaxis, SMC, PMC), and vaccination (see **Table S1**). Continuity is important to all of these interventions, as their effectiveness erodes over time, thus requiring regular re-administration. Extreme weather events threaten that continuity by delaying or degrading public health activities (*18*). Understanding the barrier this poses to achieving malaria control targets requires detailed data on infection risk during periods of likely disruption. However, limitations in surveillance systems make such data rare.

Tropical cyclones are an example of extreme weather events repeatedly shown to disrupt health systems for weeks to months following the event (*21–26*). Madagascar, our focus here, is subject to regular tropical cyclones (**Figure 1**), resulting in widespread destruction, population displacement, and damage to healthcare-related infrastructure (**Data S1**). We leverage a unique dataset comprising repeated measures of malaria infection status surrounding cyclone events (*n* = 20,718 observations) to develop a platform to explore the spectrum of recommended interventions and their robustness under climate change.

Results

1. High rates of exposure to cyclones and malaria infection

1.1 Cyclone frequency and disruptive effect in Madagascar

Cyclone frequency and severity has increased in the southern Indian Ocean (*27, 28*) with storms in Madagascar causing repeated humanitarian emergencies in recent years. Cyclones Batsirai and Freddy destroyed over 300 health facilities (*29–32*) and resulted in an estimated 112,115 and 290,000 people in need of immediate humanitarian assistance in 2022 and 2023, respectively (*33*) (**Figure 1D**). Excess rainfall for Cyclone Batsirai was made more likely by increased

83 greenhouse gas and changes in aerosol emission per climate attribution studies (*34*). Our study
84 site, Mananjary district, is located on the east coast of Madagascar, a moderate to high malaria
85 transmission area.

86
87

88 **Figure 1: The scale of cyclone-malaria interactions globally and in Madagascar**

90
Figure 1A: Blue lines show cyclone tracks from IBTrACS for all storms since 1980 (*35*, *36*). Malaria incidence
estimates are from the Malaria Atlas Project (*37*). Inset: Intensity for the 1705 recorded storms impacting currently
malaria endemic countries since 1980 (defined as storm center approaching within 60 nautical miles of landfall). For
boxplots, the midline shows the median number of storms in the period 1980-2023, lower and upper hinges
correspond to the 25th and 75th percentiles and whiskers extend 1.5 times the interquartile range. Points show the
number of storms for a given year, jittered for visibility. Categories follow the Saffir-Simpson wind scale for the
maximum wind speed observed within 60 nautical miles of landfall: Tropical Depression (TD), Tropical Storm (TS),
Hurricane Category 1-5 equivalent (H1-H5), or no wind speed data available (ND).

101
Figure 1B: Malaria burden versus cyclone exposure at the national scale. Bivariate fill color (*38*) shows average
annual person-days of exposure to tropical cyclones from 2002-2019 (*39*) vs reported national malaria incidence
(*40*). Countries with no reported malaria cases or cyclone exposure are in gray. Madagascar circled for reference.
Inset: Malaria incidence vs the number of tropical cyclones since 1980 passing within a 60 nautical mile buffer of
coastline (from IBTrACS), on a log-10 scale. Points are labeled by three-letter country codes. Among high malaria
burden countries (annual incidence > 100 per 1000 population), Madagascar (MDG, circled) and Mozambique
(MOZ) have the highest observed cyclone frequencies, followed by the Solomon Islands (SLB) and Papua New
Guinea (PNG).

Figure 1C: Tropical cyclone tracks in Africa colored by year of the storm. Inset: Tropical storm activity is highest in southeast Africa with over 47.9% of impacts in Africa overall contributed by Madagascar and Mozambique.

Figure 1D: Tropical cyclone activity for Madagascar since 1980 with the study district, Mananjary, outlined. Tracks show the estimated path of the storm center, with those passing within 60 nautical miles (111.1 km) of the Mananjary district shown in blue (others in gray). Two examples of recent cyclones impacting the Mananjary district, Cyclone Batsirai and Cyclone Freddy, are highlighted with estimated wind speed (data from IBTrACS) and estimates of the number of individuals requiring humanitarian assistance (data from EM-DAT).

1.2 Characterizing malaria infection rates

We characterize rates of malaria infection from a longitudinal cohort study of a random sample of 500 households (2954 individuals, all ages) from 10 localities (**Figure 2A**). Repeated sampling of malaria infection status by rapid diagnostic test (RDT) was performed for the period July 2021-April 2023 (mean number of samples per enrolled individual = 7.7, median = 9) (**Figure S1**). The interval between samples was short (mean = 57.4 days, median = 52 days) relative to the duration of an untreated *P. falciparum* infection (approx. 180 days (*41*, *42*)), indicating a low probability of natural infection clearance prior to the next sample. All individuals positive for malaria were treated to clear infections, allowing estimates of rates of infection and re-infection (see methods). During the course of the study, the area was impacted by Cyclone Batsirai (February 5, 2022) and Cyclone Freddy (February 21, 2023), providing estimates of malaria infection rates in the period before and for the 2 months after each event.

Figure 2: Malaria infection rates following tropical cyclones in southeast Madagascar

Figure 2A: Sampling locations for cohort study sites (MNJ.01-MNJ.10, ordered from north to south). Location of the Mananjary urban area, the capital of the Mananjary district, shown with a square. District boundary shown in white, elevation data from the Copernicus Global Digital Elevation Model (*43*).

141
Figure 2B: Raw malaria infection observations by rapid diagnostic test (RDT) by individual for the 10 follow-up
sampling time points (T01-T10) for the 10 sample localities in Mananjary District, Madagascar (MNJ.01-MNJ.10).
For visualization, each row is an individual (individuals with >50% of attendance at sampling time points shown)
with a random subsample of 100 individuals shown for each locality (full data in supplement). Inset: An overview of
sampling methods. RDT positive individuals are treated to clear infections and estimate time to re-infection. Some
infections are uncounted due a false negative RDT result (shown as circles with black outline and red fill).

148
149 **Figure 2C**: Expected proportion of individuals infected in one month by site and age category (young children = 0-
150 5y, school-aged children = 6-13y, teenagers = 14-19y, adults = 20+y). Sites are ordered by infection rate.

1.3 Malaria infection results: Force of infection and exposure following cyclones

We derive estimates of the force of infection (FOI) by age and season from ten follow-up
sampling time points (see methods) (**Figure 2B**). FOI peaked for school-aged children (**Figure S2**), consistent with the higher frequency of infection commonly observed in this group (e.g.,
(*44*)). Transmission fluctuated seasonally, with the percentage of school-aged children expected
to be infected within a month ranging, by site, from 0.8-9.1% in October to 4.9-44.3% in May
(**Figure 2C**). At the higher FOI sites (e.g., MNJ.01, MNJ.08, MNJ.09, MNJ.10), we estimate
13.5-35.6% of younger children and 20.0-49.1% of school aged children were exposed to
malaria infection in the 2 months following Cyclones Batsirai and Freddy (**Figure 2D**).
Equipped with these empirically derived estimates of the scale of the rate of infection, we can
evaluate the potential impacts of disruptions to malaria control programs during cyclone season.

164
Figure 2D: Estimated probability of an uninfected individual from a standard household becoming infected by
malaria after Cyclone Batsirai (February 5, 2022) and Cyclone Freddy (February 21, 2023) at study sites in
Mananjary district, Madagascar.

2. Malaria incidence is sensitive to brief disruptions in control

2.1 Supplemental malaria interventions in the context of disruption

Current operational plans for high malaria burden countries, including Madagascar (*45*),
recommend implementing or expanding interventions using anti-malaria drugs to reduce burden

in targeted populations, as a supplement to existing base activities (i.e., bednet distribution).

Approaches use the regular administration of either a standard first-line antimalarial with a short

half-life (protective against reinfection for 13-15 days (*46*)) to clear existing infections (mass

drug administration, MDA, or mass testing and treatment, MTaT) or a longer lasting antimalarial

(protective for 21-42 days (*47*, *48*)) to also prevent new infections (seasonal or perennial malaria

chemoprophylaxis, SMC, PMC, or intermittent preventive treatment of malaria in pregnancy,

IPTp) (see **Table S1**).

2.2 Consequences of temporal gaps in coverage

We explore the deployment of each of these supplemental interventions across the observed

range of FOIs, introducing delays to follow-up administration rounds at different points along a

seasonal cycle (see methods). Following the cessation of drug administration, protection decays

at a rate determined by the drug half-life, and infections accumulate at a rate determined by the

FOI for that locality, age group, and seasonal period (**Figure 3A**). Historical cyclone data for

Madagascar indicates the time interval with the highest risk for storm impact occurs between

January-April, contributing 81.8% of recorded tropical cyclones (**Figure 3B**). For a disruption

that results in prophylactic protection ceasing on March 1st, we estimate 5.3-11.9% of younger

children and 8.0-17.6% of school aged children are infected by day 20 in high FOI sites. This

indicates that the temporary gaps in prophylaxis administration likely in the aftermath of extreme

weather events are a major challenge for control programs aiming to minimize burden (e.g.,

maintain infection below a 1% or 10% threshold) among vulnerable children.

2.3 Incidence results are robust to variation in the force of infection

Our initial estimates of FOI indicate that prevalence rebounds rapidly when drug-based treatment

and prevention coverage ceases (e.g., reaching 10% for school-aged children within 8-92 days by

site). Future changes to malaria transmission dynamics or uncertainty in initial estimation of FOI

may result in variation in the rate of rebound. For example, undercounting infected hosts due to

imperfect RDT sensitivity will result in an underestimation of FOI (**Figure S3**). To account for

this, we evaluate a range of FOI scenarios (**Figure 3C**). Increasing FOI results in prevalence

reaching the threshold prevalence level sooner (e.g., for school aged children, reaching 10%

prevalence 3-31 days sooner for a 50% increase in FOI), demonstrating that rapid rebounds in

prevalence are robust to RDT sensitivity or other sources of underestimation. Alternatively, FOI

could be decreased by reductions in the intensity of transmission achieved by intervention rounds

prior to the disruption event. However, even in optimistic scenarios where earlier rounds of drug

administration substantially reduce transmission rates, prevalence rebounds are only slightly

delayed at high FOI sites (e.g., by 8-9 days for a 50% decrease in FOI for sites MNJ.01 and

MNJ.10). These data suggest chemoprevention-based intervention regimes with currently

available drugs are of limited efficacy in maintaining a low prevalence of infection in the face of

climate-driven disruptions for the FOI ranges likely in moderate to high transmission settings.

Figure 3: Modeled increase in malaria infection due to interrupted drug administration

Figure 3A: Schematic for estimating the risk of malaria infection over time as a function of the force of infection (FOI) and the rate of decay of prophylactic protection. For example, an interruption to the re-administration of a short and long half-life chemoprevention drug is caused by a tropical cyclone (image source: NASA, Cyclone Batsirai, February 2, 2022).

Figure 3B: Left: Calendar date of recorded tropical storms impacting Madagascar since 1980. Right: The cumulative probability of infection (i.e., the proportion of the susceptible population expected to be infected) for a 31-day interval beginning on that day, shown for school aged children (13 years of age). Differences among sites are shown as rings.

Figure 3C: Time until infection reaches a target prevalence versus the force of infection (FOI). A threshold of 10% prevalence is chosen as an example, corresponding to the WHO-defined upper limit for a low malaria burden setting. Baseline observed FOIs for March for the ten sampling localities in Mananjary district (shown with vertical dotted line) are increased or decreased by 0-99% for each site for four exemplar age groups. For reference, the dotted horizontal line shows 90 days (approx. 3 months), with the area under the horizontal line (outlined in red) corresponding to scenarios where the FOI is such that an interruption in access to chemoprevention of that duration results in infection exceeding the target threshold.

3. Vaccination under climate-driven disruption to control

3.1 Modeling vaccination

An exciting shift for malaria control is the recent availability of anti-malarial vaccines. Uniquely, vaccines have a substantially longer duration of protection (i.e., > 10 months (*49, 50*)) than currently used chemoprophylaxis drugs and the time interval relevant for seasonal, climate-driven disruptions such as tropical cyclones (i.e., 3-4 months). Additionally, vaccines using new technology such as mRNA may improve upon the efficacy against clinical disease observed so far for the recently authorized RTS,S (50-56%) (*49*) and R21 (68-75%) (*50*) vaccines. To evaluate the potential of vaccines to reduce malaria burden under climate disruptions, we quantify the reduction in burden under a range of scenarios of vaccine coverage, efficacy, and symptomatic rates of infection (see methods) (**Figure 4A**). Symptomatic rates are obtained from our cohort study where we find 37.0-53.2% of infections among children are associated with observable symptoms (**Figure S4**), in line with previous estimates of asymptomatic infection rates in moderate to high transmission settings (without vaccination) (*51*).

Figure 4: Symptomatic malaria infection probabilities during disruption under a model of vaccination

Figure 4A: Probability tree of the outcomes of malaria infection under vaccination. Parameters on branches specify the probability of either path at branching points. In the simplest analysis, the probability of symptomatic malaria infection for an age group over a time period t is determined by the probability of infection, $p(I) = 1 - e^{-\lambda t}$ where λ is the force of infection, the vaccination coverage, v, the probability of symptoms given infection, s, and any protective effect of vaccination, E. Vaccines may have efficacy against infection, E_i, or against symptoms, E_s. Note that when E_i is small, as is thought to be the case for currently available vaccines, the probability of infection is similar for vaccinated, V, and unvaccinated, V', individuals. When E_s is large, vaccinated individuals are unlikely to have symptoms upon infection.

Figure 4B: The percent reduction in the number of expected symptomatic infections across a range of vaccine efficacies. Without vaccination, the baseline probability of a symptomatic infection is $s \cdot p(I)$. Assuming $E_i = 0$, the percent reduction in symptomatic infections due to vaccination can be expressed as $(s \cdot p(I) - (S_v + S_{v'})) / (s \cdot p(I))$ which simplifies to $v \cdot E_s$. Estimates for the efficacy of the two currently available anti-malaria vaccines, RTS,S (*49*) and R21 (*50*), are shown for example. The shaded quadrant shows the scenarios with sufficient efficacy and coverage to observe a 50% reduction in symptomatic infections.

Figure 4C: For the targeted subpopulation (i.e., children), the number of symptomatic malaria infections averted by

vaccination with efficacy against symptoms E_S. Rows show results for four levels of infection rates, expressed as the

cumulative proportion of the population expected to be infected over the time interval considered. Columns show

results for three values for the probability with which infections become symptomatic.

3.2 Vaccination has the potential to reduce malaria burden following extreme weather events

When 70% of the targeted population has completed the full course for a vaccine with efficacy

68-75% (that reported for the R21 vaccine in phase 3 trials (*50*)) prior to the disruptive event, a

47.6-52.5% reduction in the expected proportion of symptomatic infections is expected (**Figure**

4B). This level of vaccine coverage is plausible given the coverage (approx. 70%) reported for

routine childhood immunization in the country (*52*, *53*). Existing estimates of vaccine efficacy

are obtained on high background rates of access to and use of bednets and other vector control

interventions (*54*). Estimates of bednet usage in Madagascar are 70-80% on the east coast of

Madagascar (*55*) and at our study sites we estimate usage was 72.6-82.6% before and 69.4-

76.4% after the two cyclone events (**Figure S5**). The number of cases averted is dependent on

the infection rate and the symptomatic rate. Using the observed parameter values for the highest

force of infection site (MNJ.10) and 50-70% coverage with the R21 vaccine as an example, we

estimate, per 1000 children, a decrease from 135.1-193.5 symptomatic infections in the 2 months

following a February cyclone to 64.1-101.4 symptomatic infections under vaccination (with

45.9-101.6 cases averted, respectively) (**Figure 4C**).

3.3 Vaccination results are robust to modest efficacy and coverage

Vaccine completeness or timeliness may be imperfect for a fraction of the population, for example due to the timing of a child's birth relative to planned vaccination dates or missed vaccination rounds. Additionally, efficacy may wane in subsequent years (*50*). To account for this variation in the degree of protection we explore a range of proportions of partially protected individuals (**Figure S6**). We also explore a range of symptomatic infection probabilities, including scenarios where symptomatic infection becomes more probable as infection derived immunity declines under lower incidence (**Figure S7**). We find vaccination can lead to a substantial reduction (i.e., approx. 50%) in the frequency of symptomatic malaria infection during disaster aftermath, results that are robust to even moderate vaccine efficacy and reasonable vaccination coverage (**Figure 4B-C**).

Discussion

1. Temporal gap-filling as a potential use case for malaria vaccination

Leveraging a unique dataset from Madagascar, we demonstrate that discontinuities in protection against malaria are a critical threat to efforts to reduce burden. Extreme weather events are a challenge to maintaining continuity. Climate change amplifies the risk of some extreme weather events (*56*), including tropical cyclones (*11*), with potentially devastating consequences relevant for health (*18*). The role of the disruptions likely to result from climate-driven disasters, and the best options for response, have been neglected in the study of climate change and malaria. Vaccination, recently available for malaria, has the unique feature of a protective effect sufficient in duration to persist through disruptive events, such as cyclones and their aftermath. To date, assessments of the cost-effectiveness of vaccination, and priority areas for vaccine deployment, do not account for the additional benefit of long lasting protection, even when moderate, against disease in times of disruption. Our findings suggest a new use case for vaccines, the deployment of vaccination in areas vulnerable to climate-driven or other disruption.

2. Limitations: Diagnostic uncertainty from reliance on RDTs and moderate vaccine efficacy

Ideally, estimates of infection rate would be determined with highly sensitive diagnostics (e.g., PCR) (*51*, *57*). Here, as in many settings with limited access to advanced health infrastructure, diagnostic options were constrained to field-deployable, rapid diagnostics (e.g., RDTs). We developed a set of sensitivity analyses to allow inference under the expected diagnostic uncertainty, with general applicability to other settings reliant on RDTs for surveillance. While RDTs are likely to miss low parasitemia infections (*58*), the extent to which these contribute to clinical burden, the target for vaccination, is unclear (*41*, *59*). Additionally, as described above, undercounting these infections minimally alters the magnitude of timing of infection rebound relevant for disaster response.

As malaria control programs prepare to scale up interventions to meet ambitious targets in coming years, investment across multiple control options must be evaluated (*60*). Benefits from covering any temporal gaps in drug administration present an important case for considering vaccination, even given current shortcomings. Limited vaccine efficacy and questions as to cost-effectiveness across varied settings (*61–63*), alongside their minimal transmission blocking activity, and restricted age range for which data are available (e.g., children < 4 years) mean that vaccines are unlikely to be deployed alone. Efforts that combine prevention approaches with different kinetics, even though they may appear redundant during periods without disturbance, are likely needed to mitigate the impact of extreme weather events. Such approaches could include reinforcement of vector control interventions such as bednets during periods of population displacement (*64–66*), strengthening health facilities to minimize damage (*67*, *68*), and creative drug distribution approaches for prophylaxis that avoid dependence on storm-vulnerable infrastructure (for example, for intermittent preventive treatment during pregnancy, IPTp).

3. The need for more research on attribution and disaggregating mechanisms of risk

A limitation of our study is reliance on estimates of aggregate risk of infection following a climate event. A challenging but important task for future investigations will be to separate the proportion of risk attributable to background and seasonal trends from the direct and indirect impacts of an extreme weather event. We hypothesize tropical cyclones alter the risk of infection through multiple mechanisms such as flooding, population displacement, and destruction of housing and clinics (*18*, *21*, *69*), that interact to modify mosquito vector abundance, host exposure to infection, and access to and use of prevention and treatment. We find the overall magnitude of infection to be high, which will dictate the burden experienced during disaster aftermath and the likely response to control measure scenarios. Obtaining further resolution, however, is limited by difficulty in deriving expectations for a non-cyclone impacted year for the study area. This is due to the scarcity of long-term data with direct estimates of malaria infection rates and the high frequency of cyclones. Both cyclone seasons of the present study (2022 and 2023) featured major storms and, at the national scale, Madagascar has experienced 48 cyclones with intensity at or greater than a category 1 hurricane in the 44 years since 1980. Additionally, storms will interact with multi-year changes in underlying drivers of malaria risk, such as periodic mass bednet campaigns, and any residual damage to infrastructure from previous years' storms. Despite these complexities around identifying a storm's possible impact on transmission potential, critically, any decline in the continuity of access to healthcare and the deployment of control measures will increase population exposure to transmission. As a result, data on total risk of infection following a climate event, even when the fraction attributable to the event is low or uncertain, will be valuable for planning and should be a priority for future surveillance efforts.

4. Implications for disease control in an era of increasingly extreme weather

The overwhelming majority of global malaria cases and deaths come from a subset of African countries (e.g., 20 African countries contribute 86% of global cases) (*1*). The context of stalled

progress in control efforts in these high burden settings, the broad effects of climate change, new
recommendations for expanded chemoprevention, and the advent of vaccination create a new
environment for malaria control. Health system and infrastructural disruptions in these high
burden areas could be a major consequence of climate change for malaria. Continuity in public
health efforts will be decisive for global efforts to reduce the burden of this pathogen. Severe
tropical cyclones, a source of likely disturbance, have marked seasonality, making the timing of
their impact predictable and enabling anticipatory interventions such as vaccination. Other forms
of disturbance (e.g., civil conflict (*70*, *71*)) may be less predictable in their timing, but could
benefit from similar analyses. In addition to malaria, other infectious diseases subject to control
regimes reliant on the regular distribution of drugs or vaccination may be vulnerable to climate-
driven disruption (*23*, *72*). This warrants increased attention on the interaction between climate
and the continuity of public health access. Our data suggest, in the malaria and tropical cyclones
case, vaccination – less sensitive to short term disruption – will be a key mitigation measure.

Acknowledgements

We thank the CRS Multisectoral Malaria Project staff for support and the CRS Madagascar office and Mananjary malaria project staff for assistance with data collection: Dr. Virginie Andreas Nambinina Ralisoa, Dr. Elanirina Andrianoelivololona, Elodie Mialinjaka Rabarijaona, Tahiniony Tonie Ludjette Rakotozafy, Johnson Rakotohasimbola, Juloce Lidovique Razafindrakoto, Hermann Raelison Paratoaly, Aymardine Flashe Kantomananjara, Simon Razafindrasoja, Arisoa Beatrice Marie Sandrine Niry, and James Hazen. We thank the High Meadows Environmental Institute (HMEI) at Princeton University for support.

Funding:

CRS SCP4 Multisectoral Malaria Project (BLR, SM, MR)

The Falcon Award for Disease Elimination - The Climate Edit by the Global Institute for Disease Elimination (GLIDE) (BLR, MR)

The Ren Che Foundation and the AWS Impact Computing Project at the Harvard Data Science Initiative (HDSI) (CDG)

The High Meadows Environmental Institute (HMEI) at Princeton University (BLR, GAV, CJEM)

Author contributions:

Conceptualization: BLR, MR, JL, BG, CDG, AW, BG, CJEM

Methodology: BLR, BG, AW, BG, CJEM

Investigation: BLR, ER, SM, MR, HR

Visualization: BLR, BG, CJEM

Funding acquisition: BLR, MR, JL, CDG

Project administration: BLR, MR, JL

Writing – original draft: BLR, BG, CJEM

Writing – review & editing: BLR, ER, JL, GAV, AW, BG, CDG, CJEM

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials. Code and date files are available at a github repository: https://github.com/bennyrice/mananjary_cyclone_code

Supplementary Materials

Materials and Methods

1. Study area: Mananjary district, southeastern Madagascar

Located in southeastern Madagascar, the Mananjary district (region Vatovavy) is a coastal district with perennial malaria transmission. The district extends approximately 60 km to the north and south of the district capital, the city of Mananjary (-21.22 S, 48.35 E) where the only hospitals in the district are located. As of 2018, 90.1% of the population of the district lives in rural *kaomina* ("communes") (Institut National de la Statistique Madagascar).

From Malaria Indicator Surveys, malaria prevalence among children 5 years or younger in the east coast of Madagascar averaged 9-18% from 2011 to 2021 (*55, 73–75*), placing the district in the WHO defined moderate transmission stratum. In terms of seasonality, higher prevalence was estimated for December to May from re-analyses of health facility data (*76, 77*), with a peak around April (*78*). Few recent epidemiological studies with active sampling are available from the region, but prevalence among households (all ages) varied from 3-46% by locality in 2017 from cross-sectional prevalence surveys (*44, 79*).

2. Extreme weather event data

Historical data on the occurrence of extreme weather events was sourced from the National Oceanic and Atmospheric Administration (NOAA) International Best Track Archive for Climate Stewardship (IBTrACS) data (accessed 21 February 2024) and EM-DAT from the Centre for Research on the Epidemiology of Disasters (CRED) (accessed 21 February 2024). See **Data S1** for the full list of weather events.

Search criteria were all storms where the eye passed within a 60 nautical mile (approx. 111 km) buffer of Madagascar's coastline. This buffer was chosen because extreme rainfall and winds may extend across the diameter of a storm. Tropical cyclones are defined as systems with maximum sustained wind speeds > 119 km/h, equivalent to a Category 1 Hurricane per the Saffir-Simpson scale used in the Atlantic basin.

3. Longitudinal sampling

3.1 Prospective cohort study

A stratified random sample of 500 households evenly distributed among 10 rural sampling clusters (communities or *tanana*) were recruited and enrolled during a baseline sample in July-October 2021. Informed consent or assent was obtained for study participants prior to enrollment. Enrolled households were followed longitudinally for 10 follow-up sampling time points. Clusters were rural (defined as more than 5km from the Mananjary district capital) and approximately 10 km apart on average.

Study procedures received approval from the Institutional Review Board (IRB) of the Harvard T.H. Chan School of Public Health (IRB#21-0111), Comité d'Ethique de la Recherche Biomédicale (CERBM) at Ministère de la Santé Publique de Madagascar (Nº019/MSANP/SG/AMM/CERBM 2021), the Mananjary district health office, local government authorities, and community leaders.

Figure S1 shows the proportion of the cohort negative, positive, and unsampled by rapid diagnostic test (RDT) per time point. The participation rate, calculated as the percentage of enrolled individuals screened, ranged from 62.5-91.0% per time point. We consider individuals with visible bands for lactate dehydrogenase (LDH) or histidine rich protein 2 (HRP2) antigens as positive. The RDTs used were the AccessBio CareStart™ Pf/Pan RDT or Abbott Diagnostics Bioline™ Malaria Ag Pf/Pan RDT per manufacturer's protocol.

3.2 Screening for malaria and symptoms

All individuals in enrolled households were screened for malaria at approximately 2-month intervals (mean interval between samples = 57.4 days, median = 52 days, n = 20,718 total observations). On site diagnosis by RDT was performed at the time of screening and a blood sample was collected on a dried blood spot (DBS) for later molecular confirmation. Individuals, or a surrogate for younger children, also completed a questionnaire on bednet usage and symptoms adapted from refs. (*80*, *81*). Sensitivity analyses were performed to account for RDT sensitivity and specificity (**Figure S3**).

Individuals positive for malaria by RDT were offered treatment by an onsite physician with artemisinin-based combination therapy (ACT) using the standard first line treatment for Madagascar (artesunate amodiaquine, AS+AQ). To observe treatment adherence, at minimum the first dose of the three-day course of treatment was observed directly by the research team. Symptomatic individuals or individuals requesting a consultation were offered a consultation with an onsite physician.

3.3. Accounting for RDT sensitivity and specificity

Due to imperfect sensitivity and specificity for RDTs, false positive and false negative RDT results may bias inference of infection rates. From a previous study, an estimated 97% of infections in the east coast region of Madagascar are by *Plasmodium falciparum* and 88% of RDT positive infections were confirmed by molecular follow-up (*82*). In studies of diagnostic performance, RDT sensitivity and specificity was estimated to be approximately 80-90% (*83*, *84*).

We use a binomial sampling approach to explore the effect of diagnostic error. We simulate 'false negatives' by drawing from a binomial distribution with probability $1 - sensitivity$ such that a fraction of RDT negatives become positive. Likewise, we simulate 'false positives' by drawing from a binomial distribution with probability equal to the *specificity* such that a fraction of RDT positives become negative. **Figure S3** shows simulations over a range of sensitivity and specificity values, where we repeat the force of infection estimation methods (described in the following section) for 500 replicates. These results illustrate minimal bias across plausible ranges of sensitivity and specificity. When the proportion of positive tests is small, accounting for possible false negatives results in an increase in the expected proportion infected. For sites with a high proportion of positive tests (i.e., approximately 50%), false negatives and false positives are similarly likely such that the expected proportion infected is minimally changed.

4. Estimating the force of infection

The interval before infection probability reaches a given threshold will hinge on target prevalence and the rate at which individuals become reinfected. To evaluate this, we define λ as the force of infection, or rate at which susceptible individuals become infected. The catalytic model (*85*) then defines the probability of evading infection up until time t as:

$$522 \qquad\qquad\qquad\qquad S(t) = exp\,(-\lambda t)$$

523 and the probability of having been infected by time t (or cumulative density function of infection) is

$$524 \qquad\qquad\qquad\qquad F(t) = 1 - exp\,(-\lambda t)$$

For malaria, infected individuals are susceptible to reinfection. The rate at which this occurs within the
time interval of interest will be negligible when the force of infection is low and/or the time interval
considered is short, estimated to be the case here (e.g., within 1-2 months). We consequently proceed with
the estimates based on $F(t)$.

Because no evidence of resistance to ACT antimalarials has been reported to date from Madagascar, we
assume the clearance rate of asexual and sexual stages of the parasite to be high. Given our direct
observation of the early doses of anti-malarial treatment for infected individuals, we assume adherence to
the treatment course is high and successful clearance of parasite infections following treatment. Thus for
individuals positive at t_i and t_{i+1} we assume that a new infection has occurred over the time interval.

The force of infection may vary according to time-invariant covariates such as geographic site, and
individual age (which can be considered approximately time-invariant at least over the time-scales
considered). We can fit these quantities using a generalized linear model fitted with a binomial likelihood
using a complementary log-log link, such that if π_i is the probability of being infected over interval i, the
link function is provided by $\eta_i = log\,(-log\,(1-\pi_i)\,)$. The cumulative density function can then be
expressed as: $F(\eta_i) = 1 - exp\,(-exp\,(\eta_i))$ from which it can be deduced that the link function must
take the form $\eta_i = \beta X_i + log(t)$ where β is a vector of coefficients, and X_i is the corresponding design
matrix of covariates for interval i. Introducing this back into the cumulative density function yields

$$542 \qquad F(t) = 1 - exp\,\bigl(-exp\,\bigl(\beta X_i + log(t)\bigr)\bigr)\, = 1 - exp\,(-(\beta X_i))\,,$$

543 indicating that the force of infection is defined by: (βX_i), as desired (*86*).

544 To additionally capture seasonal fluctuations in the force of infection (by their nature time-varying), we
545 can introduce covariates reflecting the number of days each individual at each measurement was exposed
546 to different seasons of transmission (e.g., days in each month); and we can additionally introduce smooth
547 terms to capture non-linear patterns. The extended force of infection is then provided by:

$$548 \qquad\qquad \bigl(\alpha_{site,i} + s(age_i) + \gamma_1 M_{1,i} + \gamma_2 M_{2,i} + \gamma_3 M_{3,i} + \cdots \bigr)$$

549 Where the parameter α takes a different value for each of the ten study sites; $s(age_i)$ indicates a
550 smoothed pattern across age, $M_{j,i}$ reflects the number of days an individual i was exposed to the j^{th}
551 month of the year, with γ_j the corresponding parameter converting this into effects on the force of
552 infection. Finally, we also include a random effect for household, as there was evidence of consistent
553 effects across households. The final expression for the force of infection is thus:

$$554 \qquad\qquad \bigl(\alpha_{site,i} + s(age_i) + \gamma_1 M_{1,i} + \gamma_2 M_{2,i} + \gamma_3 M_{3,i} + \cdots + H_i \bigr)$$

555 Where H_i corresponds to the random effect mapping the household of individual i.

5. Exploring the consequences of disruptions during intervention rollout

To explore the deployment of new mass chemoprevention and chemotherapy interventions, we simulate
the early phases of these interventions, when the program is building towards, but has not yet, interrupted

transmission. As a result, ongoing transmission results in the reaccumulation of infections, at a rate
determined by the force of infection, following each intervention round until transmission is interrupted
(i.e., force of infection approaches 0). We first investigate the return time, which we define as the
maximum time interval between intervention rounds for a given force of infection where malaria
prevalence is maintained below a chosen target level in the targeted population (see supplementary
methods for full details).

The required program return time Δ_c necessary to maintain prevalence below a chosen threshold z is
defined by:

$$\Delta_c = \frac{-log\,(1-z)}{\lambda}$$

The estimated force of infection captures the rate at which individuals become infected, and interventions
occur at intervals of Δ days, and clear p_t percent of infections. Conservatively, we assume clearance for
treated individuals to be 100% such that treatment successfully reduces infectiousness and treated
individuals do not participate in transmission. This is supported by the lack of evidence of resistance to
ACTs in Madagascar to date and evidence ACTs clear gametocytes (*87*). Subsequent to this intervention,
mosquito vectors infected with parasites prior to the intervention or from contact with subsets of the
population not covered by the initial round serve as a reservoir for infection. Thus 'cleared' hosts start to
become infected again from continued exposure. The longer the interval Δ, the higher the prevalence of
malaria reached within the target population.

We initially make no assumptions about feedbacks associated with transmission, and additionally
disregard recovery without treatment, since time intervals to recovery are typically long. Supporting this,
published estimates of the distribution of duration of untreated *P. falciparum* infections (mean duration
approximately 180 days (*41*, *42*)) indicate that the probability of recovery without treatment between our
longitudinal samples (mean sampling interval 57 days) and during a disruption interval (e.g., 2 months)
are very low.

583 6. Simulating additional control activities: Prophylaxis and vaccination

In addition to a primary intervention where a round of standard first line antimalarials (e.g., ACTs) is
applied to clear existing infections, various interventions that prolong the duration of protection may be
considered, from drug combinations with a longer prophylactic period to vaccination.

6.1 Mass drug administration with longer lasting chemoprevention

Including a longer-lived antimalarial (e.g., sulphadoxine-pyrimethamine, SP, or dihydroartemisinin-
piperaquine, DP) can prevent reinfection for the duration of effective prophylaxis. We define the mean
duration of prophylaxis as *D*. We first consider a mass drug administration (MDA) scenario where all
individuals, regardless of infection status, are given the ACT treatment paired with a prophylactic. When
used to prevent reinfection for clinical cases this has been termed post-discharge malaria prophylaxis
(PDMC). We also consider application in a mass test and treat (MTaT) activity where only individuals
identified as positive for malaria infection receive the chemoprevention (see **Table S1** for definitions).

Because the drugs likely to be used in PDMC, MDA, and MTaT scenarios are the same as those for
seasonal malaria chemoprevention (SMC) and intermittent preventive treatment (IPT), and thus have the
same duration of protection, the return time for all can be modeled as $D + \Delta_c$.

Table S1: World Health Organization assessed drug-based malaria interventions

Intervention	Definition
Mass Drug Administration (MDA)	Presumptive treatment of a large proportion of the population. In the simplest form, performed with a standard first-line anti-malarial (e.g., ACTs in Africa). *Aliases and derivatives*: Equivalent to MTaT when combined with diagnostic screening prior to treatment. Termed targeted drug administration (TDA) when applied to a selected subpopulation and reactive drug administration (RDA) when performed reactively for an observed infection cluster.
Post Discharge Malaria Chemoprevention (PDMC)	Addition of a longer lasting antimalarial (e.g., SP) for a prophylactic effect for treated individuals. *Aliases and derivatives*: If applied proactively for a season or year, equivalent to SMC/PMC/IPT
Mass Test and Treat (MTaT)	Testing, regardless of symptoms, a large proportion of the population and providing treatment for those testing positive. *Aliases and derivatives*: Sometimes termed Mass Screen and Treat (MSaT); targeted test and treat (TTaT) when applied to a selected subpopulation.
Intermittent Preventive Treatment (IPT)	The regular (e.g., monthly) administration of a curative antimalarial dose to curtail infections and prevent new infections. *Aliases and derivatives*: Often targeted to pregnant women (IPTp) and infants (IPTi). Equivalent to SMC/PMC.
Seasonal/Perennial Malaria Chemoprophylaxis (SMC/PMC)	Equivalent to IPT, the use of a full treatment course of antimalarials at regular intervals to reduce the risk of malaria infection. Recently recommended for scale up to more geographic areas and children over 5 *Aliases and derivatives*: SMC when targeted to the season with high transmission, PMC when year-round.

6.2 Vaccination

Vaccination with the currently available antimalarial vaccines may provide a benefit in reducing clinical cases when providing access to treatment or chemoprevention is unlikely. We use estimates of the efficacy of antimalarial vaccines against clinical malaria reported from clinical trials for the RTS,S (approx. 55%) (*49*) and R21 vaccines (approx. 75%) (*54, 88*).

In the simplest analysis, the expected number of clinical cases results from the rate of exposure, λ, giving the probability of being infected over an interval, and the probability of infection being asymptomatic (*a*) or symptomatic ($s = 1 - a$). Given no data on efficacy against infection (E_i) for currently approved vaccines, we conservatively assume $E_i = 0$. For a proportion, v, of children fully vaccinated at the beginning of cyclone season, with a vaccine of efficacy of E_s (defined as reduction in probability of symptomatic malaria), we estimate the probability of symptomatic infection among children over an interval of length t following cessation of intervention due to a disruption:

$$\Pr(\text{symptomatic infection}) = v \times (1 - e^{-\lambda t}) \times (1 - E_i) \times s \times (1 - E_s) + (1 - v) \times (1 - e^{-\lambda t}) \times s$$

See **Table S2** for parameter definitions and estimates.

617 **Table S2. Parameter definitions used in vaccination modeling**

Parameter	Definition	Range
$1 - e^{-\lambda t}$	Proportion of uninfected individuals that test positive during interval t	0.01-1.0
v	Vaccination coverage	0-100%
s	Proportion of infected individuals reporting symptoms	0.1-1.0
E_s	Efficacy against symptomatic infection for vaccination	0-100%

Supporting Information

Figure S1. Sampling for the Mananjary malaria cohort study

1A: The proportion of enrolled individuals unsampled, negative, or positive by rapid diagnostic test (RDT) per time point from baseline (T0) to final sample (T10) (all sites combined). **1B**: Distribution of the number of observations per individual (site code colors and order follow Figure 2C). **1C**: Distribution of the interval in days between samples.

643   **Figure S2. Force of infection (FOI) by age**

644

645   Observed variation in the force of infection by age for the 10 sample sites in Mananjary district, Madagascar (site

646   code colors and order follow Figure 2C). Data shown for the month of January.

647

648

649

650

651

652

653

654

655

656

657

658

659

660

Figure S3. Accounting for imperfect RDT diagnostic accuracy

From 500 simulated trials, the force of infection (FOI) per day for a given sensitivity and specificity. Sites are ordered from lowest to highest observed rate of infection. Dashed vertical lines show the mean FOI from 500 simulations. For low infection sites (e.g., Sites MNJ.06 and MNJ.04, shown at top left), reduced sensitivity increases estimates of FOI (e.g., FOI increases from an initial estimate 6.3×10^{-4} to a mean estimate of 2.7×10^{-3} for sensitivity and specificity = 90%). This indicates for low infection rate sites, our estimates for the probability of infection over a time interval are conservative, with false negative rapid diagnostic tests contributing to a higher probability of infection. For high infection sites (e.g., Sites MNJ.01 and MNJ.10, shown at bottom right) estimates of FOI are relatively invariant across the range of sensitivity and specificity values explored. False negatives and false positives largely offset such that the initial FOI estimate 7.19×10^{-3} is similar to the mean estimate (7.16×10^{-3}) from the 500 simulated trials with sensitivity and specificity at 90%.

684   **Figure S4. Symptomatic rates of infection by survey recall**

686   The percent of rapid diagnostic test (RDT) positive individuals reporting any fever within the last two weeks of
687   screening. Error bars show the 95% confidence intervals using the Clopper and Pearson (1934) method as
688   implemented in the R binom.test function.

697   **Figure S5. Bednet usage by survey recall**

699   The proportion of individuals reporting sleeping under a bednet in the last 24 hours and if reporting bednet was
700   damaged (due to holes, tears, or other damage). To the right, change in the proportion of individuals reporting not
701   using a bednet for sampling time points before and after two cyclones. Error bars show the 95% confidence intervals
702   using the Clopper and Pearson (1934) method as implemented in the R binom.test function.

715 **Figure S6. Simulating partial vaccination**

Top: Probability tree of the outcomes of malaria infection under partial vaccination. From the probability tree in
Figure 4A, the vaccinated branch can be separated into partially and fully vaccinated sub-branches. Shown here
assuming protection against infection, $E_i = 0$, and protection against symptoms for fully vaccinated individuals E_s is
greater than the mean protection against symptoms for partially vaccinated individuals $E_{partial}$. Bottom: The percent
reduction in the number of expected symptomatic infections across a range of partial vaccination rates with partial
vaccine efficacy. Two example scenarios are shown where 50% of the targeted population is fully vaccinated and 0-
50% of the population is partially vaccinated with protection varying from 0 to equaling the efficacy seen in fully
vaccinated individuals (E_s). The shaded area shows the scenarios with sufficient efficacy and coverage to observe a
50% reduction in symptomatic infections.

Figure S7. Sensitivity analyses for vaccination parameters

For the targeted subpopulation (i.e., children), the number of symptomatic malaria infections averted by vaccination with efficacy against symptoms E_S. Rows show results for four levels of infection rates, expressed as the cumulative proportion of the population expected to be infected over the time interval considered. Columns show results for three values for the probability with which infections become symptomatic.

Data S1

Tropical cyclones and related humanitarian disasters in Madagascar since 1980 (xlsx)

Data S2

Locality information for study sites in Mananjary district, Madagascar including site codes, administrative division names, and coordinates of midpoints (xslx)

Data S3

Line list data of individual rapid diagnostic test (RDT) results and sample dates (csv)

Data S4

Tropical cyclone tracks data from IBTrACS (csv)

Accessed from: https://www.ncei.noaa.gov/data/international-best-track-archive-for-climate-stewardship-ibtracs/v04r00/access/csv/

Data S5

Global estimates of average annual tropical cyclone exposure from Jing et al 2023 (*39*) (csv)

Data S6

Spatial estimates of *Plasmodium falciparum* incidence rate for 2020 from the Malaria Atlas Project (tif)

Accessed from: https://data.malariaatlas.org/

Data S7

National level estimated incidence of malaria from the World Health Organization (xlsx)

Data S8

Bednet usage questionnaire response data for the Mananjary cohort study (csv)

Data S9

Symptoms questionnaire response data for the Mananjary cohort study (csv)

References and Notes:

1. "World Malaria Report 2023" (Global Malaria Programme, World Health Organization, 2023); https://www.who.int/publications/i/item/9789240086173.

2. J. R. Poespoprodjo, N. M. Douglas, D. Ansong, S. Kho, N. M. Anstey, Malaria. *Lancet* **402**, 2328–2345 (2023).

3. E. A. Mordecai, J. M. Caldwell, M. K. Grossman, C. A. Lippi, L. R. Johnson, M. Neira, J. R. Rohr, S. J. Ryan, V. Savage, M. S. Shocket, R. Sippy, A. M. Stewart Ibarra, M. B. Thomas, O. Villena, Thermal biology of mosquito-borne disease. *Ecol. Lett.* **22**, 1690–1708 (2019).

4. O. C. Villena, S. J. Ryan, C. C. Murdock, L. R. Johnson, Temperature impacts the environmental suitability for malaria transmission by Anopheles gambiae and Anopheles stephensi. *Ecology* **103**, e3685 (2022).

5. W. M. de Souza, S. C. Weaver, Effects of climate change and human activities on vector-borne diseases. *Nat. Rev. Microbiol.*, doi: 10.1038/s41579-024-01026-0 (2024).

6. E. A. Mordecai, K. P. Paaijmans, L. R. Johnson, C. Balzer, T. Ben-Horin, E. de Moor, A. McNally, S. Pawar, S. J. Ryan, T. C. Smith, K. D. Lafferty, Optimal temperature for malaria transmission is dramatically lower than previously predicted. *Ecol. Lett.* **16**, 22–30 (2013).

7. K. P. Paaijmans, A. F. Read, M. B. Thomas, Understanding the link between malaria risk and climate. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 13844–13849 (2009).

8. S. J. Ryan, C. A. Lippi, F. Zermoglio, Shifting transmission risk for malaria in Africa with climate change: a framework for planning and intervention. *Malar. J.* **19**, 170 (2020).

9. A. S. Siraj, M. Santos-Vega, M. J. Bouma, D. Yadeta, D. Ruiz Carrascal, M. Pascual, Altitudinal changes in malaria incidence in highlands of Ethiopia and Colombia. *Science* **343**, 1154–1158 (2014).

10. C. Caminade, S. Kovats, J. Rocklov, A. M. Tompkins, A. P. Morse, F. J. Colón-González, H. Stenlund, P. Martens, S. J. Lloyd, Impact of climate change on global malaria distribution. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 3286–3291 (2014).

11. K. T. Bhatia, G. A. Vecchi, T. R. Knutson, H. Murakami, J. Kossin, K. W. Dixon, C. E. Whitlock, Recent increases in tropical cyclone intensification rates. *Nat. Commun.* **10**, 635 (2019).

12. J. B. Elsner, J. P. Kossin, T. H. Jagger, The increasing intensity of the strongest tropical cyclones. *Nature* **455**, 92–95 (2008).

13. L. Li, P. Chakraborty, Slower decay of landfalling hurricanes in a warming world. *Nature* **587**, 230–234 (2020).

14. S. Wang, R. Toumi, Recent migration of tropical cyclones toward coasts. *Science* **371**, 514–517 (2021).

15. A. M. D. Ortiz, P. L. C. Chua, D. Salvador Jr, C. Dyngeland, J. D. G. Albao Jr, R. A. Abesamis, Impacts of tropical cyclones on food security, health and biodiversity. *Bull. World*

821   *Health Organ.* **101**, 152–154 (2023).

16.   J. M. Shultz, J. Russell, Z. Espinel, Epidemiology of tropical cyclones: the dynamics of disaster, disease, and development. *Epidemiol. Rev.* **27**, 21–35 (2005).

17.   R. M. Parks, G. B. Anderson, R. C. Nethery, A. Navas-Acien, F. Dominici, M.-A. Kioumourtzoglou, Tropical cyclone exposure is associated with increased hospitalization rates in older adults. *Nat. Commun.* **12**, 1545 (2021).

18.   T. Alcayna, I. Fletcher, R. Gibb, L. Tremblay, S. Funk, B. Rao, R. Lowe, Climate-sensitive disease outbreaks in the aftermath of extreme climatic events: A scoping review. *One Earth* **5**, 336–350 (2022).

19.   K. L. Ebi, J. Vanos, J. W. Baldwin, J. E. Bell, D. M. Hondula, N. A. Errett, K. Hayes, C. E. Reid, S. Saha, J. Spector, P. Berry, Extreme Weather and Climate Change: Population Health and Health System Implications. *Annu. Rev. Public Health* **42**, 293–315 (2021).

20.   H. Nissan, I. Ukawuba, M. Thomson, Climate-proofing a malaria eradication strategy. *Malar. J.* **20**, 190 (2021).

21.   K. M. Searle, D. E. Earland, A. Francisco Bibe, A. Novela, V. Muhiro, J. L. Ferrão, Long-lasting household damage from Cyclone Idai increases malaria risk in rural western Mozambique. *Sci. Rep.* **13**, 21590 (2023).

22.   S. Devi, Cyclone Idai: 1 month later, devastation persists. *Lancet* **393**, 1585 (2019).

23.   A. Pozniak, A. Atzori, C. Marotta, F. Di Gennaro, G. Putoto, HIV continuity of care after Cyclone Idai in Mozambique. *Lancet HIV* **7**, e159–e160 (2020).

24.   Q. Fernandes, O. Augusto, S. Chicumbe, L. Anselmi, B. H. Wagenaar, R. Marlene, S. Agostinho, S. Gimbel, J. Pfeiffer, C. Inguane, D. M. Uetela, J. Crocker, I. Ramiro, B. Matsinhe, S. Tembe, N. Carimo, S. Gloyd, I. Manhiça, E. Tavede, P. Felimone, K. Sherr, Maternal and Child Health Care Service Disruptions and Recovery in Mozambique After Cyclone Idai: An Uncontrolled Interrupted Time Series Analysis. *Glob Health Sci Pract* **10** (2022).

25.   J. E. Coalson, E. J. Anderson, E. M. Santos, V. Madera Garcia, J. K. Romine, B. Dominguez, D. M. Richard, A. C. Little, M. H. Hayden, K. C. Ernst, The Complex Epidemiological Relationship between Flooding Events and Human Outbreaks of Mosquito-Borne Diseases: A Scoping Review. *Environ. Health Perspect.* **129**, 96002 (2021).

26.   J. T. Watson, M. Gayer, M. A. Connolly, Epidemics after natural disasters. *Emerg. Infect. Dis.* **13**, 1–5 (2007).

27.   IPCC, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021).

28.   J. P. Kossin, K. R. Knapp, T. L. Olander, C. S. Velden, Global increase in major tropical cyclone exceedance probability over the past four decades. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 11975–11980 (2020).

29.   "Cyclone Freddy deepens health risks in worst-hit countries" (World Health Organization

(WHO) African Region, 2023); https://www.afro.who.int/news/cyclone-freddy-deepens-
health-risks-worst-hit-countries.

30. "Mobilizing health assistance after deadly cyclones devastate Madagascar" (World Health
Organization (WHO) African Region, 2022); https://www.afro.who.int/photo-story/mobilizing-
health-assistance-after-deadly-cyclones-devastate-madagascar.

31. "In Madagascar, mobile clinics bolster health surveillance during cyclones" (World Health
Organization (WHO) Regional Office for Africa, 2023);
https://www.afro.who.int/countries/madagascar/news/madagascar-mobile-clinics-bolster-
health-surveillance-during-cyclones.

32. "Outbreaks and emergencies bulletin, week 12: 13 - 19 march 2023: Cyclone Freddy in the
WHO African Region" (12, World Health Organization (WHO) African Region, 2023);
https://www.afro.who.int/countries/togo/publication/outbreaks-and-emergencies-bulletin-
week-12-13-19-march-2023.

33. D. Delforge, V. Wathelet, R. Below, C. L. Sofia, M. Tonnelier, J. van Loenhout, N.
Speybroeck, EM-DAT: The Emergency Events Database, *Research Square* (2023).
https://doi.org/10.21203/rs.3.rs-3807553/v1.

34. FEL Otto, M Zachariah, P Wolski, I Pinto, R Barimalala, B Nhamtumbo, R Bonnet, R
Vautard, S Philip, S Kew, LN Luu, D Heinrich, M Vahlberg, R Singh, J Arrighi, L
Thalheimer, Mv Aalst, S Li, J Sun, G Vecchi, LJ Harrington, "Climate change increased
rainfall associated with tropical cyclones hitting highly vulnerable communities in
Madagascar, Mozambique & Malawi" (World Weather Attribution (WWA) Initiative, 2022);
https://www.worldweatherattribution.org/climate-change-increased-rainfall-associated-with-
tropical-cyclones-hitting-highly-vulnerable-communities-in-madagascar-mozambique-
malawi/.

35. K. R. Knapp, M. C. Kruk, D. H. Levinson, H. J. Diamond, C. J. Neumann, The International
Best Track Archive for Climate Stewardship (IBTrACS): Unifying Tropical Cyclone Data.
Bull. Am. Meteorol. Soc. **91**, 363–376 (2010).

36. K. R. Knapp, H. J. Diamond, J. P. Kossin, M. C. Kruk, C. J. Schreck, International best
track archive for climate stewardship (IBTrACS) project, version 4, NOAA National Centers
for Environmental Information (2018); https://doi.org/10.25921/82TY-9E16.

37. D. A. Pfeffer, T. C. D. Lucas, D. May, J. Harris, J. Rozier, K. A. Twohig, U. Dalrymple, C. A.
Guerra, C. L. Moyes, M. Thorn, M. Nguyen, S. Bhatt, E. Cameron, D. J. Weiss, R. E.
Howes, K. E. Battle, H. S. Gibson, P. W. Gething, malariaAtlas: an R interface to global
malariometric data hosted by the Malaria Atlas Project. *Malar. J.* **17**, 352 (2018).

38. C. P. T. G. Zehr, *Biscale: Tools and Palettes for Bivariate Thematic Mapping R Package*
(2022; https://chris-prener.github.io/biscale/authors.html).

39. R. Jing, S. Heft-Neal, D. R. Chavas, M. Griswold, Z. Wang, A. Clark-Ginsberg, D. Guha-
Sapir, E. Bendavid, Z. Wagner, Global population profile of tropical cyclone exposure from
2002 to 2019. *Nature* **626**, 549–554 (2024).

40. The Global Health Observatory, World Health Organization, Malaria incidence per 1 000
population at risk per year (estimated) - By country, World Health Organization (2023);

https://apps.who.int/gho/data/node.main.MALARIAESTINCIDENCE?lang=en.

41. H. C. Slater, A. Ross, I. Felger, N. E. Hofmann, L. Robinson, J. Cook, B. P. Gonçalves, A.
Björkman, A. L. Ouedraogo, U. Morris, M. Msellem, C. Koepfli, I. Mueller, F. Tadesse, E.
Gadisa, S. Das, G. Domingo, M. Kapulu, J. Midega, S. Owusu-Agyei, C. Nabet, R.
Piarroux, O. Doumbo, S. N. Doumbo, K. Koram, N. Lucchi, V. Udhayakumar, J. Mosha, A.
Tiono, D. Chandramohan, R. Gosling, F. Mwingira, R. Sauerwein, R. Paul, E. M. Riley, N.
J. White, F. Nosten, M. Imwong, T. Bousema, C. Drakeley, L. C. Okell, The temporal
dynamics and infectiousness of subpatent Plasmodium falciparum infections in relation to
parasite density. *Nat. Commun.* **10**, 1433 (2019).

42. E. A. Ashley, N. J. White, The duration of Plasmodium falciparum infections. *Malar. J.* **13**,
500 (2014).

43. European Space Agency, Sinergise, OpenTopography - Copernicus global digital elevation
models, (2021); https://doi.org/10.5069/G9028PQB.

44. B. L. Rice, C. D. Golden, H. J. Randriamady, A. A. N. A. Rakotomalala, M. A. Vonona, E. J.
G. Anjaranirina, J. Hazen, M. C. Castro, C. J. E. Metcalf, D. L. Hartl, Fine-scale variation in
malaria prevalence across ecological regions in Madagascar: a cross-sectional study. *BMC*
Public Health **21**, 1018 (2021).

45. "U.S. President's Malaria Initiative Madagascar Malaria Operational Plan FY 2023"
(President's Malaria Initiative, 2023); https://www.pmi.gov/resources/malaria-operational-
plans-mops/.

46. M. T. Bretscher, P. Dahal, J. Griffin, K. Stepniewska, Q. Bassat, E. Baudin, U.
D'Alessandro, A. A. Djimde, G. Dorsey, E. Espié, B. Fofana, R. González, E. Juma, C.
Karema, E. Lasry, B. Lell, N. Lima, C. Menéndez, G. Mombo-Ngoma, C. Moreira, F.
Nikiema, J. B. Ouédraogo, S. G. Staedke, H. Tinto, I. Valea, A. Yeka, A. C. Ghani, P. J.
Guerin, L. C. Okell, The duration of chemoprophylaxis against malaria after treatment with
artesunate-amodiaquine and artemether-lumefantrine and the effects of pfmdr1 86Y and
pfcrt 76T: a meta-analysis of individual patient data. *BMC Med.* **18**, 47 (2020).

47. M. Cairns, I. Carneiro, P. Milligan, S. Owusu-Agyei, T. Awine, R. Gosling, B. Greenwood,
D. Chandramohan, Duration of protection against malaria and anaemia provided by
intermittent preventive treatment in infants in Navrongo, Ghana. *PLoS One* **3**, e2227
(2008).

48. M. Cairns, A. Barry, I. Zongo, I. Sagara, S. R. Yerbanga, M. Diarra, C. Zoungrana, D.
Issiaka, A. A. Sienou, A. Tapily, K. Sanogo, M. Kaya, S. Traore, K. Diarra, H. Yalcouye, Y.
Sidibe, A. Haro, I. Thera, P. Snell, J. Grant, H. Tinto, P. Milligan, D. Chandramohan, B.
Greenwood, A. Dicko, J. B. Ouedraogo, The duration of protection against clinical malaria
provided by the combination of seasonal RTS,S/AS01E vaccination and seasonal malaria
chemoprevention versus either intervention given alone. *BMC Med.* **20**, 352 (2022).

49. RTS,S Clinical Trials Partnership, S. T. Agnandji, B. Lell, S. S. Soulanoudjingar, J. F.
Fernandes, B. P. Abossolo, C. Conzelmann, B. G. N. O. Methogo, Y. Doucka, A. Flamen,
B. Mordmüller, S. Issifou, P. G. Kremsner, J. Sacarlal, P. Aide, M. Lanaspa, J. J. Aponte, A.
Nhamuave, D. Quelhas, Q. Bassat, S. Mandjate, E. Macete, P. Alonso, S. Abdulla, N.
Salim, O. Juma, M. Shomari, K. Shubis, F. Machera, A. S. Hamad, R. Minja, A. Mtoro, A.
Sykes, S. Ahmed, A. M. Urassa, A. M. Ali, G. Mwangoka, M. Tanner, H. Tinto, U.

D'Alessandro, H. Sorgho, I. Valea, M. C. Tahita, W. Kaboré, S. Ouédraogo, Y. Sandrine, R. T. Guiguemdé, J. B. Ouédraogo, M. J. Hamel, S. Kariuki, C. Odero, M. Oneko, K. Otieno, N. Awino, J. Omoto, J. Williamson, V. Muturi-Kioi, K. F. Laserson, L. Slutsker, W. Otieno, L. Otieno, O. Nekoye, S. Gondi, A. Otieno, B. Ogutu, R. Wasuna, V. Owira, D. Jones, A. A. Onyango, P. Njuguna, R. Chilengi, P. Akoo, C. Kerubo, J. Gitaka, C. Maingi, T. Lang, A. Olotu, B. Tsofa, P. Bejon, N. Peshu, K. Marsh, S. Owusu-Agyei, K. P. Asante, K. Osei-Kwakye, O. Boahen, S. Ayamba, K. Kayan, R. Owusu-Ofori, D. Dosoo, I. Asante, G. Adjei, G. Adjei, D. Chandramohan, B. Greenwood, J. Lusingu, S. Gesase, A. Malabeja, O. Abdul, H. Kilavo, C. Mahende, E. Liheluka, M. Lemnge, T. Theander, C. Drakeley, D. Ansong, T. Agbenyega, S. Adjei, H. O. Boateng, T. Rettig, J. Bawa, J. Sylverken, D. Sambian, A. Agyekum, L. Owusu, F. Martinson, I. Hoffman, T. Mvalo, P. Kamthunzi, R. Nkomo, A. Msika, A. Jumbe, N. Chome, D. Nyakuipa, J. Chintedza, W. R. Ballou, M. Bruls, J. Cohen, Y. Guerra, E. Jongert, D. Lapierre, A. Leach, M. Lievens, O. Ofori-Anyinam, J. Vekemans, T. Carter, D. Leboulleux, C. Loucq, A. Radford, B. Savarese, D. Schellenberg, M. Sillman, P. Vansadia, First results of phase 3 trial of RTS,S/AS01 malaria vaccine in African children. *N. Engl. J. Med.* **365**, 1863–1875 (2011).

50. M. S. Datoo, A. Dicko, H. Tinto, J.-B. Ouédraogo, M. Hamaluba, A. Olotu, E. Beaumont, F. Ramos Lopez, H. M. Natama, S. Weston, M. Chemba, Y. D. Compaore, D. Issiaka, D. Salou, A. M. Some, S. Omenda, A. Lawrie, P. Bejon, H. Rao, D. Chandramohan, R. Roberts, S. Bharati, L. Stockdale, S. Gairola, B. M. Greenwood, K. J. Ewer, J. Bradley, P. S. Kulkarni, U. Shaligram, A. V. S. Hill, R21/Matrix-M Phase 3 Trial Group, Safety and efficacy of malaria vaccine candidate R21/Matrix-M in African children: a multicentre, double-blind, randomised, phase 3 trial. *Lancet* **403**, 533–544 (2024).

51. T. Bousema, L. Okell, I. Felger, C. Drakeley, Asymptomatic malaria infections: detectability, transmissibility and public health relevance. *Nat. Rev. Microbiol.* **12**, 833–840 (2014).

52. E. Rajaonarifara, M. H. Bonds, A. C. Miller, F. A. Ihantamalala, L. Cordier, B. Razafinjato, F. H. Rafenoarimalala, K. E. Finnegan, R. J. L. Rakotonanahary, G. Cowley, B. Ratsimbazafy, F. Razafimamonjy, M. Randriamanambintsoa, E. M. Raza-Fanomezanjanahary, A. Randrianambinina, C. J. Metcalf, B. Roche, A. Garchitorena, Impact of health system strengthening on delivery strategies to improve child immunisation coverage and inequalities in rural Madagascar. *BMJ Glob Health* **7** (2022).

53. F. K. Jones, K. Mensah, J.-M. Heraud, F. M. Randriatsarafara, C. J. E. Metcalf, A. Wesolowski, The Challenge of Achieving Immunity Through Multiple-Dose Vaccines in Madagascar. *Am. J. Epidemiol.* **190**, 2085–2093 (2021).

54. M. S. Datoo, H. M. Natama, A. Somé, D. Bellamy, O. Traoré, T. Rouamba, M. C. Tahita, N. F. A. Ido, P. Yameogo, D. Valia, A. Millogo, F. Ouedraogo, R. Soma, S. Sawadogo, F. Sorgho, K. Derra, E. Rouamba, F. Ramos-Lopez, M. Cairns, S. Provstgaard-Morys, J. Aboagye, A. Lawrie, R. Roberts, I. Valéa, H. Sorgho, N. Williams, G. Glenn, L. Fries, J. Reimer, K. J. Ewer, U. Shaligram, A. V. S. Hill, H. Tinto, Efficacy and immunogenicity of R21/Matrix-M vaccine against clinical malaria after 2 years' follow-up in children in Burkina Faso: a phase 1/2b randomised controlled trial. *Lancet Infect. Dis.* **22**, 1728–1736 (2022).

55. Institut National de la Statistique (INSTAT) et ICF, *Enquête Démographique et de Santé à Madagascar (EDSMD-V) 2021* (2022).

56. Intergovernmental Panel on Climate Change (IPCC), "Weather and Climate Extreme Events in a Changing Climate" in *Climate Change 2021 – The Physical Science Basis:*

Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (Cambridge University Press, 2023), pp. 1513–1766.

57. H. C. Slater, A. Ross, A. L. Ouédraogo, L. J. White, C. Nguon, P. G. T. Walker, P. Ngor, R. Aguas, S. P. Silal, A. M. Dondorp, P. La Barre, R. Burton, R. W. Sauerwein, C. Drakeley, T. A. Smith, T. Bousema, A. C. Ghani, Assessing the impact of next-generation rapid diagnostic tests on Plasmodium falciparum malaria elimination strategies. *Nature* **528**, S94–101 (2015).

58. L. Wu, L. L. van den Hoogen, H. Slater, P. G. T. Walker, A. C. Ghani, C. J. Drakeley, L. C. Okell, Comparison of diagnostics for the detection of asymptomatic Plasmodium falciparum infections to inform control and elimination strategies. *Nature* **528**, S86–93 (2015).

59. J. T. Lin, D. L. Saunders, S. R. Meshnick, The role of submicroscopic parasitemia in malaria transmission: what is the evidence? *Trends Parasitol.* **30**, 183–190 (2014).

60. "WHO guidelines for malaria" (World Health Organization, Geneva, 2023); https://www.ncbi.nlm.nih.gov/pubmed/36580567.

61. N. Schmit, H. M. Topazian, H. M. Natama, D. Bellamy, O. Traoré, M. A. Somé, T. Rouamba, M. C. Tahita, M. D. A. Bonko, A. Sourabié, H. Sorgho, L. Stockdale, S. Provstgaard-Morys, J. Aboagye, D. Woods, K. Rapi, M. S. Datoo, F. R. Lopez, G. D. Charles, K. McCain, J.-B. Ouedraogo, M. Hamaluba, A. Olotu, A. Dicko, H. Tinto, A. V. S. Hill, K. J. Ewer, A. C. Ghani, P. Winskill, The public health impact and cost-effectiveness of the R21/Matrix-M malaria vaccine: a mathematical modelling study. *Lancet Infect. Dis.*, doi: 10.1016/S1473-3099(23)00816-2 (2024).

62. H. M. Topazian, N. Schmit, I. Gerard-Ursin, G. D. Charles, H. Thompson, A. C. Ghani, P. Winskill, Modelling the relative cost-effectiveness of the RTS,S/AS01 malaria vaccine compared to investment in vector control or chemoprophylaxis. *Vaccine* **41**, 3215–3223 (2023).

63. O. O. Adeshina, S. Nyame, J. Milner, A. Milojevic, K. P. Asante, Barriers and facilitators to nationwide implementation of the malaria vaccine in Ghana. *Health Policy Plan.* **38**, 28–37 (2023).

64. World Health Organization, *Malaria Control in Humanitarian Emergencies: An Inter-Agency Field Handbook* (World Health Organization, 2013).

65. M. Burns, M. Rowland, R. N'Guessan, I. Carneiro, A. Beeche, S. S. Ruiz, S. Kamara, W. Takken, P. Carnevale, R. Allan, Insecticide-treated plastic sheeting for emergency malaria prevention and shelter among displaced populations: an observational cohort study in a refugee setting in Sierra Leone. *Am. J. Trop. Med. Hyg.* **87**, 242–250 (2012).

66. L. A. Messenger, J. Furnival-Adams, B. Pelloquin, M. Rowland, Vector control for malaria prevention during humanitarian emergencies: protocol for a systematic review and meta-analysis. *BMJ Open* **11**, e046325 (2021).

67. World Health Organization, *Operational Framework for Building Climate Resilient Health Systems* (World Health Organization, Genève, Switzerland, 2015).

68. M. Romanello, A. McGushin, C. Di Napoli, P. Drummond, N. Hughes, L. Jamart, H.

Kennard, P. Lampard, B. Solano Rodriguez, N. Arnell, S. Ayeb-Karlsson, K. Belesova, W. Cai, D. Campbell-Lendrum, S. Capstick, J. Chambers, L. Chu, L. Ciampi, C. Dalin, N. Dasandi, S. Dasgupta, M. Davies, P. Dominguez-Salas, R. Dubrow, K. L. Ebi, M. Eckelman, P. Ekins, L. E. Escobar, L. Georgeson, D. Grace, H. Graham, S. H. Gunther, S. Hartinger, K. He, C. Heaviside, J. Hess, S.-C. Hsu, S. Jankin, M. P. Jimenez, I. Kelman, G. Kiesewetter, P. L. Kinney, T. Kjellstrom, D. Kniveton, J. K. W. Lee, B. Lemke, Y. Liu, Z. Liu, M. Lott, R. Lowe, J. Martinez-Urtaza, M. Maslin, L. McAllister, C. McMichael, Z. Mi, J. Milner, K. Minor, N. Mohajeri, M. Moradi-Lakeh, K. Morrissey, S. Munzert, K. A. Murray, T. Neville, M. Nilsson, N. Obradovich, M. O. Sewe, T. Oreszczyn, M. Otto, F. Owfi, O. Pearman, D. Pencheon, M. Rabbaniha, E. Robinson, J. Rocklöv, R. N. Salas, J. C. Semenza, J. Sherman, L. Shi, M. Springmann, M. Tabatabaei, J. Taylor, J. Trinanes, J. Shumake-Guillemot, B. Vu, F. Wagner, P. Wilkinson, M. Winning, M. Yglesias, S. Zhang, P. Gong, H. Montgomery, A. Costello, I. Hamilton, The 2021 report of the Lancet Countdown on health and climate change: code red for a healthy future. *Lancet* **398**, 1619–1662 (2021).

69. M. E. Beatty, E. Hunsperger, E. Long, J. Schürch, S. Jain, R. Colindres, G. Lerebours, Y.-M. Bernard, J. G. Dobbins, M. Brown, G. G. Clark, Mosquitoborne infections after Hurricane Jeanne, Haiti, 2004. *Emerg. Infect. Dis.* **13**, 308–310 (2007).

70. T. Fürst, G. Raso, C. A. Acka, A. B. Tschannen, E. K. N'Goran, J. Utzinger, Dynamics of socioeconomic risk factors for neglected tropical diseases and malaria in an armed conflict. *PLoS Negl. Trop. Dis.* **3**, e513 (2009).

71. L. Sedda, Q. Qi, A. J. Tatem, A geostatistical analysis of the association between armed conflicts and Plasmodium falciparum malaria in Africa, 1997-2010. *Malar. J.* **14**, 500 (2015).

72. M. A. Connolly, M. Gayer, M. J. Ryan, P. Salama, P. Spiegel, D. L. Heymann, Communicable diseases in complex emergencies: impact and challenges. *Lancet* **364**, 1974–1983 (2004).

73. {Institut National de la Statistique-INSTAT/Madagascar, Programme National de lutte contre le Paludisme-PNLP/Madagascar, Institut Pasteur de Madagascar - IPM/Madagascar, and ICF International, *Enquête Sur Les Indicateurs Du Paludisme 2016 Madagascar* (Calverton, MD, USA : INSTAT, PNLP, IPM and ICF International., 2017).

74. Institut National de la Statistique-INSTAT/Madagascar, Programme National de lutte contre le Paludisme-PNLP/Madagascar, Institut Pasteur de Madagascar-IPM/Madagascar, ICF International, *Madagascar Enquête Sur Les Indicateurs Du Paludisme (EIPM) 2013* (2013).

75. Institut National de la Statistique/Madagascar, Programme National de Lutte contre le Paludisme/Madagascar, ICF International, Madagascar Enquête sur les Indicateurs du Paludisme à Madagascar (EIPMD) 2011. (2012).

76. R. Arambepola, S. H. Keddie, E. L. Collins, K. A. Twohig, P. Amratia, A. Bertozzi-Villa, E. G. Chestnutt, J. Harris, J. Millar, J. Rozier, S. F. Rumisha, T. L. Symons, C. Vargas-Ruiz, M. Andriamananjara, S. Rabeherisoa, A. C. Ratsimbasoa, R. E. Howes, D. J. Weiss, P. W. Gething, E. Cameron, Spatiotemporal mapping of malaria prevalence in Madagascar using routine surveillance and health survey data. *Sci. Rep.* **10**, 18129 (2020).

77. R. E. Howes, S. A. Mioramalala, B. Ramiranirina, T. Franchard, A. J. Rakotorahalahy, D. Bisanzio, P. W. Gething, P. A. Zimmerman, A. Ratsimbasoa, Contemporary

epidemiological overview of malaria in Madagascar: operational utility of reported routine
case data for malaria control planning. *Malar. J.* **15**, 502 (2016).

78. M. Nguyen, R. E. Howes, T. C. D. Lucas, K. E. Battle, E. Cameron, H. S. Gibson, J. Rozier,
S. Keddie, E. Collins, R. Arambepola, S. Y. Kang, C. Hendriks, A. Nandi, S. F. Rumisha, S.
Bhatt, S. A. Mioramalala, M. A. Nambinisoa, F. Rakotomanana, P. W. Gething, D. J. Weiss,
Mapping malaria seasonality in Madagascar using health facility data. *BMC Med.* **18**, 26
(2020).

79. R. Ratovoson, A. Garchitorena, D. Kassie, J. A. Ravelonarivo, V. Andrianaranjaka, S.
Razanatsiorimalala, A. Razafimandimby, F. Rakotomanana, L. Ohlstein, R.
Mangahasimbola, S. A. N. Randrianirisoa, J. Razafindrakoto, C. M. Dentinger, J.
Williamson, L. Kapesa, P. Piola, M. Randrianarivelojosia, J. Thwing, L. C. Steinhardt, L.
Baril, Proactive community case management decreased malaria prevalence in rural
Madagascar: results from a cluster randomized trial. *BMC Med.* **20**, 322 (2022).

80. C. D. Golden, E. J. G. Anjaranirina, L. C. H. Fernald, D. L. Hartl, C. Kremen, D. A. Milner Jr,
D. H. Ralalason, H. Ramihantaniarivo, H. Randriamady, B. L. Rice, B. Vaitla, S. K.
Volkman, M. A. Vonona, S. S. Myers, Cohort Profile: The Madagascar Health and
Environmental Research (MAHERY) study in north-eastern Madagascar. *Int. J. Epidemiol.*
46, 1747–1748d (2017).

81. C. D. Golden, B. L. Rice, H. J. Randriamady, A. M. Vonona, J. F. Randrianasolo, A. N.
Tafangy, M. Y. Andrianantenaina, N. J. Arisco, G. N. Emile, F. Lainandrasana, R. F. F.
Mahonjolaza, H. P. Raelson, V. R. Rakotoarilalao, A. A. N. A. Rakotomalala, A. D.
Rasamison, R. Mahery, M. L. Tantely, R. Girod, A. Annapragada, A. Wesolowski, A.
Winter, D. L. Hartl, J. Hazen, C. J. E. Metcalf, Study Protocol: A Cross-Sectional
Examination of Socio-Demographic and Ecological Determinants of Nutrition and Disease
Across Madagascar. *Front Public Health* **8**, 500 (2020).

82. B. L. Rice, C. D. Golden, E. J. G. Anjaranirina, C. M. Botelho, S. K. Volkman, D. L. Hartl,
Genetic evidence that the Makira region in northeastern Madagascar is a hotspot of malaria
transmission. *Malar. J.* **15**, 596 (2016).

83. S. Xiaodong, E. Tambo, W. Chun, C. Zhibin, D. Yan, W. Jian, W. Jiazhi, Z. Xiaonong,
Diagnostic performance of CareStart™ malaria HRP2/pLDH (Pf/pan) combo test versus
standard microscopy on falciparum and vivax malaria between China-Myanmar endemic
borders. *Malar. J.* **12**, 6 (2013).

84. G. M. Bwire, B. Ngasala, M. Kilonzi, W. P. Mikomangwa, F. F. Felician, A. A. R.
Kamuhabwa, Diagnostic performance of CareStart™ malaria HRP2/pLDH test in
comparison with standard microscopy for detection of uncomplicated malaria infection
among symptomatic patients, Eastern Coast of Tanzania. *Malar. J.* **18**, 354 (2019).

85. D. A. Griffiths, A Catalytic Model of Infection for Measles. *J. R. Stat. Soc. Ser. C Appl. Stat.*
23, 330–339 (1974).

86. N. Hens, Z. Shkedy, M. Aerts, C. Faes, P. Van Damme, P. Beutels, *Modeling Infectious*
Disease Parameters Based on Serological and Social Contact Data (Springer New York).

87. L. C. Okell, C. J. Drakeley, A. C. Ghani, T. Bousema, C. J. Sutherland, Reduction of
transmission from malaria patients by artemisinin combination therapies: a pooled analysis

of six randomized trials. *Malar. J.* **7**, 125 (2008).

88. M. S. Datoo, A. Dicko, H. Tinto, J.-B. Ouédraogo, M. Hamaluba, A. Olotu, E. Beaumont, F.
Ramos-Lopez, H. Magloire Natama, S. Weston, M. Chemba, Y. D. Compaore, D. Issiaka,
D. Salou, S. Omenda, A. Lawrie, P. Bejon, D. Chandramohan, R. Roberts, S. Bharati, L.
Stockdale, B. Greenwood, K. Ewer, J. Bradley, C. S. Poonawalla, P. S. Kulkarni, U.
Shaligram, A. V. S. Hill, A Phase III Randomised Controlled Trial Evaluating the Malaria
Vaccine Candidate R21/Matrix-M™ in African Children (2023).
https://doi.org/10.2139/ssrn.4584076.