Paving the way for precision treatment of psychiatric

symptoms with functional connectivity neurofeedback

Taylor JE[1]*[†], Oka T[1,2][†], Murakami M[1], Motegi T[1,3], Yamada T[1,4,5], Kawashima T[6], Kobayashi Y[6],

Yoshihara Y[6], Miyata J[6,7], Murai T[6], Kawato M[1], & Cortese A[1].

[1] The Department of Decoded Neurofeedback, Computational Neuroscience Laboratories, Advanced Telecommunications

Research Institute International, Kyoto, Japan

[2] The Department of Clinical Psychology, Graduate School of Human Sciences, Osaka University, Suita, Japan

[3] Saint-Pierre Hospital, Takasaki, Japan

[4] Division of Developmental Neuroscience, United Graduate School of Child Development, Osaka University, 2-2 Yamadaoka,

Suita, Osaka, 565-0871, Japan

[5] Medical Institute of Developmental Disabilities Research, Showa University, Tokyo, Japan

[6] Department of Psychiatry, Graduate School of Medicine, Kyoto University, Kyoto, Japan

[7] Department of Psychiatry, Aichi Medical University, Aichi, Japan

* correspondence: jessie.elizabeth.taylor@gmail.com

† Dr. Taylor and Dr. Oka contributed to this work equally.

Short Title: FCNef for Precision Psychiatry

Corresponding Author: Jessica Elizabeth Taylor

E-mail address: jessie.elizabeth.taylor@gmail.com

Keywords: Precision psychiatry, real-time fMRI neurofeedback, Major Depressive Disorder, Rumination

Abstract

Introduction: Major depressive disorder (MDD) remains challenging to treat, with many patients failing to respond adequately to existing therapies. Patients with MDD have heterogeneous subsets of symptoms with differing underlying neural aberrations. Treatment response may improve if treatments become more individualised. We recently showed preliminary evidence that normalisation of a neural network and a corresponding reduction in related symptoms can be achieved using a Brain Machine Interface (BMI) called real-time fMRI functional connectivity neurofeedback (FCNef). However, the robustness of this effect, and the best FCNef parameters for optimising therapeutic outcomes remained unknown. ***Methods:*** We ran additional participants, with a final dataset of N = 68, in our FCNef protocol. Functional connectivity between the dorsolateral prefrontal cortex (DLPFC) and posterior cingulate cortex/precuneus (PCC) was targeted with the goal of reducing brooding rumination symptoms. Core FCNef parameters (experimental schedule and reward) were manipulated between participants. ***Results:*** We replicated findings that normalisation of DLPFC-PCC connectivity with FCNef correlates significantly with reductions in brooding rumination, but not with changes in anxiety, which is associated with different neural circuits. The difference between these correlations was significant, highlighting the precision of this effect. Finally, we found that successful DLPFC-PCC normalisation and corresponding changes in brooding rumination depended on specific FCNef parameters. The most effective protocol involved consecutive training days with greater external reward. ***Conclusions***: These results highlight the potential of FCNef for precision medicine in psychiatry and underscore the importance of optimising parameters to enhance feasibility of BMI-based clinical interventions.

54 1. Introduction

The World Health Organization estimates that major depressive disorders will become the top
cause of global disease burden by 2030 [1]. Of patients who do receive treatment, an estimated 30-50%
do not respond fully [2,3]. Patients with the same clinical diagnosis can have heterogeneous subsets of
symptoms that relate to different underlying neural mechanisms [4]. Nonetheless, they usually receive
relatively homogenous treatment. For example, all or most clinical practice guidelines recommend
selective serotonin reuptake inhibitors as first-line treatment for depression [5]. To improve response
rates, individual differences clearly need to be considered. Future treatment may become more
individualised using Brain-Machine Interfaces (BMIs) to identify and target underlying patient neural
aberrations. However, medical regulatory approval presents serious challenges to this [6]. To date, only a
handful of BMIs have received approval from local medical regulatory agencies for human trials [7–9],
and even fewer have received full market authorisation [10,11]. A key step toward approval of a given
BMI technique by regulatory agencies is demonstrating the optimality of chosen parameters. Here, we
extend previous results to better examine the robustness of a promising form of BMI called Functional-
Connectivity Neurofeedback (FCNef) [12], and to systematically investigate a specific set of its
parameters.

FCNef is a functional magnetic resonance imaging (fMRI) neurofeedback technique in which
participants receive real-time feedback about the current state of functional connectivity between targeted
brain regions (measured as the correlation between time-courses of BOLD activity from these regions).
This feedback is used to train participants to make a targeted functional connection more positive or
negative, a result that has been demonstrated in multiple studies [12–19]. Showing promise for precision
medicine, recent FCNef studies have reported precise correspondence between normalisation of
functional connections and reductions in specifically related symptoms [12,17,18,20]. Consistent with
these results, we previously found that normalisation of functional connectivity between the dorsolateral
prefrontal cortex (DLPFC) and the precuneus/posterior cingulate cortex (PCC) that occurred with FCNef
was related to reductions in brooding rumination, but not anxiety symptoms. Importantly, these effects
persisted at least 1-2 months after FCNef [12]. Nonetheless these results were reported from only a small
sample. We have since continued to collect data and here we examine the robustness of these effects by
testing for their replication in the newly collected data. Furthermore, because combining the new and old
data provides sufficient statistical power to properly compare correlation coefficients, for the first time,
we are able to directly examine the specificity of this effect.

Given its potential use as a medical tool, FCNef should maximise health outcomes while

minimising patient burden. Here, we operationalised FCNef success as the normalisation of DLPFC/PCC

functional connectivity and a related reduction in brooding (but not anxiety) symptoms. We sought to find

parameters that would best enhance this, while also keeping participant fatigue to a minimum (because

preliminary testing in a clinical sample caused fatigue-related drop-out of one out of six patients [21]). To

accomplish our objective, we focused on the following parameters: (1) Reward schedule: During real-

time neurofeedback tasks, feedback has conventionally been provided simply as scores that reflect how

similar the induced brain activity is to the target brain activity [22–24]. However, recent evidence

suggests that target neural activity may be better reinforced during neurofeedback when external reward,

such as money, is also used [25]. Here, we manipulated how bonus money was assigned to feedback

scores so that different groups of participants could earn less/more overall external reward. (2)

Experimental schedule. Most FCNef studies [15], including our own [12,19], have required participants to

come in for multiple consecutive days of experimentation. This can be exhausting and requires motivation

and organisation skills that can be diminished in psychiatric disorders [26]. Therefore, we tested whether

a more flexible schedule, over non-consecutive days, could yield similar results to the consecutive

training schedule.

Overall, we ran 68 participants in our FCNef for depression paradigm while manipulating reward

schedule (low/high) and experimental schedule (consecutive/non-consecutive training days). Our goals

were twofold: (1) using a larger sample size to examine the robustness of our previously reported results

and more precisely examine the specificity of the FCNef effect, and (2) to fine-tune underlying

parameters.

2. Methods

2.1. Participants

Participants were screened twice using the Beck Depression Inventory-II (BDI) questionnaire

[27] (see Supplementary Methods for more detail). Only those with an average score ≥ 8, who indicated

no intention of committing suicide, who spoke Japanese, who held no current clinical diagnosis, and who

were not currently receiving treatment for a psychiatric illness were invited to participate. Overall, 69

people passed the screening and participated in the main experiment. However, experimental data of one

participant was subsequently excluded from data analysis because it was revealed, subsequent to

experimentation, that that subject held a current clinical diagnosis. Of the 68 participants whose data were

used for analyses, the average BDI score over the two screening measurements was 14.33, with a standard

deviation (STD) of 5.26. This puts them generally in the category of "mild depression" [27]. For this

reason and because these participants lacked current clinical diagnoses, we considered them "subclinical".

2.2. Experimental Conditions

We wished to examine the overall success of our paradigm and to determine whether it is

influenced by reward and experimental schedules. However, due to financial and time constraints (it takes

10 days of experimentation to screen and run just one participant in this paradigm), we could only run

participants in three experimental groups: those with consecutive days of experimentation and a high-

reward schedule (hereafter, "Consec/High-Rew"), those with consecutive days of experimentation and a

low-reward schedule (hereafter, "Consec/Low-Rew"), and those with non-consecutive days of

experimentation and a low-reward schedule (hereafter, "Non-Consec/Low-Rew") (see Table 1 for sample

size details). Participants in the three groups did not differ significantly in baseline levels of our main

measures of interest: (1) DLPFC-PCC resting-state functional-connectivity (rs-FC), (2) BDI scores, and

(3) brooding scores (measured on a subscale of the Rumination Response Scale [28,29]) (see

Supplementary Table 7). However, we found that baseline anxiety levels (measured with the trait anxiety

subscale of the State-Trait Anxiety Inventory [30]) differed significantly between the Consec/Low-Rew

and Non-Consec/Low-Rew groups (see Supplementary Table 7). This is unlikely to have had a large

impact on results because: (a) the rs-FC we targeted is not thought to relate to anxiety, and (b) results

showed that differences in FCNef success were not greatest between these two groups (as can be seen

further down).

Participant group assignments were determined by concurrent availability of participants,

experimenters, and MR machine facilities. This allocation method reflects real-world constraints in

neuroimaging research, though we acknowledge it could have introduced potential self-selection effects.

To address possible allocation bias, we conducted analyses to compare baseline demographic

measurements between groups (age, sex): No meaningful differences were found (see Supplementary

Results). It should be noted that data of 19 participants (9/21 from the Consec/High-Rew group and 10/23

from the Consec/Low-Rew group) have been reported elsewhere [12]. Further details about recruitment,

participant demographics, and payment can be found in the Supplementary Methods.

Group	Main Experiment Sample Size	1-Month Follow-up Sample Size	2-Month Follow-up Sample Size
Consec/High-Rew	21	11	9
Consec/Low-Rew	23	20	17
Non/Low-Rew	24	22	20
Total	68	53	46

Table 1. Sample sizes for three groups of participants.

This table shows the sample sizes for: (1) participants run in consecutive days of FCNef with the high-reward

schedule (the "Consec/High-Rew group"), (2) those run in consecutive days of FCNef with the low-reward schedule

(the "Consec/Low-Rew group"), and (3) those run over non-consecutive days of FCNef with the low-reward

schedule (the "Non-Consec/Low-Rew group"). No participants were run over non-consecutive days of FCNef with

the high-reward schedule (no "Non-Consec/High-Rew" group). Sample sizes are shown for the main experiment

and for one- and two-month follow-up tests. There were fewer participants for the follow-up tests for the

Consec/High-Rew group because the long-term tests were not included in the earliest stages of this experiment, so

only 12 participants from this group were invited back.

2.3. Experimental procedure, materials, and imaging data acquisition

An outline of the experimental procedure is shown in Figure 1 alongside details of the FCNef

task. These were largely the same as in our previous report [12], except for specific experimental

conditions of interest. These experimental conditions are described in the sub-section below entitled

"Differences in experimental conditions." The protocol and imaging data acquisition details are identical

to those in our previous report [12] and are summarised in the Supplementary Methods.

Details of the symptom questionnaires can be found in the Supplementary Methods, but overall

general depressive symptoms were measured with the BDI [27]. Brooding rumination symptoms were

measured with a subscale of the Rumination Response Scale (RRS) [28,29], and trait anxiety symptoms

were measured with the trait anxiety subscale of the State-Trait Anxiety Inventory (STAI-Y2) [30]. As

can be seen in Figure 1, because there would be limited clinical meaning, these symptom questionnaire

scores were not measured on all days of the main experiment. Instead, they were only measured on the

first day (Day 0) and last day (Day 4) of the main experiment and during one- and two-month follow-up

tests.

Figure 1. Experimental procedure and example FCNef trial.

a. The order of events on each day of experimentation. *Questionnaires = the Beck Depression Inventory-II [27],*

the Rumination Response Scale [28], and the State-Trait Anxiety Inventory [30]. Anatomical = T1-weighted

structural MRI. N-back = a well-known executive control task [54], used here as a functional localiser[12]. ***b. An***

example FCNef trial. *During the rest period, participants were to simply relax. During the induction period, they*

were asked to "somehow" manipulate their brain activity to get the best possible feedback. Participants were told

that different strategies of brain activity manipulation might work for different people. Unbeknown to participants

(nothing changed on screen), there was a 2s calculation period at the end of the induction period. During FCNef,

DLPFC-PCC connectivity (from the induction period) was calculated during the calculation period and this

determined the feedback presented during the feedback period. During SHAM, however, feedback was just random.

Feedback was presented on screen as a green circle and participants had been clearly instructed that the larger this

was, the more monetary reward they would receive on that trial. During FCNef, they were instructed to try to make

the green circle bigger than the red circle that was also presented on screen. The circumference of this red circle

represented the participant's baseline DLPFC-PCC connectivity (the average from SHAM). During SHAM, there

was no red circle and participants were simply instructed to try to make the green circle as big as possible. Modified

with permission from Taylor et al. (2022)[12].

2.4. Differences in experimental conditions

The following parameters were manipulated, such that some participants were run under

conditions different from those reported in our previous papers [12,19].

2.4.1. Reward schedule:

All participants received a baseline reward bonus of ¥500 on each day of the SHAM and FCNef

sessions. This is the maximum reward they could receive in the SHAM task, but they could get an

additional reward bonus in the FCNef task depending on their average FCNef scores from that day.

Importantly, two calculation methods were used to determine this additional reward bonus. Participants in

the low-reward schedule conditions could achieve an additional reward bonus of ¥100 for each average

FCNef score point over 75. Instead, participants in the high-reward schedule condition could achieve an

additional reward bonus of ¥50 for each average FCNef score point over 50. See Supplementary Table 2

for specific examples. The way in which scores were calculated in the SHAM and FCNef tasks was

identical across conditions. This was the same method as in our previous report [12] and is described in

the Supplementary Methods.

2.4.2. Experimental schedule:

Two groups of participants completed Days 0-4 over 5 consecutive days (Monday-Friday). The

third group completed Days 0-4 over 5 non-consecutive days, which took place over a period of several

weeks to months (mean of 18.5 days ± STD of 15.3 days).

2.4.3. Induction time-window:

Consistent with earlier versions of our FCNef for depression paradigm [12,19], about half of the

participants from each group completed FCNef and SHAM with a 40 sec induction time-window (see

Supplementary Table 1). Because re-analysis of pilot data suggested it would not affect results (see

Supplementary Results, Supplementary Table 3, and Supplementary Figure 1), the other half of

participants completed FCNef and SHAM with a 20 sec induction time-window. We included this

manipulation with the goal of improving our paradigm. Using a 20 sec time-window reduces the overall

experiment by about 10 mins each day, which could be crucial for patients who tire easily. Indeed, when

we checked the data of our three groups of participants, we found that our measure of FCNef success did

not change in any meaningful way that depended on the induction time-window (see Supplementary

Results and Supplementary Tables 4-6). Therefore, in the main text of this report, we did not split the data

by induction time-window because (a) it is hard to draw strong conclusions about null effects, and (b)

splitting the data further like this would make sample sizes in each cell even smaller, making it hard to

draw conclusions about other parameters of interest.

2.5. Data analyses

2.5.1. Correlations to extend previously reported results

We previously reported correlations between rs-FC and symptom changes using data from 19

participants who were run over consecutive days of FCNef [12]. Ten of these participants were run with

the high-reward schedule and the other nine were run with the low-reward schedule, but overall their

combined data showed promising results. Since then, we have collected data from 25 more participants

under the same experimental conditions (consecutive days of FCNef with low/high-reward schedules),

which gives us an overall dataset with 44 participants run under these conditions (see Table 1). Note that

this does not include the data of the Non-Consec/Low-Rew group, which has also been collected since

our previous report. This is because this group was run with a different experimental parameter to our

previous report (non-consecutive days of experimentation) and so its data cannot contribute directly to our

replication test. One goal of collecting this additional data was to assess the robustness of previously

reported results in a larger dataset. Before doing this in the 44 participants, however, we first re-ran

relevant correlations using only data of the additional 25 participants to ensure that we were not

introducing bias by adding new data to the existing pool. After this test, we then re-ran the previously

reported correlations with the larger pooled dataset. To check the adequacy of the sample size for

statistical tests, we conducted post-hoc power analyses with G*Power version 3.1.9.7 (Franz Faul, Kiel

University, Germany). The significance threshold was set at $\alpha = 0.05$, and we input relevant correlational

values to calculate statistical power.

2.5.2. Linear Mixed Effect (LME) models to examine parameters of interest

We ran multiple LME models to examine how multiple measures related to FCNef success were

impacted by our manipulated parameters (see Table 2). Dependent variables (DVs) in these models were

taken from the task (e.g. average task score) or from the resting-state scans (e.g. changes in resting-state

functional connectivity from before to after FCNef). All 3 groups of participants (N = 68) were included

in these analyses so we could best examine parameters of interest.

We were unable to run a Non/High-Rew group (see Table 1). This means we could not run full

factorial analyses to investigate all possible interactions between parameters that we manipulated,

although we had no reason to expect interactions anyway. Instead, in all models reported here, we

included "Group" as a categorical Independent Variable (IV) with 3 levels. By nature, this type of

analysis compares the variance related to the second two levels with the variance related to the first level

(the reference), so it is important to consider what data will be input at what level. Here, we decided to set

the Consec/High-Rew group as the first level because this group had parameters that we used in the initial

pilot experiment [19] and we wanted to examine whether changes to these initial parameters would

improve or worsen FCNef success.

In addition to Group, the LME models also included other relevant IVs, which differed between

models depending on the DV. In models examining how brain changes predicted symptom changes, this

was changes in resting-state functional connectivity from before to after FCNef. In other models, this was

the relevant experimental days.

For each DV, to explore the data, we ran LME models with and without the interaction term

between IVs. We then tested these against one another to see which best fit the data.

No interaction term: *'DV~X+Group+(1|Subject)'*

With interaction term: *'DV~X*Group+(1|Subject)'*

Likelihood ratio tests were used to compare models with and without interaction terms for all
DVs. If there was a significant difference between models, the best-fit model was always selected based
on the lowest Akaike Information Criterion (AIC, a measure of goodness of fit, which can be used to
assess a model's relative quality [31]). If there was no significant difference between models then the
simplest model, that without interaction, was selected as best-fit. Detailed information about the best-fit
of each of these pairs of models can be found in tables in the Supplementary Results (Supplementary
Tables 8, 9, 11-17, 19-24), with all significant main effects and interactions from the best-fit models
additionally being reported in the Results Section below.

2.5.3. Follow-up statistical testing

Significant main effects and interactions from the best-fit models, as well as effects that we had
specific hypotheses about, were followed up with t-tests or correlations. We applied False Discovery Rate
(FDR) correction [32] whenever there were multiple comparisons. We had strong directional hypotheses
that symptoms would decrease from before to after FCNef and that rs-FC would become more negative,
aligning with patterns observed in healthy individuals. Therefore, we conducted the related t-tests using a
one-tailed approach. Similarly, based on our strong directional hypothesis that symptoms would decrease
as rs-FC became more negative (indicating a positive correlation between changes in symptoms and
changes in rs-FC), we also used a one-tailed approach for the related correlation analyses.

3. Results

3.1. Extension of past results

3.1.1. Previously reported data

In our previous publication [12], we reported correlations between rs-FC and symptom changes
using combined data of 9 participants from the current Consec/High-Rew group and 10 participants from
the current Consec/Low-Rew group. We found a significant positive relationship between general
depression change and rs-FC change ($r = 0.78$, $p_{FDR} = 0.0001$, 95% CI [0.51, 0.91]) and between brooding
change and rs-FC change ($r = 0.43$, $p_{FDR} = 0.048$, 95% CI [-0.27, 0.74]), but a non-significant negative
relationship between anxiety change and rs-FC change ($r = 0.11$, $p_{FDR} = 0.32$, 95% CI [-0.36, 0.54]) (note
that we previously only reported correlation coefficients and uncorrected p-values). Because we
specifically hypothesised that changes in DLPFC-PCC rs-FC from before to after FCNef should relate to
changes in brooding but not anxiety symptoms, here we calculated Z-tests to compare these correlation
coefficients [33]. These were run with one tail due to our directional hypothesis, and no significant

difference was found ($z = 0.88$, $p = 0.16$). A post-hoc power analysis was conducted using correlation

coefficients used for this Z-test ($r = 0.43$ and $r = 0.11$) and the coefficient for the correlation between

brooding and anxiety changes ($r = -0.03$). This revealed a post-hoc power of 0.25, suggesting that the

sample size (N = 19) was statistically insufficient to detect an effect of this magnitude.

3.1.2. Newly collected data

We have since collected more data for each of these groups (12 more participants for the

Consec/High-Rew group and 13 more participants for the Consec/Low-Rew group) with the goal of

testing robustness of the aforementioned findings in a larger dataset. To ensure that we were not adding

bias or overestimating effects, before testing results for the pooled larger dataset, we first ran a replication

test using just the newly collected data (N = 25). Consistent with the aforementioned results, we found a

positive relationship between general depression change and rs-FC change ($r = 0.26$, $p_{FDR} = 0.16$, 95% CI

[-0.16, 0.60]) and between brooding change and rs-FC change ($r = 0.40$, $p_{FDR} = 0.08$, 95% CI [-0.01,

0.69]), although these did not reach significance here. We found a non-significant negative relationship

between anxiety change and rs-FC change ($r = -0.17$, $p_{FDR} = 0.80$, 95% CI [-0.53, 0.24]).

3.1.3. Pooled data

Overall, we consider the previously reported effects to be replicated in the newly collected data

because confidence intervals overlapped and effect sizes were comparable. When we next pooled these

data (total N = 44) we found a significant correlation between general depression change and rs-FC

change ($r = 0.48$, $p_{FDR} = 0.002$, 95% CI [0.21, 0.68]), and between brooding change and rs-FC change (r

$= 0.42$, $p_{FDR} = 0.004$; 95% CI [0.13, 0.64]), but not between anxiety change and rs-FC change ($r = -0.03$,

$p_{FDR} = 0.57$; 95% CI [-0.32, 0.27]). Demonstrating the specificity achieved when targeting the

DLPFC/PCC functional connection, we found a significant difference between coefficients from brooding

change/rs-FC change and anxiety change/rs-FC change correlations ($z = 2.15$, $p = 0.016$) (see Figure 2). A

post-hoc power analysis was conducted using correlation coefficients employed in this Z-test ($r = 0.42$

and $r = -0.03$) and the coefficient for the correlation between brooding and anxiety changes ($r = 0.23$).

This revealed a power of 0.79, suggesting that the sample size (N = 44) was sufficient to detect an effect

of this magnitude with high probability.

Figure 2. Comparison of rs-FC/Brooding and rs-FC/anxiety correlations

*Promising for precision medicine, changes in DLPFC-PCC resting state functional connectivity (rs-FC) from before to after FCNef were significantly correlated with changes in brooding, but not anxiety symptoms. Pooling participants from consecutive conditions (to be consistent with conditions used in our previous report) provided us with the statistical power to compare these correlation coefficients. The significant difference found, highlights the precision of FCNef targeting this functional connection. Note that this same data, but split by condition, is presented again in Figure 5a and 5b. ** = $p_{FDR} < 0.01$, * = $p < 0.05$.*

3.2. Parameter Investigations

Analyses	Formulae				AICs	
	DV	IV1	IV2	Random Intercept	+	x
FCNef Scores	Mean FCNef Score	~ FCNef Day	+ or x Group	+ (1|Subject)	2109.2	2108.7
	STD FCNef Score	~ FCNef Day	+ or x Group	+ (1|Subject)	1599.8	**1593**
Symptoms	General Depressive Scores	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	776.72	779.13
	Brooding Scores	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	681.4	682.68
	Anxiety Scores	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	913.86	914.62
	General Depressive Score Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	926.03	925.45
	Brooding Score Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	766.01	769.92
	Anxiety Score Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	1095.7	1095
rs-FC	rs-FC	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	-22.99	-23.67
	rs-FC Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	-58.77	-55.47
Symptom Change/	General Depressive Score Changes (D4)	~ rs-FC Change	+ or x Group	+ (1|Subject)	316.57	**-153.29**
rs-FC Change	Brooding Score Changes (D4)	~ rs-FC Change	+ or x Group	+ (1|Subject)	287.8	287.52
Relationship	Anxiety Score Changes (D4)	~ rs-FC Change	+ or x Group	+ (1|Subject)	408.46	410.45

Table 2. The models used to examine dependent variables from the FCNef experiment.

Models with and without interactions between independent variables (IVs) were compared to see which would best predict each dependent variable (DV). Akaike Information Criteria (AIC) are displayed in columns labelled '+' (for models without interactions) and 'x' (for models with interactions). These are highlighted in bold in cases in which likelihood ratio testing showed the corresponding model to be significantly better fit than the alternative model. If there was no significant difference, then the model without the interaction was selected as best-fit because it was the simplest. FCNef Day = Days 1-4; Subject = experimental participant; STD = standard deviation; First/last FCNef Day = Days 0 and 4; Post-Day = Day 4, and the 1- and 2-month follow-up test days; Changes = data from the day indicated in brackets (e.g. D4) minus data from Day 0; rs-FC = resting state functional connectivity between the DLPFC-PCC.

3.2.1. FCNef scores

In the total dataset (N = 68), while comparing groups, we first examined whether mean FCNef scores increased significantly across training days, because this would indicate that participants successfully modified the target functional connection in the trained direction. Additionally, we assessed whether variance in FCNef scores decreased significantly across training days, because this would indicate that participants had gained better control of the DLPFC-PCC functional connection.

The best-fit LME model to predict means of average daily FCNef scores was the model with no interaction (Table 2 and Supplementary Table 8). This model showed a significant main effect of FCNef day ($p = 0.0009$). In short, mean scores increased over training days (see Figure 3), which indicates that

participants could do the task. There was also a significant effect of the Non-Consec/Low-Rew group ($p =$

0.046): Overall scores were lower for this group than for the reference Consec/High-Rew group (see

Figure 3a), who were the only group capable of raising their scores significantly above baseline.

The best-fit model to explain variance in FCNef scores did include the interaction term (see Table

2 and Supplementary Table 9). This model showed a significant main effect of FCNef day ($p < 0.0001$),

which was qualified by a significant interaction between the Non-Consec/Low-Rew group and FCNef day

($p = 0.001$). This interaction indicates that changes in FCNef score variance over FCNef days did not

follow the same trajectory for the Non-Consec/Low-Rew group as it did for the reference Consec/Low-

Rew group; As can be seen in Figure 3b- the variance decreased over days for the reference group, but not

for the Non-Consec/Low-Rew group. No such significant interaction was found between the

Consec/Low-Rew group and FCNef Day ($p = 0.11$), which indicates that the Consec/Low-Rew group's

trajectory did not differ from that of the reference group. Indeed, the data for the Consec/Low-Rew group

lay between that of the other two groups, showing a trend for a decrease in variance over days (see Figure

3b).

Figure 3. Means and Standard Deviations of FCNef scores

(a) Mean daily FCNef scores. Investigative t-tests were conducted separately for each group to examine whether scores on each FCNef day were higher than a baseline score of 50 (which was the average score given during SHAM on Day 0). Scores improved significantly over FCNef days only for the Consec/High-Rew group. The Group Comparison inset shows the mean ± standard error of mean daily scores overlaid for the different groups. (b) Daily STDs in FCNef scores. Investigative t-tests were conducted separately for each group to examine whether variance in scores on each subsequent day of FCNef was lower than that on the Day 1 of FCNef. Score variance decreased significantly over FCNef days only for the Consec/High-Rew group, but a similar trend was found for the Consec/Low-Rew group. The Group Comparison inset shows the mean ± standard error of STD daily scores overlaid for the different groups; This highlights the Group x Day interaction found in the best-fit model for the Non-Consec/Low-Rew group relative to the reference Consec/High-Rew group. *** represents p = 0.001, ** represents $p_{FDR} < 0.01$, * represents $p_{FDR} < 0.05$, # represents $p_{FDR} < 0.1$, STD = standard deviation.

3.2.2. Self-report symptom scores

We analysed symptom questionnaire scores and compared these between the three groups as a secondary measure related to FCNef effects (Supplementary Table 10). We did not include symptom changes from before to after FCNef and into the long-term in our operationalisation of FCNef success because symptom reduction could occur for multiple reasons, including the placebo effect. Symptom reduction would be meaningful here only if specifically related to changes in the targeted brain activity.

Nonetheless, we monitored symptom changes to ensure they had not worsened over the course of the

study. Based on preliminary results [12], we predicted that symptoms should decrease from before to after

FCNef and then remain lower across post-days. Therefore, we used separate LME models to examine

initial and long-term effects.

3.2.2.1. Symptoms: The initial effect

For the best-fit models to predict both general depressive symptoms and brooding symptoms (see

Table 2 and Supplementary Tables 11 and 12), we found main effects of first/last FCNef day ($p < 0.0001$

for general depressive symptoms, $p = 0.0003$ for brooding symptoms), but no main effects of group ($ps >$

0.05). Follow-up t-tests showed that these symptoms were lower on day 4 than on day 0 ($t(67) = 4.78$,

$p_{FDR} < 0.0001$ for general depressive symptoms; $t(67) = 4.33$, $p_{FDR} < 0.0001$ for brooding symptoms). For

the best-fit models to predict anxiety symptoms (see Table 2 and Supplementary Table 13), we found a

significant main effect of first/last FCNef day ($p = 0.003$). A follow-up t-test showed that overall, anxiety

symptoms were lower on day 4 than on day 0 ($t(67) = 2.95$, $p_{FDR} = 0.007$). In this model there was also a

significant main effect of the Non-Consec/Low-Rew group ($p = 0.042$), and a trend for a main effect of

the Consec/Low-Rew group ($p = 0.0995$). Compared to the reference Consec/High-Rew group,

participants in the Non-Consec/Low-Rew group had higher anxiety and participants in the Consec/Low-

Rew group had lower anxiety (on Days 0 and 4) (see Supplementary Table 10). If anxiety has any effect

on overall FCNef success, then we should see deviations from the reference group in opposite directions

for these two groups, which we did not.

3.2.2.2. Symptoms: Changes in the long-term

Next, we examined whether any post-changes in symptoms differed or remained stable over the

long-term. For all symptom types, the best-fit model (see Table 2 and Supplementary Tables 14-17) had

no significant main effect of post-day or group ($ps > 0.05$). This may mean that after symptoms decreased

from day 0 to day 4, they remained at this new reduced level in the long-term. To examine this possibility,

we used t-tests to compare symptoms from follow-up tests with symptoms from day 0. Indeed, we found

that relative to symptoms from day 0, general depressive and brooding symptoms from the 1-month and

2-month follow-up tests were significantly reduced ($t(52) = 4.28$, $p_{FDR} = 0.0001$ for general depressive at

1-month, $t(52) = 4.28$, $p_{FDR} < 0.0001$ for brooding at 1-month, ($t(45) = 2.24$, $p_{FDR} = 0.015$ for general

depressive symptoms at 2-months, $t(45) = 5.60$, $p_{FDR} < 0.0001$ for brooding symptoms at 2-months). We

found a similar pattern, but non-significant, for anxiety symptoms ($t(52) = 4.28$, $p_{FDR} = 0.061$ at 1-month,

$t(45) = 0.92$, $p_{FDR} = 0.18$ at 2-months). Overall, these results show that general depressive and brooding

symptoms significantly decreased from before to after FCNef and remained at this new reduced level in

the long-term. Anxiety symptoms also decreased significantly from before to after FCNef and remained

in this same direction in the long-term, albeit not significantly. There was no persuasive evidence that

group, and therefore the manipulated parameters, influenced these results.

3.2.3. Resting-state functional connectivity

If FCNef is successful, then we predict that participant rs-FC should become more negative with

FCNef, which was the trained direction, and that it should remain so across the post-days. Therefore, we

used separate LME models to examine rs-FC initial and long-term effects for the three groups of

participants. Note that we measured rs-FC on all days (0-4) of the main experiment. However, we only

included data from days 0 and 4 in LME models to examine the initial effect. This is because rs-FC on

days 1-3 may be subject to homeostatic and/or compensatory mechanisms, causing overshoots [12] or

rebounds [34] in brain activity. Average rs-FC for each group for each day of experimentation can be seen

in Supplementary Table 18.

3.2.3.1. rs-FC: The initial effect

Likelihood ratio testing showed that models to predict rs-FC from days 0 and 4 did not differ

significantly, depending on whether an interaction term was included. Therefore, we selected the model

without the interaction term as best-fit, because it was simpler (see Table 2 and Supplementary Table 19).

This model showed a trend for a main effect of first/last FCNef day ($p = 0.094$), but no main effect of

group ($ps > 0.05$). Nonetheless, Figure 4 clearly shows that rs-FC on day 4 was more negative relative to

baseline for the Consec/High-Rew group, but not for the Non-Consec/Low-Rew group. rs-FC changes

from day 0 to day 4 for the Consec/Low-Rew group lay in between those of the other 2 groups (see Figure

4). Evidence for these between-group differences may be found at the LME model level if this experiment

is run with greater sample sizes; This is suggested by the fact that the model with the interaction term

trended toward being better than the model without, despite this not reaching full significance ($\chi2(2) =$

4.69, p = 0.096). The model with the interaction term had a significant first/last FCNef day x Non-

Consec/Low-Rew group interaction ($p = 0.036$) which indicates that the trajectory for rs-FC from day 0 to

day 4 differed for the Non-Consec/Low-Rew group compared to the reference Consec/Low-Rew group

(see Supplementary Table 20 and for visualisation of this see the Group Comparison inset in Figure 4).

3.2.3.2. rs-FC: Changes in the long-term

We next examined whether any post-changes in DLPFC-PCC rs-FC differed or remained stable

over time. In the best-fit model, which had no interaction term (see Table 2 and Supplementary Table 21),

no significant main effect of post-day was found. This indicates that changes from baseline remained

stable across post-days. A significant main effect of the Non-Consec/Low-Rew group was found ($p =$

0.007), indicating that post-changes in rs-FC differed for the Non-Consec/Low-Rew group and the

reference Consec/High-Rew group. Specifically, post-changes were more negative, which was the trained

direction, for the Consec/High-Rew group versus the Non-Consec/Low-Rew group (see Figure 4). Post-

changes for the Consec/Low-Rew group lay between those of the other two groups (see Figure 4).

Figure 4. Long-term changes in rs-FC.

Investigative t-tests revealed that for the Consec/High-Rew group, overall DLPFC-PCC resting state functional connectivity (rs-FC) was significantly more negative than baseline immediately after FCNef had been completed (D4) and one-month later (M1), and that it trended toward this even two-months later (M2). Nothing of significance was found for either of the other groups, although visually, the rs-FC appears to shift in the non-targeted direction (positive) for the Non-Consec/Low-Rew group. These results should be further explored using larger sample sizes for long-term data. Note that bars for each Group represent different sample sizes because not all participants came back for long-term testing (see Table 1). The Group Comparison inset shows the mean ± standard error of the change in rs-FC, over different post-days, overlaid for the different groups. Changes = data from each post-day minus data from Day 0; D = Day, M = Month, * represents $p_{FDR} < 0.05$, # represents $p_{FDR} = 0.051$.

3.2.4. Relationship between changes in self-report symptom scores and changes in rs-FC

Finally, in the three groups of participants, we examined how changes in participant DLPFC-PCC rs-FC from before to after FCNef related to changes in general depressive, brooding, and anxiety symptoms. Here, changes were defined as day 0 data subtracted from day 4 data. If the FCNef paradigm

was successful, then we would expect changes in the DLPFC-PCC rs-FC to be positively related to

changes in general depressive symptoms. This would mean that the more negative the rs-FC became, the

more depressive symptoms decreased. If, as previously hypothesised [12], the targeted functional

connection (FC) is specifically related to maladaptive symptoms of rumination, then we would expect

changes in the DLPFC-PCC rs-FC to be positively related to changes in brooding symptoms, but not to

changes in anxiety symptoms (which we used as a control). Importantly, by including Group as a factor in

the models, we examined whether these effects were impacted by manipulated parameters.

3.2.4.1. Relationships between symptom changes and rs-FC changes

The best-fit model to explain changes in general depressive scores (see Table 2 and

Supplementary Table 22) had a significant interaction between rs-FC change and group ($p = 0.0001$).

Follow-up correlations revealed a significant positive correlation between general depression change and

rs-FC change for the Consec/High-Rew group ($r = 0.66$, $p_{FDR} = 0.002$), a positive, but non-significant

relationship for the Consec/Low-Rew group ($r = 0.30$, $p_{FDR} = 0.13$), and a negative, non-significant

relationship for the Non-Consec/Low-Rew group ($r = -0.41$, $p_{FDR} = 0.98$).

The best-fit model to explain changes in brooding scores (see Table 2 and Supplementary Table

23) showed no significant main effects. However, hypothesis-driven follow-up correlations revealed a

significant positive correlation between brooding change and rs-FC change for the Consec/High-Rew

group ($r = 0.63$, $p_{FDR} = 0.003$), but nothing even approaching significance for the other two groups ($r =$

0.18, $p_{FDR} = 0.31$ for the Consec/Low-Rew group; $r = 0.05$, $p_{FDR} = 0.40$ for the Non-Consec/Low-Rew

group).

The best-fit model to explain changes in anxiety scores (see Table 2) had no significant main

effects or interactions (ps > 0.05; see Supplementary Table 22). Follow-up correlations also showed

nothing of significance ($r = 0.05$, $p_{FDR} = 0.63$ for the Consec/High-Rew group; $r = -0.08$, $p_{FDR} = 0.65$ for

the Consec/Low-Rew group; $r = 0.34$, $p_{FDR} = 0.63$ for the Non-Consec/Low-Rew group).

The relationship between symptom changes and rs-FC changes for all groups can be seen in

Figure 5. Data of one participant whose general depression change was more than 3 STD lower than the

group general depression change mean was excluded from the general depression change analysis (and

Figure 5a). Data of another participant whose brooding change was more than 3 STD lower than the

group mean was excluded from the brooding change analysis (and Figure 5b). This was because, although

inclusion of these participants did not change the results, it meant the data failed to meet a crucial

assumption underlying LME models (normal distribution of the residuals).

3.2.4.2. Specificity of the targeted functional connection

We specifically hypothesised that changes in DLPFC-PCC rs-FC from before to after FCNef should relate to changes in brooding, but not anxiety symptoms. Z-tests to compare correlation coefficients [33], run with one tail due to our directional hypothesis, demonstrated a significant difference for the Consec/High-Rew group ($z = 2.08$, $p = 0.019$), but not for the other groups ($z = 0.83$, $p = 0.21$ for the Consec/Low-Rew group; $z = 0.11$, $p = 0.46$ for the Non-Consec/Low-Rew group).

Figure 5. How changes in rs-FC relate to changes in symptomatology.

*Changes in DLPFC-PCC resting state functional connectivity (rs-FC) are plotted against **(a)** changes in General Depression Scores, **(b)** changes in Brooding scores, and **(c)** changes in Anxiety scores. Overall, promisingly for precision medicine, the Consec/High-Rew group showed significant positive correlations between changes in rs-FC and changes in **related** (General Depression and Brooding), but not unrelated (Anxiety) symptoms. These effects were in the same direction (reaching a trend with the uncorrected p-value for the General Depression change correlation) for the Consec/Low-Rew group. On the contrary, promising results were not found for the Non-Consec/Low-Rew group, for whom the relationship between rs-FC changes and General Depression score changes were actually numerically in the opposite direction to expectation. ** represents $p_{FDR} < 0.01$, # represents $p_{UNCORR} = 0.085$.*

Expected Effects	Consec/ High-Rew	Consec/ Low-Rew	Non-Consec/ Low-Rew
Increase in mean FCNef score	d = -0.58	d = -0.09	d = -0.31
Decrease in FCNef score STD	d = 0.72	d = 0.42	d = -0.19
Normalization of rs-FC	d = 0.47	d = 0.30	d = -0.11
Positive relationship between BDI changes and rs-FC changes	r = 0.66	r = 0.30	r = -0.41
Positive relationship between Brooding changes and rs-FC changes	r = 0.62	r = 0.21	r = 0.05

Large	Medium	Small	Trivial	Trivial	Small	Medium	Large

Hypothesized direction　　　Non-hypothesized direction

Table 3: Effect sizes summarized.

"Expected effects" are effects that we expect to see if FCNef training is successful. Comparisons for "Increase in

mean FCNef scores" and "Decrease in FCNef score STD" were t-tests comparing data from the first and the last

day of FCNef (Days 1 and 4). Comparisons for "Normalization of rs-FC" were t-tests comparing rs-FC from before

to after FCNef (Days 0 and 4). The two "Positive relationship…" comparisons were correlations calculated

between changes in rs-FC and changes in symptoms, where "changes" were defined as Day 4 - Day 0 data. Effect

sizes for each comparison are shown in the relevant cell. These are shown here as Cohen's d (d) for t-tests and

Pearson's correlation coefficient (r) for correlations. Following convention, these can be described on a scale

ranging from trivial (d < |0.2| or r < |0.1|), small (0.2 ≤ |d| < 0.5 or |0.1| ≤ r < |0.3|), medium (|0.5| ≤ d < |0.8| or

|0.3| ≤ r < |0.5|), to large (d ≥ |0.8| or r ≥ |0.5|). Effects in the hypothesized direction are shown in shades of blue

and effects in the non-hypothesized direction are shown in shades of orange. The larger the effect size, the stronger

the shade of the colour in the relevant cell.

4. Discussion

Overall, we found that the more participant DLPFC-PCC rs-FCs normalised over consecutive

days of Functional Connectivity Neurofeedback (FCNef), the greater their corresponding decrease in

brooding rumination, which we expected to relate to this functional connection. No such correlation was

shown for changes in DLPFC-PCC rs-FC and anxiety, which is thought to relate to different underlying

neural circuitry [35,36]. Importantly, here we found these results in new participants (run with

consecutive days of FCNef to be consistent with our past report). Combining this newly collected data

with the previously reported data provided us with sufficient statistical power for direct comparison of

these correlation coefficients and a significant difference was found. When we looked at data from three

groups of participants (see Table 3), we found the most promising results for the group run with a higher-

reward schedule to reinforce the targeted shift in functional connectivity, and an experimental schedule

with consecutive days. Weaker results were found for the group of participants that completed FCNef

over consecutive days, but who were reinforced with a lower reward schedule. FCNef did not seem to

have any effect on the group of participants who attended over non-consecutive days and that were

reinforced with a lower reward schedule.

4.1. FCNef for precision medicine.

The current results strengthen our previous finding that when FCNef is run over consecutive

days, normalisation of a target functional connection relates to a specific reduction in only related

symptoms (see Figure 2). This replication of previous results with an independent sample of newly

collected data shows their robustness and indicates that previous results were unlikely to have been

spurious [12]. Furthermore, the finding that changes in the targeted rs-FC correlated significantly more

with changes in related than unrelated symptoms highlights the precision of the FCNef technique. This

brings us one step closer to a future in which patients may one day enter a clinic, have their brains

scanned, and then have targeted treatment to normalise specific neural aberrations related to their own

subset of symptoms.

4.2. The DLPFC-PCC functional connection and rumination.

Not only do our results have implications for precision medicine in general, but also specifically

for understanding and treating brooding rumination symptoms. We previously hypothesised, based on

both data- and hypothesis-driven evidence from past studies, that the DLPFC-PCC functional connection

is likely to relate to brooding rumination [11]. Our initial study found preliminary support consistent with

this proposal. Specifically, we found that changes in connectivity between these regions relate to changes

in brooding rumination [12]. There has since been at least one other report that also showed increased

connectivity between the PCC and PFC related to ruminative symptoms [37]. We here find even stronger

support that brooding rumination relates to the DLPFC-PCC functional connection, by showing that the

correlation between changes in these is significantly stronger than the correlation between changes in the

same functional connection and a type of symptom thought to be unrelated (anxiety). Evidence from

previous studies using repetitive transcranial magnetic stimulation to target the DLPFC [38,39] and using

real-time neurofeedback to target the PCC [40], have also shown promising results for amelioration of

depressive and brooding rumination symptoms. This could be because targeting these regions affects the

functional connection between them, which if true, could mean that targeting the functional connection

itself more directly (as we did here) could be of even further advantage.

4.3. Parameter testing.

The current report extends our previous report by clarifying some of the parameters under which

FCNef for depression can best be achieved. These results should help guide the design of future

neurofeedback and other BMI studies. When selecting parameters for neurofeedback, past studies have

tended to follow convention or have gone with what seemed best in terms of cost/benefit trade-offs and/or

in terms of making things easy for participants. Often, this is the only feasible way to design a BMI study

because the cost of testing all possible parameters is enormous. However, our current results show that

certain parameters can make a difference in BMI effectiveness. This means that without knowing the

optimal parameters for a given BMI design, researchers may be finding null results simply because they

are not running their designs in the most effective way. There is no simple solution to this problem,

especially because optimal parameters may differ for different populations, target neural activities,

experimental goals, etc. Nonetheless, the current results provide initial evidence that can be used to help

future designs. Below, we discuss specific results for these parameter analyses, as well as their

implications.

Reward schedule. Participants run with the high-reward schedule had better FCNef success than

those run with the low-reward schedule (see Table 3 and Figures 3, 4, and 5). These results support the

proposal that external reward might work as reinforcement that is additional to that provided by feedback

scores during neurofeedback [41]. Based on these results, we recommend using liberal external reward in

future neurofeedback studies. Furthermore, because BMIs generally do not use external rewards for

reinforcement, this result might be worthy of consideration beyond the realm of neurofeedback.

Of course, disturbances to reward circuitry and disturbances in reward processing (usually

reductions) are commonly reported in depressive and other psychiatric disorders [42–49]. This means that

the effect of external reward on reinforcement of the target neural activity might be diminished when

neurofeedback is conducted in patients with such disorders. Here, our results with subclinical patients did

not corroborate this, but it remains worthy of further investigation in clinically depressed patients.

Experimental schedule. FCNef appears to be more effective when participants come for
consecutive, as opposed to non-consecutive, days of FCNef. All expected effects were strongest in the
consecutive condition with high reward, in the same direction (albeit with less strength) in the consecutive
condition with low reward, and weak or in the opposite direction for the non-consecutive condition (see
Table 3). Of course, the non-consecutive condition that we tested was with low reward and it would have
been nice to fully balance this by also testing the non-consecutive condition with high. reward. While
these results are therefore not fully conclusive, they do not appear promising for using non-consecutive
days of FCNef. They indicate that, although possibly more tiring for participants, consecutive days of
FCNef may be necessary to achieve positive outcomes. It is possible that consecutive days of
reinforcement are needed to drive learning effectively and/or that more non-controllable confounding
personal circumstances can occur between non-consecutive days of training (an idea that researchers
designing future BMIs and clinical treatments ought to consider).

Another point to consider is the possibility that neural plasticity related to learning might cause
dynamic rs-FC fluctuations in strength and direction that occur before settling into a new pattern (similar
to the rebound effect first documented by Kluetsch et al., 2014 [34]). If so, then our analyses may not
have fairly tested consecutive versus non-consecutive conditions. Learning could begin from Day 1; but,
post-FCNef measurements (from Day 4, and 1- and 2- months later) differed in the number of
days after Day 1 for consecutive/non-consecutive conditions (and even for different participants in the
non-consecutive condition). This may mean that measurements were taken at different
points during ongoing dynamic rs-FC fluctuations for these different conditions. If so then comparisons
between these conditions may actually have compared different snapshots of learning effects. Other
researchers using neurofeedback over non-consecutive days should also consider this possibility.

4.4. Limitations of the current design.

One limitation of our study is that it involved participants who had only subclinical levels of
depression. Nonetheless, preliminary studies using this FCNef technique with patients diagnosed with
MDD have shown promise [19,21]. Results with clinical patients may be improved if the right parameters
are employed. The most effective parameter that we found was high(er) monetary reward. We used
money because we wanted to test the effects of reward schedule with a type of reward that is well known
to strongly activate the human reward system. However, now that proof-of-concept has been provided,
this idea should be further tested more creatively with other types of reward that might be more
appropriate for a clinical setting. For example, revealing consecutive puzzle pieces for each successful

trial of neuromodulation (see Ramot et al., 2017 [16]). A second limitation of our study is the absence of a

control group or within-subject control condition. Therefore, it is possible that our target rs-FC changed

and that symptoms improved for reasons such as the placebo or Hawthorne effects. However, only

changes in relevant (depressive and brooding rumination, but not anxiety) symptoms changed parallel to

the targeted FC, which would be unlikely to occur merely from such nonspecific effects. A third

limitation of our study is the incomplete factorial design due to practical constraints, as discussed in the

Experimental Conditions section. While this limits our ability to examine potential interaction effects

between the experimental schedule and reward schedule parameters, the current design still allows us to

investigate main effects of these parameters separately, which was our main aim. Future research should

include all factorial combinations to provide a comprehensive understanding of parameter interactions.

4.5. Future directions.

Although different symptoms are likely to arise from aberrations in wider brain networks

involving multiple FCs, our current FCNef approach can directly target only one FC. In that sense,

connectome-based FCNef [50] or neurofeedback targeting an estimation of the dynamic weighted linear

sum of FCs might be more effective. Some promise for such types of neurofeedback has been found,

including when targeting an estimation of the dynamic weighted linear sum of FCs from the greater

biomarker from which we identified the DLPFC-PCC FC [T. Ogawa , personal communication, 27[th]

January, 2025; 51]. However, the authors have also reported increased difficulty for participants with

regard to the credit assignment (they report that it is difficult to target multiple functional connections and

to know what actually worked) and overall experimental interpretability [T. Ogawa, personal

communication, 27[th] January, 2025]. Furthermore, FCNef itself affects broader brain networks than just

the targeted functional connection anyway [15,16], so it remains possible that our simple FCNef approach

might be best at ameliorating symptoms without the need for added complexity and burden for patients.

Future studies should attempt to directly compare effectiveness of these types of neurofeedback.

Our current results are promising for precision medicine, but they were shown with functional

connectivity in fMRI, which can be costly and impractical (but not impossible) for real clinical treatment.

In the future, we expect FCNef to evolve further so that it may be conducted using electroencephalogram

(EEG) signatures (see Keynan et al., 2019 [52]) of target FCs or so that it may be conducted using EEG

signatures of weighted linear sums of multiple FCs. If successful, then this would allow neurofeedback

targeting functional connections to eventually be conducted with portable EEG headsets, possibly even

away from the clinic in the privacy of the patient's own home (for more detailed discussion see Taylor et

al., 2021 [53]).

4.6. Conclusion

Overall, we have extended previous results to show that normalisation of the targeted neural

network (DLPFC-PCC) correlated significantly more with reductions in symptoms thought to relate to

this neural circuitry (brooding rumination) than to changes in symptoms thought to relate to different

neural circuitry (anxiety). This highlights the precision of the FCNef technique and brings us one step

closer to a future where psychiatric treatment might be tailored to the individual patient. Here, we

additionally extended our previous work by investigating parameters under which our FCNef for

depression paradigm is most effective. We found that FCNef effectiveness changes depending on those

parameters with which it was run. Specifics and implications of some parameter-related results may be

relevant beyond neurofeedback to BMIs in general. Furthermore, some of our results highlight benefits of

testing conventional parameters. Overall, these results should be informative for design of future BMI

testing and for inspiring new interpretations of existing data. For example, previously found null results

should be considered in the context of the parameters under which the BMI was run. More broadly, by

documenting how parameter optimisation can increase beneficial outcomes and reduce patient burden, we

hope to inspire more of this in the future, with the ultimate goal of bringing optimised BMIs to the

medical clinic.

Acknowledgements

We thank Kaori Nakamura for help in scheduling and conducting the experiments, Rumi Yorizawa for

help in screening participants, and Toshinori Yoshioka for continued support with regard to experimental

scripts.

Statement of ethics

Participants all provided written informed consent on each day of screening and experimentation prior to

commencement. This research was approved by the Ethics Committee of the Review Board of Advanced

Telecommunications Research Institute International, Japan (Ethics No. 132, 172) and by the Kyoto

University Certified Review Board (YC0849) and the Committee on Medical Ethics of Kyoto University

(C0849). All experiments were performed in accordance with the guidelines and regulations of these

Ethics Committees. This research was conducted in association with research registered with the Japan

Registry of Clinical Trials (jRCTs052180169) and the University Hospital Medical Information Network

Clinical Trials Registry (UMIN000015249).

711   Conflict of interest statement

712   MK is an inventor of patents related to functional connectivity neurofeedback. The original assignee of

713   the patents is ATR, with which the authors are affiliated. We have no other conflicts of interest.

714   Funding Sources

715   This research was supported by the Japan Agency for Medical Research and Development (AMED) under

716   Grant Numbers JP18dm0307008, JP21dm0307102 and JP17dm0107044, by the Japan Society for the

717   Promotion of Science (JSPS) KAKENHI under Grant Number 24K22822, and by the Innovative Science

718   and Technology Initiative under Security Grant Number JPJ004596, ATLA, Japan.

719   Author Contributions

720   JET, TM, TY and MK designed experiments; JET, TO, MM and TM acquired data; JET analysed

721   data; JET prepared the original draft; TO, MM, MT, TY, TK, YK, YY, JM, TM, MK, and AC reviewed

722   and edited the manuscript. All authors gave final approval for submission and agreed to take

723   responsibility for the manuscript.

724   Data accessibility

725   Data and code supporting this study's findings will be publicly available on our GitHub at publication.

References

Patel V, Chisholm D, Parikh R, Charlson FJ, Degenhardt L, Dua T, et al. Addressing the burden of mental, neurological, and substance use disorders: key messages from Disease Control Priorities, 3rd edition. Lancet. 2016 Apr;387(10028):1672–85.

Fava M, Davidson KG. Definition and epidemiology of treatment-resistant depression. Psychiatr Clin North Am. 1996 Jun;19(2):179–200.

McIntyre RS, Alsuwaidan M, Baune BT, Berk M, Demyttenaere K, Goldberg JF, et al. Treatment-resistant depression: definition, prevalence, detection, management, and investigational interventions. World Psychiatry. 2023 Oct;22(3):394–412.

Herrman H, Patel V, Kieling C, Berk M, Buchweitz C, Cuijpers P, et al. Time for united action on depression: a Lancet–World Psychiatric Association Commission. Lancet. 2022 Mar;399(10328):957–1022.

Gabriel FC, de Melo DO, Fráguas R, Leite-Santos NC, Mantovani da Silva RA, Ribeiro E. Pharmacological treatment of depression: A systematic review comparing clinical practice guideline recommendations. PLoS One. 2020 Apr;15(4):e0231700.

Schalk G, Brunner P, Allison BZ, Soekadar SR, Guan C, Denison T, et al. Translation of neurotechnologies. Nat Rev Bioeng. 2024 May;2(8):637–52.

The tech: Restore [Internet]. Synchron. 2021 Dec [cited 2023 Oct 12]. Available from: https://synchron.com/technology/brain-io

Neuralink announces first-in-human clinical trial of BCI device [Internet]. [cited 2023 Oct 12]. Available from: https://www.fdanews.com/articles/211989-neuralink-announces-first-in-human-clinical-trial-of-bci-device?v=preview

Paradromics. Paradromics raises $33 million in funding, achieves breakthrough medical device designation from FDA [Internet]. PR Newswire. 2023 May [cited 2023 Oct 12]. Available from: https://www.prnewswire.com/news-releases/paradromics-raises-33-million-in-funding-achieves-breakthrough-medical-device-designation-from-fda-301827969.html?tc=eml_cleartime

10 Neurolutions Receives FDA De Novo Market Authorization for IpsiHand^{TM} [Internet]. Neurolutions.
2021 Apr [cited 2023 Oct 12]. Available from: https://www.neurolutions.com/post/neurolutions-
receives-u-s-food-and-drug-administration-de-novo-market-authorization-for-ipsihand

11 Rustamov N, Souders L, Sheehan L, Carter A, Leuthardt EC. IpsiHand Brain-Computer Interface
Therapy Induces Broad Upper Extremity Motor Recovery in Chronic Stroke. medRxiv. 2023 Aug
DOI: 10.1101/2023.08.26.23294320

12 Taylor JE, Yamada T, Kawashima T, Kobayashi Y, Yoshihara Y, Miyata J, et al. Depressive
symptoms reduce when dorsolateral prefrontal cortex-precuneus connectivity normalizes after
functional connectivity neurofeedback. Sci Rep. 2022 Feb;12(1):2581.

13 Koush Y, Rosa MJ, Robineau F, Heinen K, W Rieger S, Weiskopf N, et al. Connectivity-based
neurofeedback: dynamic causal modeling for real-time fMRI. Neuroimage. 2013 Nov;81:422–30.

14 Koush Y, Meskaldji D-E, Pichon S, Rey G, Rieger SW, Linden DEJ, et al. Learning Control Over
Emotion Networks Through Connectivity-Based Neurofeedback. Cereb Cortex. 2017
Feb;27(2):1193–202.

15 Megumi F, Yamashita A, Kawato M, Imamizu H. Functional MRI neurofeedback training on
connectivity between two regions induces long-lasting changes in intrinsic functional network. Front
Hum Neurosci. 2015 Mar;9:160.

16 Ramot M, Kimmich S, Gonzalez-Castillo J, Roopchansingh V, Popal H, White E, et al. Direct
modulation of aberrant brain network connectivity through real-time NeuroFeedback. Elife. 2017
Sep;6. DOI: 10.7554/eLife.28974

17 Tsuchiyagaito A, Misaki M, Zoubi OA, Tulsa 1000 Investigators, Paulus M, Bodurka J. Prevent
breaking bad: A proof of concept study of rebalancing the brain's rumination circuit with real-time
fMRI functional connectivity neurofeedback. Hum Brain Mapp. 2021 Mar;42(4):922–40.

18 Tsuchiyagaito A, Misaki M, Kirlic N, Yu X, Sánchez SM, Cochran G, et al. Real-Time fMRI
Functional Connectivity Neurofeedback Reducing Repetitive Negative Thinking in Depression: A
Double-Blind, Randomized, Sham-Controlled Proof-of-Concept Trial. Psychother Psychosom. 2023
Jan;92(2):87–100.

19   Yamada T, Hashimoto R-I, Yahata N, Ichikawa N, Yoshihara Y, Okamoto Y, et al. Resting-State
Functional Connectivity-Based Biomarkers and Functional MRI-Based Neurofeedback for
Psychiatric Disorders: A Challenge for Developing Theranostic Biomarkers. Int J
Neuropsychopharmacol. 2017 Oct;20(10):769–81.

20   Zhao Z, Yao S, Li K, Sindermann C, Zhou F, Zhao W, et al. Real-time functional connectivity-based
neurofeedback of amygdala-frontal pathways reduces anxiety. bioRxiv. 2018 Apr;308924.

21   Takamura M, Okada G, Kamishikiryo T, Itai E, Kato M, Motegi T, et al. Application of functional
connectivity neurofeedback in patients with treatment-resistant depression: A preliminary report.
Journal of Affective Disorders Reports. 2023 Dec;14:100644.

22   deCharms RC, Christoff K, Glover GH, Pauly JM, Whitfield S, Gabrieli JDE. Learned regulation of
spatially localized brain activation using real-time fMRI. Neuroimage. 2004 Jan;21(1):436–43.

23   deCharms RC, Maeda F, Glover GH, Ludlow D, Pauly JM, Soneji D, et al. Control over brain
activation and pain learned by using real-time functional MRI. Proc Natl Acad Sci U S A. 2005
Dec;102(51):18626–31.

24   Scharnowski F, Rosa MJ, Golestani N, Hutton C, Josephs O, Weiskopf N, et al. Connectivity
changes underlying neurofeedback training of visual cortex activity. PLoS One. 2014
Mar;9(3):e91090.

25   Sepulveda P, Sitaram R, Rana M, Montalba C, Tejos C, Ruiz S. How feedback, motor imagery, and
reward influence brain self-regulation using real-time fMRI. Hum Brain Mapp. 2016
Sep;37(9):3153–71.

26   Grahek I, Shenhav A, Musslick S, Krebs RM, Koster EHW. Motivation and cognitive control in
depression. Neurosci Biobehav Rev. 2019 Jul;102:371–81.

27   Beck, A.T., Steer, R.A., & Brown, G.K. Beck Depression Inventory (BDI-II): Manual and
Questionnaire. The Psychological Corporation.; 1996.

28   Treynor W, Gonzalez R, Nolen-Hoeksema S. Rumination Reconsidered: A Psychometric Analysis.
Cognit Ther Res. 2003 Jun;27(3):247–59.

29   Hasegawa A. Translation and initial validation of the Japanese version of the Ruminative Responses
Scale. Psychol Rep. 2013 Jun;112(3):716–26.

30   Spielberger CD, Gorsuch RL, Lushene R, Vagg PR. Manual for the State-Trait Anxiety Inventory;
Palo Alto, CA, Ed. Palo Alto: Spielberger. 1983

31   Akaike H. Information theory and an extension of the maximum likelihood principle. In: Csáki
BNP&., editor. 2nd international symposium on information theory. Akadémia Kiadó.; 1973; pp
267–81.

32   Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful approach
to multiple testing. J R Stat Soc. 1995 Jan;57(1):289–300.

33   Takeuchi RF. Comparison test of two correlation coefficient: corr_rtest(ra, rb, na, nb).
2023Available from: https://au.mathworks.com/matlabcentral/fileexchange/61398-comparison-test-
of-two-correlation-coefficient-corr_rtest-ra-rb-na-nb

34   Kluetsch RC, Ros T, Théberge J, Frewen PA, Calhoun VD, Schmahl C, et al. Plastic modulation of
PTSD resting-state networks and subjective wellbeing by EEG neurofeedback. Acta Psychiatr
Scand. 2014 Aug;130(2):123–36.

35   Williams LM. Precision psychiatry: a neural circuit taxonomy for depression and anxiety. Lancet
Psychiatry. 2016 May;3(5):472–80.

36   Williams LM. Defining biotypes for depression and anxiety based on large-scale circuit dysfunction:
a theoretical review of the evidence and future directions for clinical translation. Depress Anxiety.
2017 Jan;34(1):9–24.

37   Mısır E, Alıcı YH, Kocak OM. Functional connectivity in rumination: a systematic review of
magnetic resonance imaging studies. J Clin Exp Neuropsychol. 2023 Nov;45(9):928–55.

38   Chu SA, Tadayonnejad R, Corlier J, Wilson AC, Citrenbaum C, Leuchter AF. Rumination
symptoms in treatment-resistant major depressive disorder, and outcomes of repetitive Transcranial
Magnetic Stimulation (rTMS) treatment. Transl Psychiatry. 2023 Sep;13(1):293.

39   Richieri R, Jouvenoz D, Verger A, Fiat P, Boyer L, Lançon C, et al. Changes in dorsolateral

prefrontal connectivity after rTMS in treatment-resistant depression: a brain perfusion SPECT study.
Eur J Nucl Med Mol Imaging. 2017 Jun;44(6):1051–5.

40 Kirlic N, Cohen ZP, Tsuchiyagaito A, Misaki M, McDermott TJ, Aupperle RL, et al. Self-regulation
of the posterior cingulate cortex with real-time fMRI neurofeedback augmented mindfulness training
in healthy adolescents: A nonrandomized feasibility study. Cogn Affect Behav Neurosci. 2022
Aug;22(4):849–67.

41 Watanabe T, Sasaki Y, Shibata K, Kawato M. Advances in fMRI Real-Time Neurofeedback. Trends
Cogn Sci. 2017 Dec;21(12):997–1010.

42 Nestler EJ, Carlezon WA Jr. The mesolimbic dopamine reward circuit in depression. Biol
Psychiatry. 2006 Jun;59(12):1151–9.

43 Russo SJ, Nestler EJ. The brain reward circuitry in mood disorders. Nat Rev Neurosci. 2013
Sep;14(9):609–25.

44 Zhang W-N, Chang S-H, Guo L-Y, Zhang K-L, Wang J. The neural correlates of reward-related
processing in major depressive disorder: a meta-analysis of functional magnetic resonance imaging
studies. J Affect Disord. 2013 Nov;151(2):531–9.

45 Baskin-Sommers AR, Foti D. Abnormal reward functioning across substance use disorders and
major depressive disorder: Considering reward as a transdiagnostic mechanism. Int J Psychophysiol.
2015 Nov;98(2 Pt 2):227–39.

46 Whitton AE, Treadway MT, Pizzagalli DA. Reward processing dysfunction in major depression,
bipolar disorder and schizophrenia. Curr Opin Psychiatry. 2015 Jan;28(1):7–12.

47 Keren H, O'Callaghan G, Vidal-Ribas P, Buzzell GA, Brotman MA, Leibenluft E, et al. Reward
Processing in Depression: A Conceptual and Meta-Analytic Review Across fMRI and EEG Studies.
Am J Psychiatry. 2018 Nov;175(11):1111–20.

48 Knowland D, Lim BK. Circuit-based frameworks of depressive behaviors: The role of reward
circuitry and beyond. Pharmacol Biochem Behav. 2018 Nov;174:42–52.

49 Fox ME, Lobo MK. The molecular and cellular mechanisms of depression: a focus on reward

circuitry. Mol Psychiatry. 2019 Dec;24(12):1798–815.

50   Scheinoist D, Hsu TW, Avery EW, Hampson M, Constable, RT, Chun MM, Rosenberg MD.
Connectome-based neurofeedback: A pilot study to improve sustained attention. NeuroImage. 2020
May;15(212):116684.

51   Ichikawa N, Lisi G, Yahata N, Okada G, Takamura M, Hashimoto R-I, et al. Primary functional
brain connections associated with melancholic major depressive disorder and modulation by
antidepressants. Sci Rep. 2020 Feb;10(1):3542.

52   Keynan JN, Cohen A, Jackont G, Green N, Goldway N, Davidov A, et al. Electrical fingerprint of
the amygdala guides neurofeedback training for stress resilience. Nat Hum Behav. 2019 Jan;3(1):63–
73.

53   Jessica Elizabeth Taylor, Itamar Jalon, Toshinori Chiba, Tomokazu Motegi, Mitsuo Kawato, and
Talma Hendler. Chapter 12: Translation to the clinic and other modalities. In: Hampson M, editor.
fMRI Neurofeedback. Academic Press, Elsevier; 2021; pp 265–86.

54   Kirchner WK. Age differences in short-term retention of rapidly changing information. J Exp
Psychol. 1958 Apr;55(4):352–8.

Group	Main Experiment Sample Size	1-Month Follow-up Sample Size	2-Month Follow-up Sample Size
Consec/High-Rew	21	11	9
Consec/Low-Rew	23	20	17
Non/Low-Rew	24	22	20
Total	68	53	46

Table 1. Sample sizes for three groups of participants.

This table shows the sample sizes for: (1) participants run in consecutive days of FCNef with the high-reward schedule (the "Consec/High-Rew group"), (2) those run in consecutive days of FCNef with the low-reward schedule (the "Consec/Low-Rew group"), and (3) those run over non-consecutive days of FCNef with the low-reward schedule (the "Non-Consec/Low-Rew group"). No participants were run over non-consecutive days of FCNef with the high-reward schedule (no "Non-Consec/High-Rew" group). Sample sizes are shown for the main experiment and for one- and two-month follow-up tests. There were fewer participants for the follow-up tests for the Consec/High-Rew group because the long-term tests were not included in the earliest stages of this experiment, so only 12 participants from this group were invited back.

Analyses	Formulae				AICs	
	DV	IV1	IV2	Random Intercept	+	x
FCNef Scores	Mean FCNef Score	~ FCNef Day	+ or x Group	+ (1|Subject)	2109.2	2108.7
	STD FCNef Score	~ FCNef Day	+ or x Group	+ (1|Subject)	1599.8	**1593**
Symptoms	General Depressive Scores	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	776.72	779.13
	Brooding Scores	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	681.4	682.68
	Anxiety Scores	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	913.86	914.62
	General Depressive Score Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	926.03	925.45
	Brooding Score Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	766.01	769.92
	Anxiety Score Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	1095.7	1095
rs-FC	rs-FC	~ first/last FCNef Day	+ or x Group	+ (1|Subject)	-22.99	-23.67
	rs-FC Changes (D4, 1M, 2M)	~ Post-Day	+ or x Group	+ (1|Subject)	-58.77	-55.47
Symptom Change/	General Depressive Score Changes (D4)	~ rs-FC Change	+ or x Group	+ (1|Subject)	316.57	**-153.29**
rs-FC Change	Brooding Score Changes (D4)	~ rs-FC Change	+ or x Group	+ (1|Subject)	287.8	287.52
Relationship	Anxiety Score Changes (D4)	~ rs-FC Change	+ or x Group	+ (1|Subject)	408.46	410.45

Table 2. The models used to examine dependent variables from the FCNef experiment.

Models with and without interactions between independent variables (IVs) were compared to see which would best predict each dependent variable (DV). Akaike Information Criteria (AIC) are displayed in columns labelled '+' (for models without interactions) and 'x' (for models with interactions). These are highlighted in bold in cases in which likelihood ratio testing showed the corresponding model to be significantly better fit than the alternative model. If there was no significant difference, then the model without the interaction was selected as best-fit because it was the simplest. FCNef Day = Days 1-4; Subject = experimental participant; STD = standard deviation; First/last FCNef Day = Days 0 and 4; Post-Day = Day 4, and the 1- and 2-month follow-up test days; Changes = data from the day indicated in brackets (e.g. D4) minus data from Day 0; rs-FC = resting state functional connectivity between the DLPFC-PCC.

Expected Effects	Consec/ High-Rew	Consec/ Low-Rew	Non-Consec/ Low-Rew
Increase in mean FCNef score	$d = -0.58$	$d = -0.09$	$d = -0.31$
Decrease in FCNef score STD	$d = 0.72$	$d = 0.42$	$d = -0.19$
Normalization of rs-FC	$d - 0.47$	$d - 0.30$	$d - -0.11$
Positive relationship between BDI changes and rs-FC changes	$r = 0.66$	$r = 0.30$	$r = -0.41$
Positive relationship between Brooding changes and rs-FC changes	$r = 0.62$	$r = 0.21$	$r = 0.05$

Large	Medium	Small	Trivial	Trivial	Small	Medium	Large

Hypothesized direction   Non-hypothesized direction

Table 3: Effect sizes summarized.

"Expected effects" are effects that we expect to see if FCNef training is successful. Comparisons for "Increase in mean FCNef scores" and "Decrease in FCNef score STD" were t-tests comparing data from the first and the last day of FCNef (Days 1 and 4). Comparisons for "Normalization of rs-FC" were t-tests comparing rs-FC from before to after FCNef (Days 0 and 4). The two "Positive relationship..." comparisons were correlations calculated between changes in rs-FC and changes in symptoms, where "changes" were defined as Day 4 - Day 0 data. Effect sizes for each comparison are shown in the relevant cell. These are shown here as Cohen's d (d) for t-tests and Pearson's correlation coefficient (r) for correlations. Following convention, these can be described on a scale ranging from trivial ($d < |0.2|$ or $r < |0.1|$), small ($0.2 \leq |d| < 0.5$ or $|0.1| \leq r < |0.3|$), medium ($|0.5| \leq d < |0.8|$ or $|0.3| \leq r < |0.5|$), to large ($d \geq |0.8|$ or $r \geq |0.5|$). Effects in the hypothesized direction are shown in shades of blue and effects in the non-hypothesized direction are shown in shades of orange. The larger the effect size, the stronger the shade of the colour in the relevant cell.

a.

Localizer

Questionnaires Anatomical Resting state N-back task x 3
MRI measurements

Day 0 (SHAM)

SHAM task Resting state
MRI measurements
Questionnaires

Days 1-3

FCNef task Resting state
MRI measurements

Day 4

FCNef task Resting state
MRI measurements
Questionnaires

Follow-up Tests (1 & 2 month)

Resting state
MRI measurements
Questionnaires

Time within each session

b.

Left DLPFC

Left PCC

Correlation
↓
Score

Rest 14s   Induction 20/40s + 2s calculation   Feedback 6s

Trial