Work-related road traffic accidents: emergence of new modes of personal journey.

Analysis based on data from a register of road traffic accidents

Emmanuel Fort[a], Nicolas Connesson[a], Julien Brière[b], Amina Ndiaye[a], Blandine Gadegbeku[a],

Barbara Charbotel[a,c]

[a] Univ Lyon, Univ Gustave Eiffel, Univ Claude Bernard Lyon 1, UMRESTTE UMR T 9405, F

69622 Lyon, France.

emmanuel.fort@univ-lyon1.fr

nicolas.connesson@gmail.com

amina.ndiaye@univ-eiffel.fr

blandine.gadegbeku@univ-eiffel.fr

barbara.charbotel@univ-lyon1.fr

[b]Department of Environmental and Occupational Health, Santé Publique France, French

National Public Health Agency, Saint Maurice, France. julien.briere@santepubliquefrance.fr

[c]CRPPE de Lyon (Regional Center for Occupational and Environmental Pathologies), Hospices

Civils de Lyon, Centre Hospitalier Lyon Sud, Pierre-Bénite, France.

Corresponding author: Emmanuel Fort

Univ Lyon, Univ Gustave Eiffel, Univ Claude Bernard Lyon 1, UMRESTTE UMR T 9405, F 69622

Lyon, France emmanuel.fort@univ-lyon1.fr

Acknowledgments

We would like to thank all those involved in data collection and computerization, through

the Association pour le registre des victimes d'accidents de la circulation du Rhône (Arvac,

chaired by Étienne Javouhey) and the Université Gustave-Eiffel-TS2-Umrestte (under the

direction of Barbara Charbotel).

Funding

This work was supported by The French National Public Health Agency (Santé publique
France), and Carsat Rhône Alpes.

Declaration of Interest(s)

None

Data Statement

Data is unavailable to access to post.

Author Contributions

Emmanuel Fort: Conceptualization, Methodology, Software, Formal analysis, Writing - Original Draft

Nicolas Connesson: Writing - Review & Editing. Julien Brière: Conceptualization, Methodology,

Visualization, Investigation, Writing - Review & Editing. Amina Ndiaye: Writing - Review & Editing.

Blandine Gadegbeku: Conceptualization, Methodology, Writing - Review & Editing, Validation.:

Barbara Charbotel: Conceptualization, Funding acquisition, Writing- Reviewing and Editing,

Word count : 4,553

Abstract

Introduction: According to the 2018–2019 People Mobility Survey, work-related journeys (commuting and on-duty journeys) account for approximately 25% of all journeys. The use of non-motorized (nm) and motorized (m) personal mobility devices (PMDs) has steadily increased since their introduction into the French market in the last decade.

Objective: This study aimed to describe the characteristics of work-related road accidents and their evolution since the introduction of new PMDs in France.

Materials and methods: This was a retrospective, cross-sectional study using data from the Rhône Road Trauma Registry. Data were collected from 2015 to 2020. We included the data for the victims aged 18–70 years who were injured in work-related accidents.

Results: We identified 11,296 individuals aged 18–70 years who experienced work-related road accidents. An injury report was provided for a total of 11,277 patients. A total of 546 passengers and 78 drivers of other motorized vehicles (buses/trams, construction equipment, and tractors) were excluded from the analysis. Seven patients died at the time of the accident, and seven died after hospitalization. Of the 10,653 (94.4%) victims, there were pedestrians (5.1%), or drivers of bicycles (16.9%), scooters (3.8%), other PMDs (roller blades, skateboards, monowheels, gyropods, and hoverboards; 0.4%), motorized two-wheeler (21.4%), car (45.3%), and truck (1.5%). More than half of the scooter drivers and 80% of other PMD-drivers were men. More than 60% of other PMD-drivers and 53% of scooter drivers were under 34 years of age. Most scooter accidents occurred during commuting (95.6%). Sixty-five percent of the scooter- and 50% of other PMD-accidents did not have

antagonists. Overall, one-quarter of the victims experienced accidents without antagonists.

Most scooter users had injuries to their upper limbs (59.2%), lower limbs (46.8%), face

(21.2%) or head (17.9%).

Discussion: Most accidents were of low or moderate severity (97.5%; Maximum Abbreviated

Injury Scale, MAIS score < 3). The frequency and severity of injuries among scooter- and

other PMD users were similar. Scooter antagonists were rarely observed in pedestrian

accidents (12/575). Most scooter- and other PMD-accidents did not have antagonists.

Conclusion: Many head injuries could be prevented with more widespread use of helmets,

among scooter- and other PMD users and bicycle users.

Keywords: e-scooter; personal mobility device; mobility; road accident; work-related road

accident; injuries; regulations

What is already known on this topic

• The use of personal mobility devices (PMDs) has steadily increased in France.

• Work-related journeys (commuting and on-duty journeys) account for approximately 25% of

all journeys

What this study adds

• We describe work-related road accidents' characteristics since PMDs' introduction.

• Accidents involving scooters or other PMDs are of low severity.

How this study might affect research, practice or policy

• Many head injuries could be prevented with more widespread helmet use.

• Companies can take preventive actions to ensure that employees are better informed.

1. Introduction

While a large proportion of travel in France involves private journeys (leisure or other), work-related journeys (home-work and on-duty) account for approximately 25% according to the 2018–2019 People Mobility Survey [1]. The main modes of transportation for commuting journeys were cars (72%), public transport (12%), walking (9%), cycling (3%), and motorized two-wheelers (2%); other modes accounted for less than 1%. According to the National Transport and Mobility Survey, work-related journeys accounted for approximately 27% of all journeys in 2008 [2]. The 2016 National Survey of Environmental Behavior [3] showed that 25% of journeys were mainly for commuting. According to the Sumer surveys in France, 25% of employees drive on public roads as part of their work (Fort et al., 2016, 2019). This proportion has remained constant since the surveys were conducted (2003, 2010, and 2017).

The mode of transport to reach work depends on several factors. The distance to work, availability of transportation, car ownership, household size, parking, and cost are determining factors in choosing the travel mode(s). In addition, there is a significant difference between the types of territory; walking, cycling, and other soft modes of transport are preferred in Paris and its suburbs, as well as in large cities and urban areas (those with 100,000–10,000,000 inhabitants).

The use of non-motorized and motorized personal mobility devices (nmPMDs and mPMDs, respectively) has been steadily increasing since their introduction into the market in France in the last decade, as well as in other cities globally [6]. These modes of transport can be referred to as micromobility that combines all easy modes of transport that allow their users to make hybrid use of the vehicle either as a pedestrian or as a vehicle occupant [7]. It

represents between 8–15% of daily journeys of less than 8 km, which today account for 50–

60% of the distance traveled in China, the European Union, and the USA, and replaces 20%

of public transport journeys (in addition to bridging the gap between the first and last

kilometers) as well as bicycle, moped, scooter, or on-foot journeys [8].

The introduction of e-scooter sharing (in June 2018 in France) has accelerated this

phenomenon. It has been observed in all industrialized countries, mainly in large

metropolises [9]. The development of PMDs is growing, primarily in urban areas, for several

reasons. First, their use could replace motorized vehicles, such as cars or motorized two-

wheelers. Second, they reduce air pollution; therefore, their use is favored in cities with

reserved lanes (soft mode). In fact, it appears that most walking and public transport

journeys are replaced by trips using PMDs [10]. Electric scooters replace walking or biking for

the last mile of a journey [11], potentially reducing the overall physical activity of the

population [12]. The introduction of large low-emission zones (ZFE), where vehicle speeds

are limited to 30 km/h, is a new incentive to use PMDs. The COVID-19 pandemic has

increased the use of PMDs, particularly by encouraging social distancing [13]. Thus, public

transport users have abandoned subways, tramways, or buses, where distances between

people can be too small, for the use of PMDs. Finally, the increase in the cost of fossil energy

favors the development of PMDs. However, some authors have highlighted problems caused

by electric scooters, both regarding the safety of other vulnerable users and the impact on

the environment during the production of scooters, especially batteries, which cause

pollution [14].

In Vienna, Austria, Laa and Leth [15] showed that users were shifting to electric scooters,

followed by buses and street cars, to replace walking. Electric scooters are used in

combination with other modes of transport. A study from the USA on users of the shared-

bike service in Washington, D.C. found that 35% of occasional users reported using it as a

substitute for public transport [16]. In Belgium, nearly 50% of customers reported replacing

one or more modes of transport with electric scooters [17].

PMDs now represent a significant part of urban and peri-urban transport modes. In France,

22% of respondents have used them at least once, and 11% are regular users (at least once a

month; [18]). The electric scooters is the most-used mPMD, followed to a lesser extent by

the gyro-mot, electric skateboard, gyropod, and hoverboard. One of two daily users of

mPMD uses a self-service electric scooter. In Greece, 40% of the users mentioned using

electric scooters for work purposes [9]. These electric vehicles offer new opportunities to

users in terms of travel speed and less effort than their non-electric counterparts; however,

they cause problems with public space sharing and road safety [19].

Many regulations have been implemented and experiments have been conducted in cities

and countries to better control the use of electric scooters [6] and reduce the injuries

associated with them, such as restrictions on hiring and using electric scooters at night [20],

wearing helmets, adhering to speed limits, and traffic zones [6].

Many studies in the USA [21], the United Kingdom [22,23], Spain [24], Finland, [25] and

France [26] have found an increase in injuries associated with the use of electric-sharing

scooters [27–31].

Thus, although this increase has not yet been quantified in mobility surveys, there has been

an increase in road accidents for users. In a recent publication, the Rhône Registry reported

1,186 scooter accidents, resulting in 1,197 injured users and a 7.3-fold increase in the

number of scooter accidents between 2018 and 2019 [32].

Therefore, this study aimed to describe the characteristics of victims injured during a work-

related road accident and their evolution since the emergence of PMDs. A secondary

objective was to characterize and compare accidents involving scooters and other PMDs

with the usual user categories.

2. Materials and Methods

2.1.   Design

This retrospective cross-sectional study used data from the Rhône Road Trauma Registry.

2.2.   The Rhône Road Trauma Registry

The Rhône Road Trauma Registry has been recording data prospectively since its inception in 1995 and covers the Rhône region (1.85 million inhabitants), including one of France's largest cities (Lyon, 0.5 million inhabitants, 11,000 inhabitants/km^2) within a metropolis (Lyon Metropolis, 1.4 million inhabitants). Individuals were included in the registry if they faced a road accident involving one or more road users of nmPMDs or mPMDs in the Rhône region and consulted one of the 245 public or private healthcare facilities (including 42 emergency departments and 20 intensive care units within levels I, II, and III trauma centers), including pre-hospital primary care services and forensic medicine institutes.

The Registry collects the data on demographics of each victim, their sustained injuries, and accidents. Patient information was prospectively collected from the time of the accident to hospital discharge from prehospital emergency care, emergency departments, intensive care units, and surgical units.

2.3.   Study population

Data were collected from 2015 to 2020. The victims aged 18–70 years who were injured on the way to/from work or while working were included in the Rhône Registry database (Figure 1). Persons without an injury were excluded, as were passengers in a vehicle (n=546)

184   and drivers of buses/coaches/tractors (n=61) or other motorized vehicles (tractors,

185   construction equipment, quad bikes; n=17).

2.4.   Variables

187   The following variables were analyzed: sex, age, category of work accident (commuting or

188   mission), category of road user, type of road network, antagonist, time of accident, month of

189   accident, day of accident, use of adequate safety equipment, hospitalization, injuries and

190   death at the scene or post-accident.

2.5.   Gravity of the injuries

Each injury was coded according to the Abbreviated Injury Scale (AIS), 2005 version, which

comprises 2000 codes divided into nine body areas [33,34]. Each code has an associated AIS

severity score ranging from 1 (minor injury) to 6 (beyond treatment/death). As a patient may

have multiple injuries in the same body part, the maximum AIS (MAIS) reports the greatest

severity in each injury area. Similarly, the overall MAIS score is calculated to determine the

victims' overall severity of injuries. Finally, the New Injury Severity Score (NISS) was

calculated from the three most severe injuries and was defined as the sum of the squares of

the respective AIS [35,36]. It was then classified into 3 categories: minor trauma <9,

moderate trauma (9–15), and severe trauma >=16.

2.6.   Statistical analysis

A descriptive analysis was conducted for the overall data and by user category (for the main

categories). Seven categories of users were identified: pedestrians, bicycle drivers, scooter

drivers, drivers of other PMDs (roller blades, skateboards, hoverboards, and monowheels),

car/light commercial vehicle drivers, truck drivers, and motorized two-wheeler drivers.

An analysis strategy was defined to compare the subgroups of soft mode transport users

(pedestrians, bicyclists, scooters, and other PMDs), motor vehicle users not authorized to

use bicycle paths and lanes (cars/light commercial vehicles, trucks, and motorized two-

wheelers), and the seven user categories.

Categorical variables were compared by user category using the chi-square test, and

quantitative variables, using analysis of variance (ANOVA). All analyses were performed

using SAS software version 9.4.

3. Results

3.1.   Description of the victims

Between 2015 and 2020, 53,487 victims were recorded in the Rhône registry, of whom approximately 82% (n=43,791) were between 18 and 70 years of age. Approximately 26% (n=11,296) of accidents were related to commutes or duty journeys.

Of the 11,296 victims aged 18–70 years involved in work-related road traffic accidents, 11,277 had an injury report. Seven people died at the time of the accident and seven died after hospitalization.

We excluded 546 passengers and 78 drivers of other motorized vehicles (buses, tramways, construction equipment, and tractors) from the analysis according to the main categories of interest, resulting in 10,653 participants (94.4%). There were scooter drivers (3.8%), other PMD drivers (0.4%), pedestrians (5.1%), and drivers of bicycles (16.9%), cars or light commercial vehicles (45.3%), trucks (1.5%), and motorized two-wheelers (21.4%).

Nearly half of the scooter drivers (54.8%) and more than three-quarters of other PMD drivers were male (Table 1). Overall, most victims were male (61%), except for pedestrians (45.7%) and car occupants (47.1%).

Most PMD drivers were under 35 years of age (60% for other PMDs and 50% for scooters). Overall, the population was young, with more than 50% aged < 35 years, and 75% aged < 45 years. The average age differed significantly according to the user category.

Most work-related scooter accidents occurred during commuting (95.6%). The reason for the journey was related to the mode of travel; 79% of the truck driver victims were on duty and

20% of pedestrians were involved in the accident while on duty. Overall, 87% of the victims

were commuters.

Nearly two-thirds of scooter users and half of other-PMD users had no antagonists. Overall,

only 25% of the victims faced accidents without antagonists. By definition, all the

pedestrians had antagonists.

Of the victims, 90% were involved in accidents during the week, with no difference based on

the user category. Less than 25% of scooter and other pedal bike drivers were involved in

nighttime accidents. The accidents occurred at night for 29% of the victims, with a higher

proportion of pedestrians (35%).

The majority of scooter, other PMD, and bicycle drivers, and pedestrians were involved in

accidents in street-type traffic lanes in the city. More than 45% of car and truck drivers were

involved in accidents outside urban areas, whereas approximately 90% of accidents involving

scooters, other PMDs, and bicycles occurred in urban areas (Lyon and its metropolitan area).

The use of personal protective equipment according to the user category was the highest for

car drivers (97% wore a seatbelt), truck drivers (82% wore a seatbelt), and motorized two-

wheeler drivers (95% wore a helmet). For soft-mode users, only one-third of injured bicycle

riders wore helmets, and less than one-fifth of injured PMD and scooter users wore helmets.

3.2.   Description of the injuries

Half of the drivers of other PMDs had a single injury, whereas scooter drivers, pedestrians,

and motorized two-wheeler drivers more frequently had multiple injuries (Table 2). Of the

victims, 42% had a single injury, 33% had two injuries, 18% had three injuries, and 7% had

more than four injuries. The average number of injuries differed significantly according to

user category.

Of the victims, 7.7% were hospitalized. This proportion was higher for pedestrians, and

drivers of trucks, motorized two-wheeler, and other PMDs. One scooter driver died in an

accident. Twelve people died from their injuries (including three pedestrians and three

motorcyclists).

The most frequently affected body parts were the spine (37.6%), lower limbs (36.4%), upper

limbs (36.1%), head (14.8%), thorax (9.3%), neck (9.7%), face (8.1%), and abdomen (3.3%).

Most scooter users had injuries to their upper limbs (59.2%), lower limbs (46.8%), face

(21.2%) or head (17,9%). Spinal injuries were more common among car and truck drivers,

and pedestrians than among other users. Lower limb injuries were more common among

users of bicycles, motorized bicycles, scooters, and other PMDs. Upper extremity injuries

were more common among users of scooters, bicycles, motorized two-wheelers, and other

PMDs, and pedestrians. Head injuries were more common in pedestrians, and scooter, car,

and bicycle users. Chest injuries were more common among truck and car users,

pedestrians, and two-wheeler users. Neck injuries were more common among truck and car

users. Facial injuries were more common among scooter and bicycle users, pedestrians, and

other PMD users. Finally, abdominal injuries were more common among pedestrians and

motorized two-wheeler users.

Only 2.1% of scooter users were seriously injured (MAIS 3+). Overall, 97.5% of the victims

were slightly injured (MAIS <3). The proportion of serious injuries (MAIS 3+) was higher

among pedestrians (6.6%), other PMD users (8%), and motorized bicycle users (4.5%).

Facial injuries were minor for all accident victims, except for 4% of pedestrians who had

injuries of MAIS 3+. Among scooter and other PMD users, all upper-extremity injuries were

of low severity (MAIS <3). However, three users of other PMD had lower extremity injuries

of MAIS ≥3.

282 3.3. Evolution of work-related road accidents during 2015–2020

Since 2015, an increase in the number of work-related traffic injuries has been observed,

mainly due to an increase in the number of accidents while commuting (approximately 4%);

2020, the year of the COVID-19 pandemic, was characterized by a break in this increase

(Figure 2).

During the study period, the proportion of scooter drivers increased by more than 660%

(Figure 3). In contrast, the share of car drivers decreased by 21% between 2015 and 2020,

that of motorized two-wheeler drivers, by 24%, and that of truck drivers, by 17%.

4. Discussion

4.1.   Main findings

This original study on work-related road accidents allowed us to characterize the increase in work-related road accidents associated with new modes of travel, particularly scooters. This new mode of travel has been constantly increasing for approximately 10 years and even more since the emergence of shared electric scooters. Companies in Lyon have offered this service since 2018. Its use accelerated following the COVID-19 pandemic. The results of this study showed that for work-related trips, scooter accidents increased by 660% between 2015 and 2020, and car accidents decreased by 21% in 2020. Bagou et al. [32] found a seven-times increase in scooter accident victims between 2018 and 2019 in the Rhône Department. In contrast, the proportion of car drivers decreased in 2020 due to COVID-19 pandemic-related restrictions [37].

Most work-related road accidents were commuting accidents. Among people working in a fixed and regular place outside the home (i.e., approximately 79% of employed people), approximately 29% make two or more home-to-work trips [2]. This proportion is inversely correlated with commuting journey time. Approximately 25% of employees are exposed to driving as part of their work activities (Fort et al., 2016, 2019).

However, only a small proportion of scooter users were involved in work-related accidents. Either their use is reserved for home-work trips or people using this mode of transport have less risky behavior when it comes to a trip within the framework of their professional activity.

4.2. Characterization of users of nmPMD/mPMD who had an accident

An early review of the literature on electric scooters provided insights into the circumstances and consequences of electric scooter accidents [38,39]. The majority of these accidents occur without an antagonist and as a result of a fall, collision with an object, excessive speed, or unfavorable road conditions. A prospective study in Spain also characterized electric scooter accidents [40]: women represented 40% of the injured, and approximately 13% of the injured were under 18 years of age. In Austria, two-thirds of the injuries involved men, and 60% of the accidents occurred at night [41]. In Denmark, the majority of injured electric scooter drivers were 18-25 years old. The proportion of pedestrians injured by an electric scooter was 17% in accidents involving scooters [42]. A recent German study observed a two-thirds were men, and nighttime accidents [43]. In another study, Graef et al. (2021) observed that among patients admitted to a level 1 trauma center with a mean age of 30 years, 44% were female. In Italy, patients involved in electric scooter accidents were predominantly male (79%) with an average age of 30 years [45]. In an annual incidence study of patients with injuries related to the use of electric scooters in the USA, 60% of the victims were men and 92% were drivers [46]. In Germany, following the implementation of the first registry to examine injury patterns and collect epidemiological data on people injured while riding electric scooters, Heuer et al. (2021) described injuries in 90 patients (65% male; mean age, 35 years).

Nikiforiadis et al. [9] found that electric scooter traffic on sidewalks affected pedestrian experiences the most, ahead of uncontrolled parking on sidewalks. The speed of scooters leads to feelings of insecurity among pedestrians. However, the results of this study showed that scooter antagonists were rarely observed among injured pedestrian users (12/575).

According to a study by Shichman et al. [31], which compared accidents involving electric

scooters before and after the introduction of electric- sharing scooters, 80% of the injuries

were due to falls. Sixty percent of the scooter drivers in our study did not have an antagonist

and thus, were likely to have fallen in the accident; this proportion was 80% for drivers of

other PMD.

According to a German study, PMD users perform secondary tasks while riding [48]. More

than 13% of bicycle and electric scooter users in Germany performed secondary tasks, such

as wearing headphones or earphones and chatting with another user. However, the use of

cell phones was rare. Drivers with a delivery task using a PMD as part of their activities were

at higher risk of performing secondary tasks in addition to driving [48]. This observational

study found that delivery drivers were most likely to wear headphones or earphones;

however, they did not show significantly more traffic violations, inappropriate use of

roadway infrastructure, or at-fault crashes.

350 4.3. Injuries related to PMD

Most accidents were of low or moderate severity (97.5% MAIS < 3). In the context of work-

related accidents, the frequency of injuries among scooter users and other PMDs was similar

according to the different injury areas and severity. Work-related accidents are generally less

severe than private accidents [49].

The upper limbs, lower limbs, face and head were the main areas injured by scooter drivers.

This finding is consistent with those of other studies. In an Italian study, nearly 47% of the

injuries were fractures, with the majority being radial fractures and 25% requiring surgical

intervention [40]. In a South Korean study [50], the main anatomical region affected in

electric scooter accident victims was the face, followed by the head and upper extremities.

In an annual incidence study on patients with injuries related to electric scooter use in the

USA [46], the most common injuries were fractures and head injuries. In Germany, among

90 people injured while riding electric scooters, 32 fractures and seven ligament injuries

were reported, and head injuries were found in 38 patients [47]. Moftakhar et al. (2021)

studied the severity of injuries caused by electric scooter accidents in Austria. These

accidents occurred at high kinetic energies and caused head and upper extremity injuries. In

Germany, following the introduction of shared electric scooters in 2019, the topic of safety

emerged as a public debate. A recent study [43] reported that electric scooter accidents

were the source of serious injuries, with 28% of accident victims requiring surgical

management. In another study, Graef et al. [44] observed that more than two-thirds of

patients had extremity injuries, and half had facial injuries. The chest region exhibited the

highest AIS scores. Overall, 70% of the injuries were minor. Finally, in Italy, among patients

involved in electric scooter accidents [45], 60% had a head injury, 30% were transported to

emergency departments for life-threatening injuries, and 15% were placed in intensive care

units.

The fact that not all scooter drivers used electric scooters in our study minimized the

severity of the crashes. Shichman et al. [31] studied crashes involving electric scooters

before and after the introduction of shared electric scooters and showed a significant

increase in electric scooter-related injuries during emergency department visits after the

introduction of sharing services (six times more admissions).

Compared to non-electric scooter drivers, electric scooter drivers suffered more bruises and

lacerations on the face, required sutures, and were more often under the influence of

alcohol or drugs [42].

4.4. Safety equipment

Although seatbelt use among car and truck drivers and helmet use among motorized two-

wheeler drivers are very high in the context of work travel, the results indicate that helmet

use among users of bicycles, scooters, and other PMD is low. This has also been found in

other studies in the general population. Helmet use among electric scooter users was

anecdotal in a German study [43], only one out of 43 patients wore a helmet in another

German study [44], and 4% of victims wore a helmet in a study in Dallas, USA [51].

Helmet use for electric scooter users significantly reduces the risk of serious head and facial

injuries [52], maxillofacial fractures, and soft tissue injuries [53]. Similar to the

recommendations for helmet use for bicycle users 10 years ago, the scientific literature

confirms and recommends helmet use for electric scooter users and, in a larger set, for users

of PMDs (DiMaggio et al., 2019; Lin et al., 2020).

The non-use of helmets by electric scooter users and the high prevalence of head injuries

found in numerous studies [38,39] suggest that these types of injuries may be preventable.

Studies demonstrating the effectiveness of helmet use among cyclists can confirm this [56].

As highlighted in this literature review, several cities (Brisbane, Copenhagen, Dallas, Los

Angeles, Malaga Paris, Stockolm, and Vienna) and countries (France and Italy) (La Torre,

2003; Peng et al., 2017) have legislated this area [6]. Results from an Australian study

confirmed that helmet use among electric scooter users increased to over 60% after a

mandatory helmet law was implemented in Australia[59].

4.5. Regulations

In France, a user of rollerblades, skateboards, or scooters (without a motor) is considered a pedestrian who must ride on the sidewalk. Electrical PMDs (scooters, hoverboards, gyropods, monoroue) must be used on the bicycle path, if available. Bicycles must be on a roadway or bike path, if there is one.

The Mobility Law 2019 [60] established that electric scooters are prohibited on sidewalks (unless authorized by the mayor and unless the engine is turned off). In built-up areas, traffic is only authorized on bicycle paths or, failing that, on roads with a maximum speed limit of 50 km/h. Outside built-up areas, traffic is authorized only on bicycle paths or greenways (except for exemptions). Electric scooters are prohibited for children under 12 years of age. Wearing a helmet is mandatory for minors over 12 years of age and strongly recommended for adults. Civil insurance is mandatory. Riding an electric scooter at speeds exceeding 25 km/h is forbidden. Similar to other means of transportation, the use of headphones is prohibited. Front and rear lights are mandatory, as is wearing a retroreflective vest at night or during low visibility. A horn is mandatory. Finally, passenger transport is prohibited. The regulations mandate fines: 135 euros for driving on sidewalks, 35 euros for failure to comply with traffic laws, and 1,500 euros for exceeding the authorized speed limit [61].

The various regulations on PMDs established by European countries are recent and heterogeneous[62]. Regulations regarding helmet use, specific categories of electric scooters, authorized bike lanes, authorized pedestrian lanes, and age restrictions vary by country. However, the maximum speed limit is a common rule except in Hungary. France and Germany require liability insurance. In the USA, regulations differ by state [58,63]. Overall, many countries are yet to establish the legal requirements of this mode of travel in

detail, and more specific legislation is required. In January 2017, the Singaporean

government passed the Active Mobility Act (AMA) to regulate the use of PDMs. However, an

increase in injuries related to these new travel modes has been observed[64].

Road unsafety related to electric bikes/motorcycles is becoming increasingly problematic

owing to their growing popularity compared with other two-wheeled vehicles [65].

The majority of studies on the subject have confirmed that men and users between the ages

of 18 and 34 years are most concerned with the use of and accidents involving private or

shared PMDs.

In France, one of the few studies on this subject compared the demographic characteristics

of users sharing electric scooters with those of personal electric scooters [66]. The latter are

more likely to be men, older, and have a higher income. Therefore, the use of electric

scooters is motivated by saving time, rather than financial reasons. They also seem to have

less risky behaviors, particularly with regard to alcohol consumption, than other electric

scooter users [7]. A comparison of the frequency of use (occasional versus frequent users)

shows that frequent users have riskier behaviors, perhaps because of their habit of use.

4.6. Limitations of the study

Difficulties in correctly identifying nmPMDs were noted. The Rhône Road Trauma Registry

collection form evolved in November 2019 to better quantify electric vehicle-related

accidents. A "No/Yes" checkbox was introduced for the retrieval of mPMD with certainty,

but it does not distinguish nmPMD from missing data.

The analyzed period ended at the start of the COVID-19 pandemic. The year 2020 differed

from previous years in that car accidents decreased and scooter accidents increased. It will

be relevant to conduct another study spanning a few more years to better measure the

impact of nmPMD and mPMD on work-related traffic accidents.

5. Conclusion

This is the first study describing work-related traffic accidents that have occurred since the

emergence of PMDs.

The results observed for users of scooters and other PMDs in this study were generally

consistent with those found in the scientific literature. However, in most available studies,

the type of travel was not restricted to work-related journeys: a young and predominantly

male population, accidents mostly without antagonists, and injuries localized mainly to the

upper limbs, lower limbs, and head/face.

Despite limited data, the results suggest that accidents involving scooters or other DPEs are

of low severity. Many head injuries could be prevented with more widespread helmet use.

Similar to other modes of travel, the number of casualties can be reduced if effective actions

are taken to reduce secondary tasks during scooter journeys [48].

Finally, companies can take preventive actions so that employees using these modes of

transport for work-related journeys are better informed about the risks.

References

Ministère de la Transition écologique et de la Cohésion des territoires ,. Comment les Français voyagent-ils en 2019 ? Résultats de l'enquête mobilité des personnes. 2019.https://www.statistiques.developpement-durable.gouv.fr/comment-les-francais-voyagent-ils-en-2019-resultats-de-lenquete-mobilite-des-personnes (accessed 18 Jan 2023).

Service de l'Observation et des Statistiques ,. La mobilité des Français, panorama issu de l'enquête nationale transports et déplacements 2008. 2010. https://www.statistiques.developpement-durable.gouv.fr/sites/default/files/2018-11/La_mobilite_des_Francais_ENTD_2008_revue_cle7b7471.pdf (accessed 18 Jan 2023).

Pautard É. Les Français et la mobilité durable : quelle place pour les déplacements alternatifs à la voiture individuelle en 2016 ? 2016. https://www.statistiques.developpement-durable.gouv.fr/sites/default/files/2018-10/Datalab-essentiel-36-les-francais-et-la-mobilite-durable-septembre2016.pdf (accessed 18 Jan 2023).

Fort E, Gadegbeku B, Gat E, *et al.* Working conditions and risk exposure of employees whose occupations require driving on public roads – Factorial analysis and classification. *Accid Anal Prev* 2019;**131**:254–67. doi:10.1016/j.aap.2019.07.001

Fort E, Ndagire S, Gadegbeku B, *et al.* Working conditions and occupational risk exposure in employees driving for work. *Accid Anal Prev* 2016;**89**:118–27. doi:10.1016/j.aap.2016.01.015

Gössling S. Integrating e-scooters in urban transportation: Problems, policies, and the prospect of system change. *Transp Res D Transp Environ* 2020;**79**:102230. doi:10.1016/J.TRD.2020.102230

Christoforou Z, Gioldasis C, de Bortoli A, *et al.* Who is using e-scooters and how? Evidence from Paris. *Transp Res D Transp Environ* 2021;**92**:102708. doi:10.1016/J.TRD.2021.102708

Heineke K, Kloss B, Scurtu D, *et al.* Micromobility's 15,000-mile checkup. 2019.

Nikiforiadis A, Paschalidis E, Stamatiadis N, *et al.* Analysis of attitudes and engagement of shared e-scooter users. *Transp Res D Transp Environ* 2021;**94**:102790. doi:10.1016/J.TRD.2021.102790

6t-bureau de recherche. Usages et usagers des trottinettes électriques en free-floating en france. 2019. https://cvip.sphinxonline.net/SurveyServer/s/6t-bureau/trottinettes-freefloating/questionnaire.htm (accessed 1 Mar 2023).

Sanders RL, Branion-Calles M, Nelson TA. To scoot or not to scoot: Findings from a recent survey about the benefits and barriers of using E-scooters for riders and non-riders. *Transp Res Part A Policy Pract* 2020;**139**:217–27. doi:10.1016/J.TRA.2020.07.009

Glenn J, Bluth M, Christianson M, *et al.* Considering the Potential Health Impacts of Electric Scooters: An Analysis of User Reported Behaviors in Provo, Utah. *International Journal of Environmental Research and Public Health 2020, Vol 17, Page 6344* 2020;**17**:6344. doi:10.3390/IJERPH17176344

Fédération Française de l'Assurance (FFA) , Assurance Prévention ,, Fédération des Professionnels de la Micro-Mobilité (FP2M) ,. Usages, risques et accidentalité des Engins

de Déplacement Personnel Motorisés (EDPM), avant et après le confinement.
2020.https://www.assurance-prevention.fr/nos-etudes/engins-deplacement-personnel-
motorises-edpm-usages-risques-accidentalite (accessed 18 Jan 2023).

14 Echeverría-Su M, Huamanraime-Maquin E, Cabrera FI, *et al.* Transitioning to sustainable
mobility in Lima, Peru. Are e-scooter sharing initiatives part of the problem or the solution?
Sci Total Environ 2023;**866**. doi:10.1016/J.SCITOTENV.2022.161130

15 Laa B, Leth U. Survey of E-scooter users in Vienna: Who they are and how they ride. *J Transp*
Geogr 2020;**89**:102874. doi:10.1016/J.JTRANGEO.2020.102874

16 Buck D, Buehler R, Happ P, *et al.* Are Bikeshare Users Different from Regular Cyclists?: A First
Look at Short-Term Users, Annual Members, and Area Cyclists in the Washington, D.C.,
Region. *https://doi.org/103141/2387-13* 2013;:112–9. doi:10.3141/2387-13

17 Lefrancq M. Shared freefloating micromobility regulations and results of escooter users'
survey (summer 2019). In: *ERSCharter Webinar：e scooters*. 2019.
https://erscharter.eu/sites/default/files/ resources/presentation_martin_lefrancq.pdf
(accessed 29 Mar 2022).

18 Fédération Française de l'Assurance (FFA) , Assurance Prévention ,, Fédération
des Professionnels de la Micro-Mobilité (FP2M) ,. Usages, risques et accidentalité des EDPM.
2020. https://www.assurance-prevention.fr/file/5687/download (accessed 18 Jan 2023).

19 Zagorskas J, Burinskiene M. Challenges Caused by Increased Use of E-Powered Personal
Mobility Vehicles in European Cities. *Sustainability 2020, Vol 12, Page 273* 2019;**12**:273.
doi:10.3390/SU12010273

20 Anderson B, Rupp JD, Moran TP, *et al.* The Effect of Nighttime Rental Restrictions on E-Scooter
Injuries at a Large Urban Tertiary Care Center. *International Journal of Environmental*
Research and Public Health 2021, Vol 18, Page 10281 2021;**18**:10281.
doi:10.3390/IJERPH181910281

21 Farley KX, Aizpuru M, Wilson JM, *et al.* Estimated Incidence of Electric Scooter Injuries in the
US From 2014 to 2019. *JAMA Netw Open* 2020;**3**:e2014500.
doi:10.1001/JAMANETWORKOPEN.2020.14500

22 Hourston GJM, Ngu A, Hopkinson-Woolley J, *et al.* Orthopedic injuries associated with use of
electric scooters in the UK: A dangerous trend? Case series and review of the literature. *Traffic*
Inj Prev 2021;**22**:242–5. doi:10.1080/15389588.2021.1882676

23 Pepper T, Barker M, Smyth D, *et al.* Electric scooters: a quick way to get to the emergency
department? *Br Dent J* 2022;**232**:535–7. doi:10.1038/S41415-022-4153-6

24 Bascones K, Maio Méndez TE, Yañez Siller FA. [Translated article] E-scooter accidents: A new
epidemic. *Rev Esp Cir Ortop Traumatol* Published Online First: February 2022.
doi:10.1016/J.RECOT.2022.02.001

25 Suominen EN, Sajanti AJ, Silver EA, *et al.* Alcohol intoxication and lack of helmet use are
common in electric scooter-related traumatic brain injuries: a consecutive patient series from
a tertiary university hospital. *Acta Neurochir (Wien)* 2022;**164**:643–53. doi:10.1007/S00701-
021-05098-2

26 Hennocq Q, Schouman T, Khonsari RH, *et al.* Evaluation of Electric Scooter Head and Neck
Injuries in Paris, 2017-2019. *JAMA Netw Open* 2020;**3**.
doi:10.1001/JAMANETWORKOPEN.2020.26698

27 Badeau A, Carman C, Newman M, *et al.* Emergency department visits for electric scooter-
related injuries after introduction of an urban rental program. *Am J Emerg Med*
2019;**37**:1531–3. doi:10.1016/J.AJEM.2019.05.003

28 Beck S, Barker L, Chan A, *et al.* Emergency department impact following the introduction of an
electric scooter sharing service. *Emerg Med Australas* 2020;**32**:409–15. doi:10.1111/1742-
6723.13419

29 Mair O, Wurm M, Müller M, *et al.* E-scooter accidents and their consequences: First
prospective analysis of the injury rate and injury patterns in the urban area of a German city
with over 1 million residents. *Unfallchirurg* 2021;**124**:382–90. doi:10.1007/s00113-020-00910-
7

30 Mitchell G, Tsao H, Randell T, *et al.* Impact of electric scooters to a tertiary emergency
department: 8-week review after implementation of a scooter share scheme. *EMA -*
Emergency Medicine Australasia 2019;**31**:930–4. doi:10.1111/1742-6723.13356

31 Shichman I, Shaked O, Factor S, *et al.* Emergency department electric scooter injuries after the
introduction of shared e-scooter services: A retrospective review of 3,331 cases. World J
Emerg Med. 2022;**13**:5–10. doi:10.5847/WJEM.J.1920-8642.2022.002

32 Bagou G, Ndiaye A, Hugenschmitt D, *et al.* Traumatismes consécutifs aux accidents de
trottinettes. *Annales françaises de médecine d'urgence* 2021;**11**:144–9. doi:10.3166/AFMU-
2021-0325

33 Salottolo K, Settell A, Uribe P, *et al.* The impact of the AIS 2005 revision on injury severity
scores and clinical outcome measures. *Injury* 2009;**40**:999–1003.
doi:10.1016/J.INJURY.2009.05.013

34 Medicine A for the A of A. The Abbreviated Injury Scale 2005—Update 2008. Association for
the Advancement of Automotive Medicine Barrington, IL 2008.

35 Koch M, Nygren A, Tingvall C. Validation of the new injury impairment scale (IIS). In: AAAM,
ed. *38th AAAM conference*. Lyon, France: : AAAM 1994. 123–38.

36 Li H, Ma YF. New injury severity score (NISS) outperforms injury severity score (ISS) in the
evaluation of severe blunt trauma patients. *Chinese Journal of Traumatology* 2021;**24**:261–5.
doi:10.1016/J.CJTEE.2021.01.006

37 Laumon B, Tardy H, Ndiaye A, *et al.* Traumatologie routière pendant la première période de
confinement pour cause de pandémie de Covid-19 – Registre du Rhône. *Bull Épidémiol Hebd*
2022;**18**:308–14.http://beh.santepubliquefrance.fr/beh/2022/18/2022_18_1.html (accessed
11 Jan 2023).

38 Toofany M, Mohsenian S, Shum LK, *et al.* Systematic review Injury patterns and circumstances
associated with electric scooter collisions: a scoping review. *Inj Prev* 2021;**0**:1–10.
doi:10.1136/injuryprev-2020-044085

39   Toofany M, Mohsenian S, Shum LK, *et al.* Injury patterns and circumstances associated with
electric scooter collisions: A scoping review. Injury Prevention. 2021;**27**:490–9.
doi:10.1136/injuryprev-2020-044085

40   Coelho A, Feito P, Corominas L, *et al.* Electric scooter-related injuries: A new epidemic in
orthopedics. *J Clin Med* 2021;**10**. doi:10.3390/jcm10153283

41   Moftakhar T, Wanzel M, Vojcsik A, *et al.* Incidence and severity of electric scooter related
injuries after introduction of an urban rental programme in Vienna: a retrospective
multicentre study. *Arch Orthop Trauma Surg* 2021;**141**:1207–13. doi:10.1007/s00402-020-
03589-y

42   Blomberg SNF, Rosenkrantz OCM, Lippert F, *et al.* Injury from electric scooters in Copenhagen:
a retrospective cohort study. *BMJ Open* 2019;**9**. doi:10.1136/BMJOPEN-2019-033988

43   Störmann P, Klug A, Nau C, *et al.* Characteristics and injury patterns in electric-scooter related
accidents—A prospective two-center report from Germany. *J Clin Med* 2020;**9**.
doi:10.3390/jcm9051569

44   Graef F, Doll C, Niemann M, *et al.* Epidemiology, Injury Severity, and Pattern of Standing E-
Scooter Accidents: 6-Month Experience from a German Level I Trauma Center. *Clin Orthop*
Surg 2021;**13**:443–8. doi:10.4055/CIOS20275

45   Scquizzato T, Gamberini L, Stella F, *et al.* Analysis of road traffic accidents involving standing
electric scooters reported in newspapers in Italy. *Clin Exp Emerg Med* 2022;**9**:36–40.
doi:10.15441/CEEM.21.078

46   Trivedi TK, Liu C, Antonio ALM, *et al.* Injuries Associated With Standing Electric Scooter Use.
JAMA Netw Open 2019;**2**:e187381. doi:10.1001/JAMANETWORKOPEN.2018.7381

47   Heuer S, Landschoof S, Kornherr P, *et al.* Epidemiology and Injury Pattern of E-Scooter Injuries
- Initial Results. *Z Orthop Unfall* Published Online First: 3 May 2021. doi:10.1055/a-1419-0584

48   Huemer AK, Banach E, Bolten N, *et al.* Secondary task engagement, risk-taking, and safety-
related equipment use in German bicycle and e-scooter riders - An observation. *Accid Anal*
Prev 2022;**172**:106685. doi:10.1016/J.AAP.2022.106685

49   Charbotel B, Martin JL, Chiron M. Work-related versus non-work-related road accidents,
developments in the last decade in France. *Accid Anal Prev* 2010;**42**:604–11.

50   Choi JH, Oh SS, Kim KS, *et al.* Electric Scooter-Related Trauma in Korea. *J Korean Med Sci*
2022;**37**. doi:10.3346/JKMS.2022.37.E181

51   Trivedi B, Kesterke MJ, Bhattacharjee R, *et al.* Craniofacial Injuries Seen With the Introduction
of Bicycle-Share Electric Scooters in an Urban Setting. *Journal of Oral and Maxillofacial*
Surgery 2019;**77**:2292–7. doi:10.1016/J.JOMS.2019.07.014

52   Hamzani Y, Bar Hai D, Cohen N, *et al.* The impact of helmet use on oral and maxillofacial
injuries associated with electric-powered bikes or powered scooter: a retrospective cross-
sectional study. *Head Face Med* 2021;**17**. doi:10.1186/S13005-021-00288-W

53   al Saffar MMA, Verdonschot RJCG, Stassen HS, *et al.* The Association of Helmet Use with the
Occurrence of Maxillofacial Injuries Following Bicycle or Scooter Accidents: A Retrospective

Cohort Study. *Craniomaxillofac Trauma Reconstr* 2022;**15**:21–7.
doi:10.1177/1943387521998199

54   Dimaggio CJ, Bukur M, Wall SP, *et al.* Injuries associated with electric-powered bikes and
scooters: Analysis of US consumer product data. *Injury Prevention* Published Online First:
2019. doi:10.1136/injuryprev-2019-043418

55   Lin S, Goldman S, Peleg K, *et al.* Dental and maxillofacial injuries associated with electric-
powered bikes and scooters in Israel: A report for 2014-2019. *Dental Traumatology*
2020;**36**:533–7. doi:10.1111/edt.12562

56   Olivier J, Creighton P. Bicycle injuries and helmet use: a systematic review and meta-analysis.
Int J Epidemiol 2017;**46**:278–92. doi:10.1093/IJE/DYW153

57   [Epidemiology of scooter accidents in Italy: the effectiveness of mandatory use of helmets in
preventing incidence and severity of head trauma] - PubMed.
https://pubmed.ncbi.nlm.nih.gov/12632991/ (accessed 2 Mar 2023).

58   Peng Y, Vaidya N, Finnie R, *et al.* Universal Motorcycle Helmet Laws to Reduce Injuries: A
Community Guide Systematic Review. *Am J Prev Med* 2017;**52**:820–32.
doi:10.1016/J.AMEPRE.2016.11.030

59   Haworth NL, Schramm A. Illegal and risky riding of electric scooters in Brisbane. *Med J Aust*
2019;**211**:412–3. doi:10.5694/MJA2.50275

60   Décret n° 2019-1082 du 23 octobre 2019 relatif à la réglementation des engins de
déplacement personnel. JORF n°0249 du 25 octobre 2019 2019.
https://www.legifrance.gouv.fr/loda/id/JORFTEXT000039272656 (accessed 12 Jan 2023).

61   La nouvelle réglementation pour les Engins de Déplacements Personnels motorisés (EDPM) |
Cerema. https://www.cerema.fr/fr/actualites/nouvelle-reglementation-engins-deplacements-
personnels

62   Serra GF, Fernandes FAO, Noronha E, *et al.* Head protection in electric micromobility: A critical
review, recommendations, and future trends. *Accid Anal Prev* 2021;**163**.
doi:10.1016/J.AAP.2021.106430

63   The 2022 Comprehensive Guide to Electric Scooter Laws. https://unagiscooters.com/scooter-
articles/the-comprehensive-guide-to-electric-scooter-laws-2022/ (accessed 2 Mar 2023).

64   Lee W, Goo T, Lim W, *et al.* Hospital seeing more personal mobility device accidents and
serious injuries despite active mobility act. *J Emerg Trauma Shock* 2020;**13**:274–8.
doi:10.4103/JETS.JETS_115_19

65   Zhang X, Yang Y, Yang J, *et al.* Road traffic injuries among riders of electric bike/electric moped
in southern China. *Traffic Inj Prev* 2018;**19**:417–22. doi:10.1080/15389588.2018.1423681

66   Gioldasis C, Christoforou Z, Seidowsky R. Risk-taking behaviors of e-scooter users: A survey in
Paris. *Accid Anal Prev* 2021;**163**. doi:10.1016/J.AAP.2021.106427

Table 1: Socio-demographic and accidental characteristics of victims injured in work-related accident, by user category (source the Rhône Road Trauma Registry, 2015-2020)

	Pedestrian (n=575)	Bicycle driver (n=1 908)	Scoteer driver (n=429)	Other PMD driver (n=50)			Car driver (n=5 111)	Truck driver (n=166)	Motorized two-wheelers driver (n=2 414)		All (n=10 653)	
	n (%)	n (%)	n (%)	n (%)	p-value		n (%)	n (%)	n (%)	p-value		p-value (All)
Sex					<0,0001					<0,0001		<0,0001
Women	312 (54.3)	664 (34.8)	194 (45.2)	10 (20.0)			2703 (52.9)	3 (1.8)	277 (11.5)		4163 (39.1)	
Men	263 (45.7)	1244 (65.2)	235 (54.8)	40 (80.0)			2408 (47.1)	163 (98.2)	2137 (88.5)		6490 (60.9)	
Age categories					<0,0001					<0,0001		<0,0001
18-24 years	139 (24.2)	355 (18.6)	94 (21.9)	15 (30.0)			1208 (23.6)	27 (16.3)	678 (28.1)		2516 (23.6)	
25-34 years	140 (24.4)	636 (33.3)	132 (30.8)	15 (30.0)			1597 (31.2)	42 (25.3)	683 (28.3)		3245 (30.5)	
35-44 years	113 (19.6)	426 (22.3)	108 (25.2)	12 (24.0)			1149 (22.5)	39 (23.5)	499 (20.7)		2346 (22)	
45-54 years	108 (18.8)	331 (17.4)	77 (17.9)	4 (8.0)			832 (16.3)	34 (20.5)	388 (16.1)		1774 (16.7)	
55-64 years	69 (12.0)	157 (8.2)	17 (4.0)	4 (8.0)			317 (6.2)	21 (12.6)	164 (6.8)		749 (7.0)	
65-70 years	6 (1.0)	3 (0.2)	1 (0.2)	0 (0)			8 (0.2)	3 (1.8)	2 (0.1)		23 (0.2)	
Type of work-related journey					<0,0001					<0,0001		<0,0001
Commuting	456 (79.3)	1742 (91.3)	410 (95.6)	43 (86.0)			4565 (89.3)	35 (21.1)	2041 (84.5)		9292 (87.2)	
On duty	119 (20.7)	166 (8.7)	19 (4.4)	7 (14.0)			546 (10.7)	131 (78.9)	373 (15.5)		1361 (12.8)	
Antagonist					NA					NA		NA
None	0 (0)	909 (47.6)	281 (65.5)	26 (52.0)			548 (10.7)	48 (28.9)	1015 (42.1)		2827 (26.5)	
Pedestrian	0 (0)	29 (1.5)	3 (0.7)	1 (2.0)			7 (0.1)	0 (0)	11 (0.5)		51 (0.5)	
Other or unknown	14 (2.4)	19 (1.0)	4 (0.9)	0 (0)			15 (0.3)	0 (0)	11 (0.5)		63 (0.6)	
Car	391 (68.0)	646 (33.9)	87 (20.3)	16 (32.0)			3414 (66.8)	34 (20.5)	1139 (47.2)		5727 (53.8)	

	Pedestrian (n=575)	Bicycle driver (n=1 908)	Scoteer driver (n=429)	Other PMD driver (n=50)			Car driver (n=5 111)	Truck driver (n=166)	Motorized two-wheelers driver (n=2 414)			All (n=10 653)	
	n (%)	n (%)	n (%)	n (%)	p-value		n (%)	n (%)	n (%)	p-value			p-value (All)
Fixed obstacle	0 (0)	164 (8.6)	31 (7.2)	3 (6.0)			584 (11.4)	43 (25.9)	112 (4.6)			937 (8.8)	
Light commercial vehicle, truck, bus, train/tram, tractor	66 (11.5)	63 (3.3)	10 (2.3)	1 (2.0)			517 (10.1)	40 (24.1)	85 (3.5)			782 (7.3)	
Motorized two-wheelers	52 (9.0)	21 (1.1)	5 (1.2)	1 (2.0)			23 (0.5)	1 (0.6)	31 (1.3)			134 (1.3)	
Bicycle	52 (9.0)	56 (2.9)	8 (1.9)	2 (4.0)			3 (0.1)	0 (0)	10 (0.4)			131 (1.2)	
Rollerblade, skateboard	0 (0)	1 (0.05)	0 (0)	0 (0)			0 (0)	0 (0)	0 (0)			1 (0)	
Lighting					0,0009					0,5			0,002
Day	320 (65.4)	1067 (74.1)	218 (76.5)	31 (75.6)			3072 (70.7)	105 (73.4)	1401 (69.7)			6294 (71.0)	
Night	169 (34.6)	373 (25.9)	67 (23.5)	10 (24.4)			1274 (29.3)	38 (26.6)	610 (30.3)			2567 (29.0)	
Month					<0,0001					0,0006			<0,0001
january	60 (10,4)	141 (7,4)	32 (7,5)	4 (8)			494 (9,7)	27 (16,3)	170 (7)			928 (8.7)	
February	56 (9,7)	115 (6)	22 (5,1)	3 (6)			385 (7,5)	10 (6)	172 (7,1)			763 (7.2)	
March	49 (8,5)	122 (6,4)	27 (6,3)	4 (8)			421 (8,2)	10 (6)	189 (7,8)			822 (7.7)	
April	32 (5,6)	128 (6,7)	29 (6,8)	3 (6)			359 (7)	6 (3,6)	169 (7)			726 (6.8)	
May	30 (5,2)	149 (7,8)	34 (7,9)	3 (6)			367 (7,2)	12 (7,2)	168 (7)			763 (7.2)	
June	52 (9)	180 (9,4)	45 (10,5)	5 (10)			479 (9,4)	15 (9)	239 (9,9)			1015 (9.5)	
July	38 (6,6)	193 (10,1)	50 (11,7)	9 (18)			419 (8,2)	21 (12,6)	223 (9,2)			953 (8.9)	
August	27 (4,7)	131 (6,9)	23 (5,4)	3 (6)			254 (5)	10 (6)	141 (5,8)			589 (5.5)	
September	41 (7,1)	246 (12,9)	46 (10,7)	7 (14)			436 (8,5)	12 (7,2)	255 (10,6)			1043 (9.8)	
October	64 (11,1)	199 (10,4)	37 (8,6)	4 (8)			518 (10,1)	12 (7,2)	258 (10,7)			1092 (10.3)	
November	72 (12,5)	179 (9,4)	41 (9,6)	5 (10)			536 (10,5)	14 (8,4)	218 (9)			1065 (10.0)	
December	54 (9,4)	125 (6,6)	43 (10)	0 (0)			443 (8,7)	17 (10,2)	212 (8,8)			894 (8.4)	

	Pedestrian (n=575)	Bicycle driver (n=1 908)	Scoteer driver (n=429)	Other PMD driver (n=50)			Car driver (n=5 111)	Truck driver (n=166)	Motorized two-wheelers driver (n=2 414)			All (n=10 653)	
	n (%)	n (%)	n (%)	n (%)	p-value		n (%)	n (%)	n (%)	p-value			p-value (All)
Season													
Spring	111 (19,3)	399 (20,9)	90 (21)	10 (20)			1147 (22,4)	28 (16,8)	526 (21,8)			2311 (21.7)	
Summer	117 (20,3)	504 (26,4)	118 (27,6)	17 (34)			1152 (22,6)	46 (27,6)	603 (24,9)			2557 (24.0)	
Autunm	177 (30,7)	624 (32,7)	124 (28,9)	16 (32)			1490 (29,1)	38 (22,8)	731 (30,3)			3200 (30.0)	
Winter	170 (29,5)	381 (20)	97 (22,6)	7 (14)			1322 (25,9)	54 (32,5)	554 (22,9)			2585 (24.3)	
Urban area					<0,0001					<0,0001			<0,0001
Lyon	205 (35.6)	1257 (65.9)	325 (75.8)	26 (52.0)			730 (14.3)	15 (9.0)	700 (29.0)			3258 (30.6)	
Grand Lyon	220 (38.3)	498 (26.1)	77 (17.9)	18 (36.0)			2068 (40.5)	72 (43.4)	886 (36.7)			3839 (36.0)	
Hors Grand Lyon	66 (11.5)	88 (4.6)	10 (2.3)	2 (4.0)			1375 (26.9)	53 (31.9)	388 (16.1)			1982 (18.6)	
Rhône SAP	84 (14.6)	65 (3.4)	17 (4.0)	4 (4.0)			938 (18.3)	26 (15.7)	440 (18.2)			1574 (14.8)	

Table 2: Injury characteristics of victims injured in work-related accident, by user category (source the Rhône Road Trauma Registry, 2015-2020)

	Pedestrian (n=575)	Bicycle driver (n=1908)	Scotter driver (n=429)	Other PMD driver (n=50)			Car driver (n=5111)	Truck driver (n=166)	Motorized two-wheelers driver (n=2414)			All (n=10 653)	
	n (%)	n (%)	n (%)	n (%)	p-value		n (%)	n (%)	n (%)	p-value			p-value
Hospitalization	80 (13.9)	156 (8.2)	38 (8.9)	6 (12.0)	0,0005		216 (4.2)	29 (17.5)	301 (12.5)	<0,0001		826 (7.8)	<0,0001
Death	3 (0.5)	2 (0.1)	1 (0.2)	0 (0)	NA		2 (0.1)	1 (0.6)	3 (0.1)	NA		12 (0.1)	NA
Number of injuries					0,0001					<0,0001			<0,0001
1	186 (32.3)	759 (39.8)	163 (38.0)	25 (50.0)			2414 (47.2)	66 (39.8)	854 (35.4)			4467 (41.9)	
2	183 (31.8)	601 (31.5)	146 (34.0)	13 (26.0)			1685 (33.0)	54 (32.5)	853 (35.3)			3535 (33.2)	
3	129 (22.4)	410 (21.5)	76 (17.7)	8 (16.0)			760 (14.9)	31 (18.7)	483 (20.0)			1897 (17.8)	
4+	77 (13.4)	138 (7.2)	44 (10.3)	4 (8.0)			252 (4.9)	15 (9.0)	224 (9.3)			754 (7.1)	
MAIS					NA					NA			NA
1 Minor	451 (78.4)	1433 (75.1)	315 (73.4)	37 (74.0)			4830 (94.5)	138 (83.1)	1790 (74.1)			8994 (83.5)	
2 Moderate	86 (15.0)	428 (22.4)	105 (24.5)	9 (18.0)			223 (4.4)	22 (13.2)	516 (21.4)			1389 (12.9)	
3 Serious	21 (3.6)	37 (1.9)	6 (1.4)	3 (0.6)			44 (0.8)	4 (2.4)	86 (3.6)			201 (1.9)	
4 Severe	13 (2.3)	8 (0.4)	2 (0.5)	0 (0)			10 (0.2)	2 (1.2)	14 (0.6)			49 (0.4)	
5 Critical	3 (0.5)	2 (0.1)	0 (0)	1 (2.0)			4 (0.1)	0 (0)	8 (0.3)			18 (0.2)	
6 Maximum	1 (0.2)	0 (0)	1 (0.2)	0 (0)			0 (0)	0 (0)	0 (0)			2 (0)	
MAIS3+	38 (6.6)	47 (2.5)	9 (2.1)	4 (8.0)	<0,0001		58 (1.1)	6 (3.6)	108 (4.5)	<0,0001		270 (2.5)	<0,0001
Head injury	118 (20.5)	295 (15.5)	77 (17.9)	5 (10.0)	0,02		821 (16.1)	44 (26.5)	198 (8.2)	<0,0001		1558 (14.5)	<0,0001
Global gravity of head injury MAIS3+	11 (9.3)	12 (4.1)	2 (2.6)	1 (20.0)			15 (1.8)	1 (2.3)	8 (4.0)			50 (3.2)	<0,0001
Face injury	73 (12.7)	315 (16.5)	91 (21.2)	7 (14.0)	0,004		279 (5.5)	17 (10.2)	94 (3.9)	0,0001		876 (8.1)	<0,0001
Global gravity of face injury MAIS3+	3 (4.0)	1 (0.3)	0 (0)	0 (0)			0 (0)	0 (0)	0 (0)	NA		4 (0.4)	

	Pedestrian (n=575)	Bicycle driver (n=1908)	Scotter driver (n=429)	Other PMD driver (n=50)	p-value		Car driver (n=5111)	Truck driver (n=166)	Motorized two-wheelers driver (n=2414)	p-value		All (n=10 653)	p-value
	n (%)	n (%)	n (%)	n (%)	p-value		n (%)	n (%)	n (%)	p-value			p-value
Neck injury	34 (5.9)	53 (2.8)	18 (4.2)	0 (0)			791 (15.5)	23 (23.9)	77 (3.2)			996 (9.2)	
Global gravity of neck injury MAIS3+	1 (3.0)	0 (0)	0 (0)	0 (0)			1 (0.1)	0 (0)	1 (1.3)			3 (0.3)	
Chest injury	67 (11.6)	169 (8.9)	35 (8.2)	1 (2.0)			740 (14.5)	26 (15.7)	270 (11.2)			1308 (12.1)	
Global gravity of chest injury MAIS3+	16 (23.9)	14 (8.3)	5 (14.3)	0 (0)			31 (4.2)	5 (19.2)	52 (19.3)			123 (9.4)	
Abdomen injury	33 (5.7)	47 (2.5)	13 (3.0)	0 (0)			148 (2.9)	6 (3.6)	113 (4.7)			360 (3.3)	
Global gravity of Abdomen injury MAIS3+	8 (24.2)	6 (12.8)	0 (0)	0 (0)			12 (8.1)	1 (16.7)	8 (7.1)			35 (9.7)	
Spinal injury	119 (20.7)	198 (10.4)	39 (9.1)	4 (8.0)			3114 (60.9)	70 (42.2)	373 (15.4)			3917 (36.4)	
Global gravity of Spinal injury MAIS3+	0 (0)	2 (1.0)	0 (0)	0 (0)			6 (0.2)	1 (1.4)	5 (1.3)			14 (0.4)	
Upper limb injury	230 (40.0)	1086 (56.9)	254 (59.2)	24 (48.0)			1121 (21.9)	55 (33.1)	1161 (48.1)			3931 (36.5)	
Global gravity of Upper limb injury MAIS3+	1 (0)	2 (0.2)	0 (0)	0 (0)			3 (0.3)	1 (1.8)	5 (0.4)			12 (0.3)	
Injury of the lower limbs / pelvis	395 (68.7)	899 (47.1)	201 (46.8)	26 (52.0)			750 (14.7)	45 (27.1)	1653 (68.5)			3969 (36.8)	
Global gravity of Injury of the lower limbs / pelvis MAIS3+	18 (4.6)	23 (2.6)	4 (2.0)	3 (11.5)			17 (2.3)	1 (2.2)	49 (3.0)			115 (2.9)	

	Pedestrian (n=575)	Bicycle driver (n=1908)	Scotter driver (n=429)	Other PMD driver (n=50)			Car driver (n=5111)	Truck driver (n=166)	Motorized two-wheelers driver (n=2414)			All (n=10 653)	
	n (%)	n (%)	n (%)	n (%)	p-value		n (%)	n (%)	n (%)	p-value			p-value
New Injury Severity Score (NISS)					<0,0001					<0,0001			<0,0001
NISS < 9	523 (91.0)	1824 (95.6)	413 (96.3)	45 (90.0)			5024 (98.3)	153 (92.2)	2241 (92.8)			10343	
NISS = [9–15]	21 (3.6)	62 (3.2)	10 (2.3)	4 (8.0)			49 (1.0)	8 (4.8)	112 (4.6)			266 (2.5)	
NISS ≥ 16	31 (5.4)	22 (1.2)	6 (1.4)	1 (2.0)			38 (0.7)	5 (3.0)	61 (2.5)			164 (1.5)	

Figure 1 : Flow chart

Figure 2 : Evolution of the number of drivers and pedestrians involved in work-related road accidents, by type of journey between 2015 and 2020 (source the Rhône Road Trauma Registry)

Figure 3 : Frequencies of different user categories of drivers and pedestrians injured in work-related accidents, by year (source the Rhône Road Trauma Registry, 2015-2020)