Factors shaping vaginal microbiota long-term community dynamics in young adult women

Tsukushi Kamiya[1,*], Nicolas Tessandier[1], Baptiste Elie[1,2], Claire Bernat[2,3], Vanina Boué[2], Sophie Grasset[2], Soraya Groc[2,4], Massilva Rahmoun[2], Christian Selinger[2,5], Michael S. Humphrys[6], Marine Bonneau[7], Christelle Graf[7], Vincent Foulongne[4], Jacques Reynes[8], Vincent Tribout[8], Michel Segondy[4], Nathalie Boulle[4], Jacques Ravel[6], Carmen Lía Murall[2,9], and Samuel Alizon[1,2,*]

[1]Center for Interdisciplinary Research in Biology (CIRB), Collège de France, CNRS, INSERM, Université PSL, Paris, France
[2]MIVEGEC, CNRS, IRD, Université de Montpellier, France
[3]Institut de Génomique Fonctionnelle, Université de Montpellier, CNRS, INSERM, Montpellier, France.
[4]PCCEI, Univ. Montpellier, Inserm, EFS, Montpellier, France
[5]Current address: Swiss Tropical and Public Health Institute, Basel, Switzerland
[6]Institute for Genomic Sciences, University of Baltimore, USA
[7]Department of Obstetrics and Gynaecology, Centre Hospitalier Universitaire de Montpellier, Montpellier, France
[8]Department of Infectious and Tropical Diseases, Centre Hospitalier Universitaire de Montpellier, Montpellier, France
[9]National Microbiology Laboratory (NML), Montreal Public Health Agency of Canada (PHAC), Canada
[*]Corresponding authors: tsukushi.kamiya@college-de-france.fr, samuel.alizon@college-de-france.fr

Abstract

The vaginal microbiota is categorised into five main community state types (CST) that are known to affect women's health. Yet, there is a notable paucity of high-resolution follow-up studies lasting several months, which are required to interrogate the long-term dynamics and associations with demographic and behavioural covariates. Here, we present a high-resolution longitudinal cohort of 125 women followed for a median duration of 8.6 months, providing 11 samples per woman. Using a hierarchical Bayesian Markov model, we characterised the patterns of CST persistence and transition, simultaneously estimated the impact of 16 covariates and quantified individual variability among women. We showed that 'optimal' (CSTs I, II, and V) and 'sub-optimal' (CST III) communities are more stable in time than 'non-optimal' (CST IV) ones. Furthermore, we found that some covariates — most notably alcohol consumption — impacted the probability of shifting from one CST to another. We performed counterfactual simulations to confirm that alterations of key covariates, such as alcohol consumption, could shape the prevalence of different microbiota communities in the population. Finally, our analyses indicated that there is a relatively canalised pathway leading to the deterioration of vaginal microbiota communities, whereas the paths to recovery can be highly individualised among women. In addition to providing one of the first insights into vaginal microbiota dynamics over a year, our study showcases a novel application of a hierarchical Bayesian Markov model to clinical cohort data with many covariates. Our findings pave the way for an improved mechanistic understanding of

microbial dynamics in the vaginal environment and the development of novel preventative

and therapeutic strategies to improve vaginal health.

Introduction

Epithelia of the human body are host to a diverse array of microorganisms. These microorganisms are collectively referred to as microbiota and their compositions are tightly associated with human health. In the human vaginal environment, the description of the microbiota dates back to Albert Döderlein in 1892. Its composition has been demonstrated to impact the acquisition risk of several sexually transmitted infections (STIs) [1], fertility (especially in medically-assisted procreation procedures) [2], and general well-being [3].

Vaginal microbiota communities comprise hundreds of species. To facilitate understanding, the variation in community composition is usually reduced to a handful of categories that capture key compositional signatures, such as the dominance of certain species or species evenness. This dimensionality reduction filters out noise in the data and facilitates the identification and visualisation of key patterns and relationships. One such dimensionality reduction framework, i.e., community state types (CSTs), introduced by Ravel et al. [4], categorises vaginal microbial communities into five discrete state types. The CSTs considered 'optimal' for health are dominated by *Lactobacillus* species; *Lactobacillus crispatus*, *L. gasseri*, and *L. jensenii* for CST I, II, and V, respectively. Lactobacilli produce lactic acid and hydrogen peroxide, which creates an acidic environment that helps to inhibit the growth of harmful pathogens [5]. On the other end of the spectrum, CST IV is the primary microbial context of bacterial vaginosis (BV), which elevates the risk of STI acquisition and spontaneous preterm birth, and is associated with symptoms such as mal-

odor, discharge, and itching [6,7]. This community is characterised by a diverse assemblage of anaerobic bacterial species from the *Gardnerella*, *Prevotella*, and *Fannyhessea* genera. Finally, CST III, characterised by a dominance of *L. iners*, is considered 'sub-optimal' for women's health. While *L. iners* is a member of the *Lactobacillus* genus, it is less effective at producing lactic acid and hydrogen peroxide. As such, women with CST III tend to exhibit higher vaginal pH than CST I and are more prone to experiencing adverse health consequences, including vaginal infections [8].

The CST classification represents a snapshot of the microbiota community at the time of sampling that facilitates the examination of clinically relevant microbiota variations across time and women. The development of the modern pipeline — through meta-barcoding sequencing of 16S DNA and clustering algorithms [9] — allows for CST-typing with enhanced efficiency and reduced observer bias compared to conventional microscopy-based methods of vaginal microbiota community typing (e.g., Nugent score).

The composition of vaginal microbiota is characteristically variable over both short and long timescales [10]. For instance, vaginal microbiota shifts throughout a woman's life, with prepubescent girls and postmenopausal women exhibiting lower levels of *Lactobacillus* dominance compared to women of reproductive age, though their bacterial communities are distinct from the CST IV typically seen during reproductive years [7]. On a short timescale, daily CST fluctuations are observed in some women of reproductive age, while others remain remarkably stable across menstrual cycles, suggesting that diverse factors influence the dynamics of vaginal microbiota communities [11]. For example, menstruation

is a key driver of monthly dynamics, while clinical interventions such as antibiotics and probiotics can cause temporary perturbations [7].

A significant gap in the existing literature is the understanding of the long-term dynamics of vaginal microbiota in reproductive-aged women across several months. While some studies do follow this timespan, they focus on pregnancy-specific dynamics [12, 13], have large intervals between samples (often exceeding three months) [14], or involve modest sample sizes [15]. These limitations hinder our ability to fully understand the long-term patterns of CST stability and transitions in the general population of reproductive-aged women, and the influence of clinically relevant factors such as demography, lifestyle, sexual practices, and medication.

In this study, we introduce an original follow-up cohort of 125 women in Montpellier, France. Our cohort presents a high-resolution longitudinal follow-up study with 2,103 microbial samples, spanning a median duration of over 8.6 months and a median of 11 samples per woman. We devise a hierarchical Bayesian Markov model to estimate transition probabilities between CSTs, associations between the transitions and 16 relevant covariates, and individual variability among women.

Materials and methods

Longitudinal clinical data

The samples originated from the PAPCLEAR monocentric longitudinal cohort study, which
followed 189 women longitudinally between 2016 and 2020. The participants were recruited
through posters and leaflets circulated at the main sexually transmitted infection detection
centre (CeGIDD) at the University Hospital of Montpellier (CHU) and at and around
university campuses in the city. The inclusion criteria were to be between 18 and 25 years
old, to be living in the area of Montpellier, France, to be in good health (no chronic disease),
not to have a history of human papillomavirus (HPV) infection (e.g., genital warts or high-
grade cervical lesion), and to report at least one new sexual partner over the last 12 months.
Additional details about the protocol can be found elsewhere [16]. The longitudinal data
analysed in the present study are available at `https://doi.org/10.57745/FHQR9Z`.

The inclusion visit was performed by a gynaecologist or a midwife at the CeGIDD
outside operating hours. After an interview, several samples were collected, including
vaginal swabs with eSwabs (Coppan) in Amies preservation medium from which microbiota
barcoding was later performed. The samples were aliquoted right after the visit and stored
at -20°C, before being transferred to -70°C within a month. The participants also filled in
a detailed questionnaire, which formed the basis of epidemiological covariates analysed in
this study.

Subsequent on-site visits were scheduled every two or four months, depending on the

HPV status. In between on-site visits, women were asked to perform eight self-samples at ~~100~~

home with eSwabs in Amies medium and to keep them in their freezer. The self-samples ~~101~~

were brought back in an isotherm bag at the next visit. These were then stored with the ~~102~~

swab at -70°C until processing. ~~103~~

Microbiota metabarcoding and quantification ~~104~~

The microbiota metabarcoding was performed on $200\mu L$ of vaginal swabs specimen stored ~~105~~

at -70° in Amies medium. The DNA extraction was performed using the MagAttract ~~106~~

PowerMicrobiome DNA/RNA kit (Qiagen). Next-generation sequencing of the V3-V4 ~~107~~

region of the 16S gene [17] was performed on an Illumina HiSeq 4000 platform (150 base ~~108~~

pairs paired-end mode) at the Genomic Resource Center at the University of Maryland ~~109~~

School of Medicine. ~~110~~

The taxonomic assignment was performed using the internal software package Speci- ~~111~~

ateIT (`https://github.com/Ravel-Laboratory/speciateIT`) and the community state ~~112~~

type was determined using the VALENCIA software package [9]. To examine longitudinal ~~113~~

patterns, the present study included participants who contributed at least three samples: ~~114~~

125 women met the inclusion criterion, giving 2,103 samples in total. ~~115~~

Covariates ~~116~~

In the PAPCLEAR study, a questionnaire was given to each participant to record patient- ~~117~~

level meta-data. We initially considered the following covariates based on previously pro- ~~118~~

posed roles in influencing the vaginal milieu: 119

1st menstr. Number of years since the first menstruation: The morphology of the human vagina 120
changes throughout life and the onset of puberty marks a key event that triggers 121
cascading changes [18]. 122

Alcohol Average number of glasses of alcoholic drinks consumed per week: Chronic presence 123
of alcohol in the genital environment has been linked to a shift in the immune and 124
microbiological conditions [19]. 125

Antibio. Application of antibiotics during the study, either systemic (*Antibio. (Systemic)*) 126
or genital (*Antibio. (Genital)*): The bacterial composition responds rapidly and 127
transiently to antibiotic treatments that target bacteria either broadly or with a 128
narrow taxonomic scale [20]. 129

BMI Body mass index (BMI): Obesity has been implicated in elevating vaginal microbiota 130
diversity and promoting *Prevotella* associated with BV [21]. 131

Caucasian Identity as Caucasian ethnicity or other: Ethnicity has been linked to variation in 132
vaginal microbiota compositions in several studies [4]. However, causal mechanisms 133
remain an open question. 134

Cigarettes Cigarette smoking: Smoking has been implicated in the development of BV due to its 135
anti-estrogenic effects and the presence of harmful substances such as benzo[a]pyrene 136
diol epoxide (BPDE). [22]. 137

Horm. contra. Use of hormonal contraception during the study: The vaginal hormonal landscape is 138
affected by the use of hormonal contraceptives [23]. 139

Lubricant Use of lubricant during the study: Personal lubricants contain various chemicals that 140
differentially impact the growth of vaginal microbes in-vitro [24]. 141

Menstr. cup Use of menstrual cups during the study: The vaginal microenvironment may be 142
altered by the use of menstrual cups both physically and chemically. An elevated 143
risk of fungal infections has been reported [25]. 144

Partners Cumulative number of sexual partners: The genital microbiome can be transferred 145
between sexual partners [26]. Such an external input could destabilise the resident 146
community. 147

Red meat Average number of meals that include red meat consumption per week: Diet alters 148
the vaginal environment for microbes. An unhealthy diet, linked to a high proportion 149
of red meat consumption, has been linked to an elevated risk of BV [27]. 150

Regular condom Regular use of condoms during sexual intercourse: Condom use can modify the vagi- 151
nal microenvironment by altering the exchange of microbes between partners [28]. 152

Regular sport Engaging in regular sporting activities, over 50% of the time: Physical activities 153
influence immune responses, with leisure-time physical activity associated with a 154
reduced risk of suspected bacterial infections compared to sedentary behaviour [29]. 155

Stress   Average stress level reported from 0 (min) to 3 (max): Stress hormones may disrupt [156] vaginal flora, for instance, by inhibiting glycogen production, which is the primary [157] fuel for lactobacilli [30]. [158]

Tampon   Use of tampons during the study: The use of internal menstrual health products [159] like tampons directly alters the vaginal environment, although negative effects from [160] tampon use are seldom reported [31]. [161]

Vag. product   Use of vaginal cream/tablet/capsule/gel/wipe during the study: Women frequently [162] use over-the-counter vulvovaginal treatments that contain a variety of chemical components. [163] However, the clinical effectiveness of these products in preventing BV is [164] seldom systematically evaluated [32]. [165]

Chlamydia   Tested positive for chlamydia. [166]

Female/male affinity   Affinity to female/male partner: Genital microbiome transfers during sexual activity [167] are anticipated to vary based on the genders of the partners [33]. [168]

Pregnancy   History of pregnancy: Pregnancy significantly changes the cervicovaginal environment, [169] with increased estrogen from the ovaries and placenta leading to higher vaginal [170] glycogen. This supports the growth of *Lactobacillus* species [34]. [171]

Spermicide   Use of spermicide during the study: Spermicides are chemicals that prevent sperm [172] from reaching an egg, but their use can change the vaginal microflora, potentially [173] increasing the risk of genitourinary infections [35]. [174]

Vag. douching Use of vaginal douching during study: Vaginal douching, the practice of washing inside the vagina with a liquid solution, has been shown to increase the risk of disturbing the natural balance of vaginal flora [36].

Out of the covariates initially considered above, we excluded six (*Chlamydia*, *Female affinity*, *Male affinity*, *Pregnancy*, *Spermicide* and *Vag. douching*) as data were severely skewed towards the most common value (> 90% of data). During the study, any use of antibiotics was recorded with the date and we distinguished systemic (*Antibio. (Systemic)*) and genital topical (*Antibio. (Genital)*) applications, corresponding to 'Gynecological anti-infectives and antiseptics' ('G01' ATC codes), which consisted of metronidazole treatments, and 'Antibacterials for systemic use' ('J01' ATC codes), which were more diverse. Since the exact dates of treatment were recorded, *Antibio. (Systemic)* and *Antibio. (Genital)* were included as time-inhomogenous covariates in the model. All other covariates were considered time-homogeneous meaning that the variation is among women, and static through time because the precise timing of changes in the covariate values was unknown.

To facilitate the comparison of covariate effects, we centred and scaled continuous variables [37] and deviation-coded binary variables. These transformations ensure that all covariates are modelled in a comparable scale and the intercept is located at a "representative reference value" of the modelled population: i.e., the population mean for continuous and the theoretical mid-point for binary values. Four continuous covariates (i.e., *Alcohol*, *BMI*, *Partners*, and *Red meat*) were log-transformed before scaling due to their right-

skewed distribution. We found no strong correlations among the covariates included in the analysis (Supplementary Information S1).

Modelling

Markov model

Markov models are statistical models used to represent systems that transition between discrete states over time. These models are 'memoryless', meaning that the probability of transition to another state depends on the current state, but not its historical path. In clinical research, these models are often used to predict the transitions among health states (e.g., health, illness and remission), and the propensity to transition between these states is estimated from longitudinal follow-up data. Clinical follow-up data are typically modelled using the continuous-time Markov model [38], in which the probability of transition over a given interval depends on the instantaneous transition intensity and the amount of time spent in the current state.

Vaginal microbiota state transitions are classically studied using continuous-time Markov models [12–14, 39, 40]. Our application of the continuous-time Markov model differs from those of the existing literature in its hierarchical Bayesian formulation, which allowed us to quantify individual variability among women (as unobserved heterogeneity, or random effects) and to estimate many covariate effects simultaneously (through the use of weakly informative priors).

Transition intensities

Transition intensities refer to the instantaneous rate of moving from state i to state j in a participant p (e.g., CST I to CST IV), a process that may be affected by a vector of covariates, X. Taking the form of a proportional hazards model, these rates can be expressed as:

$$q_{p,i,j} = \text{Exp}(\mu_{p,i,j} + \beta_{i,j} \, X), \tag{1}$$

where $\mu_{p,i,j}$ is the intercept and $\beta_{i,j}$ is the coefficient expressing the impact of a covariate(s). This intercept is further defined by the equation,

$$\mu_{p,i,j} = (\hat{\mu}_{i,j} + s_{p,i,j}) \cdot \mu_{sd} + \bar{\mu}, \tag{2}$$

where $\bar{\mu}$ and μ_{sd} are the prior mean and standard deviation of the intercept such that $\hat{\mu}_{i,j} \cdot \mu_{sd} + \bar{\mu}$ constitutes the non-centred parameterisation of the population-level intercept, $\mu_{i,j}$ and is assumed to be normally distributed, i.e., $\hat{\mu}_{i,j} \sim \mathcal{N}(0,1)$.

Additionally, we allowed for unobserved heterogeneity in μ, i.e., $s_{p,i,j}$, where

$$s = \text{diag}(sd_s) \cdot L_s \cdot z_s. \tag{3}$$

We sampled from the corresponding weakly informative priors, namely $sd_s \sim t_4(0,1)$,

$L_s \sim \text{LKJCorrCholesky}(2)$, and $z_s \sim \mathcal{N}(0, 1)$, as recommended by the Stan development ₂₁₉

community [41, 42]. ₂₂₀

For regression coefficients, the Student-t distributions with degrees of freedom 4 to 7 are ₂₂₁

recommended as generic, weakly informative, priors [42]: we sampled β from $\beta \sim t_4(0, 1)$, ₂₂₂

which places a comparatively wide tail within the recommendation. As all of our covariates ₂₂₃

have been proposed to impact vaginal microbiota communities *a priori* (see above), we did ₂₂₄

not strongly regularise the priors, for example, through the use of horseshoe priors [43]. ₂₂₅

We note that all covariates were modelled simultaneously, such that the interpretation of ₂₂₆

each coefficient is conditional upon other covariates included and accounts for the influence ₂₂₇

of other factors. We assumed that the covariates affect the transitions symmetrically (i.e., ₂₂₈

$\beta_{j,i} = -\beta_{i,j}$), meaning that the influence of a covariate on the affinity (or aversion) towards ₂₂₉

a particular CST is consistent, regardless of the direction of the transition. ₂₃₀

Collectively, the transition intensities form the Q matrix, in which the sum of intensities

across a row, i.e., all transitions from a particular state, is defined to be zero, such that we

have the following equation for the diagonal entries:

$$q_{i,i} = -\sum_{j \neq i} q_{i,j}. \tag{4}$$

Transition probabilities and likelihood

Taking the matrix exponential of the Q matrix for each participant, we compute the matrix P such that:

$$P = \text{Exp}\big((t_{k+1} - t_k)\,Q\big), \tag{5}$$

where k represents the sample identity for a given individual. The P matrix contains the transition probabilities between two observations (at k and $k+1$) and $t_{k+1}-t_k$ indicates the elapsed time between two observations.

Finally, the probability of observing a given state at the next sampling event (i.e., at $k+1$) is modelled by the categorical distribution:

$$y_{k+1} \sim \text{Categorical}\big(P[y_k,]\big) \tag{6}$$

where $P[y_k,]$ is the $y_k{}^{\text{th}}$ row of the P matrix containing the probabilities of transition from the state observed at k.

Model fitting

We used a Bayesian approach to fit the above continuous-time Markov model to lon- gitudinal data of vaginal microbiota CSTs. Our model was written in Stan 2.26.1 and fitted through the RStan interface 2.32.3 [44]. The Stan programme is available at https:

One participant lacked information on the years since their initial menstruation. We imputed missing values using the mice package [45] and generated 10 imputed datasets to be fitted separately. For each imputed dataset, we fitted the model in parallel using four independent chains, each with $10,000$ sampled iterations and $1,000$ warm-up iterations. The MCMC samples from separate runs (i.e., based on differently imputed data) were subsequently combined for inference.

We confirmed over $1,000$ effective samples per imputed dataset and ensured convergence of independent chains ($\hat{R} < 1.01$) for all parameters [46]. We carried out a posterior predictive check by comparing the observed and predicted CST frequency. We also quantified the posterior z-score and posterior contraction to examine the accuracy and precision of posterior distributions and the relative strength of data to prior information [47] (Supplementary Information S2).

Counterfactual predictions

We took advantage of the parameterised model to simulate the population-level outcomes of each covariate, assuming that all covariates, but a focal one, are at the representative reference value (as described above) and then varying the focal parameter within the range of values observed in the studied cohort. The model predictions were generated by randomly drawing 100 samples from the posterior distributions and simulating the Markov model for each sampled parameter set. We focused on the CST frequency as the outcome

of interest. ₂₆₁

Results and Discussion ₂₆₂

CSTs in the cohort ₂₆₃

Our longitudinal dataset from the PAPCLEAR cohort represents one of the largest analysed ₂₆₄ to date in the context of the vaginal microbiota. Detailed participant characteristics are ₂₆₅ presented in Table 1. Briefly, the participants were between 18 and 25 years old and the ₂₆₆ majority of the 2,103 samples (73.7%) were self-collected at home, the rest being collected ₂₆₇ during on-site visits (Fig. 1a). The median follow-up duration was 8.64 months and the ₂₆₈ most common intervals between analysed samples were seven and 28 days (Fig. 1a & b). ₂₆₉ On average, each of the 125 participants contributed 11 samples (Fig. 1c). ₂₇₀

The metabarcoding analysis on 16S RNA with the VALENCIA algorithm [9] was used ₂₇₁ to assign each sample to a CST. The vaginal microbiota communities were variable across ₂₇₂ women and over time (Fig. 1d). As CSTs I, II, and V are all dominated by lactobacilli ₂₇₃ and considered 'optimal' in terms of health, yet the latter two are rare ($\sim$4% of all sam- ₂₇₄ ples combined), we pooled the three optimal communities for further investigation. Over- ₂₇₅ all, optimal communities were the most frequent, representing 44.5% of samples, followed ₂₇₆ by 'sub-optimal' (CST III) at 35.2% and 'non-optimal' communities (CST IV) at 20.4% ₂₇₇ (Fig. 1e and Table 1). ₂₇₈

Figure 1: **Summary of vaginal microbiota samples analysed in the PAPCLEAR study.** a) Intervals between sampling events for clinical (i.e., on-site) and home samples. b) Follow-up duration per participant. c) Number of samples analysed per participant. d) Vaginal microbiota Community State Types (CST) over time in 125 participants. e) Frequency of the optimal (CSTs I, II, and V), sub-optimal (CST III) and non-optimal (CST IV) communities in all samples.

We implemented a continuous-time Markov model to capture the CST dynamics. Simula- 280

tions based on the estimated parameters of our model (i.e., a posterior predictive check) 281

confirmed that it accurately captures the observed CST prevalence (Fig. 2a). The optimal, 282

Table 1: Summary profile of vaginal microbiota samples and covariates in the PAPCLEAR study. Q1 and Q3 refer to first (25%) and third (75%) quantiles. Level = 1 indicates the presence of a binary condition. See Materials and Methods for the covariate definitions.

	Level	Summary
Samples (Participants)		2103 (125)
CST (%)	I	847 (40.3)
	II	39 (1.85)
	III	740 (35.2)
	IV-A	54 (2.57)
	IV-B	342 (16.3)
	IV-C	32 (1.52)
	V	49 (2.33)
Sample type (%)	On-site	553 (26.3)
	Home	1550 (73.7)
Sampling interval (median (Q1,Q3))		21 (7, 28)
Follow-up duration (median (Q1,Q3))		8.64 (5.36, 14.0)
Samples per subject (median (Q1,Q3))		11 (7, 16)
Covariates		
Identifying as 'Caucasian' (%)	1	102 (81.6)
BMI (median (Q1,Q3))		21.19 (19.78, 23.46)
Alcohol (median (Q1,Q3))		3.14 (1.40, 5.07)
Smoker (%)	1	36 (28.8)
Stress level (from 0 to 3, median (Q1,Q3))		1.41 (1.00, 1.75)
Regular sport practice (%)	1	61 (48.8)
Red meat consumption (times per week, median (Q1,Q3))		0.50 (0.16, 1.00)
Years since 1st menstruation (median (Q1,Q3))		9 (7, 10)
Hormonal contraception (%)	1	32 (25.6)
Menstrual cup user (%)	1	46 (36.8)
Vaginal product user (%)	1	73 (58.4)
Tampon user (%)	1	89 (71.2)
Lifetime number of partners (median (Q1,Q3))		5 (3, 11)
Lubricant use (%)	1	58 (46.4)
Regular condom use by partner (%)	1	23 (18.4)
Male affinity (%)	1	124 (99.2)
Chlamydia infection at inclusion (%)	1	7 (5.6)
Pregnancy during follow-up (%)	1	4 (3.2)
Vaginal douching (%)	1	4 (3.2)
Spermicide user (%)	1	1 (0.8)
Female affinity (%)	1	10 (8.0)
Systemic antibiotic treatment (%)	1	65 (52.0)
Genital antibiotic treatment (%)	1	30 (24.0)

Figure 2: **Prevalence and transition probabilities among vaginal microbiota community state types (CSTs).** a) Observed (bars) and predicted prevalence (crosses) of CSTs I (II, V), III and IV. The model predictions were generated by drawing 100 random samples from the posterior distributions and simulating the Markov model for each sampled parameter set. b) Mean estimated weekly transition probabilities of CSTs I (II, V), III and IV. The arrow thickness indicates the persistence or transition probability.

CST I(II, V), and sub-optimal, CST III, communities showed a high degree of stability, with weekly probabilities to remain in the current state estimated at 87% (95% credibility interval (95CrI): 78 - 93%) and 81% (95CrI: 68 - 90%), respectively (Fig. 2b). In contrast, the weekly persistence probability of the non-optimal CST IV was 60% (95CrI: 35 - 80%, Fig. 2b). These transition probabilities translate into sojourn times (i.e., the expected time spent in a given state before moving to another) in CST I (II, V), III and IV of 6.9 days (95CrI: 2.9 - 13.6 days), 4.23 days (95CrI: 1.8 - 8.4 days) and 1.6 days (95CrI: 0.58 - 3.8 days), respectively.

The reported persistence and transition probabilities in the literature vary widely based on the cohort characteristics. For example, focusing on women during pregnancy, DiGiulio et al. [12] estimated that the four *Lactobacillus*-dominated CSTs (CSTs I, II, III, and V) were more stable than CST IV. Notably, both CST I and II showed 98% probability of weekly persistence. The enhanced persistence of *Lactobacillus*-dominated communities during pregnancy owes to specific vaginal conditions during pregnancy including the up-regulation of oestrogen and progesterone that facilitates lactobacilli [12, 48].

In addition, the temporal dynamics of vaginal microbiota are notably different in women with BV. In contrast to pregnant women, those experiencing symptomatic BV generally exhibit less stable vaginal microbiota communities. In the cohort of Ravel et al. [49], which focused on women with symptomatic BV, Brooks et al. [39] found significantly lower stability across all CSTs. The probability of these CSTs persisting ranged from 38% to 48%, with CST I persisting only 46% of the time over a week.

Among studies that focused on non-pregnant, healthy young women — with no partic-ular emphasis on BV — the analysis by Brooks et al. [39] of the Chaban et al. cohort [15] (N = 27; Canada) estimated weekly persistence probabilities of 75% for CST I, 78% for III, 60% for IV-A, and 88% for V. In the Gajer et al. dataset [11] (N = 32; USA) analysed again by Brooks et al., [39], CST I, II and III demonstrated 72%, 84% and 77% weekly persis-tence probabilities, respectively. In this dataset, CST IV sub-categories showed markedly different stability with CST IV-A with weekly persistence of 38% and CST IV-B with persistence of 82%. A third study, Munoz et al. [14] (N = 88; South Africa), reported

the stability of vaginal microbiota in women in a three-month time frame using a differ- 312
ent microbiota classification system consisting of four categories predominantly associated 313
with: *L. crispatus* (similar to CST I), *L. iners* (similar to CST III), *G. vaginalis* (similar 314
to CST IV), or *Prevotella* spp. (similar to CST IV). They found similar persistence for 315
CST I and CST IV-like communities ranging from 51 to 53% over three months while the 316
CST III-like community was more stable at 62% over the same period. Recasting in the 317
three-month time scale, our estimates show the same extent of stability for CST I(II, V) at 318
51% (95% CrI: 29-72%) while CST III (38%, 95% CrI: 19-61%) and CST IV (15%, 95% CrI: 319
5-34%) were less stable. Taken together, our estimates of vaginal microbiota community 320
stability are within the range of values reported in other cohorts. However, the dynamics 321
of vaginal microbiota communities are likely geographically variable even among healthy 322
young women. 323

Covariate effects on transitions 324

The Bayesian approach, which can accommodate vaguely informative priors on the co- 325
variate effects, allows for the simultaneous inclusion of many covariates (as hazard ratios; 326
Eq. 1) which would otherwise prove difficult in Markov models [38]. We identified 16 co- 327
variates based on previously proposed roles in influencing the vaginal milieu and assumed 328
that covariates have a symmetrical effect on CST transitions: e.g., the magnitude of a given 329
covariate effect on the transition from CST I to III is identical to that on the transition 330
from CST III to I. We identified alcohol consumption as the strongest and most consistent 331

effect while several other covariates were identified as possible drivers of CST transitions. 332

Alcohol consumption 333

The estimated hazard ratios on community transitions indicate that alcohol consumption 334

favoured the sub-optimal (CST III) community over optimal (CST I(II, V)) with 97% 335

probability (Fig. 3). Because of our symmetry assumption, this can mean that alcohol 336

consumption increases the pace of transition from CST I(II, V) to CST III or reduces that 337

in the opposite direction by the same magnitude. Alcohol consumption also tended to 338

favour CST IV over CST III, although with a lower credibility level (with 73% probability 339

of the hazard ratio $\neq 1$, Fig. 3). 340

To examine how these effects translate to the population level, we carried out counter- 341

factual simulations in which all participant characteristics were set to the representative 342

value observed in the studied cohort, except for alcohol consumption, which ranged from 343

non-drinking to the level of the heaviest drinking observed in our cohort (19 drinks per 344

week). The simulations demonstrated that the expected prevalence of the optimal (CST I 345

(II, V)) community was 18% (95% CrI of 9 to 27%) higher in a hypothetical population 346

of non-drinkers compared to that of average-level drinkers who consumed three drinks per 347

week (Fig. 4; Supplementary Information S3). In turn, the prevalence of the optimal com- 348

munity was 19% (95% CrI of 10 to 29%) higher in the population of average-level drinkers 349

than in the heaviest drinkers. As the optimal community declined with alcohol consump- 350

tion, the prevalence of the non-optimal (CST IV) community was found to be 9% (95% CrI 351

Figure 3: **Estimated covariate effects on community transition rates.** With the symmetry assumption, there are three sets of covariate effects on transitions. The impact of covariates on community transition rates was estimated for a given set of community states as the log hazard ratio, β. The figure shows the posterior distributions of $\exp(\beta)$, the hazard ratio for the three sets of transition sets, and the corresponding 16 covariates. The numbers on the right-hand side of each panel indicate the probability that the estimated effect is different from the hazard ratio of 1 (i.e., the proportion of posterior distributions sampled on the dominant side of the effect).

of 2 to 15%) higher among average drinkers compared to non-drinkers. Therefore, while the strongest impact of alcohol on community transitions appears to be between the optimal (CST I, (II, V)) and sub-optimal (CST III) communities, an additional, non-zero impact on the sub-optimal to non-optimal (CST IV) transition means that alcohol consumption ultimately promotes non-optimal communities at the expense of optimal ones. As the effects of covariates are estimated simultaneously, potential confounding factors, including the number of partners, condom use and smoking, are controlled for in our findings.

Alcohol consumption may impact the vaginal microbiota through a variety of mecha-

Figure 4: **Counter-factual simulations predict population-level consequences of covariates.** Based on estimated hazard ratios (Fig. 3), the population-level impact (i.e., the prevalence of CST I (II,V), III and IV) was simulated for each covariate. The vertical dashed lines indicate the intercept used in estimation: i.e., the population mean for continuous and midpoint for binary variables. For continuous variables, the range of values explored was determined by the minimum and maximum values reported in the PAP-CLEAR study.

nisms. Physiologically, the chronic presence of alcohol in the genital environment has been 360

linked to a shift in immune and microbiological conditions [19]. In addition, alcohol is a 361

known modifier of sexual behaviour, which in turn has been demonstrated to increase the risk of BV, linked to CST IV [50]. Finally, alcohol alters the microbial profile in other body parts, which in turn could cross over to the vaginal milieu. For example, *Prevotella*, a genus commonly found in CST IV communities, is enriched in the oral microbiota of drinkers [51]. Similarly, others postulate the effect of alcohol on the gut microbiota may have a concurrent influence on the vaginal microbiota [52].

While there remains a lack of consensus among existing studies (briefly reviewed by Froehle et al. [52]), cohort and cross-sectional studies from diverse geographical contexts (namely, Australia, Denmark, Sweden, Thailand, Tanzania, Uganda and USA) have previously reported an association between alcohol consumption and BV [52–60]. In addition to corroborating these findings, our Markov model offers a novel insight into the ecology of microbial communities underlying these observations: alcohol consumption destabilises the optimal (CST I (II, V)) communities towards sub-optimal (CST III), which opens the gate for the deterioration towards non-optimal (CST IV), associated with BV. To the authors' knowledge, there have been no alcohol cessation studies reporting its impact on vaginal microbiota. Such studies are necessary to establish causal links, similar to those conducted on the effects of smoking [61], douching [62], and antibiotics [20] on vaginal microbiota compositions.

Potential effects of other covariates Other factors with possible effects on transitions (i.e., with more than 80% probability of hazard ratio $\neq 1$) included the use of vaginal intimate hygiene products, number of sexual partners and self-reported 'Caucasian' identity. **Vaginal hygiene products:** The use of vaginal hygiene products appeared to have multifaceted effects. Between CST I(II, V) and CST III, their use was positively linked to maintaining or transitioning to CST I(II, V) with 90% probability. For the CST I(II, V) and CST IV pair, it tended to favour a shift towards CST IV, with 85% probability. Finally, between CST III and CST IV, their use was more likely to support the persistence or a shift towards CST III, also with 90% probability. The circular effects suggest that women may experience different effects of the products marketed for 'vaginal intimate hygiene' depending on associations with certain CSTs. Nonetheless, the circular effects on community transitions meant that there was no noticeable impact at the population level in our counterfactual simulations (Fig. 4). **Number of sexual partners:** A higher number of sexual partners was also found to potentially favour CST IV over CST III, increasing the risk of maintaining (or transi- tioning to) CST IV with 89% probability of the hazard ratio $\neq 1$. The association between CST IV and the lifetime number of partners is consistent with the hypothesis that exter- nal importation of microbes could alter the dynamics of vaginal microbiota and is in line with earlier work [63]. Population-level simulations predict that an increasing number of 28

sexual partners tends to reduce the prevalence of the sub-optimal (CST III) community. 400

For example, CST III was 13% (95% CrI of 2 to 21%) less common in a hypothetical pop- 401

ulation with the highest number of partners than one conforming to the average number. 402

The decrease was accompanied by a tendency for the other CSTs to increase, although the 403

trend was less clear for individual CSTs (Fig. 4). 404

Causasian identity: It is worth noting that our cohort was not designed to achieve 405

comprehensive coverage of self-reported ethnic identity, with over 80% identifying as Cau- 406

casians (Table 1). Nonetheless, identifying oneself as a 'Caucasian' tended to favour CST III 407

over CST I(II, V) with 86% probability. European studies focusing on the role of ethnicity 408

are rare. However, a North American study has observed a qualitatively opposite trend: 409

CST III communities are comparably rare in women who identify themselves as Caucasian 410

compared to those identifying as Asian, Black and Hispanic (26.8 versus 42.7, 31.4 and 411

36.1%, respectively [4]). While previous studies have revealed differences in vaginal mi- 412

crobiota compositions among ethnic groups, the relative importance of biological, societal, 413

and environmental factors remains an open question [4, 64–66]. 414

Antibiotics: Notably, we found little association between antibiotic consumption and 415

CST transitions, neither for local treatment for BV (genital application of metronidazole) 416

nor systemic treatment (antibiotic treatment via oral intake). Such a lack of effect in 417

our study may be because the changes in the vaginal microbiota compositions following an 418

antibiotic treatment take place in a shorter time scale than our sampling intervals: the most 419

common sampling intervals were either 7 or 28 days (Table 1). In comparison, Brooks et 420

al. [39] found rapid CST transition following BV medication in the cohort of Ravel et al. [49], 421

which involved daily sampling. On a longer time scale, the re-emergence of BV-associated 422

communities following treatments is a well-documented clinical challenge [67–69]. 423

Unobserved individual variability in community transition 424

While we incorporated 16 covariates into our Markov model, some variations among women 425

remain unaccounted for. To quantify these, we estimated the extent of individual variabil- 426

ity (i.e., unobserved heterogeneity, or random effects) in community transitions for each 427

transition pair using a hierarchical Bayesian approach (Eq.2 & 3). 428

The highest variability was observed among women in the transitions involving 'recov- 429

ery' to an optimal (CST I (II,V)) from a non-optimal (CST IV) state (Fig. 5). On the 430

other hand, inverse transitions from non-optimal to optimal exhibited some of the lowest 431

individual variability. The same is true, although to a lesser extent, for the shifts from 432

sub-optimal (CST III) to optimal. These findings suggest that there are relatively lim- 433

ited pathways leading to the deterioration of vaginal microbiota communities, whereas the 434

routes to recovery can be more individualised and the source of this variation remains to 435

be fully elucidated. 436

The presence of individual-level random effects indicates that a considerable part of 437

the variability remains unaccounted for by the 16 covariates in this study. One possible 438

cause is that our study left out key drivers of the vaginal milieu. For example, while 439

menstrual cycles have been demonstrated to influence daily and weekly transitions [11], 440

Figure 5: **Individual-level variability in vaginal microbiota community state type (CST) transitions.** a) The population average (thick black) and individual (thinner colours) weekly transition probabilities. Between-women individual variation for transitions to b) a more optimal and c) a less optimal state. Colours indicate the type of transition between CSTs.

they were omitted from our analysis because the timing of menstruation was ambiguous   441

in the PAPCLEAR study. Secondly, further resolution on individual variability may be   442

gained by incorporating time-varying covariates, which could accommodate changes in   443

participant behaviours during the follow-up. In continuous-time Markov models, time-   444

varying covariates are assumed piecewise constant, meaning they are constant between   445

sampling events [38]. Such an assumption is convenient as covariate values are rarely   446

known between sampling events. Without precise knowledge of the timing of the covariate   447

changes, however, it is unclear whether the previous covariate value (at $t-1$) or the new   448

covariate value (at t) should influence the transition. Consequently, our analysis focused on static covariates, with the exception of antibiotic treatments for which the exact application dates were known. Aggregating participant behaviours as static covariates eliminated the uncertainty of covariate dynamics, albeit at a potentially lost opportunity for further precision.

Limitations and opportunities

A potential limitation from a clinical methodological perspective is that the majority of samples were collected at home during the PAPCLEAR study. While home sampling could introduce variability, the participants were provided with detailed instructions to minimise the difference in swabbing techniques between on-site and home samples, and we verified consistency in sampling dates by having participants fill out online questionnaires during sampling.

Another possible limitation of our study is the resolution of microbiota community classification. We focused on three CST groups with varying health implications: optimal (CST I(II,V)), sub-optimal (III), and non-optimal (IV). This decision stemmed from the fact that detailed classifications in a Markov model would increase the number of possible transitions, and it would be difficult to estimate transitions between rare types. However, significant functional differences may exist within these CSTs. For instance, the VALENCIA algorithm classifies subcategories within some CSTs [9], and Brooks et al. demonstrated that CST IV-B is more stable than CST IV-A [39]. Further precision of mi-

crobiota classification may be obtained through the metagenomics approach. For instance, 469
identifying lineage replacements in women with stable CSTs and investigating the impact 470
of antibiotic treatments on the prevalence of resistance genes could yield insights into the 471
within-species dynamics of vaginal microbes. 472

Finally, a promising direction for future research is the joint analysis of CST dynamics 473
and sexually transmitted infections such as HPV. Previous studies have found a weak 474
association between CST IV and HPV detection risk [70]. However, these studies tested 475
the CST effect after estimating transition rates and pooled all high-risk and low-risk HPVs, 476
making it difficult to identify coinfections or reinfections. The PAPCLEAR cohort, with 477
genotype-specific follow-ups, could provide new insights into the link between CST and 478
HPV infection, potentially identifying causal relationships. 479

Conclusion 480

We showcased a novel application of a hierarchical Bayesian Markov model to original 481
clinical cohort data of vaginal microbiota dynamics. Our approach facilitated the simulta- 482
neous estimation of several covariate effects on community transitions and the identification 483
of unobserved variability in these transitions. Our work paves the way for an improved 484
ecological understanding of microbial dynamics within the vaginal environment and the 485
development of preventative and therapeutic strategies to improve vaginal health. 486

Ethics

This study has been approved by the Comité de Protection des Personnes (CPP) Sud Méditerranée I (reference number 2016-A00712-49); by the Comité Consultatif sur le Traitement de l'Information en matière de Recherche dans le domaine de la Santé (reference number 16.504); by the Commission Nationale Informatique et Libertés (reference number MMS/ABD/ AR1612278, decision number DR-2016–488), by the Agence Nationale de Sécurité du Médicament et des Produits de Santé (reference 20160072000007), and is registered at ClinicalTrials.gov under the ID NCT02946346.

Funding

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 648963, to SA). The authors acknowledge further support from the Fondation pour la Recherche Medicale (to TK), the Agence Nationale de la Recherche contre le SIDA (ANRS-MIE, to NT), and the MemoLife Labex (to BE).

Acknowledgements

The authors thank Olivier Supplisson for his helpful feedback. We acknowledge the ISO 9001 certified IRD i-Trop HPC (member of the South Green Platform) at IRD Montpellier for providing HPC resources that have contributed to the research results reported

within this article (bioinfo.ird.fr and www.southgreen.fr). DNA extracts were (partly) performed through the genotyping and sequencing facilities of ISEM (Institut des Sciences de l'Evolution-Montpellier) and Labex Centre Méditerranéen Environnement Biodiversité.

References

[1] van de Wijgert JHHM. The vaginal microbiome and sexually transmitted infections are interlinked: Consequences for treatment and prevention. PLOS Medicine. 2017;14(12):e1002478.

[2] Haahr T, Zacho J, Bräuner M, Shathmigha K, Skov Jensen J, Humaidan P. Reproductive outcome of patients undergoing in vitro fertilisation treatment and diagnosed with bacterial vaginosis or abnormal vaginal microbiota: a systematic PRISMA review and meta-analysis. BJOG. 2019;126(2):200–207.

[3] Bilardi JE, Walker S, Temple-Smith M, McNair R, Mooney-Somers J, Bellhouse C, et al. The burden of bacterial vaginosis: women's experience of the physical, emotional, sexual and social impact of living with recurrent bacterial vaginosis. PLOS ONE. 2013;8(9):e74378.

[4] Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, et al. Vaginal microbiome of reproductive-age women. Proceedings of the National Academy of Sciences. 2011;108:4680–4687.

[5] France MT, Fu L, Rutt L, Yang H, Humphrys MS, Narina S, et al. Insight into the ecology of vaginal bacteria through integrative analyses of metagenomic and meta-transcriptomic data. Genome Biology. 2022;23(1):66.

[6] Coudray MS, Madhivanan P. Bacterial vaginosis—A brief synopsis of the literature. European Journal of Obstetrics & Gynecology and Reproductive Biology. 2020 Feb;245:143–148.

[7] France M, Alizadeh M, Brown S, Ma B, Ravel J. Towards a deeper understanding of the vaginal microbiota. Nature Microbiology. 2022;7(3):367–378.

[8] Petrova MI, Reid G, Vaneechoutte M, Lebeer S. *Lactobacillus iners*: Friend or Foe? Trends in Microbiology. 2017;25(3):182–191.

[9] France MT, Ma B, Gajer P, Brown S, Humphrys MS, Holm JB, et al. VALENCIA: a nearest centroid classification method for vaginal microbial communities based on composition. Microbiome. 2020;8(1):166.

[10] Cancelo-Hidalgo MJ, Coello LB. Genitourinary syndrome of the menopause: vaginal health and microbiota. In: Cano A, editor. Menopause: a comprehensive approach. Cham; 2017. p. 91–107.

[11] Gajer P, Brotman RM, Bai G, Sakamoto J, Schütte UM, Zhong X, et al. Temporal dynamics of the human vaginal microbiota. Science Translational Medicine. 2012;4(132):132ra52–132ra52.

[12] DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, et al. Temporal and spatial variation of the human microbiota during pregnancy. Proceedings of the National Academy of Sciences. 2015;112(35):11060–11065.

[13] Serrano MG, Parikh HI, Brooks JP, Edwards DJ, Arodz TJ, Edupuganti L, et al. Racioethnic diversity in the dynamics of the vaginal microbiome during pregnancy. Nature Medicine. 2019;25(6):1001–1011.

[14] Munoz A, Hayward MR, Bloom SM, Rocafort M, Ngcapu S, Mafunda NA, et al. Modeling the temporal dynamics of cervicovaginal microbiota identifies targets that may promote reproductive health. Microbiome. 2021;9(1):1–12.

[15] Chaban B, Links MG, Jayaprakash TP, Wagner EC, Bourque DK, Lohn Z, et al. Characterization of the vaginal microbiota of healthy Canadian women through the menstrual cycle. Microbiome. 2014;2:1–12.

[16] Murall CL, Rahmoun M, Selinger C, Baldellou M, Bernat C, Bonneau M, et al. Natural history, dynamics, and ecology of human papillomaviruses in genital infections of young women: protocol of the PAPCLEAR cohort study. BMJ Open. 2019;9(6):e025129.

[17] Frank JA, Reich CI, Sharma S, Weisbaum JS, Wilson BA, Olsen GJ. Critical evaluation of two primers commonly used for amplification of bacterial 16S rRNA genes. Applied and Environmental Microbiology. 2008;74(8):2461.

[18] Farage M, Maibach H. Lifetime changes in the vulva and vagina. Archives of Gynecology and Obstetrics. 2006;273:195–202.

[19] Loganantharaj N, Nichols WA, Bagby GJ, Volaufova J, Dufour J, Martin DH, et al. The effects of chronic binge alcohol on the genital microenvironment of simian immunodeficiency virus-infected female rhesus macaques. AIDS Research and Human Retroviruses. 2014;30(8):783–791.

[20] Mayer BT, Srinivasan S, Fiedler TL, Marrazzo JM, Fredricks DN, Schiffer JT. Rapid and profound shifts in the vaginal microbiota following antibiotic treatment for bacterial vaginosis. Journal of Infectious Diseases. 2015;212(5):793–802.

[21] Si J, You HJ, Yu J, Sung J, Ko G. *Prevotella* as a hub for vaginal microbiota under the influence of host genetics and their association with obesity. Cell Host & Microbe. 2017;21(1):97–105.

[22] Brotman RM, He X, Gajer P, Fadrosh D, Sharma E, Mongodin EF, et al. Association between cigarette smoking and the vaginal microbiota: a pilot study. BMC Infectious Diseases. 2014;14(1):1–11.

[23] Achilles SL, Austin MN, Meyn LA, Mhlanga F, Chirenje ZM, Hillier SL. Impact of contraceptive initiation on vaginal microbiota. American Journal of Obstetrics and Gynecology. 2018;218(6):622–e1.

[24] Laniewski P, Owen KA, Khnanisho M, Brotman RM, Herbst-Kralovetz MM. Clinical and personal lubricants impact the growth of vaginal lactobacillus species and colonization of vaginal epithelial cells: an in vitro study. Sexually Transmitted Diseases. 2021;48(1):63–70.

[25] Tessandier N, Uysal IB, Elie B, Selinger C, Bernat C, Boué V, et al. Does exposure to different menstrual products affect the vaginal environment? Molecular Ecology. 2023;32(10):2592–2601.

[26] Vodstrcil LA, Twin J, Garland SM, Fairley CK, Hocking JS, Law MG, et al. The influence of sexual activity on the vaginal microbiota and *Gardnerella vaginalis* clade diversity in young women. PLOS ONE. 2017;12(2):e0171856.

[27] Noormohammadi M, Eslamian G, Kazemi SN, Rashidkhani B. Association between dietary patterns and bacterial vaginosis: a case–control study. Scientific Reports. 2022;12(1):12199.

[28] Hutchinson KB, Kip KE, Ness RB. Condom use and its association with bacterial vaginosis and bacterial vaginosis-associated vaginal microflora. Epidemiology. 2007;p. 702–708.

[29] Pape K, Ryttergaard L, Rotevatn TA, Nielsen BJ, Torp-Pedersen C, Overgaard C, et al. Leisure-time physical activity and the risk of suspected bacterial infections. Medicine and Science in Sports and Exercise. 2016;48(9):1737–1744.

[30] Amabebe E, Anumba DO. Psychosocial stress, cortisol levels, and maintenance of vaginal health. Frontiers in Endocrinology. 2018;p. 568.

[31] Klebanoff MA, Nansel TR, Brotman RM, Zhang J, Yu KF, Schwebke JR, et al. Personal hygienic behaviors and bacterial vaginosis. Sexually Transmitted Diseases. 2010;37(2):94.

[32] Van Kessel K, Assefi N, Marrazzo J, Eckert L. Common complementary and alternative therapies for yeast vaginitis and bacterial vaginosis: a systematic review. Obstetrical & Gynecological Survey. 2003;58(5):351–358.

[33] Ma ZS. Microbiome transmission during sexual intercourse appears stochastic and supports the red queen hypothesis. Frontiers in Microbiology. 2022;12:789983.

[34] Juliana NC, Peters RP, Al-Nasiry S, Budding AE, Morré SA, Ambrosino E. Composition of the vaginal microbiota during pregnancy in women living in sub-Saharan Africa: a PRISMA-compliant review. BMC Pregnancy and Childbirth. 2021;21(1):1–15.

[35] Gupta K, Hillier SL, Hooton TM, Roberts PL, Stamm WE. Effects of contraceptive method on the vaginal microbial flora: a prospective evaluation. Journal of Infectious Diseases. 2000;181(2):595–601.

[36] Brotman RM, Klebanoff MA, Nansel TR, Andrews WW, Schwebke JR, Zhang J, et al. A longitudinal study of vaginal douching and bacterial vaginosis—a marginal structural modeling analysis. American Journal of Epidemiology. 2008;168(2):188–196.

[37] Gelman A. Scaling regression inputs by dividing by two standard deviations. Statistics in Medicine. 2008;27(15):2865–2873.

[38] Christopher H Jackson. Multi-State Models for Panel Data: The msm Package for R. Journal of Statistical Software. 2011;38(8):1–29.

[39] Brooks JP, Buck GA, Chen G, Diao L, Edwards DJ, Fettweis JM, et al. Changes in vaginal community state types reflect major shifts in the microbiome. Microbial Ecology in Health and Disease. 2017;28(1):1303265.

[40] Fettweis JM, Serrano MG, Brooks JP, Edwards DJ, Girerd PH, Parikh HI, et al. The vaginal microbiome and preterm birth. Nature Medicine. 2019;25(6):1012–1021.

[41] Stan Development Team. Stan Functions Reference; 2023. Accessed: 2024-02-24. Available from: `https://mc-stan.org/docs/functions-reference/index.html`.

[42] Stan Development Team. Prior Choice Recommendations; 2023. Accessed: 2024-02-15. `https://github.com/stan-dev/stan/wiki/Prior-Choice-Recommendations`.

[43] Piironen J, Vehtari A. Sparsity information and regularization in the horseshoe and other shrinkage priors. Electronic Journal of Statistics. 2017;11(2):5018–5051.

[44] Stan Development Team. RStan: the R interface to Stan; 2023. R package version 2.32.3. Available from: `https://mc-stan.org/`.

[45] van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R. Journal of Statistical Software. 2011;45(3):1–67.

[46] Stan Development Team. The Stan Core Library; 2018. Version 2.18.0. Available from: http://mc-stan.org/17.

[47] Betancourt M. Towards a principled Bayesian workflow; 2020. Available from: https://betanalpha.github.io/assets/case_studies/principled_bayesian_workflow.html.

[48] Odogwu NM, Onebunne CA, Chen J, Ayeni FA, Walther-Antonio MR, Olayemi OO, et al. *Lactobacillus crispatus* thrives in pregnancy hormonal milieu in a Nigerian patient cohort. Scientific Reports. 2021;11(1):18152.

[49] Ravel J, Brotman RM, Gajer P, Ma B, Nandy M, Fadrosh DW, et al. Daily temporal dynamics of vaginal microbiota before, during and after episodes of bacterial vaginosis. Microbiome. 2013;1(1):1–6.

[50] Fethers KA, Fairley CK, Hocking JS, Gurrin LC, Bradshaw CS. Sexual risk factors and bacterial vaginosis: a systematic review and meta-analysis. Clinical Infectious Diseases. 2008;47(11):1426–1435.

[51] Liao Y, Tong XT, Jia YJ, Liu QY, Wu YX, Xue WQ, et al. The effects of alcohol drinking on oral microbiota in the Chinese population. International Journal of Environmental Research and Public Health. 2022;19(9):5729.

[52] Froehle L, Ghanem KG, Page K, Hutton HE, Chander G, Hamill MM, et al. Bacterial vaginosis and alcohol consumption: a cross-sectional retrospective study in Baltimore, Maryland. Sexually transmitted diseases. 2021;48(12):986–990.

[53] Smart S, Singal A, Mindel A. Social and sexual risk factors for bacterial vaginosis. Sexually Transmitted Infections. 2004;80(1):58–62.

[54] Thorsen P, Vogel I, Molsted K, Jacobsson B, Arpi M, Møller BR, et al. Risk factors for bacterial vaginosis in pregnancy: a population-based study on Danish women. Acta Obstetricia et Gynecologica Scandinavica. 2006;85(8):906–911.

[55] Shoubnikova M, Hellberg D, Nilsson S, Mårdh PA. Contraceptive use in women with bacterial vaginosis. Contraception. 1997;55(6):355–358.

[56] Rugpao S, Sriplienchan S, Rungruengthanakit K, Lamlertkittikul S, Pinjareon S, Werawatakul Y, et al. Risk factors for bacterial vaginosis incidence in young adult Thai women. Sexually Transmitted Diseases. 2008;35(7):643–648.

[57] Baisley K, Changalucha J, Weiss HA, Mugeye K, Everett D, Hambleton I, et al. Bacterial vaginosis in female facility workers in north-western Tanzania: prevalence and risk factors. Sexually Transmitted Infections. 2009;85(5):370–375.

[58] Francis SC, Looker C, Vandepitte J, Bukenya J, Mayanja Y, Nakubulwa S, et al. Bacterial vaginosis among women at high risk for HIV in Uganda: high rate of recurrent diagnosis despite treatment. Sexually Transmitted Infections. 2016;92(2):142–148.

[59] Cu-Uvin S, Hogan JW, Warren D, Klein RS, Peipert J, Schuman P, et al. Prevalence of lower genital tract infections among human immunodeficiency virus (HIV)—seropositive and high-risk HIV-seronegative women. Clinical Infectious Diseases. 1999;29(5):1145–1150.

[60] French AL, Adeyemi OM, Agniel DM, Evans CT, Yin MT, Anastos K, et al. The association of HIV status with bacterial vaginosis and vitamin D in the United States. Journal of Women's Health. 2011;20(10):1497–1503.

[61] Brotman RM, He X, Gajer P, Fadrosh D, Sharma E, Mongodin EF, et al. Association between cigarette smoking and the vaginal microbiota: a pilot study. BMC Infectious Diseases. 2014;14(1):471.

[62] Brotman RM, Ghanem KG, Klebanoff MA, Taha TE, Scharfstein DO, Zenilman JM. The effect of vaginal douching cessation on bacterial vaginosis: a pilot study. American Journal of Obstetrics and Gynecology. 2008 Jun;198(6):628.e1–628.e7.

[63] Sobel JD, Sobel R. Current and emerging pharmacotherapy for recurrent bacterial vaginosis. Expert Opinion on Pharmacotherapy. 2021;22(12):1593–1600.

[64] Zhou X, Brown CJ, Abdo Z, Davis CC, Hansmann MA, Joyce P, et al. Differences in the composition of vaginal microbial communities found in healthy Caucasian and black women. ISME J. 2007;1(2):121–133.

[65] Fettweis JM, Brooks JP, Serrano MG, Sheth NU, Girerd PH, Edwards DJ, et al. Differences in vaginal microbiome in African American women versus women of European ancestry. Microbiology. 2014;160(Pt 10):2272.

[66] Borgdorff H, Veer Cvd, Houdt Rv, Alberts CJ, Vries HJd, Bruisten SM, et al. The association between ethnicity and vaginal microbiota composition in Amsterdam, the Netherlands. PLOS ONE. 2017;12(7):e0181135.

[67] Lambert JA, John S, Sobel JD, Akins RA. Longitudinal analysis of vaginal microbiome dynamics in women with recurrent bacterial vaginosis: recognition of the conversion process. PLOS ONE. 2013;8(12):e82599.

[68] Srinivasan S, Liu C, Mitchell CM, Fiedler TL, Thomas KK, Agnew KJ, et al. Temporal variability of human vaginal bacteria and relationship with bacterial vaginosis. PLOS ONE. 2010;5(4):e10197.

[69] Armstrong E, Hemmerling A, Joag V, Huibner S, Kulikova M, Crawford E, et al. Treatment success following standard antibiotic treatment for bacterial vaginosis is not associated with pre-treatment genital immune or microbial parameters. Open Forum Infectious Diseases. 2023 Jan;p. ofad007.

[70] Brotman RM, Shardell MD, Gajer P, Tracy JK, Zenilman JM, Ravel J, et al. Interplay Between the Temporal Dynamics of the Vaginal Microbiota and Human Papillomavirus Detection. Journal of Infectious Diseases. 2014;210(11):1723–1733.

[71] Team SD. Stan Functions Reference; 2024. Accessed: 2024-02-05. `https://mc-stan.` 709
`org/docs/functions-reference`. 710

Supplementary Information

Competing interests

SAlizon is a recommender for PCI Evolutionary Biology and PCI Ecology.

JReynes reports personal fees from Gilead (consulting and payment or honoraria for lec-
tures, presentations, speaker's bureaus, manuscript writing, or educational events), Janssen
(payment or honoraria for lectures, presentations, speaker's bureaus, manuscript writing,
or educational events), Merck (payment or honoraria for lectures, presentations, speaker's
bureaus, manuscript writing, or educational events), Theratechnologies (payment or hon-
oraria for lectures, presentations, speaker's bureaus, manuscript writing, or educational
events), and ViiV Healthcare (consulting and payment or honoraria for lectures, presen-
tations, speaker's bureaus, manuscript writing, or educational events) and support for
attending meetings and/or travel from Gilead and Pfizer, outside of the submitted work.

JRavel is co-founder of LUCA Biologics, a biotechnology company focusing on translat-
ing microbiome research into live biotherapeutics drugs for women's health. He is Editor-
in-Chief at *Microbiome*.

None of the other authors report any conflict of interest.

S1: Pairwise correlations between covariates

There were no strong correlations among covariates, with the strongest correlation found
between BMI and stress ($r = 0.41$; Fig. S1).

Figure S1: Correlation between covariates. Pairwise Pearson's correlation coefficients between covariates

S2: Assessment of posterior accuracy, precision and prior contraction

We leveraged the properties of posterior distributions to identify potential model fitting problems that might manifest from our model assumptions. To examine the accuracy and precision of posterior distributions, we first generated simulated observations based on the estimated posterior mean parameters. We then refitted our model to the simulated observations (i.e., secondary fitting) to compute the posterior z-score for each parameter, which measures how closely the posterior recovers the parameters of the data generating process [47]:

$$z = \frac{\mathbb{E}_{\text{sim}} - \mathbb{E}_{\text{post}}}{\sigma_{\text{sim}}},$$

where $\mathbb{E}_{\text{post}}$ denotes the posterior mean of the fit to the actual data that we consider the [731] 'true' parameter. $\mathbb{E}_{\text{sim}}$ and σ_{sim} denote the mean and standard deviation of the posterior [732] distribution of the secondary fitting. The smaller the z-score, the closer the bulk of the [733] posterior is to the true parameter [47]. In contrast, large z-values may be indicative of [734] overfitting and, or poor prior specifications [47]. [735]

To examine the influence of the likelihood function in relation to prior information, we computed the posterior contraction, k:

$$k = 1 - \frac{\sigma_{\text{post}}^2}{\sigma_{\text{prior}}^2}$$

where σ_{post}^2 and σ_{prior}^2 correspond to the variance of posterior and prior distributions, respectively. The k values close to zero indicate that data contain little information (i.e., rendering priors strongly informative). Conversely, values close to 1 indicate that data are much more informative than the prior [47].

We found that most of our model parameters and hyperparameters — were estimated with accuracy, precision, and identifiability, with the absolute posterior z-scores below three (Fig. S2). Some individual variations in transition rates, sd_s showed a tendency towards overfitting (the absolute posterior z-scores above three). Thus, caution might be warranted when interpreting the extent of between-women variation in CST transition rates, a small number of z-scores exceeding the absolute number of three is unlikely to be a cause of concern [47]. We found that the posterior distributions for covariate coefficients, β, contracted by over 75% compared to the prior distribution, for all but one covariate effect, meaning that the covariate coefficients were well-identified from data (Fig. S2). Although we used generic priors recommended by Stan [71], the L_s parameters that define correlations among between-woman variation showed limited posterior contraction (i.e., $\leq \sim 0.25$), indicating that these parameters are poorly informed by data. As such, we refrain from making biological inferences about these correlations.

Figure S2: Accuracy, precision and identifiability of estimated parameters. Posterior z-score (y-axis) measures how closely the posterior recovers the parameters of the true data-generating process and posterior contraction (x-axis) evaluates the influence of the likelihood function over the prior, respectively. Smaller absolute posterior z-scores indicate that the posterior accurately recovers the parameters of the data-generating process: the absolute value beyond three to four may indicate substantial bias [47]. The posterior contraction values close to one indicate that data are much more informative than the prior. The estimated parameters are represented by a filled dot.

S3: Predicted difference in community state type (CST) prevalence at various counterfactual scenarios.

Our counterfactual simulations predicted that alcohol consumption and the number of partners are factors that impact the population-level outcome in terms of the prevalence of different community state types. The full list of comparisons is available in Fig. S3.

Figure S3: Difference in community state type (CST) prevalence at predicted various counterfactual scenarios. The differences were calculated from posterior samples simulated at 0 and 1 for binary variables and at the population maximum and minimum values recorded by the PAPCLEAR for continuous variables (left panel). Additional differences were computed between the population maximum and mean (middle panel) and the population mean and minimum for continuous variables (right panel).