The Efficacy of Psychological Interventions for Child and Adolescent PTSD: A Network Meta-Analysis

Thole H. Hoppen*[1], Ph.D., Lena Wessarges[1], B.Sc., Marvin Jehn[1], M.Sc., Julian Mutz[2], Ph.D., Ahlke Kip[1], Ph.D., Pascal Schlechter, Ph.D. [1], Richard Meiser-Stedman[3], Ph.D., & Nexhmedin Morina[1], Ph.D.

[1]Institute of Psychology, University of Münster, Münster, Germany

[2]Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom

[3]Department of Clinical Psychology and Psychological Therapies, Norwich Medical School, University of East Anglia, Norwich, United Kingdom

Short Title: Efficacy of psychological interventions for pediatric PTSD

*Corresponding Author
Thole H. Hoppen, Ph.D.
Institute of Psychology
University of Münster
Fliednerstr. 21
48149 Münster
Germany
Tel: +492518339415
Fax: +492518331331
e-Mail: thoppen@uni-muenster.de

28 **Key Points**
29
30 **Question:** How efficacious are psychological treatments for pediatric PTSD?
31
Findings: Trauma-focused cognitive behavior therapies (TF-CBTs) are currently the most
evaluated treatment for pediatric PTSD (74% of included studies). Data for other
interventions are emerging. At short-term, TF-CBTs, Eye Movement Desensitization and
Reprocessing (EMDR), non-trauma-focused interventions, and multi-disciplinary treatments
(MDTs) all significantly reduced pediatric PTSD relative to no treatment. TF-CBTs produced
the strongest short- and long-term treatment effects. EMDR and MDTs had insufficient long-
term data.
39
40 **Meaning:** TF-CBTs should be the first-line treatment recommendation for pediatric PTSD.
41 While data for other treatment approaches emerged with some promising findings, more data
42 (including long-term data) are needed to draw firmer conclusions.

Abstract

Importance: Pediatric post-traumatic stress disorder (PTSD) is a common and debilitating mental disorder. Yet, a comprehensive network meta-analysis examining the efficacy of psychological interventions is lacking.

Objective: To synthesize all available evidence on psychological interventions for pediatric PTSD in a comprehensive network meta-analysis.

Data Sources: PsycINFO, MEDLINE, Web of Science, and PTSDpubs were searched from inception to January 2nd 2024 and 74 related systematic reviews were screened.

Study Selection: Two independent raters screened publications for eligibility. Inclusion criteria were: Randomized controlled trial (RCT) with ten or more patients per arm examining a psychological intervention for pediatric PTSD compared to a control group in children and adolescents (< 19 years) with full or subthreshold PTSD.

Data Extraction and Synthesis: PRISMA guidelines were followed to synthesize and present evidence. Two independent raters extracted data and assessed risk of bias with Cochrane criteria. Random effects network meta-analyses were run.

Main Outcome and Measures: Standardized mean differences (Hedges' g) in PTSD severity.

Results: In total, 70 RCTs (N = 5,528 patients) were included. Most RCTs (74%) examined trauma-focused cognitive behavior therapy interventions (TF-CBTs). At treatment endpoint, TF-CBTs, EMDR, multi-disciplinary treatments (MDTs), and non-trauma-focused interventions (non-trauma-focused interventions) were all efficacious in reducing PTSD when compared to passive control conditions, with large pooled effects ($gs \geq 0.86$, all $ps <$ .001) in the random effects network meta-analysis. TF-CBTs produced the strongest short-term effects relative to both passive and active control conditions and across all sensitivity analyses. In a sensitivity analysis including only trials with parent involvement, TF-CBTs were significantly more efficacious in reducing PTSD than non-trauma-focused interventions ($g = 0.35$, $p = .026$). Results for mid-term (up to 5 months posttreatment) and long-term data (6-24 months posttreatment) were very similar.

Conclusions and Relevance: The present network meta-analysis is the most comprehensive summary of psychological treatments for pediatric PTSD to this date. Results confirm that TF-CBTs can efficaciously reduce PTSD symptom severity in children and adolescents in the short-, mid-, and long-term. More long-term data are needed for EMDR, MDTs, and non-trauma-focused interventions. Results of TF-CBTs are encouraging and disseminating these results may help reduce common treatment barriers.

One to two-thirds of children and adolescents from the general population report

exposure to at least one traumatic event in their lifetimes.[1-4] While most children and

adolescents react resiliently to trauma, about one-fifth develop post-traumatic stress disorder

(i.e., pediatric PTSD).[4,5] Pediatric PTSD is a common, impairing,[4] and often chronic[6] mental

disorder characterized by vivid re-experiencing of trauma, avoidance of trauma-related

stimuli, changes in cognitions and emotions, and hyperarousal symptoms.[7] Given the high

prevalence and disease burden of pediatric PTSD,[8-10] the examination and implementation of

efficacious treatments constitutes a public health priority.

International treatment guidelines recommend cognitive behavior therapies with a

trauma focus (TF-CBTs, e.g., TF-CBT[11], prolonged exposure[12], cognitive processing

therapy[13]) as first-line treatment for pediatric PTSD.[14-19] Research on other psychological

interventions such as eye movement desensitization and reprocessing (EMDR) or non-

trauma-focused interventions is also emerging. In recent years, number of published RCTs

has increased substantially. To inform clinical practice about the relative efficacy of all

treatment approaches, a comprehensive network meta-analysis (NMA) is required.

NMAs integrate data from both direct (i.e., comparison of arms within a given RCT)

and indirect comparisons (i.e., comparisons of arms across RCTs), which enables conclusions

about relative effects of all interventions based on all data.[20] Three NMAs of psychological

interventions for pediatric PTSD have been published.[21-23] However, four omissions in

previous works need to be addressed. First, a comprehensive NMA is needed. Caro et al.[23]

only focused on pediatric PTSD relating to sexual abuse (rather than any trauma type).

Mavranezouli et al.[22] analyzed follow-up data only up to four months posttreatment and

cannot discern long-term efficacy. Xiang et al.[21] included data from 56 RCTs published until

2020, compared to 70 RCTs published until 2024 in the present work. Second, no previous

NMA included a sensitivity analysis of low risk of bias evidence. Low quality evidence may

bias results in quantitative synthesis.[24] Third, no NMA has performed sensitivity analyses

concerning delivery format (e.g., individual delivery only or treatments with parent/caregiver

involvement only) and efficacy may differ by delivery formats. Fourth, efficacy might also

differ by age group (i.e., children vs. adolescents), and no NMA has yet performed an age-

based sensitivity analysis. To enhance our understanding of relative treatment efficacy of

interventions, the present work addresses these four omissions.

116   **Methods**

117   We followed Preferred Reporting Items for Systematic Reviews and Meta-analyses

(PRISMA) 2015 guidelines.[25] The systematic literature search, data extraction, and risk of

bias assessment were carried out independently by at least two authors. Disagreements were

discussed between at least three authors (THH/LW/AK/NM). To address missing data, we

sent data requests to corresponding authors, with a reminder one month later. The objectives

and methods of the present NMA were pre-registered in the PROSPERO database (ID:

CRD42020206290). We defined the main research question (Population, Intervention,

Comparison, Outcome, and Study; PICOS) as follows: In children and adolescents with full

or subthreshold PTSD (P), how do psychological interventions (I), compared to passive

control conditions, active control conditions, or amongst different categories of interventions

(C), perform in terms of lowering PTSD symptom severity (O) in randomized controlled

trials (S)?

Identification and Selection of Studies

Search Strategy

132   For the timespan from inception to April 21st 2022, we relied on the literature search

of our previous work[26] including 57 eligible RCTs. We conducted a new search with identical

search strategy on January 2[nd] 2024 covering literature published between April 21[st] 2022 and

January 2[nd] 2024. Appendix A outlines the full search string. Briefly, we searched four large

bibliometric health sciences databases: PsycINFO, MEDLINE, Web of Science, and

PTSDpubs with various search terms for PTSD (ptsd OR ptss OR post-traumatic stress OR

posttraumatic stress) and treatment (trial* OR treatment* OR intervention*) in all-field

searches. Consistent with our previous search, no restrictions were applied to languages or

publication formats. We also screened 74 recently published related systematic reviews

(Appendix B) and the reference lists of included trials.

Eligibility Criteria

In line with our previous work,[26] we included trials that met all of the following

inclusion criteria: 1) RCT, 2) investigating the efficacy of a psychological intervention for

pediatric PTSD compared to a control condition, 3) all participants had PTSD complaints

(full or sub-threshold PTSD), 4) sample mean age < 19.0 years, and 5) outcome data reported

for at least ten participants per arm.

Quality Assessment

Risk of bias was independently assessed by two authors (THH/NM) based on eight

dichotomous quality criteria reported by Cuijpers and colleagues.[24] These eight criteria

originated partly from the Cochrane Collaboration criteria[27] and authoritative criteria for

evidence-based psychological interventions.[28] In the present study, RCTs were defined as

high-quality trials (i.e., with low risk of bias) when fulfilling at least six of the eight quality

criteria (for quality criteria and quality ratings per trial see Appendices C & D, respectively).

Initial agreement between independent raters was good (91.35%).

Data Extraction

Trial characteristics (e.g., treatment delivery format), sample characteristics (e.g.,

mean age), and PTSD outcome data were extracted independently by at least two authors

(THH/LW/AK/NM). When applicable, intent-to-treat (ITT) data were prioritized over

completer data. When applicable, outcome data from clinician-based measures were

prioritized over self-report-based data.

Categorization of Psychological Interventions and Control Conditions

The present study compared four categories of psychological interventions based on

the number of trials available: 1) TF-CBTs (i.e., any CBT-based intervention with a trauma

focus, such as TF-CBT,[11] prolonged exposure,[12] or cognitive processing therapy[13]), 2)

EMDR, 3) non-trauma-focused interventions (i.e., any intervention without a trauma focus),

and 4) multi-disciplinary treatments (MDTs, i.e., treatments that combine techniques from at

least two of the aforementioned categories of interventions, such as intensive multimodal

group program[29]). Other trauma-focused interventions (i.e., interventions with a trauma focus

but neither belonging to TF-CBTs or EMDR, such as expressive supportive groups[30]) were

planned as fifth category, but lacked evidence (kes < 4). In the present work, k denotes the

number of RCTs, whereas *kes* denotes the number of direct comparisons, which differ due to

multi-arm trials. Control conditions were divided into passive control conditions (e.g., wait-

list control) and active control conditions (e.g., treatment-as-usual, for all categorizations see,

Appendix E).

Categorization of Assessment Timepoints

To evaluate the short, mid, and long-term efficacy and consistent with previous

research,[31] we distinguished between three periods: 1) *post-treatment* (i.e., short-term), which

we defined as assessments at treatment endpoint, 2) *mid-term*, which we defined as

assessments of up to or equal to five months after the treatment endpoint, and 3) *long-term*,

which we defined as assessments longer than five months after the treatment endpoint. When

several assessments fell into the mid-term and long-term categories, the longest assessment

was chosen.

Outcomes

The primary outcome of interest was the standardized mean differences (Hedges' g)[32]

in PTSD symptom severity between comparator groups.

Statistical Analysis

Random effects NMAs were conducted given that high heterogeneity in outcomes

was expected.[32] Level of statistical significance was set to p-values below 0.05 (two-sided)

for all analyses, including Egger's test. Analyses were performed in R (version 4.1.1)[33] with

the netmeta package.[34] Effect sizes (Hedges' g) were first calculated at the study-level[35] and

then pooled and compared between all comparison dyads in NMA.[32] Following Cohen,[36] g

was interpreted as small (0.20), medium (0.50), and large effect size (0.80). We only included

intervention categories with minimally sufficient evidence base (i.e., $kes \geq 4$)[37] to ensure that

effects were also estimated via a sufficient number of direct comparisons.[38]

For the transitivity assumption, we examined whether the distribution of various trial

and sample characteristics was similar across comparison dyads and performed sensitivity

analyses. We analyzed inconsistency between direct and indirect evidence[38] globally (in the

network) and locally (per comparison dyad)[25,39] with the net splitting method[40] and inspected

net heat plots[41]. We performed inconsistency-corrected analyses when applicable. We

calculated outlier-adjusted NMAs whenever (≥ 1) outliers were detected (i.e., ≥ 3.3 standard

deviations above or below the pooled g)[42]. To examine the influence of small-study effects,

we performed Egger's test[43] and inspected comparison-adjusted funnel plots[44] (i.e.,

comparisons of interventions to passive and active control conditions only). We calculated

the I^2 and τ^2 statistics as estimates of overall heterogeneity,[45] and Q_{het} and Q_{inc} as estimates of

heterogeneity within and between comparison dyads.[46] We also calculated surface under the

cumulative ranking (SUCRA; 50,000 resamples), allowing for a ranking by efficacy. To

visualize distribution of available evidence, we build network graphs. In addition to main

NMAs (across all data), we performed four sensitivity NMAs: 1) only high-quality trials, 2)

only trials delivering treatment(s) individually, 3) only trials involving parents/caregivers in

treatment, and 4) only trials with sample mean age < 12 years (i.e., involving mainly

children) as well as only trials with sample mean age ≥ 12 years (i.e., involving mainly

adolescents).

222   **Results**

Study Selection Process

The new search wave covered 8,845 electronic records with ten additional eligible

RCTs. Thus, a total of 70 independent RCTs were eligible for the present purposes. Figure 1

details the study synthesis process.

Study Characteristics

The 70 RCTs reported data of 5,528 participants (for trial characteristics and their

references, see Appendices F & G, respectively). Apart from one dissertation,[47] all RCTs

were peer-reviewed. Only Dorsey et al.[48] reported on more than one RCT (i.e., four RCTs).

In total, 41 RCTs (59% of trials) delivered interventions individually and 29 RCTs (41% of

trials) involved parents or primary caregivers in treatments. Across trials reporting this

information, mean number of total sessions was 10 and total duration of treatments (i.e., total

sessions times length) was on average 11 hours ($SD = 6.5$ hours). In total, 40 RCTs (57% of

trials) assessed follow-up data (range = 1mo-24mo posttreatment). ITT PTSD data were

reported in 44 RCTs (63% of trials). In total, 41 RCTs (59%) were conducted in high-income

countries and the remaining 29 RCTs (41%) in low and middle-income countries. However,

two RCTs (conducted in the US and Germany) exclusively involved refugees originating

from low-income countries.[49,50] Across trials reporting this information ($k = 52$ or 74%), 90%

of the participants met diagnostic criteria for PTSD at baseline. Fifty-seven trials (81%)

involved mixed gender samples, whereas nine RCTs (13%) included only females and four

RCTs (6%) only males. Across all trials, 60% of participants identified as females. Average

age across trials was 12.21 years ($SD = 3.08$). In terms of trauma history, 31 RCTs (44%)

included a sample with varying trauma histories. In the other trials, only participants with a

particular trauma history were included, such as sexual assault ($k = 10$, 14%) or parental

death ($k = 6$, 9%).

Network Meta-analyses of Short, Mid, and Long-term Efficacy

Assumptions

Assumptions were mostly met. Apart from two analyses, no inconsistencies were

observed. In the main NMA on mid-term efficacy, significant inconsistency (i.e., between

direct and indirect evidence) was found for MDTs and a corrected analysis (i.e., without

MDTs) was performed. In the sensitivity NMA on mid-term efficacy of treatments with

individual delivery only, significant inconsistencies were found for all interventions

(precluding corrected re-analysis) and results are thus not reported. The distribution of

sample and methodological characteristics across comparison dyads is presented in Appendix

H.

Network Graphs

Figure 2 shows the network graphs for the NMAs on short-, mid-, and long-term

efficacy. Most available trials assessed TF-CBTs. Only TF-CBTs had enough accumulated

evidence to be analyzed across all three timepoints.

Network Meta-analysis of Short-term Efficacy

Table 1 provides all short-term results including sensitivity analyses. At treatment

endpoint, all therapy categories – TF-CBTs, EMDR, MDTs, and non-trauma-focused

interventions – were efficacious in treating pediatric PTSD compared to passive control

conditions, with *g*s ranging from 0.86 for non-trauma-focused interventions to 1.06 for TF-

CBTs (see Appendix I for the corresponding forest plot). Compared to active control

conditions, only TF-CBTs ($g = 0.55$, $p < .001$) and MDTs ($g = 0.43$, $p = .013$) were

efficacious (Appendix J). Differences in efficacy between treatment categories were not

significant, with few or no direct comparisons for most comparison dyads. Heterogeneity in

this main analysis of the short-term efficacy was large within and between comparison dyads

($\tau^2 = 0.14$, $I^2 = 68.9\%$; $Q_{total} = 196.06$, $df = 61$, $p < .001$; $Q_{het} = 173.76$, $df = 50$, $p < .001$;

$Q_{inc} = 21.40$, $df = 11$, $p = .030$). No significant inconsistencies were detected in the net

splitting method (Appendices K & L). No evidence for small-study effects was found

(Appendix M). Two outliers[51,52] investigating TF-CBTs were detected. Outlier-adjusted

analysis excluding these trials produced similar results (Appendix N).

Sensitivity Analyses for Short-term Efficacy

In high-quality trials only, the results for comparisons to passive control conditions

were similar, with effect sizes ranging from 0.80 for non-trauma-focused interventions to

1.05 for TF-CBTs. Only TF-CBTs ($g = 0.53$, $p < .001$) and EMDR ($g = 0.43$, $p = .047$)

produced significant short-term effects compared to active control conditions. In the

sensitivity analysis concerning trials with individual treatment delivery only, results were

similar to the main analysis with most favorable outcomes for TF-CBT. In the sensitivity

analysis concerning only trials with parent/caregiver involvement, results were also similar

for the comparison to passive control conditions, with all categories with sufficient evidence

for synthesis (TF-CBT, MDTs, & non-trauma-focused interventions) producing significant

effects. Compared to active control conditions, however, only TF-CBTs ($g = 0.42$, $p < .001$)

produced a significant effect. Moreover, TF-CBTs with parent/caregiver involvement were

more efficacious than non-trauma-focused interventions with caregiver involvement ($g =$

0.35, $p = .026$). In the sensitivity analysis of samples with mean age < 12 (mostly children),

results were very similar with all intervention categories with sufficient evidence (TF-CBTs,

EMDR, & non-trauma-focused interventions) yielding significant reductions in PTSD

compared with passive controls ($gs \geq 0.78$). Yet, only TF-CBTs produced a significant effect

compared to active controls ($g = 0.55$, $p < .001$). In the sensitivity analysis of samples with

mean age $\geq$ 12 (mostly adolescents), results were also very similar with TF-CBTs, EMDR,

and MDTs yielding significant reductions in PTSD compared with passive controls ($gs \geq$

0.93). Only TF-CBTs ($g = 0.53$, $p < .001$) and MDTs ($g = 0.46$, $p = .042$) produced

significant effects compared to active controls.

Network Meta-analyses of Mid and Long-term Efficacy

Table 2 provides all results. For non-trauma-focused interventions, too few direct

comparisons were available. At mid-term (up to 5 months posttreatment), TF-CBTs, EMDR,

and MDTs were efficacious compared to passive control conditions with *gs* ranging from

0.59 for MDTs to 0.95 for EMDR (Appendix O). Compared to active control conditions, only

EMDR ($g = 0.52$, $p = .032$) and TF-CBTs ($g = 0.45$, $p = .002$) produced significant mid-term

effects (Appendix P). Heterogeneity in this main analysis concerning mid-term efficacy was

large within and between comparison dyads ($\tau^2 = 0.15$, $I^2 = 66.4\%$; $Q_{total} = 68.49$, $df = 23$, $p <$

.001; $Q_{bet} = 49.40$, $df = 16$, $p < .001$; $Q_{inc} = 19.85$, df = 7, $p = .006$). There was no evidence

for small-study effects (Appendix Q). Significant inconsistency was detected for MDTs in the

net splitting method (Appendices R & S). Results remained similar to those of the main

analysis in a re-analysis excluding MDTs (Appendix T).

In the long-term (6-to-24 months posttreatment), only TF-CBTs and non-trauma-

focused interventions had sufficient evidence for synthesis. Compared to passive control

conditions, both TF-CBTs ($g = 0.76$, $p = .002$) and non-trauma-focused interventions ($g =$

0.71, $p = .014$) produced significant effects (Appendix U). Both TF-CBTs ($g = 0.55$, $p <$

.001) and non-trauma-focused interventions ($g = 0.50$, $p = .016$) were also efficacious when

compared to active control conditions (Appendix V). Heterogeneity in this main analysis

concerning long-term efficacy was large within and between comparison dyads ($\tau^2 = 0.11$, I^2

= 67.6%; $Q_{total} = 46.30$, $df = 15$, $p < .001$; $Q_{bet} = 32.33$, $df = 10$, $p < .001$; $Q_{inc} = 14.02$, df =

5, $p = .016$). No inconsistencies (Appendices W & X) and no evidence for small-study effects

(Appendix Y) were found.

Sensitivity Analyses for Mid- and Long-term Efficacy

At mid-term, sensitivity analysis on high-quality trials could be conducted with TF-

CBTs and EMDR only. EMDR ($g = 1.15$, $p < .001$) and TF-CBTs ($g = 1.06$, $p < .001$)

produced large treatment effects compared to passive control conditions. Compared to active

control conditions, however, only TF-CBTs ($g = 0.33$, $p = .029$) produced a significant

treatment effect. The sensitivity analysis concerning trials with individual treatment delivery

only was infeasible given detected inconsistency for all interventions. The sensitivity analysis

of trials with parent/caregiver involvement produced similar results to the main analysis. Yet,

TF-CBTs were found more efficacious than MDTs ($g = 0.67$, $p = .042$). The sensitivity

analysis of trials with mean age below 12 was infeasible due to lacking evidence. The

sensitivity analysis of trials with mean age ≥ 12 (mainly adolescents) produced similar results

to the main analysis. TF-CBT ($g = 0.76$, $p < .001$) and MDTs ($g = 0.70$, $p = .034$) both

produced significant effects relative to passive controls. Yet, only TF-CBT ($g = 0.54$, $p <$

$.001$) was found significantly superior to active controls.

At long-term, only TF-CBTs and non-trauma-focused interventions had sufficient

accumulated data. Passive control conditions were also lacking. Compared to active control

conditions in high-quality trials, only TF-CBTs produced a significant moderate-sized

treatment effect ($g = 0.53$, $p < .001$). Sensitivity analysis concerning trials with individual

treatment delivery only produced similar results to the main analysis. Sensitivity analysis

concerning trials with parent/caregiver involvement was infeasible due to lacking long-term

evidence (kes < 4). The sensitivity analysis of trials with mean age ≥ 12 was infeasible due to

lacking evidence. The sensitivity analysis of trials with mean age < 12 (mainly involving

children) produced similar results to the main analysis, with only TF-CBT (but not non-

trauma-focused interventions) producing significant effects compared to active control

conditions ($g = 0.54$, $p = .003$).

Ranking of Efficacy

Table 3 shows SUCRA rankings. TF-CBTs were the highest ranking category of

interventions at all timepoints and all analyses, except at mid-term when it was second to

EMDR.

Discussion

The present work synthesized data from 70 RCTs in a comprehensive NMA. TF-

CBTs are currently the most evaluated and the most efficacious category of treatments for

PTSD in children and adolescents, followed by (in this order) EMDR, MDTs, and non-

trauma-focused interventions. This supports recommendations of international treatment

guidelines for pediatric PTSD, including the International Society for Traumatic Stress

Studies[18] and the National Institute of Clinical Excellence.[17] Our review confirms and

extends previous NMAs[21-23] and pairwise meta-analyses.[26,53,54] TF-CBTs were efficacious in

treating pediatric PTSD relative to different comparators (passive and active control

conditions), across time (short-, mid-, and long-term follow-up), and when restricting

analyses to trials with high-quality, utilizing an individual delivered intervention, with parent

involvement, and to samples involving mainly children or adolescents, respectively. These

results are important for the training of therapist and treatment implementation in clinical

practice and might help in reducing common treatment barriers.

Our review also reveals remaining gaps in the literature. While short and mid-term

efficacy of EMDR was supported, our review highlights the lack of evidence concerning

long-term follow-up data. The present results therefore support some international treatment

guidelines[17,19] that list EMDR as second-line treatment recommendation. Data for two other

categories of interventions – MDTs and non-trauma-focused interventions – are emerging.

More data (including long-term data), however, are needed to robustly investigate the

(relative) efficacy of these two categories as well as EMDR. For the time being, only the

efficacy of TF-CBTs can be robustly supported given that about three quarters of the

available data concern TF-CBTs. There was some evidence for TF-CBTs producing

significantly higher effects than non-trauma-focused interventions. Yet, more data is needed

to draw firmer conclusions with regards to relative efficacy between categories of

interventions.

Limitations

Five limitations should be noted. First, the categories of non-trauma-focused interventions,

MDTs, and other trauma-focused interventions are heterogenous with regards to theoretical

foundations of included interventions. However, there is no solid ground for further sub-

categorization given the low number of available RCTs. As more RCTs accumulate, more

homogenous categorizations will become feasible. Second, we found evidence for

inconsistency in the NMA regarding mid-term efficacy data. However, results remained

similar in a consistency-corrected re-analysis. Third, our age group sensitivity analyses

provide approximate estimations of treatment effects for children and adolescents, as the

categorization was based on sample mean age. An individual patient data meta-analysis

would allow for a solid differentiation of effects for children and adolescents, which was

beyond the scope of the present work. Fourth, while the distinction between passive and

active control conditions is a strength of the present work, active control condition such as

treatment-as-usual can comprise very different elements, depending on the context. Future

research might be able to disentangle this heterogenous comparator group in a more fine-

grained way. Fifth, relatively low rates of reported ITT data are concerning and trialists are

encouraged to report ITT data (including ITT means and SDs) in future trials.

Conclusions

There is robust evidence that pediatric PTSD can be effectively treated by

psychological treatments, in particular TF-CBTs. A large evidence base for TF-CBTs

supports efficacy at short, mid, and long-term, relative to both passive and active control

conditions and in high-quality trials. A comparably thin evidence base also supports the short

and mid-term efficacy of EMDR and to a lesser extent MDTs and non-trauma-focused

interventions. More high-quality trials including long-term assessments of PTSD are needed.

Statements

Author Contributions: THH, NM, LW, AK, MJ, and JM had full access to all of the data in
the study and take responsibility for the integrity of the data and the accuracy of the data
analysis.
Concept and design: THH and NM.
Acquisition, analysis, or interpretation of data: All authors.
Drafting of the manuscript: THH and RM.
Critical review of the manuscript for important intellectual content: All authors.
Statistical analysis: THH, MJ, LW, JM.
Obtained funding: Not applicable (no funding).
Administrative, technical, or material support: THH.
Supervision: THH and NM.

Conflict of Interest Disclosures:
THH, MJ, LW, AK, PS and NM declare no competing interests. JM is funded by the National
Institute for Health and Care Research (NIHR) Maudsley Biomedical Research Centre at
South London and Maudsley NHS Foundation Trust and King's College London. The views
expressed are those of the authors and not necessarily those of the NHS, the NIHR or the
Department of Health and Social Care. RM occasionally receives payment (from universities
and private training providers) for training therapists in the delivery of cognitive therapy for
PTSD for children and adolescents; is a co-investigator or chief investigator on four National
Institute for Health and Care Research (NIHR)-funded or Medical Research Council-funded
clinical trials of psychological therapies, particularly cognitive therapy for PTSD in children
and young people; and was the chair of a steering committee for a trial addressing the online
treatment of PTSD in adults. RM institution (University of East Anglia) has received
payment through the following research grants: "Addressing the trauma-related distress of
young people in care: a randomised feasibility trial across social-care and mental health
services" (NIHR RfPB NIHR200586); "Internet-delivered Cognitive Therapy (iCT) for
young people with Post Traumatic Stress Disorder (PTSD)" (MRC DPFS MR/P017355/1);
"Supporting services to deliver trauma-focused cognitive behavioural therapy for care-
experienced young people: a pilot implementation study", NIHR Applied Research
Collaboration West; "Cognitive Behavioural Therapy for the treatment of post-traumatic
stress disorder (PTSD) in youth exposed to multiple traumatic stressors: a phase II
randomised controlled trial" (NIHR CDF-2015-08-073). RM institution part owns the
intellectual property for an online guided self-help version of cognitive therapy for PTSD for
children and young people as a result of RM involvement in one of these trials.

Funding/Support: This research received no specific grant from any funding agency,
commercial or not-for-profit sectors.

Role of Funder/Sponsor: Not applicable.

Data Sharing Statement: The datasets and the R script are available on request via email to
Dr. Thole H. Hoppen (thoppen@uni-muenster.de).

Additional Contributions: We thank the authors of the included studies for their work and
for responding to data request emails and the provision of data.

References

1. Copeland WE, Keeler G, Angold A, Costello EJ. Traumatic events and posttraumatic

stress in childhood. *Arch Gen Psychiatry*. 2007;64(5):577-584.

2. Saunders BE, Adams ZW. Epidemiology of traumatic experiences in childhood. *Child

Adolesc Psychiatr Clin*. 2014;23(2):167-184.

3. Danese A, Smith P, Chitsabesan P, Dubicka B. Child and adolescent mental health amidst

emergencies and disasters. *Br J Psychiatry*. 2020;216(3):159-162.

4. Lewis SJ, Arseneault L, Caspi A, et al. The epidemiology of trauma and post-traumatic

stress disorder in a representative cohort of young people in England and Wales. *Lancet

Psychiatry*. 2019;6(3):247-256.

5. Alisic E, Zalta AK, van Wesel F, et al. Rates of post-traumatic stress disorder in trauma-

exposed children and adolescents: meta-analysis. *Br J Psychiatry*. 2014;204(5):335-340.

doi:10.1192/bjp.bp.113.131227.

6. Widom CS. Posttraumatic stress disorder in abused and neglected children grown up. *Am

J Psychiatry*. 1999;156(8):1223-1229.

7. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders:

DSM-5*. 5. ed. Washington, DC: American Psychiatric Publishing; 2013.

8. Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and

consequences of child maltreatment in high-income countries. *Lancet*.

2009;373(9657):68-81.

9. Lansford JE, Dodge KA, Pettit GS, Bates JE, Crozier J, Kaplow J. A 12-year prospective

study of the long-term effects of early child physical maltreatment on psychological,

behavioral, and academic problems in adolescence. *Arch Pediatr Adolesc Med*.

2002;156(8):824-830.

10. Guiney H, Caspi A, Ambler A, et al. Childhood sexual abuse and pervasive problems

across multiple life domains: Findings from a five-decade study. *Dev Psychopathol*.

2024;36(1):219-235.

11. Cohen JA, Deblinger E, Mannarino AP, Steer RA. A multisite, randomized controlled

trial for children with sexual abuse–related PTSD symptoms. *Journal of the American*

Academy of Child & Adolescent Psychiatry. 2004;43(4):393-402.

12. Foa EB, McLean CP, Capaldi S, Rosenfield D. Prolonged exposure vs supportive

counseling for sexual abuse-related PTSD in adolescent girls: A randomized clinical trial.

JAMA. 2013;310(24):2650-2657.

13. Rosner R, Rimane E, Frick U, et al. Effect of developmentally adapted cognitive

processing therapy for youth with symptoms of posttraumatic stress disorder after

childhood sexual and physical abuse: A randomized clinical trial. *JAMA Psychiatry*.

2019;76(5):484-491.

14. Bandelow B, Allgulander C, Baldwin DS, et al. World Federation of Societies of

Biological Psychiatry (WFSBP) guidelines for treatment of anxiety, obsessive-compulsive

and posttraumatic stress disorders – Version 3. Part II: OCD and PTSD. *World J Biol*

Psychiatry. 2023;24(2):118-134. doi:10.1080/15622975.2022.2086296.

15. World Health Organization. *Guidelines on mental health promotive and preventive*

interventions for adolescents: helping adolescents thrive. Geneva: World Health

Organization; 2020.

16. Cohen JA, Issues, The Work Group On Quality, AACAP Work Group on Quality Issues.

Practice parameter for the assessment and treatment of children and adolescents with

posttraumatic stress disorder. *J Am Acad Child Adolesc Psychiatry*. 2010;49(4):414-430.

17. National Institute for Health and Care Excellence. Post-traumatic stress disorder (NICE

guideline NG116): [B] Evidence reviews for psychological, psychosocial and other non-

pharmacological interventions for the treatment of PTSD in children and young people.

https://www.nice.org.uk/guidance/ng116/resources/posttraumatic-stress-disorder-pdf-

66141601777861.

18. Bisson JI, Berliner L, Cloitre M, et al. The international society for traumatic stress

studies new guidelines for the prevention and treatment of posttraumatic stress disorder:

Methodology and development process. *J Trauma Stress*. 2019;32(4):475-483.

19. Phelps AJ, Lethbridge R, Brennan S, et al. Australian guidelines for the prevention and

treatment of posttraumatic stress disorder: Updates in the third edition. *Aust N Z J*

Psychiatry. 2022;56(3):230-247.

20. Cipriani A, Higgins JPT, Geddes JR, Salanti G. Conceptual and technical challenges in

network meta-analysis. *Ann Intern Med*. 2013;159(2):130-137.

21. Xiang Y, Cipriani A, Teng T, et al. Comparative efficacy and acceptability of

psychotherapies for post-traumatic stress disorder in children and adolescents: a

systematic review and network meta-analysis. *BMJ Ment Health*. 2021;24(4):153-160.

22. Mavranezouli I, Megnin☐Viggars O, Daly C, et al. Research Review: Psychological and

psychosocial treatments for children and young people with post☐traumatic stress

disorder: a network meta☐analysis. *J Child Psychol Psychiatry*. 2020;61(1):18-29.

23. Caro P, Turner W, Caldwell DM, Macdonald G. Comparative effectiveness of

psychological interventions for treating the psychological consequences of sexual abuse

in children and adolescents: a network meta☐analysis. *Cochrane Database Syst Rev*.

2023;(6):CD013361.

24. Cuijpers P, van Straten A, Bohlmeijer E, Hollon SD, Andersson G. The effects of

psychotherapy for adult depression are overestimated: a meta-analysis of study quality

and effect size. *Psychol Med*. 2010;40(2):211-223. doi:10.1017/S0033291709006114.

25. Hutton B, Salanti G, Caldwell DM, et al. The PRISMA extension statement for reporting

of systematic reviews incorporating network meta-analyses of health care interventions:

checklist and explanations. *Ann Intern Med*. 2015;162(11):777-784.

26. Hoppen TH, Meiser-Stedman R, Jensen TK, Birkeland MS, Morina N. Efficacy of

psychological interventions for post-traumatic stress disorder in children and adolescents

exposed to single versus multiple traumas: meta-analysis of randomised controlled trials.

Br J Psychiatry. 2023;222(5):196-203.

27. Higgins JPT, Green S, eds. *Cochrane handbook for systematic reviews of interventions*.

Repr. with corr. Chichester: Wiley-Blackwell; 2009. Cochrane book series.

28. Chambless DL, Hollon SD. Defining empirically supported therapies. *J Consult Clin*

Psychol. 1998;66(1):7-18. doi:10.1037//0022-006x.66.1.7.

29. Roque-Lopez S, Llanez-Anaya E, Álvarez-López MJ, et al. Mental health benefits of a 1-

week intensive multimodal group program for adolescents with multiple adverse

childhood experiences. *Child Abuse & Neglect*. 2021;122:105349.

30. Shechtman Z, Mor M. Groups for children and adolescents with trauma-related

symptoms: Outcomes and processes. *International Journal of Group Psychotherapy*.

2010;60(2):221-244.

31. Hoppen TH, Jehn M, Holling H, Mutz J, Kip A, Morina N. The efficacy and acceptability

of psychological interventions for adult PTSD: A network and pairwise meta-analysis of

randomized controlled trials. *J Consult Clin Psychol*. 2023;91(8):445-461.

32. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to meta-analysis*.

John Wiley & Sons; 2011.

33. R Core Team. R: A Language and Environment for Statistical Computing. http://www.R-

project.org/.

34. Balduzzi S, Rücker G, Nikolakopoulou A, et al. netmeta: An r package for network meta-
analysis using frequentist methods. *J Stat Soft*. 2023;106(2):1-40.

35. Lipsey MW, Wilson DB. *Practical meta-analysis*. [Nachdr.]. Thousand Oaks, Calif.:
SAGE Publ; 2009. Applied social research methods series; 49.

36. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hoboken: Taylor
and Francis; 2013. http://gbv.eblib.com/patron/FullRecord.aspx?p=1192162.

37. Hoppen TH, Meiser-Stedman R, Kip A, Birkeland MS, Morina N. The efficacy of
psychological interventions for adult post-traumatic stress disorder following exposure to
single versus multiple traumatic events: a meta-analysis of randomised controlled trials.
Lancet Psychiatry. 2024;11(2):112-122.

38. Donegan S, Williamson P, D'Alessandro U, Tudur Smith C. Assessing key assumptions
of network meta‐analysis: a review of methods. *Res Synth Methods*. 2013;4(4):291-323.

39. *Netmeta: Network meta-analysis using frequentist methods*; 2021.

40. Dias S, Welton NJ, Caldwell DM, Ades AE. Checking consistency in mixed treatment
comparison meta-analysis. *Stat Med*. 2010;29(7-8):932-944.

41. Krahn U, Binder H, König J. A graphical tool for locating inconsistency in network meta-
analyses. *BMC Medical Research Methodology*. 2013;13(1):1-18.

42. Tabachnick BG, Fidell LS. *Using multivariate statistics*. 6. ed., internat. ed. Boston,
Mass.: Pearson; 2013. Always learning.

43. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple,
graphical test. *BMJ*. 1997;315(7109):629-634.

44. Chaimani A, Salanti G. Using network meta‐analysis to evaluate the existence of
small‐study effects in a network of interventions. *Res Synth Methods*. 2012;3(2):161-
176.

45. DerSimonian R, Laird N. Meta-Analysis in Clinical Trials. *Controlled Clinical Trials*.

1986;7:177-188.

46. Cochran WG. The combination of estimates from different experiments. *Biometrics*.

1954;10(1):101-129.

47. Schauer E. *Trauma treatment for children in war: Build-up of an evidence-based large-*

scale mental health intervention in north-eastern Sri Lanka. University of Konstanz.

48. Dorsey S, Lucid L, Martin P, et al. Effectiveness of task-shifted trauma-focused cognitive

behavioral therapy for children who experienced parental death and posttraumatic stress

in Kenya and Tanzania: A randomized clinical trial. *JAMA Psychiatry*. 2020;77(5):464-

473.

49. Schottelkorb AA, Doumas DM, Garcia R. Treatment for childhood refugee trauma: A

randomized, controlled trial. *International Journal of Play Therapy*. 2012;21(2):57-73.

50. Ruf M, Schauer M, Neuner F, Catani C, Schauer E, Elbert T. Narrative exposure therapy

for 7□to 16□year□olds: A randomized controlled trial with traumatized refugee children.

Journal of Traumatic Stress. 2010;23(4):437-445.

51. McMullen J, O'callaghan P, Shannon C, Black A, Eakin J. Group trauma□focused

cognitive□behavioural therapy with former child soldiers and other war□affected boys in

the DR Congo: A randomised controlled trial. *J Child Psychol Psychiatry*.

2013;54(11):1231-1241.

52. Barron I, Freitas F, Bosch CA. Pilot randomized control trial: Efficacy of a group-based

psychosocial program for youth with PTSD in the Brazilian favelas. *J Child Adolesc*

Trauma. 2021;14(3):335-345.

53. Bastien RJ-B, Jongsma HE, Kabadayi M, Billings J. The effectiveness of psychological

interventions for post-traumatic stress disorder in children, adolescents and young adults:

A systematic review and meta-analysis. *Psychol Med*. 2020;50(10):1598-1612.

54. Hoppen TH, Morina N. Is high-quality of trials associated with lower treatment efficacy?

A meta-analysis on the association between study quality and effect sizes of

psychological interventions for pediatric PTSD. *Clin Psychol Rev*. 2020;78:101855.

doi:10.1016/j.cpr.2020.101855.

Table 1. Short-term efficacy of psychological interventions for pediatric PTSD

Reference group	Psych. interv.	kes (N)	SMD	[95% CI]	p	I^2 (τ^2)
Main analysis (i.e., across all data, irrespective of trial quality and trauma history)						
relative to PCC	TF-CBTs	18 (1,145)	**1.06**	**[0.86 – 1.26]**	**< .001**	68.9
	EMDR	7 (297)	**0.86**	**[0.54 – 1.18]**	**< .001**	***
	non-TF	1 (40)	**0.88**	**[0.53 – 1.23]**	**< .001**	
	MDTs	4 (270)	**0.95**	**[0.62 – 1.28]**	**< .001**	(0.14)
	ACC	1 (22)	**0.52**	**[0.24 – 0.79]**	**< .001**	
relative to ACC	TF-CBTs	21 (2,048)	**0.55**	**[0.36 – 0.74]**	**< .001**	
	EMDR	0 (0)	0.35	[-0.04 – 0.73]	.078	
	non-TF	0 (0)	**0.36**	**[0.01 – 0.72]**	**.042**	
	MDTs	4 (146)	**0.43**	**[0.09 – 0.77]**	**.013**	
relative to EMDR	TF-CBTs	3 (185)	0.20	[-0.14 – 0.55]	.240	
	non-TF	0 (0)	0.02	[-0.43 – 0.47]	.933	
	MDTs	0 (0)	0.09	[-0.36 – 0.53]	.704	
relative to non-TF	TF-CBTs	9 (631)	0.19	[-0.11 – 0.48]	.224	
	MDTs	0 (0)	0.07	[-0.38 – 0.51]	.767	
relative to MDTs	TF-CBTs	2 (72)	0.12	[-0.21 – 0.45]	.485	
Sensitivity analysis: high quality trials[a] only						
relative to PCC	TF-CBTs	10 (882)	**1.05**	**[0.83 – 1.28]**	**< .001**	67.8
	EMDR	3 (145)	**0.95**	**[0.57 – 1.34]**	**< .001**	***
	non-TF	0 (0)	**0.80**	**[0.37 – 1.23]**	**< .001**	
	MDTs	3 (218)	**0.91**	**[0.54 – 1.28]**	**< .001**	(0.10)
	ACC	0 (0)	**0.53**	**[0.23 – 0.82]**	**< .001**	
relative to ACC	TF-CBTs	13 (1,792)	**0.53**	**[0.33 – 0.72]**	**< .001**	
	EMDR	0 (0)	**0.43**	**[0.01 – 0.85]**	**.047**	
	non-TF	0 (0)	0.28	[-0.14 – 0.69]	.189	
	MDTs	1 (46)	0.38	[-0.03 – 0.80]	.070	
relative to EMDR	TF-CBTs	3 (185)	0.10	[-0.28 – 0.48]	.602	
	non-TF	0 (0)	-0.15	[-0.67 – 0.37]	.567	
	MDTs	0 (0)	-0.04	[-0.56 – 0.47]	.869	
relative to non-TF	TF-CBTs	5 (322)	0.25	[-0.11 – 0.61]	.171	
	MDTs	0 (0)	0.11	[-0.42 – 0.64]	.687	
relative to MDTs	TF-CBTs	1 (50)	0.14	[-0.25 – 0.53]	.469	
Sensitivity analysis: individual treatment delivery only						
relative to PCC	TF-CBTs	10 (563)	**1.07**	**[0.79 – 1.35]**	**< .001**	63.0
	EMDR	4 (150)	**1.02**	**[0.62 – 1.41]**	**< .001**	***
	non-TF	1 (40)	**0.88**	**[0.49 – 1.27]**	**< .001**	
	ACC	0 (0)	**0.42**	**[0.03 – 0.82]**	**.033**	(0.12)
relative to ACC	TF-CBTs	10 (766)	**0.65**	**[0.37 – 0.92]**	**< .001**	
	EMDR	0 (0)	**0.59**	**[0.11 – 1.07]**	**.016**	
	non-TF	0 (0)	**0.46**	**[0.06 – 0.86]**	**.024**	
Relative to EMDR	TF-CBTs	3 (185)	0.06	[-0.34 – 0.45]	.778	
	non-TF	0 (0)	-0.13	[-0.62 – 0.35]	.590	
relative to non-TF	TF-CBTs	9 (631)	0.19	[-0.10 – 0.48]	.196	
Sensitivity analysis: treatments with parent involvement only						
relative to PCC	TF-CBTs	5 (364)	**1.07**	**[0.76 – 1.38]**	**< .001**	55.5
	non-TF	0 (0)	**0.72**	**[0.28 – 1.16]**	**.001**	***
	MDTs	3 (188)	**0.94**	**[0.59 – 1.29]**	**< .001**	(0.07)
	ACC	0 (0)	**0.65**	**[0.28 – 1.02]**	**< .001**	
relative to ACC	TF-CBTs	9 (852)	**0.42**	**[0.19 – 0.66]**	**< .001**	
	non-TF	0 (0)	0.07	[-0.32 – 0.46]	.719	
	MDTs	2 (56)	0.29	[-0.11 – 0.69]	.159	
relative to non-TF	TF-CBTs	5 (476)	**0.35**	**[0.04 – 0.66]**	**.026**	
	MDTs	0 (0)	0.22	[-0.26 – 0.70]	.378	
relative to MDTs	TF-CBTs	2 (72)	0.14	[-0.23 – 0.50]	.465	
Sensitivity analysis: samples with mean age < 12 (i.e., mostly children)						
relative to PCC	TF-CBTs	4 (252)	**1.08**	**[0.66 – 1.49]**	**< .001**	69.2

	EMDR	3 (89)	**0.86**	**[0.35 – 1.36]**	**< .001**	***
	non-TF	0 (0)	**0.78**	**[0.24 – 1.32]**	**.004**	(0.12)
	ACC	0 (0)	**0.52**	**[0.03 – 1.01]**	**.037**	
relative to ACC	TF-CBTs	10 (1,104)	**0.55**	**[0.29 – 0.81]**	**< .001**	
	EMDR	0 (0)	0.34	[-0.29 – 0.96]	.290	
	non-TF	0 (0)	0.26	[-0.17 – 0.69]	.236	
relative to EMDR	TF-CBTs	1 (52)	0.22	[-0.35 – 0.78]	.451	
	non-TF	0 (0)	-0.08	[-0.74 – 0.58]	.823	
relative to non-TF	TF-CBTs	6 (507)	0.29	[-0.05 – 0.63]	.094	
Sensitivity analysis: samples with mean age ≥ 12 (i.e., mostly adolescents)						
	TF-CBTs	13 (854)	**1.09**	**[0.82 – 1.37]**	**< .001**	73.6
relative to PCC	EMDR	3 (147)	**0.93**	**[0.42 – 1.45]**	**< .001**	***
	MDTs	1 (82)	**1.02**	**[0.53 – 1.52]**	**< .001**	(0.19)
	ACC	1 (22)	**0.56**	**[0.18 – 0.95]**	**.004**	
	TF-CBTs	11 (944)	**0.53**	**[0.24 – 0.82]**	**< .001**	
relative to ACC	EMDR	0 (0)	0.37	[-0.22 – 0.96]	.222	
	MDTs	4 (146)	**0.46**	**[0.02 – 0.90]**	**.042**	
relative to EMDR	TF-CBTs	2 (133)	0.16	[-0.36 – 0.68]	.550	
	MDTs	0 (0)	0.09	[-0.59 – 0.77]	.791	
relative to MDTs	TF-CBTs	2 (72)	0.07	[-0.39 – 0.53]	.773	

Note. ACC = active control conditions (e.g. = treatment-as-usual); EMDR = eye movement desensitization and reprocessing; kes = number of direct comparisons for the given comparison; MDTs = multidisciplinary treatments; N = total number of participants; non-TF = non-trauma-focused psychological interventions; PCC = passive control conditions (e.g. = waitlist); Psych. interv. = psychological interventions; SMD = standardized mean differences (i.e. = Hedges' g); TF-CBTs = trauma-focused cognitive behaviour therapies. **Bold** print highlights statistical significance of findings. A positive (negative) SMD indicates superior (inferior) efficacy of the given psychological intervention relative to the given reference group.
[a]Meeting at least six of eight trial quality criteria (Cuijpers et al., 2010).
*** p < .001, ** p < .01, * p < .05, corresponding to the respective Q-statistic as a measure of heterogeneity in outcomes.

Table 2. Mid-term efficacy (top) and long-term efficacy (bottom) of psychological interventions for pediatric PTSD

Reference group	Psych. interv.	kes (N)	SMD	[95% CI]	p	I^2 (τ^2)
Mid-term efficacy (i.e., ≤ 5 months follow-up) - main analysis						
relative to PCC	TF-CBTs	9 (389)	**0.87**	**[0.58 – 1.17]**	**< .001**	66.4
	EMDR	2 (86)	**0.95**	**[0.48 – 1.41]**	**< .001**	***
	MDTs	1 (52)	**0.59**	**[0.03 – 1.15]**	**.039**	(0.15)
	ACC	2 (74)	**0.42**	**[0.06 – 0.79]**	**.024**	
relative to ACC	TF-CBTs	9 (813)	**0.45**	**[0.17 – 0.73]**	**.002**	
	EMDR	1 (74)	**0.52**	**[0.04 – 1.00]**	**.032**	
	MDTs	3 (79)	0.17	[-0.36 – 0.69]	.535	
relative to EMDR	TF-CBTs	2 (125)	-0.07	[-0.51 – 0.37]	.753	
	MDTs	0 (0)	-0.35	[-1.02 – 0.31]	.298	
relative to MDTs	TF-CBTs	1 (20)	0.28	[-0.26 – 0.83]	.309	
Sensitivity analysis: high quality trials[a] only						
relative to PCC	TF-CBTs	4 (223)	**1.06**	**[0.64 – 1.49]**	**< .001**	71.9
	EMDR	1 (23)	**1.15**	**[0.56 – 1.75]**	**< .001**	***
	ACC	1 (52)	**0.73**	**[0.24 – 1.21]**	**.004**	(0.14)
relative to ACC	TF-CBTs	7 (781)	**0.33**	**[0.03 – 0.63]**	**.029**	
	EMDR	1 (74)	0.43	[-0.08 – 0.94]	.102	
relative to EMDR	TF-CBTs	2 (125)	-0.09	[-0.58 – 0.39]	.702	
Sensitivity analysis: treatments with parent involvement only						
relative to PCC	TF-CBTs	3 (128)	**1.09**	**[0.56 – 1.63]**	**< .001**	71.1
	MDTs	1 (52)	0.43	[-0.25 – 1.10]	.216	***
	ACC	0 (0)	**0.73**	**[0.11 – 1.35]**	**.021**	(0.14)
relative to ACC	TF-CBTs	5 (619)	0.37	[-0.02 – 0.75]	.060	
	MDTs	2 (43)	-0.30	[-0.95 – 0.34]	.357	
relative to MDTs	TF-CBTs	1 (20)	**0.67**	**[0.02 – 1.32]**	**.042**	
Sensitivity analysis: samples with mean age ≥ 12 (i.e., mostly adolescents)						
relative to PCC	TF-CBTs	6 (252)	**0.76**	**[0.43 – 1.09]**	**< .001**	44.9
	MDTs	0 (0)	**0.70**	**[0.05 – 1.34]**	**.034**	*
	ACC	2 (74)	0.22	[-0.18 – 0.61]	.282	(0.07)
relative to ACC	TF-CBTs	7 (492)	**0.54**	**[0.26 – 0.83]**	**< .001**	
	MDTs	3 (79)	0.48	[-0.05 – 1.01]	.076	
relative to MDTs	TF-CBTs	1 (20)	0.06	[-0.52 – 0.64]	.831	
Long-term efficacy (i.e., 6-24 months follow-up) - main analysis						
relative to PCC	TF-CBTs	3 (118)	**0.76**	**[0.27 – 1.26]**	**.002**	67.6
	non-TF	1 (40)	**0.71**	**[0.15 – 1.27]**	**.014**	***
	ACC	1 (51)	0.21	[-0.31 – 0.73]	.431	(0.11)
relative to ACC	TF-CBTs	9 (920)	**0.55**	**[0.30 – 0.81]**	**< .001**	
	non-TF	1 (45)	**0.50**	**[0.09 – 0.93]**	**.016**	
relative to non-TF	TF-CBTs	5 (343)	0.06	[-0.29 – 0.40]	.754	
Sensitivity analysis: high quality trials[a] only						
relative to ACC	TF-CBTs	8 (887)	**0.53**	**[0.27 – 0.80]**	**< .001**	70.7
	non-TF	1 (45)	0.46	[-0.03 – 0.95]	.065	***
relative to non-TF	TF-CBTs	3 (151)	0.07	[-0.37 – 0.52]	.744	(0.11)
Sensitivity analysis: individual treatment delivery only						
relative to PCC	TF-CBT	3 (118)	**0.78**	**[0.34 – 1.22]**	**< .001**	46.2
	non-TF	1 (40)	**0.64**	**[0.13 – 1.15]**	**.014**	
	ACC	1 (51)	0.17	[-0.33 – 0.67]	.503	(0.07)
relative to ACC	TF-CBT	5 (280)	**0.61**	**[0.28 – 0.94]**	**< .001**	
	non-TF	0 (0)	**0.47**	**[0.01 – 0.92]**	**.043**	
relative to non-TF	TF-CBT	5 (343)	0.14	[-0.18 – 0.46]	.384	
Sensitivity analysis: samples with mean age < 12 (i.e., mostly children)						
relative to PCC	TF-CBTs	1 (26)	1.00	[-0.11 – 2.11]	.077	78.5
	non-TF	0 (0)	0.83	[-0.37 – 2.03]	.176	***
	ACC	0 (0)	0.46	[-0.71 – 1.63]	.441	(0.15)

relative to ACC	TF-CBTs	5 (673)	**0.54**	**[0.18 – 0.90]**	**.003**
	non-TF	1 (45)	0.37	[-0.16 – 0.91]	.173
relative to non-TF	TF-CBTs	3 (266)	0.17	[-0.29 – 0.63]	.472

Note.   ACC = active control conditions (e.g. = treatment-as-usual); EMDR = eye movement desensitization and reprocessing; kes = number of direct comparisons for the given comparison; MDTs = multidisciplinary treatments; N = total number of participants; non-TF = non-trauma-focused psychological interventions; PCC = passive control conditions (e.g. = waitlist); Psych. interv. = psychological interventions; SMD = standardized mean differences (i.e. = Hedges' g); TF-CBTs = trauma-focused cognitive behaviour therapies. **Bold** print highlights statistical significance of findings. A positive (negative) SMD indicates superior (inferior) efficacy of the given psychological intervention relative to the given reference group.
[a]Meeting at least six of eight trial quality criteria (Cuijpers et al., 2010).

*** p < .001, ** p < .01, * p < .05, corresponding to the respective Q-statistic as a measure of heterogeneity in outcomes.

Table 3. Efficacy rankings of psychological interventions for pediatric PTSD

	SUCRA @short-term - all data (main analysis)	SUCRA @short-term - outlier-adjusted	SUCRA @short-term - high-quality trials only	SUCRA @short-term - individual delivery of treatments only	SUCRA @short-term - treatments with caregiver involvement only	SUCRA @short-term - samples with mean age < 12 (mostly children)	SUCRA @short-term - samples with mean age ≥ 12 (mostly adolescents)	SUCRA @mid-term - all data (main analysis)	SUCRA @mid-term - MDTs deleted due to inconsistency	SUCRA @mid-term - high-quality trials only	SUCRA @mid-term - treatments with caregiver involvement only	SUCRA @mid-term - samples with mean age ≥ 12 (mostly adolescents)	SUCRA @long-term - all data (main analysis)	SUCRA @long-term - high-quality trials only	SUCRA @long-term - individual delivery of treatments only	SUCRA @long-term - samples with mean age < 12 (mostly children)
TF-CBTs	0.90	0.89	0.87	0.88	0.94	0.87	0.83	0.81	0.80	0.77	0.98	0.86	0.87	0.81	0.93	0.91
EMDR	0.58	0.61	0.71	0.77	NA	0.69	0.64	0.87	0.86	0.87	NA	NA	NA	NA	NA	NA
MDTs	0.70	0.72	0.66	NA	0.74	NA	0.74	0.50	NA[a]	NA	0.36	0.78	NA	NA	NA	NA
Non-TF	0.60	0.57	0.52	0.60	0.46	0.60	NA	NA	NA	NA	NA	NA	0.79	0.67	0.73	0.68
ACC	0.21	0.21	0.23	0.25	0.36	0.34	0.28	0.32	0.34	0.36	0.62	0.31	0.26	0.02[b]	0.25	0.29
PCC	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.04	0.05	0.07	NA	0.09	0.12

Note. ACC = active control conditions (e.g. = treatment-as-usual); @long-term = long-term efficacy (i.e., assessments 6-24 months after treatment endpoint); @mid-term = mid-term efficacy (i.e., assessments ≤ 5 months after treatment endpoint); @short-term = short-term efficacy(i.e., assessments at treatment endpoint); EMDR = eye movement desensitization and reprocessing; high-quality trials only = sensitivity-analysis exclusively involving high-quality trials (i.e., trials fulfilling at least six of eight quality criteria); MDTs = multi-disciplinary treatments; NA = not applicable given insufficient accumulated evidence (i.e., kes < 4); Non-TF = non-trauma-focused psychological interventions (i.e., psychological interventions without a trauma focus); PCC = passive control conditions (i.e., waitlist); SUCRA = surface under the cumulative ranking (i.e., efficacy ranked by means of surface under the cumulative ranking with 50,000 resamples); TF-CBTs = trauma-focused cognitive behaviour therapies.

[a]Excluded from this analysis given that significant inconsistency was detected.

[b]Reference group for this analysis given insufficient number of direct comparisons (k < 4) for PCC.

Figure 1. PRISMA flow chart depicting the study selection process

Figure 2

Network graphs for the main analyses concerning short-term (left), mid-term (middle), and long-term (right) efficacy

Note. ACC = active control conditions (e.g., = treatment-as-usual); EMDR = eye movement desensitization and reprocessing; MDT = multi-disciplinary treatments (e.g., bounce back); non-TF-PIs = non-trauma-focused psychological interventions (e.g., non-directive supportive therapy); PCC = passive control conditions (e.g., = waitlist); TF-CBTs = trauma-focused cognitive behaviour therapies (e.g., prolonged exposure).
Nodes sizes are proportional to the number of included participants per dyad and the thickness of lines is proportional to the number of direct comparisons (see also number of direct comparisons inserted for each comparison dyad).