Immune-developmental processes contribute to schizophrenia risk: insights from a genetic overlap

study with height

Cato Romero[1], Christiaan de Leeuw[2], Marijn Schipper[2], Bernardo de A.P.C. Maciel[2], Martijn P. van den

Heuvel[1,2], Rachel M. Brouwer[2], August B. Smit[3], Frank Koopmans[3], Danielle Posthuma[2,4], & Sophie van

der Sluis[1]

Affiliations:

[1]Department of Child and Adolescent Psychiatry and Psychology, Section Complex Trait Genetics,

Amsterdam Neuroscience, Vrije Universiteit Medical Center, Amsterdam UMC, Amsterdam, the

Netherlands

[2]Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Amsterdam

Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

[3]Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive

Research, Neuroscience Campus Amsterdam, VU University, Amsterdam, the Netherlands

[4]Department of Clinical Genetics, Section Complex Trait Genetics, Amsterdam Neuroscience, Vrije

Universiteit Medical Center, Amsterdam University Medical Centre, Amsterdam 1081 HZ, the

Netherlands

¯Correspondence should be addressed to: Sophie van der Sluis: Department of Complex Trait Genetics,

Vrije Universiteit Amsterdam, Research building VU, De Boelelaan 1100, 1081 HZ, Amsterdam, The

Netherlands. Phone: +31 20 598 6833, s.vander.sluis@vu.nl

Main text word count: 3,982 words

Short title: Immune-developmental processes contribute to SCZ risk

Keywords: *schizophrenia; GWAS; psychiatric genetics; comorbidity; genetic overlap; development*

Background

Shorter stature has been phenotypically linked to increased prevalence of schizophrenia (SCZ), but the

nature of this association is unknown.

Methods

Using genome-wide genetic data, we studied the SCZ-height relationship on a genetic level. Applying

novel genetic methods and tools, we analyzed gene-sets, tissue-types, cell-types, local genetic correlation,

conditional genetic analyses, and fine-mapping of effector-genes to scrutinize the SCZ-height

relationship.

Results

We identified 142 genes statistically associated with both SCZ and height and found enrichment in

immune response gene-sets. Genetic annotations implicated the pituitary and specifically mesenchymal

stem cells for height and thyrotropic cells for SCZ. While the global SCZ-height genetic correlation was

nonsignificant, 9 genomic regions showed robust local genetic correlations (7 negative, 6 in the MHC-

region). The shared genetic signal for SCZ and height within the 6 MHC-regions was partially explained

by mutual genetic overlap with white blood cell count, particularly lymphocytes. Fine-mapping

prioritized 3 shared effector-genes (*GIGYF2*, *HLA-C*, and *LIN28B*) involved in immune response

sensitivity and development of immune and pituitary cell-types.

Conclusions

Overall, our findings suggest an involvement during height-development of thyrotropic cells and immune

response sensitivity contributing towards risk of SCZ.

Introduction

Shorter stature has been linked to a small but consistent association with schizophrenia (SCZ) [1–9], with some studies showing a 15% increase in SCZ prevalence in males of below-average height[10]. While adult height shows consistent associations with SCZ prevalence, evidence for an association with birth length is inconsistent[4,11], prompting further investigation into the nature of the SCZ-height relationship. The association between adult height and hormonal imbalance[12], brain size[13], infections[14], nutrition[15], and socioeconomic status[16] alludes to a developmental risk component in the SCZ-height relationship, as factors influencing physical growth may interact with SCZ pathogenesis. Despite availability of large-scale genome-wide association studies (GWAS), few studies have investigated the genetic SCZ-height relationship directly. Current genetic evidence suggest certain SCZ-linked deletions (e.g., 22q11.2, 1q21.1, and 3q29) reduce height[17]. However, previous genome-wide association studies (GWAS) observed sign-discordant as well as sign-concordant shared SCZ-height signal[18], and some SCZ-linked duplications (e.g., 16p11.2) associate with an *increase* in height[19]. These findings point to the existence of a complex genetic relationship between SCZ and height that can contribute to the observed negative phenotypic association. Leveraging larger GWAS datasets and novel methods, we aimed to further investigate the extent and nature of genetic overlap between SCZ and height, and to identify biological processes underlying this relationship (**Supplementary Figure 1** displays the study design).

Methods and Materials

Genome-wide association study of height using UK Biobank

We conducted GWASs on height using the UK Biobank (UKB) data. Prior to analyses, we excluded variants with high missingness (>0.05), low imputation score (INFO value <0.9) and low minor allele frequency (MAF < 0.01), leaving 8,515,957 genetic variants. Subjects were filtered on genetic relatedness and European ancestry, resulting in an available sample size of 382,754 participants (54% female). Covariates were standardized and included age, sex, array and the first 20 principal components of population structure. This study was conducted under UKB application number 16406. Based on cohort descriptions, there is no sample overlap between the SCZ, UKB-height and GIANT-height samples.

Global genetic correlations and SNP-based heritability estimation

We applied bivariate Linkage-Disequilibrium Score (LDSC) regression[20] to estimate the genome-wide genetic correlation (r_g) between SCZ and height, utilizing precomputed LD scores from the 1000 Genomes data (for each of the European, East-Asian, Latino, and African ancestries, separately) and filtering GWAS summary statistics to HapMap3 SNPs (**Data availability**). LDSC regression, requiring only GWAS summary statistics, provides r_g-estimates unbiased by sample overlap.

MiXeR estimation of shared genetic architecture beyond genetic correlations

To estimate shared genetic architecture, MiXeR[21] analyzes GWAS summary datasets using a Gaussian

mixture model, assuming a "mixture" of causal and non-causal variants that each follow a distinct

Gaussian distribution. The model accounts for factors such as LD structure, genomic inflation from

cryptic relatedness, MAF, GWAS sample size, and sample overlap. In this study, we used bivariate

MiXeR to analyze SCZ and height in both full and sex-stratified samples. To prevent bias in polygenicity

estimates caused by complex LD, the MHC region was excluded from all GWAS prior to analysis. All

AIC and BIC fit indices were positive, indicating sufficient fit.

Overlapping SNPs, genes, and gene-set enrichment analyses

To identify genome-wide significant (GWS) SNPs shared between SCZ and height, we filtered GWAS

summary statistics of SCZ (7,193,791 SNPs), UKB-height (8,515,957 SNPs) and GIANT-height

(2,550,858 SNPs) to 832,763 overlapping SNPs. SNPs meeting the GWS threshold ($P < 5e-8$) in SCZ and

UKB-height were extracted, with semi-replication using GIANT-height ($\alpha_{BON} = 1.23 \times 10^{-4}$). Fisher's

exact tests evaluated whether overlap exceeded chance levels. LD structure was addressed by clumping

shared GWS SNPs using PLINK 1.9[22], defining independent loci (LD $r^2 < 0.2$, 250kb). MHC lead SNPs

were further clumped using progressively larger (100kb) window sizes to account for long-range LD.

Effect concordance was assessed through lead SNP beta coefficients. LD correlation structure was

obtained through 10K UKB genotypes (**Data availability**).

SNP-to-gene annotation used Multi-marker Analysis of GenoMic Annotation (MAGMA)[23],

which calculated gene-based P-values accounting for LD between SNPs (19,427 genes; GRCh37). Shared

genes were defined by Bonferroni-corrected significance ($\alpha_{BON} = .05/19,427 = 2.57 \times 10^{-6}$). Functional

enrichment of shared genes was analyzed in Functional Mapping and Annotation of Genome-Wide

Association Studies (FUMA)[24], testing 34,550 MSigDB[25] v6.2 gene-sets with correction for multiple

testing ($\alpha_{BON} = .05/34,550 = 1.45 \times 10^{-6}$). MAGMA also assessed gene-set and gene-property annotations,

testing 50 hallmark gene-sets, 53 tissue-types (GTEx v8[26]), and 25 pituitary cell-types[27] with combined

Bonferroni correction ($\alpha_{BON} = .05/128 = 3.90 \times 10^{-4}$)

Local genetic correlation analysis of SCZ and height

Local Analysis of [co]Variant Annotation (LAVA) identifies genomic regions with correlated genetic

signals between traits[28]. LAVA requires GWAS summary statistics, estimated sample overlap (using

bivariate LDSC intercepts), an external LD reference (1000 Genomes v3; **Data Availability**), and

predefined genomic regions. Using LAVA's default genomic region definitions, 2,517 genomic regions

were defined for European data. LAVA's workflow involves two steps: filtering regions with sufficient

genetic signal in both traits (default local h^2_{SNP}; $P < 1 \times 10^{-4}$) and estimating bivariate local r_g in selected

regions. Of the 2,517 regions, 816 showed sufficient genetic signal for both SCZ and height. Bivariate

local r_g P-values were Bonferroni-corrected for multiple testing ($\alpha_{BON\square} = 3.13 \times 10^{-5}$; two-sided)

Fine-mapping and gene prioritization in regions associated between SCZ and UKB-height

Fine-mapping of the 9 genomic regions that significantly correlated between SCZ and UKB-height was

conducted using FINEMAP[29], with LD structure estimated from 100K unrelated UKB subjects. Due to

the complexity of fine-mapping the MHC region, FINEMAP was restricted to model 1 causal SNP per

locus. To prioritize genes from FINEMAP's 95% credible sets, we used FLAMES[30] (Fine-mapped Locus

Assessment Model of Effector geneS; v1.0.0), which predicts effector genes mediating trait associations

based on locus-specific SNP-to-gene links and gene network convergence. Genes with FLAMES

confidence scores > 0.05 were considered likely effector genes. Shared effector genes between SCZ and

UKB-height were defined as those with FLAMES scores > 0.05 for both traits. We ran FLAMES using

the default settings, i.e., only annotating SNPs to coding genes mapped within 750kb of the SNPs.

For *HLA-C*, we assessed expression and regulation in dorsolateral prefrontal cortex proteomics

data from 47 SCZ cases versus 49 controls (Koopmans et al., under review, PRIDE dataset ID:

PXD058441).

Conditional local genetic correlation analysis in LAVA

We used conditional local genetic analysis in LAVA to assess whether the genetic overlap between SCZ

and UKB-height in correlated regions was influenced by other covariates. This method tests whether the

marginal genetic correlation ($r_g^{marginal}$) between two traits (Y and X_1) significantly differs from the

conditional estimate ($r_g^{conditional}$) after adjusting for a covariate (X_2). If $r_g^{marginal} \neq r_g^{conditional}$, the covariate

significantly impacts the genetic covariance. Significance was determined by the conditional genetic

covariance (Cov_g; details in Supplementary Note 9.). Defined as a regression model, either SCZ or height

must be chosen as the outcome (Y). Based on Mendelian Randomization SCZ was chosen as the outcome.

To be assessed, a covariate required significant local univariate genetic signal ($P < 1 \times 10^{-4}$) and

genetic correlation with SCZ and height ($\alpha_{BON} = .05/9 = 5.55 \times 10^{-3}$). Applying these filters, 25 out of 44

covariates could be assessed. We tested 25 covariates across 9 loci, correcting for multiple testing ($\alpha_{BON} =$

$.05/(9 \times 25) = 2.06 \times 10^{-4}$). Covariates with significant effects were further replicated in SCZ and Giant-

height analyses. Sample overlap estimates for LAVA were calculated using LDSC.

Replication in non-European ancestries

We attempted to replicate our findings in East-Asian, African, and Latino ancestry GWAS[31,32] data. First,

we tested the overlap of 20 lead SNPs, 142 MAGMA genes, and gene-property results

(pituitary/mesenchymal cells for height; pituitary/thyrotropic cells for SCZ), applying Bonferroni

corrections ($\alpha_{BON}=\alpha/nr.$ *Lead SNPs* $= .05/20 = 2.50 \times 10^{-3}$; MAGMA genes: $\alpha_{BON}=\alpha/nr.$ *genes* $=$

$.05/142 = 3.52 \times 10^{-4}$). Due to insufficient sample sizes in Latino ($N_{SCZcases}$ = 1,234; N_{height} = 58,709)

and African ($N_{SCZcases}$ = 6,152; N_{height} = 168,193) GWASs, only East-Asian ancestry GWASs ($N_{SCZcases}$ =

14,004; N_{height} = 363,856) were viable for local genetic correlation replication. To account for differences

in LD structure, we used the LAVA partitioning algorithm on 1000 Genomes (v3) East-Asian LD

reference. We focused on the 9 genomic regions significantly associated with SCZ and height in

European analyses and identified 11 overlapping East-Asian regions for replication.

12   **Results**

Phenotypic association, Height GWAS and shared genetic architecture with SCZ

Using data from the UK Biobank (UKB), we confirmed the reported association between SCZ and shorter

height in both sexes, although the lower genetic liability of SCZ cases in UKB limits the generalizability

of our findings[33] (**Supplementary Notes 1–2; Supplementary Figures 2A–C; Supplementary Table**

1). Next, we sought to explore how common genetic variation accounts for this observed phenotypic

relationship. We obtained SCZ GWAS summary statistics of European (EUR) ancestry individuals from

the Psychiatric Genetics Consortium[31] (N_{eff}[34] = (4 x N_{cases} x $N_{controls}$)/N_{total} = 157,066). We conducted a

GWAS on height using EUR-ancestry data in the UKB (UKB-height; **Methods**). Although larger height

GWAS exist, their sparse SNP coverage (~1 million SNPs) adversely affects genetic overlap analyses

(e.g., a substantial proportion of LAVA loci would fail). The UKB data, with its denser coverage (~8

million SNPs), ensures more robust and comprehensive analyses of genetic overlap with SCZ.

(**Supplementary Table 2** displays all SCZ and height GWAS data used in this study). Using LDSC

regression[20] (**Methods**), we estimated the genome-wide genetic correlation between SCZ and UKB-

height to be near-zero and non-significant (r_g = -.01, SE = .02, *P* = .72), which is in line with previous

reports[18]. MiXeR[21], unlike LDSC, quantifies shared genetic architectures between traits beyond the global

genetic correlation. Bivariate MiXeR (**Methods**) estimated ~1,100 shared SNPs (SE = 200) contributing

to shared genetic architecture between SCZ and UKB-height (12% and 35% of all SCZ and UKB-height

MiXeR estimated SNPs, respectively; **Figure 1A**-**D**). The sign-concordance of shared SNPs was 48%

(SD = 1.0%), consistent with a near-zero global genetic correlation.

Genetic variants and functional annotations shared between SCZ and height

We investigated specific GWS SNPs, genes and biological annotations shared between SCZ and UKB-height and then used an independent height dataset (GIANT-height[35]; $N_{GIANT-height}$ = 253,288) for semi-replication. That is, for the replication we used the same SCZ GWAS summary statistics (as such data are not available for a sample of similar size and ancestry) but different height data. Analyses were conducted with and without the MHC region (chromosome 6 28,477,797 – 33,448,354 in hg19) to address possible inflation in overlap due to high gene-density and structural complexity. Among the GWS SNPs from SCZ (2,469; 176 independent risk loci) and UKB-height (23,567; 1,226 independent risk loci), we observed 408 mutually associated SNPs, out of which 268 SNPs were semi-replicated and sign-concordant (α_{BON}=.05/408 = 1.23 × 10^{-4}) in GIANT-height, which is above the expected chance level (OR = 3.68, Fisher's exact test P = 1.18 x 10^{-32}). Clumping yielded 20 independent risk loci (5 within the MHC), and 55% of their lead SNPs showed concordant effects (**Supplementary Table 3**). Next, we performed genome-wide gene-based association analysis using MAGMA (**Methods**)[23]. We identified 558 GWS genes for SCZ and 2,883 for UKB-height. Of these, 205 overlapped, and 142 replicated in GIANT-height (α_{BON}=.05/205 = 2.44 × 10^{-4}), which is again significantly above chance level (OR = 2.01, Fisher's exact test P-value = 2.87 × 10^{-11}, without MHC genes OR = 1.89, Fisher's exact test P = 5.33 × 10^{-8}, **Supplementary Table 4**). Significant SNP and/or gene-overlap was also seen between height and other psychiatric disorders. Specifically, SNP and gene-overlap with height was significant for bipolar disorder[36] (BIP; N_{eff} = 49,366), SNP-overlap was significant for insomnia disorder[37] (N_{eff} = 313,722) and autism spectrum disorder[38] (N_{eff} = 44,393), but overlap was not significant for attention deficit/hyperactivity disorder[39] (N_{eff} = 48,990), alcohol use disorder[40] (N_{eff} = 145,999), major depressive disorder[41] (N_{eff} = 156,547), and anorexia nervosa[42] (N_{eff} = 52,027; **Supplementary Table 5**). Out of the genes shared between BIP and height, 90% (N = 18) were also shared between SCZ and height, suggesting concordance in SCZ-BIP overlap with height.

Convergence of the 142 genes that were associated to both SCZ and height onto gene-sets was investigated using GENE2FUNC in FUMA (**Methods**)[24]. Gene-sets were obtained from MSigDB[25] ($N_{MSigDB gene-sets}$ = 34,550). Significant enrichment was found for 7 gene-sets (α_{BON} =.05/34,550 = 1.45 x 10^{-6}; **Figure 1E**), including 4 positional gene-sets (*chr16p11, chr3p21, chr6p21*, and *chr6p22*), and 3 functional gene-sets (*sodium phosphate symporter activity, interferon gamma signaling* and *proximal deletion syndrome*; although *chr6p21* and *interferon gamma signaling* were non-significant when MHC genes were excluded; **Supplementary Table 6**). Gene-set enrichment analysis of mutual GWS genes is an incomplete assessment of functional convergence as different genes can partake in the same biological processes or be expressed in the same biological tissue. To investigate the existence of additionally significant gene-sets, as well as tissue enrichment, we conducted gene-set and gene-property analysis in MAGMA on both SCZ and height (using the biologically-established and non-redundant hallmark gene-

sets $N_{hallmark\ gene-sets}$ = 50 from MsigDB v6.2[25,43] and $N_{tissue-types}$ = 53 from GTEx v8.0[26]) with the entire set

of 19,427 MAGMA genes. While no additional gene-sets were significant, gene-property analysis of

tissue-types found *the pituitary* as significantly associated with both SCZ and UKB-height. Follow-up

cell-type analysis within the pituitary ($N_{cell-types}$ = 25 from Zhang et al.[27]) found significant enrichment for

mesenchymal stem cells (MSC) for UKB- and GIANT-height, and *thyrotrophic cells* (TC) for SCZ. All

gene-property results were significant after correcting for multiple testing and after excluding the MHC

region (α_{BON}=.05/(50 + 53 + 25) = 3.90 × 10^{-4}; **Supplementary Table 7**). In summary, multiple

risk loci and genes are associated with both phenotypes, implicating shared biological processes between

these traits.

Investigating genetic correlations within genomic regions

Shared genetic architecture with concordant and discordant SNP effects suggest the possibility of

significantly shared *local* or regional genetic signal. To evaluate shared local signal, we conducted local

genetic correlation analysis using LAVA (**Methods**)[28]. To filter out clearly non-associated loci, only

genomic regions with suggestive (LAVA default $P < 1$ x 10^{-4}) univariate genetic signal in both SCZ and

height were considered for estimation of local bivariate genetic correlations, resulting in 816 out of 2,517

predefined, roughly independent genomic regions. Subsequently, bivariate genetic correlation was

evaluated in these 816 regions and 16 regions showed significant SCZ-height local correlation after

correcting for multiple testing (α_{BON} = .05/816 = 6.13 x 10^{-5}; **Table 1**; **Figure 1F**). Semi-replication with

the independent GIANT-height data evinced significant associations in 9 of these 16 regions (α_{BON} =

.05/16 = 3.13 x 10^{-3}; **Supplementary Figure 3A-B**). Among these 9 semi-replicated significant regions, 2

were positively and 7 were negatively correlated. Of these 7, 6 were localized within the MHC region on

chromosome 6 (all but two regions (chromosome 6 104,945,107 – 106,052,133 and chromosome 6

107,319,938 – 109,273,876) overlap with the aforementioned 20 shared lead SNPs).

We sought to investigate multiple additional aspects of the local SCZ-height genetic relationship,

including the extent of sex-specific effects, non-linear SCZ-height associations across height strata, and

evidence against indirect genetic effects (e.g., assortative mating, population stratification, and gene-

environment interactions). In summary, we detected 6 additional local genetic correlations between SCZ

and height specific to males using UKB-height data, of which only 1 semi-replicated (chromosome 15

89,385,687 – 90,632,694) in male-stratified GIANT-height data[44]. No female-specific local genetic

correlations were observed, and concordant effects between sexes were confirmed using MiXeR on sex-

stratified SCZ-height overlap (**Supplementary Note 3; Supplementary Table 8**). Within the 9 genomic

regions in which we observed local genetic correlations, the association between SCZ and height

appeared to be linear, except for one region (chromosome 6 107,319,938 – 109,273,876) where the effect

was primarily driven by the 20% shortest individuals (**Supplementary Note 4; Supplementary Table 9**).

Using a within-sibling GWAS on height[45], which can better control for indirect genetic effects, 4 of the 9

robust regions were semi-replicated, lending evidence to associations being based on direct (rather than

indirect) genetic effects in these regions (**Supplementary Note 5**; **Supplementary Table 10**).

Next, we used FLAMES (**Methods**)[30] to prioritize which genes most likely underlie the local

genetic SCZ-height correlations in the 9 robust regions: three genes — *GIGYF2*, *HLA-C* and *LIN28B* —

converged between SCZ and UKB-height as the most likely genes underlying the genetic correlation in

three genomic regions (chromosome 2 233,360,693 – 234,089,948, chromosome 6 31,250,173 –

31,320,268, and chromosome 6 104,945,107 – 106,052,133, respectively; **Table 2, Supplementary**

Figure 4; **Supplementary Note 6; Supplementary Table 11**). Gene-prioritization of MHC in SCZ has

previously not supported a role for *HLA-C*[46]. However, look-up of *HLA-C* in SCZ differentially expressed

proteins in dorsal lateral prefrontal cortex (dlPFC) found *HLA-C* to be significantly differentially

regulated ($\log^2$(FC) = -0.89, *Q*-value = 0.02 (5% FDR, 5,242 proteins tested); Koopmans et al., under

review; **Methods**) in 47 SCZ cases as compared to 49 controls. In the remaining 6 regions no or different

genes were prioritized between SCZ and height.

Assessing involvement of external traits on correlated genomic regions

We assessed whether the 9 local genetic SCZ-height correlations could be further understood through

shared genetic covariance with other traits, that is, whether the genetic relationship between SCZ and

height is mediated by other traits. To this end, we conducted conditional local genetic correlation analyses

in LAVA (**Methods; Supplementary Note 7**). GWAS summary statistics of potential covariates were

selected based on their known epidemiological relation to SCZ or height to: 1) investigate whether the

genetic SCZ-height overlap is shared with related traits (e.g., BIP and body-mass index), 2) assess

involvement of potentially confounding factors (e.g., social deprivation), and 3) assess potential

biological implications in the relationship between SCZ and height (e.g., (sex-)hormones, and

metabolites; full list of all 44 selected covariates and their sources in **Supplementary Table 12**).

Subsequently, covariates were filtered for having significant univariate genetic signal (LAVA default *P* <

1×10^{-4}) and significant genetic correlation with both SCZ and UKB-height in at least 1 of the 9 genomic

regions ($\alpha_{BON} = 0.05/9 = 5.55 \times 10^{-3}$). After filtering, 25 covariates remained.

Specified as a mediation model, conditional genetic analyses require the appointment of a

predictor and an outcome variable. Generalized Summary-data-based Mendelian Randomization

(GSMR)[47] uses genetic information to determine, for genetically and phenotypically correlated traits, the

most likely direction of causation. Here, we used GSMR to determine whether SCZ or UKB-height

should be set as outcome variable in our local genetic mediation analyses. GSMR yielded stronger

evidence, both for p-value and standardized effect size, for UKB-height predicting SCZ (B_{height} = -.024,

SE = .006, P = 2.00 x 10^{-5}) than for SCZ predicting UKB-height (B_{SCZ} = -.020, SE = .008, P = .02), hence

in our mediation analyses, the effects of the 25 covariates were assessed with SCZ as the outcome

variable and UKB-height as predictor (**Supplementary Note 8**). Among related or confounding traits,

only conditioning on BIP resulted in a large (sub-threshold) significant decrease in the genetic correlation

between SCZ and UKB-height within 4 loci (**Figure 2**). Among traits potentially sharing biological

processes with both SCZ and height, conditioning on the genetic signal of white blood cell subtype counts

(lymphocyte and neutrophil counts), significantly decreased the genetic correlation between SCZ and

UKB-height in 6 out of 9 genomic regions, all residing within the MHC region. Lastly, conditioning on

the genetic effects of intra-cranial volume strongly affected the SCZ-height relation on chromosome 6

outside MHC. All the effects of covariates remained significant in semi-replication using GIANT-height

GWAS summary statistics (**Supplementary Table 13; Supplementary Figure 5-6** shows results from

model with height as outcome).

To follow-up the genetic relation between SCZ, height and white blood cell count, we conducted

analyses in UKB data to assess whether phenotypic relationships mirror the observed genetic relations.

Results indeed showed a negative phenotypic relation of white blood cell count (in 10^9 cells/liter) with

UKB-height (in cm; β = -.13, $SE_β$ = 2.01 x 10^{-3}, B = -.03, SE_B = 4.66 x 10^{-4}, T = -63.71, P < 2.00 x 10^{-16}),

and a positive relation with SCZ diagnosis (β = .02, $SE_β$ = 1.46 x 10^{-3}, B = .82, SE_B = .06, T = 14.39, P <

2.00 x 10^{-16}), mirroring previous reports[48,49].

Replication in non-European ancestry

Lastly, to evaluate whether the genetic overlap between SCZ and height observed in EUR-ancestry is

evident across other ancestries, we analyzed GWAS data for East-Asian (EAS), African (AFR), and

Latino (LAT) ancestry. Global genetic correlations were non-significant for EAS and LAT but positive

for AFR (r_g = .29, SE = .13, P = .02). Replication of EUR-ancestry lead SNPs, genes, and annotations

showed limited significance in EAS data and none in AFR or LAT data (see **Supplementary Note 9;**

Supplementary Tables 3, 4, 7). Using EAS-based summary statistics for SCZ (N_{eff} = 30,523) and height

(N = 363,856), none of the nine EUR-significant genomic regions replicated (**Supplementary Note 10;**

Supplementary Table 14). While small SCZ samples affect the power of these cross-ancestry replication

attempts, top SCZ EUR prioritized SNPs did show lower MAF in EAS ancestry (**Table 2,**

Supplementary Table 15) suggesting potential ancestry-specific genetics effects.

33   **Discussion**

We investigated the SCZ-height genetic relationship via multiple post-GWAS methods, replicating 6 of 8

previously reported shared lead SNPs[18], and observe 14 addition lead SNPs, 142 shared genes, and

several shared biological properties. Our findings paint a complex genetic picture, related to immune

response and pituitary-driven development. The shared gene-set *sodium phosphate symporter activity* is

involved in processes like nutrient absorption, cellular signaling, and skeletal development[50]. Although

SCZ and height were enriched in different cell-types in the pituitary, these cell types are known to

interact: thyroid stimulating hormone secreted by pituitary thyrotropic cells regulates mesenchymal cell

differentiation[51]. Shared prioritized genes, *LIN28B*, *GIGYF2*, and *HLA-C*, appear to modulate

inflammation and thyroid function, potentially explaining how SCZ's dysregulated thyroid hormones can

impact growth. Collectively, findings align with *the developmental risk factor model of psychosis*,

suggesting that disruptions in growth- and immune-related processes may heighten susceptibility to

SCZ[52]. For full discussion, see **Supplementary Note 11**.

Our data indicate that SCZ-height genetics may be more pronounced than in other psychiatric

disorders. Gene and SNP overlap was observed for BIP, and BIP showed significant conditional effects in

4 loci, but most of this overlap was shared with SCZ, highlighting the strong genetic correlation (.67)

between BIP and SCZ[53] rather than BIP-specific components, an interpretation substantiated by non-

significant associations between BIP and height in epidemiological literature[54]. However, statistical power

needs to increase for other psychiatric disorders before we can verify the specificity of SCZ's genetic

association to height.

The aforementioned biological processes represent a plausible hypothesis concerning the

relationship between SCZ and height; however, caution is needed regarding the generalizability of this

hypothesis. While our results suggest that the hypothesis generalizes to both sexes, results did not

replicate in non-EUR ancestries. To our knowledge, no epidemiological study has been published on the

SCZ-height association in non-EUR samples, and our non-EUR UKB sample lacked statistical power to

detect a mean height difference (**Supplementary Note 9**). Replication is hampered by power issues

relating to smaller non-EUR (especially SCZ) GWAS samples, and lower MAF of the EUR lead SNPs in

non-EUR ancestries (**Table 2; Supplementary Table 15**). While variation in MAF between ancestries

could suggest that the SCZ-height relationship is specific to, or more pronounced in, people of EUR

ancestry, additional epidemiological and genetic research in non-EUR ancestry is required to test this

hypothesis.

The MHC region, frequently appearing in our results, poses methodological challenges for

GWAS studies. While we replicated results without the MHC where possible, LAVA and FLAMES

results are affected by the strong LD in this region: long-range LD can cause interdependence among

nonadjacent loci that is not accounted for in LAVA, rendering the independence of the 6 MHC loci

uncertain. Additionally, while we used FINEMAP to prioritize SNPs in LAVA-defined regions, specialized methods (SNP2HLA and HLA*SNP) are better suited for fine-mapping the MHC but require individual-level data. Nevertheless, our prioritization of *HLA-C* for SCZ is supported by significant downregulation in dlPFC SCZ proteomic data, suggesting *HLA-C* may contribute to the MHC-SCZ association in addition to other, previously implicated genes.

In summary, despite a near-zero global genetic correlation, shared genetic signals between SCZ and height is evident. By leveraging the genetic overlap with height, our findings converge on the involvement of thyrotropic cells and immune response sensitivity in SCZ and highlights development-specific biological processes contributing towards the etiology of SCZ.

Acknowledgements

C.R., B.A.P.C.M., M.P.vd.H., M.S., R.M.B., A.B.S., D.P. and S.v.d.S were funded by NWO Gravitation:

BRAINSCAPES: A Roadmap from Neurogenetics to Neurobiology (grant no. 024.004.012 [to D.P.]).

F.K. was supported by the Simons Foundation, SFARI Award 882976. D.P. and C.d.L. were funded by a

European Research Council advanced grant (no. ERC-2018-AdG GWAS2FUNC 834057 [to

D.P.]). M.P.v.d.H. is supported by NWO VIDI Grant 452-16-015 and the ERC Consolidator of the

European Research Council Grant 101001062.

The analyses were carried out on the Genetic Cluster Computer, which is financed by the

Netherlands Scientific Organization (NWO: 480-05-003), by the VU University, Amsterdam, the

Netherlands, and by the Dutch Brain Foundation, and is hosted by the Dutch National Computing and

Networking Services SurfSARA. This research has been conducted using the UK Biobank resource under

application number 16406. We thank the numerous participants, researchers, and staff from many studies

who collected and contributed to the data. We particularly express our gratitude to all UKB participants

that have been so generous to share their data for analysis.

Supplementary Note Figure 6 and **Supplementary Figure 1** was created with BioRender.com

with permission to publish.

Author Contributions: C.R.: Conceptualization, formal analyses, methodology, visualization, writing.

C.d.L.: Feedback, formal analysis, methodology. M.S.: Feedback, formal analysis. B.A.P.C.M.:

Feedback, visualization. M.P.vd.H.: Feedback, resources. R.M.B.: Feedback, resources. A.B.S.:

Resources. F.K.: Resources. D.P.: Feedback, funding acquisition, resources. S.vd.S.: Conceptualization,

methodology, project administration, writing, supervision.

Disclosures and Competing Interests:

C.d.L is funded by Hoffman-La Roche. The other authors declare no competing interests.

Locus number	CHR	Start	Stop	Rho	*P*-value	Semi-replication *P*-value
410	2	233360693	234089948	0.56	5.50×10^{-8}	3.57×10^{-5}
970(*)	6	30316328	30932188	-0.45	8.74×10^{-5}	1.34×10^{-5}
971(*)	6	30932189	31143244	-0.46	4.08×10^{-8}	8.20×10^{-7}
972(*)	6	31143245	31250172	-0.66	1.06×10^{-14}	8.13×10^{-6}
973(*)	6	31250173	31320268	-0.64	2.12×10^{-12}	3.61×10^{-5}
974(*)	6	31320269	31577220	-0.64	4.04×10^{-13}	1.66×10^{-6}
981(*)	6	32636612	32682213	-0.51	5.53×10^{-8}	3.93×10^{-5}
1050	6	104945107	106052133	0.64	1.03×10^{-7}	6.13×10^{-5}
1052	6	107319938	109273876	-0.41	2.58×10^{-6}	7.00×10^{-7}

1   *Table 1. Genomic regions showing significant local genetic correlation between SCZ and height.*

2   *Note: All semi-replicated results showed sign concordance with the main analysis. (*) indicates region*

3   *within the MHC region on chromosome 6.*

Locus	Trait	Flames Gene	Flames Score	Top SNP	PIP	nSNP in set	SNP on Gene	EUR MAF	EAS MAF	AFR MAF	LAT MAF
410	Height	*GIGYF2*	.05	rs283477	.12	75	sQTL, eQTL, intron variant	.31	.36	.29	.30
	SCZ	*GIGYF2*	.11	rs778371	.16	46	sQTL	.30	.11	.24	.31
970(*)	Height	*SFTA2*	.21	rs2253718	.93	6	eQTL	.36	.07	.41	.20
	SCZ	*DDX39B*	.12	rs1264347	.10	69		.12	.01	.04	.03
971(*)	Height	*MUC22*	.24	rs9262549	.99	1	Missense variant	.30	.28	.12	.21
	SCZ	*TRIM39*	.04	rs1634726	.36	11		.12	.00	.02	.03
972(*)	Height	*HLA-C*	.16	rs2524080	.74	5	sQTL, eQTL	.29	.19	.16	.22
	SCZ	*HCG27*	.04	rs1793894	.73	3	eQTL	.13	.003	.09	.06
973(*)	Height	*HLA-C*	.06	rs2524054	.45	5	sQTL, eQTL	.27	.19	.09	.17
	SCZ	*HLA-C*	.10	rs2923008	.13	31	sQTL, eQTL	.12	.004	.02	.04
974(*)	Height	*HLA-B*	.02	rs2523578	.36	3	sQTL, eQTL	.26	.03	.20	.13
	SCZ	*ATP6V1G2*	.03	rs2596495	.11	15	DHS-promoter	.13	.004	.11	.06
981(*)	Height	*HLA-DQB1*	.04	rs9275109	.50	5	sQTL, eQTL, Hi-C	.37	.15	.36	.20
	SCZ	*HLA-DQA1*	.03	rs1617322	.10	15	sQTL, eQTL	.13	.06	.08	.06
1050	Height	*LIN28B*	.34	rs314265	.17	11	eQTL	.23	.23	.41	.25
	SCZ	*LIN28B*	.14	rs457286	.07	52	eQTL, intron variant	.44	.33	.77*	.33
1052	Height	*FOXO3*	.24	rs9400239	.16	20	Intron variant	.32	.30	.77*	.40
	SCZ	*SNX3*	.05	rs11964401	.09	42		.49	.36	.83*	.49

Table 2. Top prioritized SNPs within SCZ-height correlated LAVA loci and allele frequencies

Note: Flames score above 0.05 is regarded as sufficient. MAF for multiple ancestries are based on

gnomADv2.1.1. Which allele is the minor allele is based on the European ancestry reference panel, MAF

*values with * indicate that this allele is the major allele in the respective population, not minor allele.*

PIP = Posterior Inclusion Probability; nSNP in set refers to the number of SNPs in the 95% credible set,

sQTL: splicing quantitative trait locus, eQTL: expression quantitative trait locus, HI-C: chromatin

conformation capture, DHS-promoter: DN-ase I hypersensitive site-promoter.

Figure 1. Shared genetic overlap between SCZ and height

1A) Venn-diagram of estimated MiXeR overlap of shared SNPs between SCZ and UKB-height. Jaccard index (ratio of overlap to total area) is 9.65%. Estimates are in 1000 SNPs, i.e., 1.1 refers to 1,100 shared SNPs with SE = 200. B-C) Conditional Q-Q plot of expected versus observed -log10(P) compares the distribution of p-values for SNP associations with SCZ | UKB-height SNPs (B) and UKB-height | SCZ SNPs (C) against the uniform null expectation (x-axis). The diagonal dashed line represents the null hypothesis, where deviation by observed p-values represent significant association for SNPs with P < 0.001, 0.01, 0.1, and 1, respectively. D) Negative log-likelihood plot as function of polygenic overlap. Lowest point of line indicates number of shared variants with optimal fit. E) Flowchart summarizing genetic overlap between SCZ and height regarding specific SNPs, independent risk loci, shared MAGMA genes, gene-set enrichment, tissue and cell type results. Gene-sets with * are not significant after exclusion of the MHC region F) Local genetic correlations between SCZ and UKB-height over all loci in the genome. Y-axis is defined as -log10(P)*sign of rho. Red lines represent local genetic correlations that replicated between SCZ and GIANT-height.

Figure 2. Results from conditional local genetic correlation analyses

Results of the conditional local genetic correlation analyses for covariates that showed (nominal)

significant conditional effects. Left side of the plot shows covariate name and category; top side gives

genomic region definitions. Circle sizes represent the estimated covariate effect. The covariate effect is

the genetic covariance between the covariate trait on the one hand and the covariance shared between

SCZ-height on the other. Grey circles represent theoretical complete overlap between covariate and SCZ-

height, while colored circles represent the observed covariate – SCZ-height overlap within each of the 9

genomic regions. To illustrate, for Total white blood cell count in region chromosome 6 32,636,612 –

32,682,213 (top panel, rightmost circle), all genetic covariance shared between SCZ and height is also

shared with white blood cell count, while in the other regions the genetic signal is only partly, yet often

significantly, shared. Colors indicate significance of the covariate trait estimate: light yellow = not

significant, light orange = nominally significant, orange = significant corrected for number of genomic

regions tested (9), and dark red = significant corrected for number of genomic regions and number of

covariates tested (9 x 25). This figure uses absolute covariate effect, for the sign of the covariate effect,

*please see **Supplementary table 13**.*

Data availability

GWAS summary statistics were downloaded from <u>link</u> for SCZ and <u>link</u> for GIANT-height.

All GWAS summary statistics are based on Human Genome Build 37 (GRCh37/hg19).

Precomputed LD scores and HapMap 3 reference file were obtained from: <u>Link</u>

1000 Genomes reference data: <u>Link</u>

LD correlation structure was obtained through the LD information available from 10K UKB genotypes:

<u>https://www.uk10k.org</u>

Gene information was obtained from the GeneCard database (v5.12.0 Build 702;

<u>https://www.genecards.org</u>).

Software Availability

FLAMES(v1.0.0) <u>https://github.com/Marijn-Schipper/FLAMES</u>

MAGMA (v.1.10) gene-based and gene-property analysis: <u>https://ctg.cncr.nl/software/magma</u>

LAVA(v0.1.0) local genetic correlations: <u>https://ctg.cncr.nl/software/lava</u>

PLINK (v1.9) <u>https://www.cog-genomics.org/plink/</u>

G*power**: <u>Link</u>**

References

1. Burchard, E. M. L. *Physique and Psychosis: An Analysis of the Postulated Relationship Between Bodily Constitution and Mental Disease Syndrome*. (Johns Hopkins Press, 1936).

2. Kemali, D., Polani, N., Polani, P. E. & Amati, A. A dermatoglyphic study of 219 Italian schizophrenic males. *Clin. Genet.* **9**, 51–60 (1976).

3. Nopoulos, P., Flaum, M., Arndt, S. & Andreasen, N. Morphometry in schizophrenia revisited: height and its relationship to pre-morbid function. *Psychol. Med.* **28**, 655–663 (1998).

4. Gunnell, D. *et al.* The association of fetal and childhood growth with risk of schizophrenia. Cohort study of 720,000 Swedish men and women. *Schizophr. Res.* **79**, 315–322 (2005).

5. Niemi, L. T., Suvisaari, J. M., Haukka, J. K. & Lönnqvist, J. K. Childhood growth and future development of psychotic disorder among Helsinki high-risk children. *Schizophr. Res.* **76**, 105–112 (2005).

6. Perrin, M. A., Chen, H., Sandberg, D. E., Malaspina, D. & Brown, A. S. Growth trajectory during early life and risk of adult schizophrenia. *Br. J. Psychiatry* **191**, 512–520 (2007).

7. Laurens, K. R. *et al.* Common or distinct pathways to psychosis? A systematic review of evidence from prospective studies for developmental risk factors and antecedents of the schizophrenia spectrum disorders and affective psychoses. *BMC Psychiatry* **15**, 205 (2015).

8. Jørgensen, T. S. H. *et al.* Cognitive ability, education, height and body mass index in relation to risk of schizophrenia and mortality following its diagnosis. *Eur. J. Epidemiol.* **39**, 893–904 (2024).

9. Garcia-Rizo, C. *et al.* Anthropometry in antipsychotic-naïve first-episode psychosis patients: An exploratory approach to the role of environmental early life events in two independent samples. *Schizophr. Res.* **266**, 216–226 (2024).

10.   Zammit, S. *et al.* Height and body mass index in young adulthood and risk of

schizophrenia: a longitudinal study of 1 347 520 Swedish men. *Acta Psychiatr. Scand.* **116**,

378–385 (2007).

11.   Latham, K. & Kirkpatrick, B. Meta-analysis of adult height and birth length in

schizophrenia. *Schizophr. Res.* **195**, 110–114 (2018).

12.   Laron, Z. Insulin-like growth factor 1 (IGF-1): a growth hormone. *Mol. Pathol.* **54**, 311–

316 (2001).

13.   Taki, Y. *et al.* Correlation among body height, intelligence, and brain gray matter volume

in healthy children. *NeuroImage* **59**, 1023–1027 (2012).

14.   Stephensen, C. B. Burden of infection on growth failure. *J. Nutr.* **129**, 534S-538S (1999).

15.   Perkins, J. M., Subramanian, S. V., Davey Smith, G. & Özaltin, E. Adult height,

nutrition, and population health. *Nutr. Rev.* **74**, 149–165 (2016).

16.   Thompson, K., Portrait, F. & Schoonmade, L. The height premium: A systematic review

and meta-analysis. *Econ. Hum. Biol.* **50**, 101273 (2023).

17.   Owen, D. *et al.* Effects of pathogenic CNVs on physical traits in participants of the UK

Biobank. *BMC Genomics* **19**, 867 (2018).

18.   Bacanu, S.-A., Chen, X. & Kendler, K. S. The genetic overlap between schizophrenia and

height. *Schizophr. Res.* **151**, 226–228 (2013).

19.   Niarchou, M. *et al.* Psychiatric disorders in children with 16p11.2 deletion and

duplication. *Transl. Psychiatry* **9**, 8 (2019).

20.   Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from

polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

21. Frei, O. *et al.* Bivariate causal mixture model quantifies polygenic overlap between complex traits beyond genetic correlation. *Nat. Commun.* **10**, 2417 (2019).

22. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

23. Leeuw, C. A. de, Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. *PLOS Comput. Biol.* **11**, e1004219 (2015).

24. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

25. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739–1740 (2011).

26. The Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene regulation in humans | Science. https://www.science.org/doi/full/10.1126/science.1262110.

27. Zhang, S. *et al.* Single-cell transcriptomics identifies divergent developmental lineage trajectories during human pituitary development. *Nat. Commun.* **11**, 5275 (2020).

28. Werme, J., van der Sluis, S., Posthuma, D. & de Leeuw, C. A. An integrated framework for local genetic correlation analysis. *Nat. Genet.* **54**, 274–282 (2022).

29. Benner, C. *et al.* FINEMAP: efficient variable selection using summary data from genome-wide association studies. *Bioinformatics* **32**, 1493–1501 (2016).

30. Schipper, M. *et al.* Gene prioritization in GWAS loci using multimodal evidence. 2023.12.23.23300360 Preprint at https://doi.org/10.1101/2023.12.23.23300360 (2024).

31. Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* **604**, 502–508 (2022).

32. Yengo, L. *et al.* A saturated map of common genetic variants associated with human height. *Nature* **610**, 704–712 (2022).

33. Legge, S. E. *et al.* Genetic and Phenotypic Features of Schizophrenia in the UK Biobank. *JAMA Psychiatry* **81**, 681–690 (2024).

34. Grotzinger, A. D., Fuente, J. de la, Privé, F., Nivard, M. G. & Tucker-Drob, E. M. Pervasive Downward Bias in Estimates of Liability-Scale Heritability in Genome-wide Association Study Meta-analysis: A Simple Solution. *Biol. Psychiatry* **93**, 29–36 (2023).

35. Wood, A. R. *et al.* Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat. Genet.* **46**, 1173–1186 (2014).

36. Mullins, N. *et al.* Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nat. Genet.* **53**, 817–829 (2021).

37. Jansen, P. R. *et al.* Genome-wide analysis of insomnia in 1,331,010 individuals identifies new risk loci and functional pathways. *Nat. Genet.* **51**, 394–403 (2019).

38. Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum disorder. *Nat. Genet.* **51**, 431–444 (2019).

39. Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat. Genet.* **51**, 63–75 (2019).

40. Kranzler, H. R. *et al.* Genome-wide association study of alcohol consumption and use disorder in 274,424 individuals from multiple populations. *Nat. Commun.* **10**, 1499 (2019).

41. Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat. Genet.* **50**, 668–681 (2018).

42. Watson, H. J. *et al.* Genome-wide association study identifies eight risk loci and implicates metabo-psychiatric origins for anorexia nervosa. *Nat. Genet.* **51**, 1207–1214 (2019).

43. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst.* **1**, 417–425 (2015).

44. Randall, J. C. *et al.* Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. *PLoS Genet.* **9**, e1003500 (2013).

45. Howe, L. J. *et al.* Within-sibship genome-wide association analyses decrease bias in estimates of direct genetic effects. *Nat. Genet.* **54**, 581–592 (2022).

46. Sekar, A. *et al.* Schizophrenia risk from complex variation of complement component 4. *Nature* **530**, 177–183 (2016).

47. Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat. Commun.* **9**, 224 (2018).

48. Liemburg, E. J., Nolte, I. M., Klein, H. C. & Knegtering, H. Relation of inflammatory markers with symptoms of psychotic disorders: a large cohort study. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **86**, 89–94 (2018).

49. Shimizu, Y. *et al.* Short stature is an inflammatory disadvantage among middle-aged Japanese men. *Environ. Health Prev. Med.* **21**, 361–367 (2016).

50. Stelzer, G. *et al.* The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence Analyses: The GeneCards Suite. in *Current protocols in bioinformatics / editoral board, Andreas D. Baxevanis ... [et al.]* vol. 54 1.30.1-1.30.33 (2016).

1   51.   Bagriacik, E. U., Yaman, M., Haznedar, R., Sucak, G. & Delibasi, T. TSH-induced gene

2   expression involves regulation of self-renewal and differentiation-related genes in human

3   bone marrow-derived mesenchymal stem cells. *J. Endocrinol.* **212**, 169–178 (2012).

4   52.   Murray, R. M., Bhavsar, V., Tripoli, G. & Howes, O. 30 Years on: How the

5   Neurodevelopmental Hypothesis of Schizophrenia Morphed Into the Developmental Risk

6   Factor Model of Psychosis. *Schizophr. Bull.* **43**, 1190–1196 (2017).

7   53.   Romero, C. *et al.* Exploring the genetic overlap between twelve psychiatric disorders.

8   *Nat. Genet.* **54**, 1795–1802 (2022).

9   54.   Okholm, G. T. *et al.* Body mass index and height in young adult men in relation to

10   subsequent risk of mood disorder. *Eur. J. Epidemiol.* **36**, 1065–1074 (2021).