1    # TITLE PAGE

2    **Manuscript Title-** Optimal Systolic Blood pressure control after thrombectomy in acute
3    ischemic stroke-a systematic review and meta-analysis
4    **Short title**-Optimal Systolic blood pressure after thrombectomy
5    **Authors and affiliations**

| Author | Qualification | Affiliation | Email IDs |
|---|---|---|---|
| Baikuntha Panigrahi | MD,DM | Department of Neurology, All India Institute of Medical Sciences, New Delhi 110029,India | baikunthap199@gmail.com |
| Rohit Bhatia* | MD,DM | Department of Neurology, All India Institute of Medical Sciences, New Delhi 110029,India | rohitbhatia71@yahoo.com |
| Partha Haldar | MD | Centre for Community Medicine, All India Institute of Medical Sciences, New Delhi 110029,India | parthahaldar@outlook.com |
| Risha Sarkar | MA | Department of Neurology, All India Institute of Medical Sciences, New Delhi 110029,India | risha98sarkar@gmail.com |
| Imnameren Longkumer | PhD | Department of Neurology, All India Institute of Medical Sciences, New Delhi 110029,India | imna01lkr@gmail.com |

**\*Corresponding author**
Dr.Rohit Bhatia
Professor
Room 603. 6th Floor
Department of Neurology
Neurosciences Centre
All India Institute of Medical Sciences
New Delhi 110029. India
Mail-rohitbhatia71@yahoo.com

**Manuscript Metadata**
Abstract word count-319 words
Manuscript Total Word count-4048 words
Tables-2
Figures-6
References-22

**Optimal Systolic Blood pressure control after thrombectomy in acute**
**ischemic stroke-a systematic review and meta-analysis**
3

## Abstract
**Introduction**
Although endovascular thrombectomy(EVT) is the standard of care for acute large vessel
occlusions, ideal systolic blood pressure (SBP) target post procedure has remained elusive.
With recent publications of randomised controlled trials (RCTs) comparing intensive BP
control to a less intensive strategy, there is a need to conduct a systematic review and meta-
analysis on existing studies for a timely update. Our study aimed to address the question of
whether in adult patients of acute ischemic stroke who undergo EVT does an intensive blood
pressure control as compared to a less intensive blood pressure control/conventional control
up to 24 hours after procedure lead to a good functional outcome as defined by modified
Rankin score(mRS) of 0-2 at 90days.
15
**Methods**
We used the PRISMA guidelines for this systematic review. Databases were searched for
English language articles using predefined search terms till Sep 15 2023. Risk of bias was
assessed using the New Castle Ottawa scale(NOS) for observational studies, and the revised
Cochrane risk-of-bias 2 (ROB2) tool for RCTs. The primary outcome was mRS 0-2 at
90days.
22
**Results**
Initial search revealed 9883 records. 8857 records entered title and abstract screening out of
which 72 full text studies were assessed for eligibility and twelve studies (5439 patients-eight
observational and four RCTs) were included in the final analysis. The primary outcome was
not significantly different between both the arms(RR-1.16 ; 95%CI-0.98-1.37; p=0.08).There
were no significant differences in the mortality at 90days and the risk of symptomatic
intracranial haemorrhage(RR-death:0.83;95%CI-0.68-1.02;p=0.08,RR-symptomatic
intracranial haemorrhage:0.84;95%CI-0.61-1.16;p=0.29). Fewer patients required
decompressive surgery in the intensive arm(RR-0.40; 95%CI-0.25-0.66; p=0.0003). A
separate analysis of observational studies and RCTs for the primary outcome showed that
observational studies favoured intensive control (RR-1.34;95%CI-1.20-1.48; p<0.00001)
whereas data from RCTs favoured conventional control(RR-0.82;95%CI-0.72-0.93;p=0.003).
35
36 **Conclusion**
37 The results of this meta-analysis suggest that neither intensive nor conventional SBP control
38 resulted in better functional outcomes but intensive control had a significant effect by
39 reducing the need for decompression. Further RCT's will help strengthen evidence.

# **ABBREVIATIONS**

**AIS-**Acute ischemic stroke

**EVT**-Endovascular thrombectomy

**LVO**-Large vessel occlusion

**BP**-Blood Pressure

**SBP**-Systolic Blood Pressure

**RCT**-Randomised Controlled trial

**PRISMA**- Preferred Reporting Items for Systematic Reviews and Meta-Analyses

**NOS**-New Castle Ottawa Scale

**ROB2**-Cochrane Risk of Bias-2

**mRS**-Modified Rankin Scale

**ICH**-Intracranial haemorrhage

**sICH**-Symptomatic Intracranial Haemorrhage

**RR**-Relative Risk/Risk ratio

**AHA/ASA-**American Heart Association/American Stroke Association

**LOE**-Level of evidence

**COR**-Class of recommendation

**PICO**-Patients/Intervention/Control or Comparator/Outcomes

**MeSH**-Medical Subject headings

**NIHSS**-National Institute of Health Stroke Scale

**CONSORT**- Consolidated Standards of Reporting Trials

**GRADE**- Grading of Recommendations, Assessment, Development, and Evaluations

**Introduction**

Endovascular thrombectomy with or without thrombolysis is currently the standard of care in patients with acute ischemic stroke due to a large vessel occlusion.[1] However, even after successful radiological evidence of reperfusion a proportion of patients do not achieve the desired level of functional independence.[2] This is dependent on numerous modifiable, non-modifiable and unknown factors. Among these, blood pressure (BP) is a major modifiable factor which influences the outcomes after a successful reperfusion.[2,3] The interaction between BP and outcomes is extremely complex as other critical factors including the collateral status, size of the penumbra and degree of recanalization play a role in the final outcome.[4] However, the optimal blood pressure after thrombectomy is not clear. The 2019 American Heart Association/American Stroke Association(AHA/ASA) guidelines for acute ischemic stroke recommend that it is reasonable to maintain BP at ≤180/105 mm Hg during and for 24 hours after the procedure (LOE-B-NR;COR-IIa)[1]. It is well acknowledged in the guideline that the data on optimal blood pressure target is sparse and the data supporting this guideline was drawn from blood pressure management protocols in the thrombectomy trials. Since a majority of the patients enrolled in the six hour time window trials had already received thrombolysis, the post thrombolysis blood pressure targets applied to them. Additional data was obtained from the DAWN and ESCAPE trial protocols.[5,6] The DAWN protocol recommended the systolic BP (SBP) to be maintained at <140mmHg in the first 24hrs after the procedure.[6] However, the ESCAPE protocol recommended the SBP to be maintained at >150mmHg to maintain the collateral perfusion.[5]

Since the last recommendations, multiple randomised and observational data has become available with conflicting results which makes it imperative to perform a systematic review and meta-analysis to carefully address this important question. We conducted this meta-analysis to address the question of whether in adult patients with an acute ischemic stroke due to  large vessel occlusion who undergo thrombectomy does an intensive blood pressure control (Target SBP cutoff as defined by the respective study or an upper limit of  SBP target for defining intensive control as <140mmHg) as compared to a less intensive blood pressure control/conventional control up to 24 hours after the procedure lead to a good functional outcome as defined by  modified Rankin score (mRS) of 0-2 at 90 days.

**Methods**

*Search strategy*

We used the standard PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines for conducting and reporting our review (Checklist in *Supplementary material-1*). The protocol was registered on PROSPERO (https://www.crd.york.ac.uk/prospero/#myprospero) vide registration number-CRD42023463173. An amendment was submitted on 28 January 2024 to define blood pressure targets as per the respective study definition. A comprehensive preliminary search was carried out by three authors (B.P, R.S and I.L) for English Language articles from inception until September 15, 2023, using databases which included PubMed, SCOPUS, EMBASE, Google Scholar, Web of Science and hand searching. The MeSH terms included for the search string were – "Acute stroke", OR "Ischemic stroke" OR "cerebrovascular accident" OR "cerebral infraction" AND "thrombectomy" OR "endovascular therapy" OR "mechanical aspiration" OR "endovascular technique" OR "endovascular procedure" OR "stent retriever" AND "blood pressure" OR "blood pressure monitor" OR "high blood

pressure" OR "low blood pressure" OR "hypertension". The search results obtained from the various sources were imported to the Covidence software (https://www.covidence.org). Duplicates were removed using the software automatically.

*Screening and Eligibility criteria*

Titles and abstracts were reviewed independently by four authors (B.P, RB, R.S, I.L ) to evaluate for inclusion and screening of full text. The review question and the PICO statements are summarised in the form of a table (available as *Supplementary material-2*). The inclusion criteria of the studies were-Randomized Controlled Trials (RCTs), observational prospective studies, observational retrospective studies with at least 20 patients, patients included in the study must have had a baseline mRS-0-1,included patients with Acute Ischemic Stroke (AIS) due to large vessel occlusion (LVO) involving either the anterior or the posterior circulation and treated with thrombectomy using any technique with or without bridging thrombolytic agent, systemic blood pressure monitoring during and at least 24 hours after the procedure and functional outcomes using the mRS at 90days should have been reported. Intensive control was defined as aggressive control of SBP to a predefined target of <140mmHg or as described in the respective study with the use of anti-hypertensive drugs for up to 24 hours after a successful thrombectomy irrespective of whether the patient received thrombolysis or not . Less Intensive or Conventional blood pressure control was defined as a liberal SBP control strategy where SBP was not lowered until it was >180mmHg as per guidelines. We excluded case series, case reports, studies without a separate comparison outcome data between intensive and less intensive blood pressure groups, studies comparing blood pressure control in AIS due to LVO treated only with Intravenous thrombolysis (IVT) without Endovascular therapy (EVT) and studies without a primary efficacy outcome data. The primary efficacy outcome was the proportion of patients who had a mRS of 0-2 at 90days. Other safety outcomes that were analysed were death at 90 days, development of symptomatic intracranial haemorrhage(sICH) and requirement of decompressive surgery. Any discrepancies were resolved by R.B. Full texts were retrieved for further consideration for inclusion in the study. Individual patient data was not available for the study.

*Data Extraction*

The data was extracted on MS Excel (Ver 16.66.1).This included: the name of the journal, year of publication, 1st author's name, study type- retrospective, prospective study or randomized trial (monocentric or multicentric), number of patients, proportion of patients with a history of hypertension, median NIHSS, stroke localization, intravenous thrombolysis use (if provided), recanalization rates, procedure time, number of patients in the intensive and conventional treatment arm as per the respective study cut-offs, BP parameters measured at least for 24 hours post procedure (mean, minimum, maximum according to the study), rates of symptomatic intracranial haemorrhage (ICH) at 24-36 hours as per the respective study definition, 3- month mRS according to peri-procedural BP.

*Risk of Bias Assessment*

The Newcastle Ottawa Scale (NOS) was used by B.P (reviewed by P.H) to assess the risk of bias in the observational studies. The Risk of Bias (ROB2) tool was used by B.P (reviewed by P.H) for assessing risk of bias in randomized controlled trials.

*Strategy for Data Analysis*

Statistical analyses were performed using Review Manager 5.4.1 software by B.P and reviewed by P.H. Random-effects models was used in view of expected heterogeneity to be present due to known differences in stroke patient characteristics and symptom definitions.

The overall effect estimate, and 95% CI was used to generate Forest plots. Heterogeneity was assessed by the Higgin $I^2$ statistics. The primary and all secondary efficacy and safety outcomes were assessed by risk ratios (RR). Sensitivity analysis using the leave one study out analysis was done. GRADE pro GDT was used for certainty of evidence assessment.

**Results**

*Study selection and evaluation*
An initial database search using the MeSH terms yielded 9883 records which included 8958 from PubMed, 1045 from Google Scholar, 545 from Scopus, 288 from EMBASE and 129 from Web of Science. 1026 duplicates were removed (Figure-1). 8857 studies entered the title and abstract screening of which 8713 studies were found to be irrelevant. 72 studies were assessed for full text eligibility and 60 studies were excluded (reasons in the CONSORT-Figure 1). A total of twelve studies were included in the meta-analysis.

*Study Characteristics*
Of the twelve studies involving 5439 patients included in the final analysis, eight were observational and four were RCTs.[7–18] Observational study data for 3868 patients and data from RCTs involving total of 1571 patients was also extracted separately. We used the published data only for conducting the analysis. We did not have access to individual patient data in any of the studies included in the meta-analysis. Five out of the eight observational studies used a SBP cut-off of 140 mmHg to define intensive control.[11,13,15–17] Two(Chang et al-2019 and P.Upadhyaya et al-2023) studies used a cut-off of 130 mmHg and one study(Eva.A.Mistry et al) used a cut-off of 158mmHg for defining intensive control.[12,14,18] In case of the RCTs the ENCHANTED2/MT used a cut-off SBP of 120mmHg to define its intensive arm.[8] The BP-TARGET on the other hand used a range of SBP between 100-129mmHg to define its intensive arm.[7] There were 3 SBP groups in the BEST-II(140,160,180).[9] For combining the effects we clubbed the <160mmHg and <180mmHg groups into the less intensive group and the <140mmHg as the intensive arm. The OPTIMAL-BP used a SBP cut-off of 140mmHg.[10] The baseline characteristics and attributes of each study included in this meta-analysis are summarised in the Table-1 and Table-2.

*Risk of Bias*
- Observational Studies
  The Newcastle Ottawa Scale results are summarised in the *Supplementary material-3*. We found all studies to be of good methodological quality with a median of 8 stars.

- RCTs
  The ROB2 tool demonstrated a "high risk" of bias in the deviations from the intended intervention domain (Domain-2) in three out of four included RCTs. One study had "some concerns" in this domain. The other domains examined had a low risk of bias. The detailed analysis is provided in the *Supplement (Supplementary material-3)*.

*Outcomes*
*Primary Outcome mRS 0-2*
Pooled data of all the twelve studies involving 5398 patients (reported outcomes) with 2983 patients in the intensive blood pressure control group and 2415 patients in the conventional control group demonstrated that the primary outcome of mRS-0-2 was seen in 1508/2983 (50.55%) patients in the intensive arm as compared to 1106/2415 (45.8%) in the conventional

arm and was not significantly different between both the arms(RR-1.16 ; 95%CI-0.98-1.37; p=0.08) as seen in Figure-2.[7–18]

Separate analysis of data from eight observational studies showed that mRS 0-2 was seen in 1171 out of 2235 in the intensive control group (52.39%) as compared to 674 out of 1633 patients in the conventional group (41.23%) favouring intensive blood pressure control for a good outcome(RR-1.34; 95%CI-1.20-1.48; p<0.00001) as shown in Figure-2.[11–17]

Similarly, a separate analysis of data from 4 RCTs showed that the primary outcome was seen in 337/748 (45.05%) patients in the intensive arm as compared to 432/782 (55.24%) in the conventional arm thus favouring the conventional arm for a good outcome (RR-0.82; 95%CI-0.72-0.93; p=0.003) as shown in Figure-2.[7–10]

*Death at 90 days*

Outcomes for death at 90days were available for 2985 patients in the intensive arm and 2417 patients in the conventional arm from twelve studies. A total of 487/2985(16.3%) patients died in the intensive arm as compared to 485/2417(20.06%) in the conventional arm and the difference between both the arms was not significant (RR-0.83; 95%CI-0.68-1.02; p=0.08) as shown in Figure-3.[7–18]

A separate analysis of data from eight observational studies showed that 373/2235 (15.97%) patients died in the intensive arm as compared to 375/1633 (22.96%) in the conventional arm. The difference between both the arms was significant with lower risk of dying in the intensive arm (RR-0.72; 95%CI-0.59-0.87; p=0.0008) as shown in Figure-3.[11–17].

Data from the four RCTs showed that 114/750 (15.2%) patients died in the intensive arm as compared to 110/784 (14.03%) patients in the conventional arm and the difference between both the arms was not statistically significant (RR-1.13; 95%CI-0.88-1.44; p=0.34) as shown in Figure-3.[7–10]

*Symptomatic ICH*

Data on symptomatic ICH (sICH) was available for nine out of twelve studies involving 2385 patients in the intensive arm and 1797 patients in the conventional arm. A total of 130/2385(5.45%) patients had sICH (as per respective study definition) in the intensive arm compared to 127/1797(7.07%) in the conventional arm and the difference between the two groups was not statistically significant(RR-0.84; 95%CI-0.61-1.16; p=0.29) as shown in Figure-4.[7–11,13,16–18]

A separate analysis of data from five observational studies showed that sICH occurred in 74/1632 (4.53%) patients in the intensive arm compared to 76/1010 (7.52%) patients in the conventional arm and the risk of sICH was significantly lower in the intensive arm compared to the conventional arm (RR-0.64; 95%CI-0.43-0.96; p=0.03) as shown in Figure-4.[11,13,16–18]

Data from the four RCTs showed that symptomatic ICH occurred in 56/753 (7.44%) patients in the intensive arm compared to 51/787 (6.48%) patients in the conventional arm with no significant difference between the two groups (RR-1.13; 95%CI-0.78-1.63; p=0.52) as shown in Figure-4.[7–10]

*Decompressive surgery*

Data on decompressive surgery was available from four out of twelve studies for 1684 patients in the intensive arm as compared to 959 patients in the conventional arm. A total of 44/1684(2.61%) patients in the intensive arm required decompression as compared to 54/959(5.63%) patients in the conventional arm. There was significantly higher requirement of decompressive surgery in the conventional management arm compared to the intensive arm (RR-0.40; 95%CI-0.25-0.66; p=0.0003) as shown in Figure-5.[7,13,16,18]

A separate analysis of only observational study data from three studies showed that 43/1526 (2.82%) patients required decompressive surgery in the intensive arm compared to 54/797 (6.76%) patients in the conventional arm with the chances of needing a decompression higher in the conventional arm than intensive arm (RR-0.39; 95%CI-0.25-0.59; p<0.00001) as shown in Figure-5.[13,18,19]

*Optimal Systolic Blood Pressure cut-offs*
We further explored the various SBP cut-offs (SBP at which anti-hypertensive medications would be started and the target SBP up to 24hours after EVT) used across the studies and segregated them into groups of studies using a cut-off of 140mmHg to define intensive control and 130mmHg to define intensive control.

*Primary outcome for a SBP cut-off of 140mmHg*
Pooled analysis of seven studies(n=2156 patients in intensive arm and 1488 patients in conventional arm) for the primary outcome of mRS 0-2 at 90days with a cut-off of 140mmHg to define intensive control did not show in any significant differences between the intensive or less intensive/conventional control(RR-1.14; 95%CI-0.96-1.36; p=0.14) as shown in the Figure-6.[9–11,13,15–17]
However a separate analysis of five observational studies (n=1964 patients in the intensive arm and 1265 patients in conventional arm) only using the 140mmHg cut-off for the primary outcome demonstrated a significant chance of favourable outcome in the intensive arm(RR-1.26; 95%CI-1.13-1.40; p<0.0001) as shown in Figure-6.[11,13,15–17]
A similar analysis for two RCTs (n=192 patients in intensive arm and 223 patients in the conventional arm) with the SBP cut-off of 140mmHg did not show a significant difference between the two groups(RR-0.79; 95%CI-0.61-1.04; p=0.09) as shown in Figure-6.[9,10]
Among the other outcomes death at 90days was significantly lower in the intensive control arm in the observational data involving five studies (n=1964 in intensive arm and n=1265 in conventional arm) with the cut-off of 140mmHg (RR-0.76; 95%CI-0.60-0.97; p=0.03) as shown in Supplemental Figure-2. The details about the other outcomes of sICH , decompressive surgery at different cut-offs can be found in the *Supplemental material-4 Figures-1,2,3.*

*Primary outcome for a SBP cut-off of 130mmHg*
An analysis of three studies(n=244 in intensive arm and 254 in conventional arm) with respect to the primary outcome using a target SBP cut-off of 130mmHg did not show any significant differences between the two groups(RR-1.27; 95%CI-0.92-1.74; p=0.15) as shown in Figure-6.[7,14,18]
The details about the other outcomes of sICH and decompressive surgery and different cut-offs can be found in the *Supplemental material-4;Figures-1,2,3.*
We could not explore other lower(one study) and higher cut-offs(two studies) due to limited number of studies in these groups and non-availability of individual patient data.

*Leave-out-one study Influence analysis*
Leaving out the ENCHANTED-2/MT and OPTIMAL-BP results resulted in a change in the outcome of the pooled data such that the intensive arm had a higher odds of having a better functional outcome as compared to conventional BP management.[8,10] Leave out one study analysis did not lead to any change in the primary outcome in a separate analysis of only observational studies. Similar to the combined analysis, leaving out the ENCHANTED-2/MT and OPTIMAL-BP results resulted in a change in the outcome of the RCT pooled data but the difference was in the outcome such that neither arm was preferred[8,10]. Leaving out the other

two RCTs did not change outcomes. The details are provided in the supplement *(Supplementary material-5)*.

*GRADE*
The GRADEpro GDT (https://www.gradepro.org) was used to assess the certainty of evidence and was found to be moderate to low in case of observational studies and moderate in case of the RCTs. The summary of the GRADE findings is presented in the *Supplementary material-6.*

**Discussion**
Our systematic review and meta-analysis involving eight observational studies and four RCTs analysed the effects of an intensive as compared to a conventional/less intensive blood pressure control up to 24 hours of thrombectomy in all strokes (eleven studies on anterior circulation and one on posterior circulation stroke). The results suggest that neither intensive nor conventional management of blood pressure post thrombectomy resulted in any difference in the functional outcomes i.e. mRS-0-2 at 90days. Similarly the risk of death and sICH were also similar in both groups. A significantly fewer patients required decompressive surgery in the intensive arm as compared to the conventional arm. Separate analysis of data from observational studies and data from RCTs revealed discordant results. The primary outcome i.e. achieving a mRS 0-2 at 90 days was higher in the intensive arm as compared to the conventional arm in observational studies whereas the conventional management was favoured from RCT data. Mortality at 90 days was also lower in the intensive arm as compared to the conventional arm in the observational data and both groups were comparable as per pooled data from the RCTs. Similarly the observational data favoured intensive control with lower rates of symptomatic ICH as compared to a conventional blood pressure control. The pooled data of the RCTs did not favour any particular arm. Decompressive surgery was more likely to be required in the conventional arm as compared to the intensive arm.

The optimal blood pressure after endovascular therapy has remained elusive ever since the procedure was introduced. Numerous hypotheses have been suggested favouring and negating intensive and conventional management. Arguments in favour of intensive management primarily derived from observational studies and stemmed from the fact that intensive control reduced the rates of symptomatic ICH, reduced cerebral hyper perfusion thereby reducing cerebral edema and improving outcomes.[13,19] However with the robust RCTs like the ENCHANTED-2/BP-TARGET not favouring the intensive arm an alternate school of thought has emerged.[7,8] Firstly, higher SBP might be associated with maintenance of perfusion to distal branches of the blood vessel after successful recanalization. A more intensive SBP control would lead to focal areas of hypoperfusion leading to infarct progression and poor outcomes. Secondly, a successful recanalization after EVT may have persistent venous postcapillary thrombosis and an intensive BP reduction could be detrimental and further promote infarct and ischemia progression.[20]

Blood pressure is a dynamic entity subject to cerebral autoregulation which in turn is unique to a patient. Blood pressure variability post EVT have also been observed to be associated with poor outcomes.[21] It is therefore important to avoid wide fluctuations in the blood pressure.

The pooled data did not show any difference in outcomes with the type of BP management strategy used although there was a trend towards better outcomes in the intensive arm which

could be due to the reduced risk of reperfusion injury and lower risk of haemorrhagic transformation.[11,16] The requirement of decompressive surgery was also significantly lower in the intensive arm. This could also be explained by the fact that lower SBP probably reduced the risk of reperfusion injury and subsequent infarct expansion and cerebral edema.[16] The separate analysis of observational and RCT data showed discordant results. This discordance in the observational and randomised data could be attributed to multiple factors. First, randomised trials are conducted in a controlled environment with strict BP targets and had patients for longer periods of time in the intended target. The higher rates of hypotensive events in the intensive group patients in the RCTs might have been a potential confounder for the poor outcomes in this group. On the other hand the BP in observational studies is more like the real world with some studies using the peak SBP and some using the mean SBP over a period of 24 hrs after the procedure. Since BP fluctuates throughout the day and the mean BP is unlikely to reflect these wide fluctuations and it is difficult to ascertain whether these so called 'intensive' arm patients were actually intensively treated.

Second, RCTs used drugs to both increase and decrease the blood pressure to set targets and it is possible that these drugs could have been a potential confounder with respect to unknown pleotropic effects of the anti-hypertensive drug.[8] Such data was not available for all the observational studies.

Third, although the observational studies selected had a net low risk of bias (using the NOS tool) it is not possible to completely eliminate selection and observer's bias inherent in all observational studies.

Fourth, the observational cut-offs used were different for different studies. We used the definition of intensive as defined by the study which is the reason for the significant heterogeneity. A total of five out of eight observational studies namely Goyal et al, Anadani et al(2019), Cernik et al, Anadani et al (2020) and Han et al used a SBP cut off of 140mmHg as a definition for intensive control.[13,15,17,19,22] On the other hand 2 studies by Chang et al and P Upadhyay et al used a lower cut-off of 130mmHg.[14] A much liberal cut-off of 158mmHg was used for defining intensive control in the Mistry et al (2019) study.[12]

The cut-offs used in the RCTs for defining intensive was also different across studies with the ENCHANTED-2/MT and BP-TARGET defining the upper limit of SBP for the intensive arm at 120mmHg and the BEST-II dividing the BP into three groups of 140,160 and 180 and the OPTIMAL-BP using a upper limit SBP of 140mmHg.[7–9] Even though we did a separate analysis using different cut-offs of 140mmHg and 130mmHg to define the intensive arm there was no difference in the results as the observational data favoured the intensive arm. The difference was seen in the RCT data as removing ENCHANTED-2/MT and BP-TARGET trials resulted in no significant difference in the two arms.[7,8] A lower cut off of 130mmHg also did not result in any significant differences between the arms. Access to individual patient data of these studies could have allowed us to study higher BP targets to identify the SBP sweet spot across studies by pooling the blood pressure recordings and subsequently dichotomising the outcomes.

Another important aspect to note is the heterogeneity in perfusion deficit of the patients recruited in the study. In the RCT data the ENCHANTED2/MT had reported a considerable number of patients with a significant perfusion deficit before the thrombectomy and such cases are extremely prone to reinfarction and infarct progression if the blood pressure is lowered.[8]

Further studies involving more patients and more uniform BP cut-offs will provide better answers in the future. The findings of our study based on RCT data which are considered Level-1 evidence favour a conventional management of blood pressure with target SBP of 140-180mmHg in line with the recommendations from the AHA/ASA.

**Strengths**

The major strengths of this meta-analysis is the comprehensive search methodology and an inclusive study pooling data of all the published studies till date. It also includes data of one study with posterior circulation strokes.

**Limitations**

The major limitation of this study is that it uses only data that was reported. A study with individual patient data could have led to better identification of the true cut-off SBP for achieving a good outcome and we could have explored higher targets in better detail. We also could not explore posterior circulation strokes separately as there was only one study exploring this.[17] We could have performed meta-regression by adjusting for co-variates to predict whether age, recanalization grade or the admission blood pressure had an effect on the final outcome.

**Conclusion**

The results of this meta-analysis suggest that although the combined data does not favour either intensive or conventional control, observational data favours lowering the systolic blood pressure after thrombectomy and was associated with higher chances of functional recovery, reduced mortality and reduced risk of symptomatic ICH. Data from RCTs however favours a less intensive blood pressure control after thrombectomy in terms of achieving functional recovery but has no effects on mortality. Further well conducted RCTs using homogenous BP cut-offs and larger sample sizes can help generate Level-1 evidence required for recommending one particular SBP management strategy over the other.

**Acknowledgements**

Nil

**Funding**

Nil

**Disclosures and Conflicts of Interest**

B.P, R.B, P.H, R.S and I.L report no disclosures and no conflicts of interest.

**Consent**

Not required as this is a systematic review and meta-analysis.

**Ethical Statement**

No ethical clearance was required as this is a systematic review and meta-analysis

**Contributions**

Baikuntha Panigrahi: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data. Rohit Bhatia: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data. Partha Haldar: Drafting/revision of the manuscript for content, including medical writing for content; Analysis or interpretation of data. Risha Sarkar: Major role in the acquisition of data. Imnameren Longkumer: Major role in the acquisition of data.

**Supplemental Material(S)**
**S1-**PRISMA checklist
**S2**-Review Question and PICO Statements
**S3**-Risk of Bias analysis
**S4**- Outcomes analysed using different SBP cut-offs
**S5**-Leave one study out analysis
**S6**-GRADE assessment of certainty of evidence
Supplemental Figures-S3

**References**

1. Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the Early Management of Patients With Acute Ischemic Stroke: 2019 Update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2019;50(12):e344-e418. doi:10.1161/STR.0000000000000211

2. Rabinstein AA, Albers GW, Brinjikji W, Koch S. Factors that may contribute to poor outcome despite good reperfusion after acute endovascular stroke therapy. *International Journal of Stroke*. 2019;14(1):23-31. doi:10.1177/1747493018799979

3. Katsanos AH, Malhotra K, Ahmed N, et al. Blood pressure after endovascular thrombectomy and outcomes in patients with acute ischemic stroke: An individual patient data meta-analysis. *Neurology*. 2022;98(3):e291-e301.

4. Hong L, Cheng X, Lin L, et al. The blood pressure paradox in acute ischemic stroke. *Ann Neurol*. 2019;85(3):331-339.

5. Goyal M, Demchuk AM, Menon BK, et al. Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med*. 2015;372(11):1019-1030. doi:10.1056/NEJMoa1414905

6. Nogueira RG, Jadhav AP, Haussen DC, et al. Thrombectomy 6 to 24 Hours after Stroke with a Mismatch between Deficit and Infarct. *N Engl J Med*. 2018;378(1):11-21. doi:10.1056/NEJMoa1706442

7. Mazighi M, Richard S, Lapergue B, et al. Safety and efficacy of intensive blood pressure lowering after successful endovascular therapy in acute ischaemic stroke (BP-TARGET): a multicentre, open-label, randomised controlled trial. *Lancet Neurol*. 2021;20(4):265-274. doi:10.1016/S1474-4422(20)30483-X

8. Yang P, Song L, Zhang Y, et al. Intensive blood pressure control after endovascular thrombectomy for acute ischaemic stroke (ENCHANTED2/MT): a multicentre, open-label, blinded-endpoint, randomised controlled trial. *Lancet*. 2022;400(10363):1585-1596. doi:10.1016/S0140-6736(22)01882-7

9. Mistry EA, Hart KW, Davis LT, et al. Blood Pressure Management After Endovascular Therapy for Acute Ischemic Stroke: The BEST-II Randomized Clinical Trial. *JAMA*. 2023;330(9):821. doi:10.1001/jama.2023.14330

10. Nam HS, Kim YD, Heo J, et al. Intensive vs conventional blood pressure lowering after endovascular thrombectomy in acute ischemic stroke: The OPTIMAL-BP randomized clinical trial. *JAMA*. 2023;330(9):832-842.

11. Goyal N, Tsivgoulis G, Pandhi A, et al. Blood pressure levels post mechanical thrombectomy and outcomes in large vessel occlusion strokes. *Neurology*. 2017;89(6):540-547. doi:10.1212/WNL.0000000000004184

12. Mistry EA, Sucharew H, Mistry AM, et al. Blood pressure after endovascular stroke therapy (BEST): Final results of a prospective multicenter cohort validation study. *Stroke*. 2019;50((Mistry E.A.; Sucharew H.; Mistry A.M.; Mehta T.; Arora N.; De Los Rios La Rosa F.; Starosciak A.K.; EI James S.; Barnhill N.R.; Patel K.; Assad S.; Tarboosh A.T.; Dakay K.; Cruz A.; Wagner J.; Fortuny E.; Bennett A.; James R.; Jagadeesan B.; Streib C.; O). doi:10.1161/str.50.suppl_1.94

13. Anadani M, Orabi MY, Alawieh A, et al. Blood pressure and outcome after mechanical thrombectomy with successful revascularization. *Stroke*. 2019;50(9):2448-2454.

14. Chang JY, Han MK. Postthrombectomy systolic blood pressure and clinical outcome among patients with successful recanalization. *Eur Neurol*. 2019;81(5-6):216-222.

15. Cernik D, Sanak D, Divisova P, et al. Impact of blood pressure levels within first 24 hours after mechanical thrombectomy on clinical outcome in acute ischemic stroke patients. *Journal of NeuroInterventional Surgery*. 2019;11(8):735-739. doi:10.1136/neurintsurg-2018-014548

16. Anadani M, Arthur AS, Tsivgoulis G, et al. Blood Pressure Goals and Clinical Outcomes after Successful Endovascular Therapy: A Multicenter Study. *Annals of Neurology*. 2020;87(6):830-839. doi:10.1002/ana.25716

17. Han B, Sun X, Liu R, et al. Impact of the perioperative blood pressure on clinical outcome after thrombectomy in acute basilar artery occlusion. *J Stroke Cerebrovasc Dis*. 2021;30(3):105590.

18. Upadhyaya P, Mehta A, Luby M, et al. Good outcome associated with blood-brain barrier disruption and lower blood pressure after endovascular therapy. *Journal of Stroke and Cerebrovascular Diseases*. 2023;32(1). doi:10.1016/j.jstrokecerebrovasdis.2022.106870

19. Anadani M, Arthur AS, Alawieh A, et al. Blood pressure reduction and outcome after endovascular therapy with successful reperfusion: a multicenter study. *J Neurointerv Surg*. 2020;12(10):932-936.

20. Deng G, Chu YH, Xiao J, et al. Risk Factors, Pathophysiologic Mechanisms, and Potential Treatment Strategies of Futile Recanalization after Endovascular Therapy in Acute Ischemic Stroke. *Aging Dis*. 2023;14(6):2096-2112. doi:10.14336/AD.2023.0321-1

21. Nepal G, Shrestha GS, Shing YK, Muha A, Bhagat R. Systolic blood pressure variability following endovascular thrombectomy and clinical outcome in acute ischemic stroke: A meta-analysis. *Acta Neurol Scand*. 2021;144(4):343-354. doi:10.1111/ane.13480

22. Goyal N, Tsivgoulis G, Iftikhar S, et al. Admission systolic blood pressure and outcomes in large vessel occlusion strokes treated with endovascular treatment. *Journal of NeuroInterventional Surgery*. 2017;9(5):451-454. doi:10.1136/neurintsurg-2016-012386

**Table-1-Baseline characteristics of the study population of RCTs**

| Characteristic | Mazighi 2021[7] | Yang 2022[8] | Mistry 2023[9] | Nam 2023[10] |
|---|---|---|---|---|
| Total number of participants randomised | 324 | 821 | 120 | 306 |
| Number of patients in Intensive Arm (ITT) | 158 | 407 | 40 | 155 |
| Number of patients in less intensive arm (ITT) | 162 | 409 | 80 | 150 |
| Age {Mean} Intensive Arm | | 68 | | 73.2 |
| SD Age in Intensive Arm | | 12 | | 12.1 |
| Age {Mean} Less Intensive Arm | | 67 | | 72.9 |
| SD Age in Less Intensive Arm | | 12 | | 10.8 |
| Age {Median} Intensive arm | 77 | | 74 | |
| IQR Age Intensive Arm | 66-85 | | 66.8-84.8 | |
| Age{Median-Less Intensive} | 76 | | | |
| IQR Age Less Intensive Arm | 62-82 | | | |
| Males n; Intensive Arm | 81 | 249 | 12 | 92 |
| % Males in Intensive Arm | 51 | 61 | 30 | 59.4 |
| Males n;Less Intensive Arm | 72 | 257 | 39 | 88 |
| Males(%) in Less Intensive Arm | 45 | 63 | 48.75 | 59.9 |
| Hypertension n; Intensive Arm | 110 | 267 | 32 | 121 |
| Hypertension(%) Intensive Arm | 70 | 66 | 80 | 78.1 |
| Hypertension n; Less Intensive Arm | 113 | 261 | 60 | 110 |
| Hypertension % Less Intensive Arm | 71 | 64 | 75 | 74.8 |
| Smoker n; Intensive Arm | 19 | | 8 | 39 |
| Smokers(%) Intensive Arm | 14 | | 20 | 25.2 |
| Smoker n; Less Intensive Arm | 24 | | 22 | 29 |
| Smoker % Less Intensive Arm | 16 | | 27.5 | 19.7 |
| Diabetes n; Intensive Arm | 34 | 81 | 12 | 65 |
| Diabetes(%) Intensive Arm | 22 | 20 | 30 | 41.9 |
| Diabetes n; Less Intensive Arm | 33 | 82 | 28 | 62 |
| Diabetes % Less intensive Arm | 21 | 20 | 35 | 42.2 |
| Atrial Fibrillation n Intensive Arm | | 84 | 19 | 77 |
| Atrial Fibrillation(%) Intensive Arm | | 21 | 47.5 | 49.7 |
| Atrial Fibrillation n; Less Intensive Arm | | 98 | 34 | 69 |
| Atrial Fibrillation(%) Less Intensive Arm | | 24 | 42.5 | 46.9 |
| Prior Stroke history n; Intensive Arm | 25 | 107 | | 36 |
| Prior Stroke History(%) Intensive Arm | 16 | 26 | | 23.2 |

| | | | | |
|---|---|---|---|---|
| Prior Stroke history; n Less Intensive Arm | 21 | 139 | | 30 |
| Prior Stroke % Less Intensive Arm | 13 | 34 | | 20.4 |
| Median NIHSS at presentation Intensive Arm | 18 | 15 | 16 | 13 |
| Median NIHSS at presentation Less Intensive Arm^ | 17 | 15 | 18^(160)+14^(180) | 12 |
| Site M1 MCA n; Intensive Arm | 117 | | 31 | |
| %M1 MCA-Intensive | 74 | | 77.5 | |
| M1 MCA n; Less Intensive Arm | 119 | | 48 | |
| M1 MCA (%) Less Intensive Arm | 75 | | 60 | |
| ICA n; Intensive Arm | | | 7 | |
| ICA % Intensive Arm | | | 17.5 | |
| ICA n; Less Intensive Arm | | | 14 | |
| ICA (%) Less Intensive Arm | | | 17.5 | |
| IV Thrombolysis use n; Intensive Arm | 85 | 132 | 17 | 44 |
| IV Thrombolysis use %; Intensive Arm | 54 | 32 | 42.5 | 28.4 |
| IV Thrombolysis use n; Less Intensive Arm | 83 | 115 | 37 | 54 |
| IV Thrombolysis use (%) Less Intensive Arm | 52 | 28 | 46.25 | 36.7 |
| Mean SBP at presentation Intensive Arm | 155 | 158.1 | 150 | |
| SD SBP- Intensive Arm | 26 | 25 | 22.5 | |
| Mean SBP at presentation; Less Intensive Arm^ | 152 | 158.7 | 151^/146^(160/180) | |
| SD SBP-Less Intensive Arm | 25 | 23 | 25.3^(160)/23^(180) | |
| Mean DBP at presentation-Intensive Arm | 86 | 89.4 | 82.6 | |
| SD DBP-Intensive Arm | 18 | 16 | 14.5 | |
| Mean DBP at presentation-Less Intensive Arm | 85 | 89.5 | 89^(160)/86.6^(180) | |
| SD DBP-Less Intensive Arm | 15 | 15 | 18.4^(160)/19.6^(180) | |
| Onset to Puncture time in minutes-Intensive Arm | | | | 388 |
| Onset to Puncture time in minutes-Less intensive Arm | | | | 357 |
| Onset to recanalization time in minutes-Intensive | 285 | | | 421 |
| Onset to recanalization time in minutes-Less Intensive | 297 | | | 399 |
| Recanalization mTICI 3 (%) Intensive Arm | 56 | | 45 | |
| Recanalization mTICI 3 (%) Less Intensive Arm | 52 | | 42.5 | |
| Recanalization mTICI 2b (%) Intensive Arm | 44 | | 40 | |
| Recanalization mTICI 2b (%) Less Intensive Arm | 48 | | 42.5 | |
| Mean procedure Time in minutes - Intensive Arm | | 54 | | 30 |
| Mean procedure Time in minutes-Less Intensive Arm | | 60 | | 31 |
| Last Known well in minutes (SD) Intensive Arm | | 558 | 238 | |
| Last Known well in minutes (SD) Less Intensive Arm | | 516 | 222 | |

1    ^*No of patients with each individual group in brackets
2
3
4

9 **Table-2-Baseline characteristics of the study population of Observational Studies**

| Characteristic | Goyal et al[11] | Eva A. Mistry et al[12] | Anadani et al[13] | Chang et al[14] | Cernik D et al[15] | Anadani et al[16] | Han et al[17] | P. Upadhyaya et al[18]. |
|---|---|---|---|---|---|---|---|---|
| Year | 2017 | 2019 | 2019 | 2019 | 2019 | 2020 | 2021 | 2023 |
| BP Cut-offs | 140 | 158 | 140 | 130 | 140 | 140 | 140 | 130 |
| No of patients-total | 217 | 485 | 1245 | 90 | 690 | 1019 | 187 | 103 |
| n-Intensive | 10 | 195 | 942 | 48 | 376 | 540 | 96 | 44 |
| n-Less intensive | 130 | 290 | 259 | 42 | 314 | 479 | 83 | 59 |
| Age(Mean)-Total | 62 | | 69 | 72.3 | 71 | | 60 | |
| Age(SD)-Total | 14 | | 14 | 11.8 | 13 | | 10 | |
| Age(Mean)_intensive | 61.2 ± 14.2 | | | 70.1 | | 70 | | |
| Age(Mean)-Less Intensive | | | | 74.9 | | | | |
| Age(Median) | | 69 | | | | | | 66 |
| Age(IQR) | | 57-79 | | | | | | 57-75 |
| Males(n) | 108 | 236 | 614 | 54 | 352 | 508 | 157 | 40 |
| Males(%) | 50 | 48.66 | 49.3 | 60 | 51 | 49.8 | 84 | 38.8 |
| Males(Intensive)-n | 2 | | | 31 | | 259 | | 18 |
| % males-Intensive | 20 | | | 64.6 | | 47.96 | | 40.9 |
| Males(Less Intensive)-n | 68 | | | 23 | | 249 | | 22 |
| %males(Less Intensive) | 52.3 | | | 54.8 | | 51.98 | | 37.3 |
| Admission NIHSS-Baseline-Total | 16(12-21) | 16(11-20) | 16(IQR-7) | 14.7±5.5 | 17 | | 22(10-34) | 19 |
| Admission NIHSS-Intensive | 17 | 15 | | 14.2±5.3 | | 15 | | 20 |
| Admission NIHSS-Less Intensive* | | 17 | | 15.3±5.8 | | 17*(160) 16*(180) | | 19 |
| Hypertension-Total-n | 166 | 364 | 797 | 43 | 525 | 644 | 133 | 63 |
| Hypertension%-Total | 77 | 75 | 64 | 49.4 | 76 | 62.59 | 71.1 | 61 |
| Hypertension-Intensive(n) | 6 | 122 | | 20 | | 326 | | 29 |
| Hypertension(%)-Intensive | 60 | 62.5 | | 43.5 | | 60 | | 66 |
| Hypertension-Less Intensive-n | 100 | 242 | | 23 | | 318 | | 34 |
| Hypertension(%)-Less Intensive | 76.9 | 83.4 | | 56.1 | | 66.4 | | 59 |
| Diabetes(Total)-n | 70 | 139 | 406 | 25 | 188 | 356 | 51 | 17 |
| Diabetes(Total)-% | 32 | 29 | 33 | 28.7 | 27 | 34.94 | 27.3 | 17 |
| Diabetes(Intensive)-n | 3 | 51 | | 12 | | 192 | | 7 |
| Diabetes(Intensive)% | 30 | 26 | | 26.1 | | 36 | | 16 |
| Diabetes(Less Intensive)-n | 44 | 88 | | 13 | | 164 | | 10 |
| Diabetes(Less Intensive%) | 33.8 | 30 | | 31.7 | | 34.2 | | 18 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Afib-Total-n | 54 | 176 | 462 | 51 | 299 | | | 31 |
| Afib-Total(%) | 25 | 36 | 37 | 58 | 43 | | | 30 |
| Afib(Intensive)-n | 4 | 63 | | 29 | | 186 | | 14 |
| Afib(Intensive(%) | 40 | 32 | | 61.7 | | 34.4 | | 33 |
| Afib(Less Intensive)-n | 29 | 113 | | 22 | | 203 | | 17 |
| Afib(Less Intensive(%) | 22.3 | 39 | | 53.7 | | 42.4 | | 30 |
| Prior Stroke-Total(n) | | | | | | | 37 | 13 |
| Prior Stroke-Total% | | | | | | | 19.8 | 13 |
| Prior Stroke-Intensive(n) | | | | | | | | 6 |
| Prior Stroke-Intensive% | | | | | | | | 14 |
| Prior Stroke-Less Intensive(n) | | | | | | | | 7 |
| Prior Stroke-Less Intensive% | | | | | | | | 12 |
| IVtPA-Total(n) | 141 | 231 | 590 | 54 | 519 | 566 | 36 | 62 |
| IVtPA-Total(%) | 65 | 48 | 55 | 60 | 75 | 55.5 | 19.3 | 60 |
| IVtPA-Intensive(n) | 9 | 107 | | 29 | | 278 | | 27 |
| IVtPA-Intensive(%) | 90 | 55 | | 60.4 | | 51 | | 61 |
| IVtPA-Less Intensive | 84 | 124 | | 25 | | 288 | | 35 |
| IV tPA-Less intensive% | 64.6 | 42 | | 59.5 | | 60 | | 59 |
| ICA-Total(n) | 33 | 129 | 172 | 26 | 160 | 150 | | 16 |
| ICA-Total(%) | 15 | 27 | 14.2 | 28.9 | 23 | 14.7 | | 16 |
| ICA-Intensive(n) | | 49 | | 11 | | 66 | | 8 |
| ICA-Intensive% | | 25 | | 22.9 | | 11 | | 18 |
| ICA-Less Intensive-n | | 80 | | 15 | | 84 | | 8 |
| ICA-Less Intensive(%) | | 27 | | 35.7 | | 17.5 | | 14 |
| MCA-M1-Total(n) | 104 | 309 | 714 | 64 | 567 | 564 | | 42 |
| MCA-M1-Total(%) | 48 | 64 | 57 | 71.1 | 82 | 55.35 | | 41 |
| M1-Intensive-n | | 108 | | 37 | | 315 | | 19 |
| M1 Intensive% | | 55.3 | | 77.1 | | 59 | | 43 |
| M1 Less Intensive(n) | | 148 | | 27 | | 249 | | 23 |
| M1 Less Intensive(%) | | 51 | | 64.3 | | 51.98 | | 39 |
| Post Circulation-Total-n | 38 | | | | | | 187 | |
| Post Circulation-% | 18 | | | | | | 100 | |

1  *No of patients with each individual group in brackets

1
2
3

4
5 **Figure-1-CONSORT diagram of the study**

6
7
8

## Combined Analysis(n=12)

| Study or Subgroup | Intensive Management Events | Total | Conventional Management Events | Total | Weight | Risk Ratio M–H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 6.2% | 1.36 [0.88, 2.11] | 2017 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 9.7% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 9.9% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 6.4% | 1.65 [1.08, 2.50] | 2019 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 9.2% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 10.3% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 6.7% | 1.22 [0.82, 1.81] | 2021 |
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 8.8% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 10.3% | 0.78 [0.69, 0.89] | 2022 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 7.3% | 1.39 [0.97, 1.99] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 6.4% | 0.97 [0.64, 1.48] | 2023 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 8.8% | 0.72 [0.57, 0.92] | 2023 |
| **Total (95% CI)** | | **2983** | | **2415** | **100.0%** | **1.16 [0.98, 1.37]** | |
| Total events | 1508 | | 1106 | | | | |

Heterogeneity: Tau² = 0.06; Chi² = 70.46, df = 11 (P < 0.00001); I² = 84%
Test for overall effect: Z = 1.74 (P = 0.08)

## Observational Studies(n=8)

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M–H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 4.9% | 1.36 [0.88, 2.11] | 2017 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 14.4% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 18.2% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 19.9% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 5.4% | 1.65 [1.08, 2.50] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 24.3% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 5.9% | 1.22 [0.82, 1.81] | 2021 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 7.0% | 1.39 [0.97, 1.99] | 2023 |
| **Total (95% CI)** | | **2235** | | **1633** | **100.0%** | **1.34 [1.20, 1.48]** | |
| Total events | 1171 | | 674 | | | | |

Heterogeneity: Tau² = 0.01; Chi² = 11.16, df = 7 (P = 0.13); I² = 37%
Test for overall effect: Z = 5.42 (P < 0.00001)

## RCTs(n=4)

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M–H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 20.9% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 48.9% | 0.78 [0.69, 0.89] | 2022 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 21.6% | 0.72 [0.57, 0.92] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 8.6% | 0.97 [0.64, 1.48] | 2023 |
| **Total (95% CI)** | | **748** | | **782** | **100.0%** | **0.82 [0.72, 0.93]** | |
| Total events | 337 | | 432 | | | | |

Heterogeneity: Tau² = 0.00; Chi² = 4.00, df = 3 (P = 0.26); I² = 25%
Test for overall effect: Z = 2.99 (P = 0.003)

**Figure-2-Forest plot for the primary outcome of mRS 0-2 at 90 days.**

**Legend-** The estimated risk ratio-RR is represented by a square and the corresponding 95% CI are shown for the primary outcome(mRS-0-2) at 90 days. The square's size is proportional to the study's weight in the meta-analysis implying studies with larger squares have greater weight. The diamond beneath is the overall pooled estimate and its width denotes the 95%CI around the combined effect size.

**Combined data(n=12)**

| Study or Subgroup | Intensive Events | Total | Conventional Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 0 | 10 | 30 | 130 | 0.5% | 0.20 [0.01, 2.98] | 2017 |
| Anadani et al 2019 | 150 | 942 | 67 | 259 | 15.3% | 0.62 [0.48, 0.79] | 2019 |
| Cernik D et al 2019 | 88 | 376 | 101 | 314 | 15.6% | 0.73 [0.57, 0.93] | 2019 |
| Chang et al 2019 | 5 | 48 | 6 | 42 | 2.8% | 0.73 [0.24, 2.22] | 2019 |
| Eva A. Mistry et al 2019 | 20 | 179 | 59 | 267 | 9.6% | 0.51 [0.32, 0.81] | 2019 |
| Anadani et al 2020 | 84 | 540 | 91 | 479 | 14.8% | 0.82 [0.63, 1.07] | 2020 |
| Han et al 2021 | 21 | 96 | 12 | 83 | 6.6% | 1.51 [0.79, 2.89] | 2021 |
| Mazighi et al 2021 | 29 | 152 | 21 | 153 | 8.7% | 1.39 [0.83, 2.33] | 2021 |
| Yang et al 2022 | 66 | 406 | 61 | 408 | 13.4% | 1.09 [0.79, 1.50] | 2022 |
| Mistry et al 2023 | 7 | 37 | 20 | 76 | 5.1% | 0.72 [0.33, 1.55] | 2023 |
| Nam et al 2023 | 12 | 155 | 8 | 147 | 4.3% | 1.42 [0.60, 3.38] | 2023 |
| P. Upadhyaya et al 2023 | 5 | 44 | 9 | 59 | 3.3% | 0.74 [0.27, 2.07] | 2023 |
| **Total (95% CI)** | | 2985 | | 2417 | 100.0% | 0.83 [0.68, 1.02] | |
| Total events | 487 | | 485 | | | | |

Heterogeneity: Tau² = 0.05; Chi² = 22.93, df = 11 (P = 0.02); I² = 52%
Test for overall effect: Z = 1.76 (P = 0.08)

Favours intensive   Favours conventional

**Observational Studies(n=8)**

| Study or Subgroup | Intensive Arm Events | Total | Conventional Arm Events | Total | Weight | Odds Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 0 | 10 | 30 | 130 | 0.5% | 0.20 [0.01, 2.98] | 2017 |
| Chang et al 2019 | 5 | 48 | 6 | 42 | 2.8% | 0.73 [0.24, 2.22] | 2019 |
| Eva A. Mistry et al 2019 | 20 | 179 | 59 | 267 | 12.0% | 0.51 [0.32, 0.81] | 2019 |
| Anadani et al 2019 | 150 | 942 | 67 | 259 | 24.9% | 0.62 [0.48, 0.79] | 2019 |
| Cernik D et al 2019 | 88 | 376 | 101 | 314 | 25.7% | 0.73 [0.57, 0.93] | 2019 |
| Anadani et al 2020 | 84 | 540 | 91 | 479 | 23.5% | 0.82 [0.63, 1.07] | 2020 |
| Han et al 2021 | 21 | 96 | 12 | 83 | 7.3% | 1.51 [0.79, 2.89] | 2021 |
| P. Upadhyaya et al 2023 | 5 | 44 | 9 | 59 | 3.3% | 0.74 [0.27, 2.07] | 2023 |
| **Total (95% CI)** | | 2235 | | 1633 | 100.0% | 0.72 [0.59, 0.87] | |
| Total events | 373 | | 375 | | | | |

Heterogeneity: Tau² = 0.02; Chi² = 10.48, df = 7 (P = 0.16); I² = 33%
Test for overall effect: Z = 3.35 (P = 0.0008)

Favours Intensive Control   Favours Conventional

**RCTs(n=4)**

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 29 | 152 | 21 | 153 | 22.7% | 1.39 [0.83, 2.33] | 2021 |
| Yang et al 2022 | 66 | 406 | 61 | 408 | 58.9% | 1.09 [0.79, 1.50] | 2022 |
| Mistry et al 2023 | 7 | 37 | 20 | 76 | 10.3% | 0.72 [0.33, 1.55] | 2023 |
| Nam et al 2023 | 12 | 155 | 8 | 147 | 8.0% | 1.42 [0.60, 3.38] | 2023 |
| **Total (95% CI)** | | 750 | | 784 | 100.0% | 1.13 [0.88, 1.44] | |
| Total events | 114 | | 110 | | | | |

Heterogeneity: Tau² = 0.00; Chi² = 2.29, df = 3 (P = 0.51); I² = 0%
Test for overall effect: Z = 0.95 (P = 0.34)

Favours Intensive   Favours Conventional

**Figure-3-Forest plot for the outcome of death at 90 days**

**Legend-**The estimated risk ratio-RR is represented by a square and the corresponding 95% CI are shown for the outcome death at 90 days. The square's size is proportional to the study's weight in the meta-analysis implying studies with larger squares have greater weight. The diamond beneath is the overall pooled estimate and its width denotes the 95%CI around the combined effect size.

## Combined data(n=9)

| Study or Subgroup | Intensive Events | Total | Conventional Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 1 | 10 | 7 | 130 | 2.4% | 1.86 [0.25, 13.64] | 2017 |
| Anadani et al 2019 | 37 | 942 | 20 | 259 | 18.2% | 0.51 [0.30, 0.86] | 2019 |
| Anadani et al 2020 | 16 | 540 | 29 | 479 | 15.9% | 0.49 [0.27, 0.89] | 2020 |
| Han et al 2021 | 19 | 96 | 16 | 83 | 15.9% | 1.03 [0.57, 1.86] | 2021 |
| Mazighi et al 2021 | 17 | 154 | 11 | 157 | 12.6% | 1.58 [0.76, 3.25] | 2021 |
| Yang et al 2022 | 23 | 407 | 25 | 409 | 17.4% | 0.92 [0.53, 1.60] | 2022 |
| Mistry et al 2023 | 2 | 37 | 3 | 72 | 3.1% | 1.30 [0.23, 7.43] | 2023 |
| Nam et al 2023 | 14 | 155 | 12 | 149 | 12.3% | 1.12 [0.54, 2.34] | 2023 |
| P. Upadhyaya et al 2023 | 1 | 44 | 4 | 59 | 2.1% | 0.34 [0.04, 2.90] | 2023 |
| **Total (95% CI)** | | 2385 | | 1797 | 100.0% | 0.84 [0.61, 1.16] | |
| Total events | 130 | | 127 | | | | |

Heterogeneity: Tau² = 0.08; Chi² = 12.13, df = 8 (P = 0.15); I² = 34%
Test for overall effect: Z = 1.07 (P = 0.29)

## Observational Studies(n=5)

| Study or Subgroup | Intensive Control Events | Total | Conventional Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 1 | 10 | 7 | 130 | 3.9% | 1.86 [0.25, 13.64] | 2017 |
| Anadani et al 2019 | 37 | 942 | 20 | 259 | 34.1% | 0.51 [0.30, 0.86] | 2019 |
| Anadani et al 2020 | 16 | 540 | 29 | 479 | 29.2% | 0.49 [0.27, 0.89] | 2020 |
| Han et al 2021 | 19 | 96 | 16 | 83 | 29.3% | 1.03 [0.57, 1.86] | 2021 |
| P. Upadhyaya et al 2023 | 1 | 44 | 4 | 59 | 3.4% | 0.34 [0.04, 2.90] | 2023 |
| **Total (95% CI)** | | 1632 | | 1010 | 100.0% | 0.64 [0.43, 0.96] | |
| Total events | 74 | | 76 | | | | |

Heterogeneity: Tau² = 0.05; Chi² = 5.37, df = 4 (P = 0.25); I² = 26%
Test for overall effect: Z = 2.15 (P = 0.03)

## RCTs(n=4)

| Study or Subgroup | Intensive Arm Events | Total | Conventional Arm Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 17 | 154 | 11 | 157 | 25.8% | 1.58 [0.76, 3.25] | 2021 |
| Yang et al 2022 | 23 | 407 | 25 | 409 | 44.9% | 0.92 [0.53, 1.60] | 2022 |
| Nam et al 2023 | 14 | 155 | 12 | 149 | 24.9% | 1.12 [0.54, 2.34] | 2023 |
| Mistry et al 2023 | 2 | 37 | 3 | 72 | 4.5% | 1.30 [0.23, 7.43] | 2023 |
| **Total (95% CI)** | | 753 | | 787 | 100.0% | 1.13 [0.78, 1.63] | |
| Total events | 56 | | 51 | | | | |

Heterogeneity: Tau² = 0.00; Chi² = 1.34, df = 3 (P = 0.72); I² = 0%
Test for overall effect: Z = 0.65 (P = 0.52)

**Figure-4- Forest plot for the outcome of symptomatic ICH(sICH)**

**Legend-**The estimated risk ratio-RR is represented by a square and the corresponding 95% CI are shown for the outcome sICH. The square's size is proportional to the study's weight in the meta-analysis implying studies with larger squares have greater weight. The diamond beneath is the overall pooled estimate and its width denotes the 95%CI around the combined effect size.

**Combined data(n=4)**

| | Intensive | | Conventional | | | Risk Ratio | | Risk Ratio |
|---|---|---|---|---|---|---|---|---|
| Study or Subgroup | Events | Total | Events | Total | Weight | M–H, Random, 95% CI | Year | M–H, Random, 95% CI |
| Anadani et al 2019 | 31 | 942 | 19 | 259 | 53.6% | 0.45 [0.26, 0.78] | 2019 | |
| Anadani et al 2020 | 11 | 540 | 34 | 479 | 40.9% | 0.29 [0.15, 0.56] | 2020 | |
| Mazighi et al 2021 | 1 | 158 | 0 | 162 | 2.3% | 3.08 [0.13, 74.93] | 2021 | |
| P. Upadhyaya et al 2023 | 1 | 44 | 1 | 59 | 3.2% | 1.34 [0.09, 20.85] | 2023 | |
| | | | | | | | | |
| Total (95% CI) | | 1684 | | 959 | 100.0% | 0.40 [0.25, 0.66] | | |
| Total events | 44 | | 54 | | | | | |

Heterogeneity: Tau$^2$ = 0.04; Chi$^2$ = 3.43, df = 3 (P = 0.33); I$^2$ = 13%
Test for overall effect: Z = 3.60 (P = 0.0003)

0.01  0.1  1  10  100
Favours Intensive  Favours Conventional

**Observational Studies(n=3)**

| | Intensive | | Conventional | | | Risk Ratio | | Risk Ratio |
|---|---|---|---|---|---|---|---|---|
| Study or Subgroup | Events | Total | Events | Total | Weight | M–H, Random, 95% CI | Year | M–H, Random, 95% CI |
| Anadani et al 2019 | 31 | 942 | 19 | 259 | 57.9% | 0.45 [0.26, 0.78] | 2019 | |
| Anadani et al 2020 | 11 | 540 | 34 | 479 | 39.8% | 0.29 [0.15, 0.56] | 2020 | |
| P. Upadhyaya et al 2023 | 1 | 44 | 1 | 59 | 2.4% | 1.34 [0.09, 20.85] | 2023 | |
| | | | | | | | | |
| Total (95% CI) | | 1526 | | 797 | 100.0% | 0.39 [0.25, 0.59] | | |
| Total events | 43 | | 54 | | | | | |

Heterogeneity: Tau$^2$ = 0.00; Chi$^2$ = 1.85, df = 2 (P = 0.40); I$^2$ = 0%
Test for overall effect: Z = 4.43 (P < 0.00001)

0.01  0.1  1  10  100
Favours Intensive Control  Favours Conventional

2
**Figure-5- Forest plot for the outcome of decompressive surgery.**
**Legend-**The estimated risk ratio-RR is represented by a square and the corresponding 95%
CI are shown for the outcome requirement of decompressive surgery. The square's size is
proportional to the study's weight in the meta-analysis implying studies with larger squares
have greater weight. The diamond beneath is the overall pooled estimate and its width
denotes the 95%CI around the combined effect size.

## Combined Analysis of n=7 studies(Intensive Control Cut-off=140mmHg)

| Study or Subgroup | Intensive Management Events | Total | Conventional Management Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 9.1% | 1.36 [0.88, 2.11] | 2017 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 17.9% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 18.5% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 0.0% | 1.65 [1.08, 2.50] | 2019 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 0.0% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 19.7% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 10.2% | 1.22 [0.82, 1.81] | 2021 |
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 0.0% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 0.0% | 0.78 [0.69, 0.89] | 2022 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 0.0% | 1.39 [0.97, 1.99] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 9.5% | 0.97 [0.64, 1.48] | 2023 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 15.2% | 0.72 [0.57, 0.92] | 2023 |
| **Total (95% CI)** | | 2156 | | 1488 | 100.0% | 1.14 [0.96, 1.36] | |
| Total events | 1093 | | 656 | | | | |

Heterogeneity: Tau² = 0.04; Chi² = 23.28, df = 6 (P = 0.0007); I² = 74%
Test for overall effect: Z = 1.49 (P = 0.14)

Favours [conventional]  Favours [intensive]

## Observational Studies(n=5)-Intensive Control Cut-off=140mmHg

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 5.3% | 1.36 [0.88, 2.11] | 2017 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 0.0% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 24.4% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 27.4% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 0.0% | 1.65 [1.08, 2.50] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 36.4% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 6.4% | 1.22 [0.82, 1.81] | 2021 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 0.0% | 1.39 [0.97, 1.99] | 2023 |
| **Total (95% CI)** | | 1964 | | 1265 | 100.0% | 1.26 [1.13, 1.40] | |
| Total events | 1015 | | 540 | | | | |

Heterogeneity: Tau² = 0.00; Chi² = 5.47, df = 4 (P = 0.24); I² = 27%
Test for overall effect: Z = 4.32 (P < 0.0001)

Favours Conventional  Favours Intensive

## RCTs(n=2)-Intensive Control Cut-off=140mmHg

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 0.0% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 0.0% | 0.78 [0.69, 0.89] | 2022 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 67.8% | 0.72 [0.57, 0.92] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 32.2% | 0.97 [0.64, 1.48] | 2023 |
| **Total (95% CI)** | | 192 | | 223 | 100.0% | 0.79 [0.61, 1.04] | |
| Total events | 78 | | 116 | | | | |

Heterogeneity: Tau² = 0.01; Chi² = 1.39, df = 1 (P = 0.24); I² = 28%
Test for overall effect: Z = 1.67 (P = 0.09)

Favours Conventional  Favours Intensive

## Combined Analysis of studies(n=3) using a SBP cut-off of 130mmHg (mRS 0-2@90d) for intensive control

| Study or Subgroup | Intensive Management Events | Total | Conventional Management Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 0.0% | 1.36 [0.88, 2.11] | 2017 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 0.0% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 0.0% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 27.9% | 1.65 [1.08, 2.50] | 2019 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 0.0% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 0.0% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 0.0% | 1.22 [0.82, 1.81] | 2021 |
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 40.1% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 0.0% | 0.78 [0.69, 0.89] | 2022 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 32.0% | 1.39 [0.97, 1.99] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 0.0% | 0.97 [0.64, 1.48] | 2023 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 0.0% | 0.72 [0.57, 0.92] | 2023 |
| **Total (95% CI)** | | 244 | | 254 | 100.0% | 1.27 [0.92, 1.74] | |
| Total events | 127 | | 113 | | | | |

Heterogeneity: Tau² = 0.05; Chi² = 5.45, df = 2 (P = 0.07); I² = 63%
Test for overall effect: Z = 1.44 (P = 0.15)

Favours [conventional]  Favours [intensive]

**Figure-6**-**mRS 0-2 at 90 days with different SBP cut-off to define intensive control**

**Legend-** The estimated risk ratio-RR is represented by a square and the corresponding 95% CI are shown for the primary outcome(mRS-0-2) at 90 days. The square's size is proportional to the study's weight in the meta-analysis implying studies with larger squares have greater weight. The diamond beneath is the overall pooled estimate and its width denotes the 95%CI around the combined effect size.

## Combined Analysis(n=12)

| Study or Subgroup | Intensive Management Events | Total | Conventional Management Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 6.2% | 1.36 [0.88, 2.11] | 2017 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 9.7% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 9.9% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 6.4% | 1.65 [1.08, 2.50] | 2019 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 9.2% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 10.3% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 6.7% | 1.22 [0.82, 1.81] | 2021 |
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 8.8% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 10.3% | 0.78 [0.69, 0.89] | 2022 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 7.3% | 1.39 [0.97, 1.99] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 6.4% | 0.97 [0.64, 1.48] | 2023 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 8.8% | 0.72 [0.57, 0.92] | 2023 |
| **Total (95% CI)** | | 2983 | | 2415 | 100.0% | 1.16 [0.98, 1.37] | |
| Total events | 1508 | | 1106 | | | | |

Heterogeneity: Tau$^2$ = 0.06; Chi$^2$ = 70.46, df = 11 (P < 0.00001); I$^2$ = 84%
Test for overall effect: Z = 1.74 (P = 0.08)

Favours [conventional]  Favours [intensive]

## Observational Studies(n=8)

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 4.9% | 1.36 [0.88, 2.11] | 2017 |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 14.4% | 1.59 [1.28, 1.98] | 2019 |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 18.2% | 1.45 [1.22, 1.73] | 2019 |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 19.9% | 1.28 [1.09, 1.50] | 2019 |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 5.4% | 1.65 [1.08, 2.50] | 2019 |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 24.3% | 1.13 [1.00, 1.28] | 2020 |
| Han et al 2021 | 38 | 96 | 27 | 83 | 5.9% | 1.22 [0.82, 1.81] | 2021 |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 7.0% | 1.39 [0.97, 1.99] | 2023 |
| **Total (95% CI)** | | 2235 | | 1633 | 100.0% | 1.34 [1.20, 1.48] | |
| Total events | 1171 | | 674 | | | | |

Heterogeneity: Tau$^2$ = 0.01; Chi$^2$ = 11.16, df = 7 (P = 0.13); I$^2$ = 37%
Test for overall effect: Z = 5.42 (P < 0.00001)

Favours Conventional  Favours Intensive

## RCTs(n=4)

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 20.9% | 0.98 [0.76, 1.26] | 2021 |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 48.9% | 0.78 [0.69, 0.89] | 2022 |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 21.6% | 0.72 [0.57, 0.92] | 2023 |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 8.6% | 0.97 [0.64, 1.48] | 2023 |
| **Total (95% CI)** | | 748 | | 782 | 100.0% | 0.82 [0.72, 0.93] | |
| Total events | 337 | | 432 | | | | |

Heterogeneity: Tau$^2$ = 0.00; Chi$^2$ = 4.00, df = 3 (P = 0.26); I$^2$ = 25%
Test for overall effect: Z = 2.99 (P = 0.003)

Favours Conventional  Favours Intensive

## Combined data(n=12)

| Study or Subgroup | Intensive Events | Total | Conventional Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 0 | 10 | 30 | 130 | 0.5% | 0.20 [0.01, 2.98] | 2017 |
| Anadani et al 2019 | 150 | 942 | 67 | 259 | 15.3% | 0.62 [0.48, 0.79] | 2019 |
| Cernik D et al 2019 | 88 | 376 | 101 | 314 | 15.6% | 0.73 [0.57, 0.93] | 2019 |
| Chang et al 2019 | 5 | 48 | 6 | 42 | 2.8% | 0.73 [0.24, 2.22] | 2019 |
| Eva A. Mistry et al 2019 | 20 | 179 | 59 | 267 | 9.6% | 0.51 [0.32, 0.81] | 2019 |
| Anadani et al 2020 | 84 | 540 | 91 | 479 | 14.8% | 0.82 [0.63, 1.07] | 2020 |
| Han et al 2021 | 21 | 96 | 12 | 83 | 6.6% | 1.51 [0.79, 2.89] | 2021 |
| Mazighi et al 2021 | 29 | 152 | 21 | 153 | 8.7% | 1.39 [0.83, 2.33] | 2021 |
| Yang et al 2022 | 66 | 406 | 61 | 408 | 13.4% | 1.09 [0.79, 1.50] | 2022 |
| Mistry et al 2023 | 7 | 37 | 20 | 76 | 5.1% | 0.72 [0.33, 1.55] | 2023 |
| Nam et al 2023 | 12 | 155 | 8 | 147 | 4.3% | 1.42 [0.60, 3.38] | 2023 |
| P. Upadhyaya et al 2023 | 5 | 44 | 9 | 59 | 3.3% | 0.74 [0.27, 2.07] | 2023 |
| **Total (95% CI)** | | **2985** | | **2417** | **100.0%** | **0.83 [0.68, 1.02]** | |
| Total events | 487 | | 485 | | | | |

Heterogeneity: Tau² = 0.05; Chi² = 22.93, df = 11 (P = 0.02); I² = 52%
Test for overall effect: Z = 1.76 (P = 0.08)

Favours intensive — Favours conventional

## Observational Studies(n=8)

| Study or Subgroup | Intensive Arm Events | Total | Conventional Arm Events | Total | Weight | Odds Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 0 | 10 | 30 | 130 | 0.5% | 0.20 [0.01, 2.98] | 2017 |
| Chang et al 2019 | 5 | 48 | 6 | 42 | 2.8% | 0.73 [0.24, 2.22] | 2019 |
| Eva A. Mistry et al 2019 | 20 | 179 | 59 | 267 | 12.0% | 0.51 [0.32, 0.81] | 2019 |
| Anadani et al 2019 | 150 | 942 | 67 | 259 | 24.9% | 0.62 [0.48, 0.79] | 2019 |
| Cernik D et al 2019 | 88 | 376 | 101 | 314 | 25.7% | 0.73 [0.57, 0.93] | 2019 |
| Anadani et al 2020 | 84 | 540 | 91 | 479 | 23.5% | 0.82 [0.63, 1.07] | 2020 |
| Han et al 2021 | 21 | 96 | 12 | 83 | 7.3% | 1.51 [0.79, 2.89] | 2021 |
| P. Upadhyaya et al 2023 | 5 | 44 | 9 | 59 | 3.3% | 0.74 [0.27, 2.07] | 2023 |
| **Total (95% CI)** | | **2235** | | **1633** | **100.0%** | **0.72 [0.59, 0.87]** | |
| Total events | 373 | | 375 | | | | |

Heterogeneity: Tau² = 0.02; Chi² = 10.48, df = 7 (P = 0.16); I² = 33%
Test for overall effect: Z = 3.35 (P = 0.0008)

Favours Intensive Control — Favours Conventional

## RCTs(n=4)

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 29 | 152 | 21 | 153 | 22.7% | 1.39 [0.83, 2.33] | 2021 |
| Yang et al 2022 | 66 | 406 | 61 | 408 | 58.9% | 1.09 [0.79, 1.50] | 2022 |
| Mistry et al 2023 | 7 | 37 | 20 | 76 | 10.3% | 0.72 [0.33, 1.55] | 2023 |
| Nam et al 2023 | 12 | 155 | 8 | 147 | 8.0% | 1.42 [0.60, 3.38] | 2023 |
| **Total (95% CI)** | | **750** | | **784** | **100.0%** | **1.13 [0.88, 1.44]** | |
| Total events | 114 | | 110 | | | | |

Heterogeneity: Tau² = 0.00; Chi² = 2.29, df = 3 (P = 0.51); I² = 0%
Test for overall effect: Z = 0.95 (P = 0.34)

Favours Intensive — Favours Conventional

## Combined data(n=9)

| Study or Subgroup | Intensive Events | Total | Conventional Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 1 | 10 | 7 | 130 | 2.4% | 1.86 [0.25, 13.64] | 2017 |
| Anadani et al 2019 | 37 | 942 | 20 | 259 | 18.2% | 0.51 [0.30, 0.86] | 2019 |
| Anadani et al 2020 | 16 | 540 | 29 | 479 | 15.9% | 0.49 [0.27, 0.89] | 2020 |
| Han et al 2021 | 19 | 96 | 16 | 83 | 15.9% | 1.03 [0.57, 1.86] | 2021 |
| Mazighi et al 2021 | 17 | 154 | 11 | 157 | 12.6% | 1.58 [0.76, 3.25] | 2021 |
| Yang et al 2022 | 23 | 407 | 25 | 409 | 17.4% | 0.92 [0.53, 1.60] | 2022 |
| Mistry et al 2023 | 2 | 37 | 3 | 72 | 3.1% | 1.30 [0.23, 7.43] | 2023 |
| Nam et al 2023 | 14 | 155 | 12 | 149 | 12.3% | 1.12 [0.54, 2.34] | 2023 |
| P. Upadhyaya et al 2023 | 1 | 44 | 4 | 59 | 2.1% | 0.34 [0.04, 2.90] | 2023 |
| **Total (95% CI)** | | **2385** | | **1797** | **100.0%** | **0.84 [0.61, 1.16]** | |
| Total events | 130 | | 127 | | | | |

Heterogeneity: Tau$^2$ = 0.08; Chi$^2$ = 12.13, df = 8 (P = 0.15); I$^2$ = 34%
Test for overall effect: Z = 1.07 (P = 0.29)

Risk Ratio M-H, Random, 95% CI — scale: 0.01 0.1 1 10 100 — Favours Intensive / Favours Conventional

## Observational Studies(n=5)

| Study or Subgroup | Intensive Control Events | Total | Conventional Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 1 | 10 | 7 | 130 | 3.9% | 1.86 [0.25, 13.64] | 2017 |
| Anadani et al 2019 | 37 | 942 | 20 | 259 | 34.1% | 0.51 [0.30, 0.86] | 2019 |
| Anadani et al 2020 | 16 | 540 | 29 | 479 | 29.2% | 0.49 [0.27, 0.89] | 2020 |
| Han et al 2021 | 19 | 96 | 16 | 83 | 29.3% | 1.03 [0.57, 1.86] | 2021 |
| P. Upadhyaya et al 2023 | 1 | 44 | 4 | 59 | 3.4% | 0.34 [0.04, 2.90] | 2023 |
| **Total (95% CI)** | | **1632** | | **1010** | **100.0%** | **0.64 [0.43, 0.96]** | |
| Total events | 74 | | 76 | | | | |

Heterogeneity: Tau$^2$ = 0.05; Chi$^2$ = 5.37, df = 4 (P = 0.25); I$^2$ = 26%
Test for overall effect: Z = 2.15 (P = 0.03)

Risk Ratio M-H, Random, 95% CI — scale: 0.01 0.1 1 10 100 — Favours Intensive / Favours Conventional

## RCTs(n=4)

| Study or Subgroup | Intensive Arm Events | Total | Conventional Arm Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 17 | 154 | 11 | 157 | 25.8% | 1.58 [0.76, 3.25] | 2021 |
| Yang et al 2022 | 23 | 407 | 25 | 409 | 44.9% | 0.92 [0.53, 1.60] | 2022 |
| Nam et al 2023 | 14 | 155 | 12 | 149 | 24.9% | 1.12 [0.54, 2.34] | 2023 |
| Mistry et al 2023 | 2 | 37 | 3 | 72 | 4.5% | 1.30 [0.23, 7.43] | 2023 |
| **Total (95% CI)** | | **753** | | **787** | **100.0%** | **1.13 [0.78, 1.63]** | |
| Total events | 56 | | 51 | | | | |

Heterogeneity: Tau$^2$ = 0.00; Chi$^2$ = 1.34, df = 3 (P = 0.72); I$^2$ = 0%
Test for overall effect: Z = 0.65 (P = 0.52)

Risk Ratio M-H, Random, 95% CI — scale: 0.01 0.1 1 10 100 — Favours Intensive / Favours Conventional

## Combined data(n=4)

| Study or Subgroup | Intensive Events | Total | Conventional Events | Total | Weight | Risk Ratio M–H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Anadani et al 2019 | 31 | 942 | 19 | 259 | 53.6% | 0.45 [0.26, 0.78] | 2019 |
| Anadani et al 2020 | 11 | 540 | 34 | 479 | 40.9% | 0.29 [0.15, 0.56] | 2020 |
| Mazighi et al 2021 | 1 | 158 | 0 | 162 | 2.3% | 3.08 [0.13, 74.93] | 2021 |
| P. Upadhyaya et al 2023 | 1 | 44 | 1 | 59 | 3.2% | 1.34 [0.09, 20.85] | 2023 |
| | | | | | | | |
| Total (95% CI) | | 1684 | | 959 | 100.0% | 0.40 [0.25, 0.66] | |
| Total events | 44 | | 54 | | | | |

Heterogeneity: Tau$^2$ = 0.04; Chi$^2$ = 3.43, df = 3 (P = 0.33); I$^2$ = 13%
Test for overall effect: Z = 3.60 (P = 0.0003)

Risk Ratio M–H, Random, 95% CI
Favours Intensive   Favours Conventional

## Observational Studies(n=3)

| Study or Subgroup | Intensive Events | Total | Conventional Events | Total | Weight | Risk Ratio M–H, Random, 95% CI | Year |
|---|---|---|---|---|---|---|---|
| Anadani et al 2019 | 31 | 942 | 19 | 259 | 57.9% | 0.45 [0.26, 0.78] | 2019 |
| Anadani et al 2020 | 11 | 540 | 34 | 479 | 39.8% | 0.29 [0.15, 0.56] | 2020 |
| P. Upadhyaya et al 2023 | 1 | 44 | 1 | 59 | 2.4% | 1.34 [0.09, 20.85] | 2023 |
| | | | | | | | |
| Total (95% CI) | | 1526 | | 797 | 100.0% | 0.39 [0.25, 0.59] | |
| Total events | 43 | | 54 | | | | |

Heterogeneity: Tau$^2$ = 0.00; Chi$^2$ = 1.85, df = 2 (P = 0.40); I$^2$ = 0%
Test for overall effect: Z = 4.43 (P < 0.00001)

Risk Ratio M–H, Random, 95% CI
Favours Intensive Control   Favours Conventional

## Combined Analysis of n=7 studies(Intensive Control Cut-off=140mmHg)

| Study or Subgroup | Intensive Management Events | Total | Conventional Management Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year | Risk Ratio M-H, Random, 95% CI |
|---|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 9.1% | 1.36 [0.88, 2.11] | 2017 | |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 17.9% | 1.45 [1.22, 1.73] | 2019 | |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 18.5% | 1.28 [1.09, 1.50] | 2019 | |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 0.0% | 1.65 [1.08, 2.50] | 2019 | |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 0.0% | 1.59 [1.28, 1.98] | 2019 | |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 19.7% | 1.13 [1.00, 1.28] | 2020 | |
| Han et al 2021 | 38 | 96 | 27 | 83 | 10.2% | 1.22 [0.82, 1.81] | 2021 | |
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 0.0% | 0.98 [0.76, 1.26] | 2021 | |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 0.0% | 0.78 [0.69, 0.89] | 2022 | |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 0.0% | 1.39 [0.97, 1.99] | 2023 | |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 9.5% | 0.97 [0.64, 1.48] | 2023 | |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 15.2% | 0.72 [0.57, 0.92] | 2023 | |
| **Total (95% CI)** | | **2156** | | **1488** | **100.0%** | **1.14 [0.96, 1.36]** | | |
| Total events | 1093 | | 656 | | | | | |

Heterogeneity: Tau² = 0.04; Chi² = 23.28, df = 6 (P = 0.0007); I² = 74%
Test for overall effect: Z = 1.49 (P = 0.14)

Favours [conventional]   Favours [intensive]

## Observational Studies(n=5)-Intensive Control Cut-off=140mmHg

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year | Risk Ratio M-H, Random, 95% CI |
|---|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 5.3% | 1.36 [0.88, 2.11] | 2017 | |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 0.0% | 1.59 [1.28, 1.98] | 2019 | |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 24.4% | 1.45 [1.22, 1.73] | 2019 | |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 27.4% | 1.28 [1.09, 1.50] | 2019 | |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 0.0% | 1.65 [1.08, 2.50] | 2019 | |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 36.4% | 1.13 [1.00, 1.28] | 2020 | |
| Han et al 2021 | 38 | 96 | 27 | 83 | 6.4% | 1.22 [0.82, 1.81] | 2021 | |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 0.0% | 1.39 [0.97, 1.99] | 2023 | |
| **Total (95% CI)** | | **1964** | | **1265** | **100.0%** | **1.26 [1.13, 1.40]** | | |
| Total events | 1015 | | 540 | | | | | |

Heterogeneity: Tau² = 0.00; Chi² = 5.47, df = 4 (P = 0.24); I² = 27%
Test for overall effect: Z = 4.32 (P < 0.0001)

Favours Conventional   Favours Intensive

## RCTs(n=2)-Intensive Control Cut-off=140mmHg

| Study or Subgroup | Intensive Control Events | Total | Conventional Control Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year | Risk Ratio M-H, Random, 95% CI |
|---|---|---|---|---|---|---|---|---|
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 0.0% | 0.98 [0.76, 1.26] | 2021 | |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 0.0% | 0.78 [0.69, 0.89] | 2022 | |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 67.8% | 0.72 [0.57, 0.92] | 2023 | |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 32.2% | 0.97 [0.64, 1.48] | 2023 | |
| **Total (95% CI)** | | **192** | | **223** | **100.0%** | **0.79 [0.61, 1.04]** | | |
| Total events | 78 | | 116 | | | | | |

Heterogeneity: Tau² = 0.01; Chi² = 1.39, df = 1 (P = 0.24); I² = 28%
Test for overall effect: Z = 1.67 (P = 0.09)

Favours Conventional   Favours Intensive

## Combined Analysis of studies(n=3) using a SBP cut-off of 130mmHg (mRS 0-2@90d) for intensive control

| Study or Subgroup | Intensive Management Events | Total | Conventional Management Events | Total | Weight | Risk Ratio M-H, Random, 95% CI | Year | Risk Ratio M-H, Random, 95% CI |
|---|---|---|---|---|---|---|---|---|
| Goyal et al 2017 | 7 | 10 | 67 | 130 | 0.0% | 1.36 [0.88, 2.11] | 2017 | |
| Anadani et al 2019 | 485 | 942 | 92 | 259 | 0.0% | 1.45 [1.22, 1.73] | 2019 | |
| Cernik D et al 2019 | 202 | 376 | 132 | 314 | 0.0% | 1.28 [1.09, 1.50] | 2019 | |
| Chang et al 2019 | 32 | 48 | 17 | 42 | 27.9% | 1.65 [1.08, 2.50] | 2019 | |
| Eva A. Mistry et al 2019 | 96 | 179 | 90 | 267 | 0.0% | 1.59 [1.28, 1.98] | 2019 | |
| Anadani et al 2020 | 283 | 540 | 222 | 479 | 0.0% | 1.13 [1.00, 1.28] | 2020 | |
| Han et al 2021 | 38 | 96 | 27 | 83 | 0.0% | 1.22 [0.82, 1.81] | 2021 | |
| Mazighi et al 2021 | 67 | 152 | 69 | 153 | 40.1% | 0.98 [0.76, 1.26] | 2021 | |
| Yang et al 2022 | 192 | 404 | 247 | 406 | 0.0% | 0.78 [0.69, 0.89] | 2022 | |
| P. Upadhyaya et al 2023 | 28 | 44 | 27 | 59 | 32.0% | 1.39 [0.97, 1.99] | 2023 | |
| Mistry et al 2023 | 17 | 37 | 36 | 76 | 0.0% | 0.97 [0.64, 1.48] | 2023 | |
| Nam et al 2023 | 61 | 155 | 80 | 147 | 0.0% | 0.72 [0.57, 0.92] | 2023 | |
| **Total (95% CI)** | | **244** | | **254** | **100.0%** | **1.27 [0.92, 1.74]** | | |
| Total events | 127 | | 113 | | | | | |

Heterogeneity: Tau² = 0.05; Chi² = 5.45, df = 2 (P = 0.07); I² = 63%
Test for overall effect: Z = 1.44 (P = 0.15)

Favours [conventional]   Favours [intensive]