

1

2

Prevalence of Social Anxiety Disorder and its associated

3

factors among Foreign-born Undergraduate Students in

4

Türkiye: A Cross-Sectional Study

5

6

Lujain Alnemr^{1¶}, Abdelaziz H. Salama^{1,2¶*}, Salma Abdelrazek^{1&}, Hussein Alfakeer^{1&}, Mohamed

7

Ali Alkhateeb¹, Perihan Torun³

8

9

¹ Hamidiye International School of Medicine, University of Health Sciences, Istanbul, Türkiye

10

11

² School of Medicine, Johns Hopkins University, Maryland, United States of America

12

13

³ Department of Public Health, Hamidiye International School of Medicine, University of Health Sciences, Istanbul, Türkiye

14

15

16

17

18

*Corresponding author

19

20

E-mail: asalama5@jh.edu

21

22

¶ These authors contributed equally to this work.

23

24

& These authors also contributed equally to this work.

25

26

Abstract

27

Social anxiety disorder (SAD) is prevalent among university students, yet data on its

28

severity among foreign-born international undergraduate students in Türkiye remains

29

limited. This study aims to determine the prevalence of SAD and its associated factors

30

within this population. A cross-sectional study was conducted using a Google Form

31

survey distributed across various universities from September 17, 2023, to February 1,

32

2024. The survey comprised two sections: sociodemographic information and 17 items of

33 the Social Phobia Inventory (SPIN), which measures the frequency and intensity of social
34 anxiety symptoms. Data analysis included descriptive statistics and inferential analysis,
35 multiple regression, and binomial logistic regression. Out of 506 participants, 455 were
36 included in the study. Results revealed that 39.1% exhibited no or very mild symptoms of
37 SAD, while 23.7% experienced mild symptoms, 21.3% faced moderate symptoms, and
38 11.6% and 4.2% presented with severe to very severe symptoms, respectively. Factors
39 such as gender ($p < 0.0001$), previous academic failures ($p = 0.013$), family history of
40 mental health issues ($p = 0.009$), exercise frequency ($p < 0.0001$), and perceptions of
41 relationships with classmates ($p < 0.0001$) were significantly associated with SAD.
42 Females showed a higher probability of SAD compared to males (OR = 1.976).
43 Individuals engaging in over 90 minutes of exercise per week were less likely to have
44 SAD (OR = 0.383), and occasional smokers had a lower risk of SAD compared to non-
45 smokers (OR = 0.422). Our study uncovered a notably elevated prevalence of Social
46 Anxiety Disorder (SAD) among foreign-born undergraduate students in Türkiye. Factors
47 such as being female, having a family history of mental illnesses, studying in a stressful
48 environment, experiencing academic failure, and engaging in less frequent exercise were
49 associated with noticeable symptoms of SAD. These findings emphasize the urgent need
50 for heightened efforts in recognizing and addressing SAD within this population.

51

52

53

54

55

56

57 **Introduction**

58 Anxiety disorders are the most common group of mental illnesses in the world and the
59 ninth most health-related cause of disability, according to the World Health Organization
60 (WHO) [1,2]. Social Anxiety Disorder (SAD), also referred to as social phobia, is
61 classified according to the Diagnostic and Statistical Manual of Mental Disorders V
62 (DSM-V) as an anxiety disorder that presents with an intense fear of being in social
63 situations where one may be under scrutiny by others [3]. SAD usually starts during late
64 childhood and continues throughout life [4, 5]. People with SAD endure severe emotional
65 and/or physical symptoms and may also show signs of shyness and discomfort when in
66 social situations, even though they crave other people's company [6].

67

68 Long-term effects of SAD include low self-esteem, decreased quality of life, and
69 academic underachievement [7-9]. Among university students, the prevalence of SAD is
70 very prominent [10-14]. Moreover, enrolling in a university forces students to deal with
71 new life experiences, such as adjusting to new life circumstances, engaging in university
72 society, and fulfilling the demands of life. These factors are recognized to impact
73 students' mental health and their participation in risky behaviors affecting their health,
74 such as smoking, drinking alcohol, eating poorly, or being physically inactive [15].

75

76

77 Students seeking education abroad face significant difficulties adjusting due to the large
78 variations in social and cultural norms [16]. They may experience emotional issues,
79 including loss, separation, travel stress, social pressure, exacerbation of any underlying or
80 hidden disorders,
81 pharmaceutical complications, and unexpected situations [17,18]. These challenges may,
82 therefore, result in anxiety, anorexia, and general avoidance of social interactions [19,20].

83 In a study done on Chinese international students, the researchers found that students
84 faced challenges regarding language, academics, psycho-socio-cultural matters, finances,
85 and political issues which had a negative impact on life satisfaction among the students
86 [21]. Foreign-born students were found to face many difficulties in language and
87 communication, and they can experience isolation and discrimination [22]. These results
88 underline the importance of more research on the mental health of foreign students.
89 Although studies worldwide assessed the prevalence of SAD in different populations,
90 they did not give enough attention to population diversity, nor did they differentiate
91 between foreign-born and non-foreign-born students.

92

93 The population of international students pursuing university education in Türkiye is
94 steadily increasing. According to the latest update from the Council of Higher Education
95 (YÖK) in 2023, it is estimated that there are 260,316 international undergraduate students
96 in the country [23]. This indicates that Türkiye provides an excellent opportunity to reach
97 a respectable number of foreign students from different backgrounds. Previous studies
98 highlighted that SAD is already prevalent among Turkish university students [24,25].
99 Furthermore, emotional, academic, and financial difficulties were determined to be the
100 most prevalent concerns faced by Turkish students [26]. Another study conducted in
101 2015 showed that particularly international students in Türkiye commonly experience
102 psychosocial challenges such as being separated from their families, fear of making
103 mistakes, feeling disappointed in their choice to study in Türkiye, social pressure,
104 difficulty expressing thoughts, and struggling with criticism [27].

105 These various obstacles that international students face make them susceptible to mental
106 health disorders. However, to our knowledge, no study has looked into the prevalence of
107 SAD in this population. This study, therefore, aims to investigate the prevalence of SAD
108 among foreign-born undergraduate students in Turkish universities and its associated
109 factors.

110

111 **Methods**

112 **Study design**

113 This study employs a cross-sectional design, utilizing a self-administered questionnaire
114 distributed to foreign-born undergraduate students in Türkiye via Google Forms. To
115 reduce the possibility of non-response bias, we required that all questions be answered
116 before submission. The survey, designed to maximize participation and guarantee
117 anonymity, was conducted from September 17, 2023, to February 1, 2024. As there is no
118 formal list representing international students in Turkish universities, we tried to reach
119 international students from various universities through WhatsApp, Facebook, and
120 Telegram groups that connect international students. Administrators of these groups were
121 requested to send periodic reminders to enhance response rates. Additionally, we
122 approached associations concerned with international students' affairs to promote the
123 survey and send reminders about it.

124

125 **Sample size**

126 Based on the most recent data from the Council of Higher Education (YÖK), in 2023,
127 there were 260,316 non-Turkish undergraduate students in Türkiye [23]. Utilizing
128 EpiInfo 7.2.5.0, we determined the sample size required to represent this population.
129 Considering a population size of 260,316, with $p=0.21$ and $q=0.79$, along with a 95%
130 confidence interval and a 5% margin of error, we calculated a minimum required sample
131 size of 255 participants.

132

133 **Participants and survey**

134 The study included undergraduate students who are foreign-born non-Turkish, aged 18 or
135 older and currently enrolled in Turkish universities. The survey is structured into two
136 sections: the first part collects basic demographic and socioeconomic information,
137 presented in both Arabic and English. The second part utilizes the Social Phobia
138 Inventory (SPIN), a validated tool for evaluating symptoms of social anxiety and phobia.

139 The SPIN is administered in both its original English version and its validated Arabic
140 translation, ensuring an accurate assessment of social anxiety symptoms among
141 participants. The questionnaire was restricted to Arabic and English. This choice was
142 driven by the notable representation of Arabic-speaking students in Türkiye, with five of
143 the top 10 international student nationalities in Türkiye being Arabic-speaking [28].
144 Additionally, the decision was influenced by the assumption that many students seeking
145 education outside their home countries have sufficient English language skills to answer
146 the questionnaire.

147

148 SPIN, with its 17 questions, measures the frequency and intensity of social anxiety
149 symptoms over the preceding week. Responses are captured on a 5-point Likert scale
150 from 0 (not at all) to 4 (extremely), allowing for a comprehensive symptom severity score
151 ranging from 0 to 68. The scoring system classifies severity into five categories: none or
152 very mild (less than 20), mild (21-30), moderate (31-40), severe (41-50), and very severe
153 (51 or more), enabling detailed analysis of social anxiety levels among the surveyed
154 population.

155

156 **Quality of measure**

157 The Social Phobia Inventory (SPIN), originally developed in English by Connor et al.
158 (2000) [29], has undergone successful translation and validation into Arabic by Sfeir et
159 al. (2022) [30]. The Arabic version has demonstrated high reliability and validity for
160 evaluating social anxiety symptoms among Arabic-speaking populations. In two separate
161 ways, the Arabic SPIN exhibited Cronbach's alpha values of 0.92 and 0.90, indicating
162 strong internal consistency comparable to the original scale. Moreover, its convergent
163 validity and gender-invariant measurement of social phobia further confirm its efficacy.
164 Consequently, the Arabic SPIN emerges as a robust tool for detecting social phobia in
165 Arabic-speaking individuals.

166

167 **Statistical analysis**

168 The statistical analysis of this study encompasses descriptive, inferential analysis,
169 multiple regression, and logistic regression. Initially, descriptive statistics, including
170 frequencies and percentages, will outline the participant characteristics of
171 sociodemographic and academic data. The Shapiro-Wilk test assesses the normality of
172 the Social Phobia Inventory (SPIN) scores. For non-normally distributed data, the Mann-
173 Whitney U and Kruskal-Wallis H tests will examine differences in SPIN scores across
174 demographic variables. Associations between demographic factors and SPIN categories
175 (ranging from “None or Very Mild” to “Very Severe”) will be explored using the non-
176 parametric chi-square test.

177 Further, multiple regression analysis will predict SPIN scores (Ranging from “0” to
178 “68”) based on variables such as gender, smoking status, previous failures in university,
179 family history of mental health issues, exercise habits, and classmate relationships.
180 Additionally, a binomial logistic regression will evaluate the impact of various
181 demographic factors, including gender, monthly budget, smoking status, field of study,
182 GPA, family history, Turkish language proficiency, living arrangements, exercise habits,
183 and classmate relationships, on the presence of social phobia, dichotomized into
184 “Absent” (No or very mild SPIN) and “Present” (Mild to very severe SPIN). All
185 statistical analyses will be conducted using IBM’s SPSS Statistics package for Windows
186 (Version 25.0).

187

188 **Ethical approval**

189 Prior to commencing the cross-sectional study, the research team obtained ethical
190 approval from the Hamidiye Scientific Research Ethics Committee [31]. Participants
191 were briefed on the study's aims and objectives, and digital consent was obtained from
192 them before they completed the survey. Consent was indicated by participants selecting a
193 designated checkbox.

194

195

196 **Results**

197 **Participant characteristics**

198 We initially collected 506 responses but narrowed them down to 455 eligible participants
199 after excluding those who did not meet the undergraduate criterion (37 respondents), one
200 due to Turkish nationality, and 13 for incomplete surveys. The majority of participants
201 were Syrian students (59.6%); second to them were Egyptian students at 11.4%. Gender
202 distribution was nearly balanced (47.9% females, 52.1% males), and over half were aged
203 18–21 (52.1%). Most were single (94.1%), reported an average monthly budget (48.1%),
204 and were majoring in medical sciences (45.9%). The social classmate environment was
205 predominantly described as friendly and relaxed (56.3%). The significant majority
206 reported being non-smokers (85.7%) and abstaining from alcohol (97.6%). Living
207 arrangements varied, with 39.6% residing with their parents. **(Table 1).**

Table 1, Participants Characteristics		n (%)
Age	18-21	237(52.1)
	22-25	181(39.8)
	30-34	27(5.9)
	35 or above	10(2.2)
	Total	455(100)
Gender	Female	218(47.9)
	Male	237(52.1)
	Total	455(100)
Marital status	Single	428(94.1)
	Married	24(5.3)
	Widowed	1(0.2)
	Divorced	2(0.4)
	Total	455(100)
Monthly budget	Low	122(26.8)
	Average	219(48.1)
	Above average	49(10.8)
	Very low	61(13.4)
	High	4(0.9)
	Total	455(100)
Do you smoke?	No	390(85.7)
	Occasionally	33(7.3)
	Daily	32(7)

	Total	455(100)
Do you drink alcoholic drinks?	No	444(97.6)
	Occasionally	11(2.4)
	Daily	0
	Total	455(100)
What is the type of your university?	Government	359(78.9)
	Foundation / Private	96(21.1)
	Total	455(100)
Which subject are you currently studying?	Medical Field	209(45.9)
	Engineering	165(36.3)
	Social Sciences	72(15.8)
	Natural Sciences	8(1.8)
	Total	454(99.8)
	Missing data	1(0.2)
What is your current academic year in school?	Preschool	10(2.2)
	Year1	97(21.3)
	Year2	127(27.9)
	Year3	98(21.5)
	Year4	96(21.1)
	Year5	26(5.7)
	Year6	1(0.2)
	Total	455(100)
What is your most recent GPA	Less than 2.5	118(25.9)
	2.5-2.9	170(37.4)
	3.0-3.4	118(25.9)
	3.5-4.0	49(10.8)
	Total	455(100)
Have you previously failed a year in college?	No, I have not failed a year.	373(82)
	Yes, I have failed a year / years.	82(18)
	Total	455(100)
Do you have Family history of mental issues?	No, we don't.	380(83.5)
	Yes, we have.	75(16.5)
	Total	455(100)
Do you use psychiatric medication?	No, I don't.	415(91.2)
	Yes, I do.	40(8.8)
	Total	455(100)
What do you think is your level of Turkish language?	Poor	29(6.4)
	Fair	79(17.4)
	Good	175(38.5)
	Very good	172(37.8)
	Total	455(100)
At present whom do you live with?	Alone	43(9.5)
	With Parents	180(39.6)
	With housemates	90(19.8)
	With relatives	11(2.4)
	In a dormitory	131(28.8)
	Total	455(100)
Do you exercise?	No	144(31.6)
	Yes, occasionally.	190(41.8)
	Yes, less than 90 minutes per week.	27(5.9)
	Yes, more than 90 minutes per week.	94(20.7)
	Total	455(100)
How would you define the climate between you and your classmates?	Friendly and relaxed.	256(56.3)
	Competitive and stimulating	32(7.0)
	Competitive and hostile	29(6.4)

	I don't have an opinion yet.	124(27.3)
	I don't have friends.	13(2.9)
	Total	454(99.8)
	Missing	1(0.2)

208

209 Prevalence of social phobia

210 The study reveals the prevalence of social phobia among participants as follows: 39.1%
 211 exhibited “no or very mild symptoms,” 23.7% experienced “mild symptoms,” 21.3%
 212 faced “moderate symptoms,” while “Severe” to “very severe symptoms” were present in
 213 11.6% and 4.2% of participants, respectively. This distribution underscores the varying
 214 degrees of social anxiety’s impact on individuals (**Fig 1**).

215

216 Fig 1. Prevalence of SPIN categories

217 Factors associated with social anxiety disorder (SAD)

218 The inferential analysis yielded significant findings related to SAD. Gender showed
 219 statistical significance in both group difference and association tests ($p < 0.0001$). Family
 220 history of mental health issues was significant in both group difference tests ($p = 0.001$)
 221 and association tests ($p = 0.009$). Exercise habits were also significant in both tests ($p <$
 222 0.0001), as were the type of relationship with classmates ($p < 0.0001$). Smoking status
 223 demonstrated statistical significance only in the group difference test ($p = 0.028$), and
 224 prior academic failure was significant only in the association test ($p = 0.013$).
 225 Several factors did not show a statistically significant association with SAD, including
 226 age, marital status, monthly budget, alcohol consumption, type of university, subject of
 227 study, recent GPA, psychiatric medication usage, level of Turkish language proficiency,
 228 and living arrangements (**Table 2**).

229

Table 2, Factors associated with Social Anxiety Disorder (SAD).

	SPIN Score.	Group Difference Test	Association Test
	N (%)		
	455(100)	Mean (SD)	<i>P-value</i>
			<i>P-value</i>

Age	18-21	237(52.1)	26.3(14)	0.56	0.61
	22-25	181(39.8)	25.6(13.2)		
	30-34	27(5.9)	24.9(15.9)		
	35 or above	10(2.2)	20.4(15.14)		
	Total	455(100)			
Gender	Female	218(47.9)	28.89(14.27)	<0.0001*	<0.0001*
	Male	237(52.1)	22.96(12.82)		
	Total	455(100)			
Marital status	Single	428(94.1)	26(13.8)	0.651	0.575
	Married	24(5.3)	23.2(13.7)		
	Widowed	1(.2)			
	Divorced	2(.4)	20(28.3)		
	Total	455(100)			
Monthly budget	Low	122(26.8)	27.7(13.5)	0.646	0.799
	Average	219(48.1)	26.2(14)		
	Above average	49(10.8)	25.36(13.84)		
	Very low	61(13.4)	24.5 (13.58)		
	High	4(.9)	23(20.34)		
	Total	455(100)			
Do you smoke?	No Smoker	390(85.7)	26.25(13.77)	0.028*	0.461
	Smoker (Occasionally)	33(7.3)	19.48(14.25)		
	Smoker (Daily)	32(7)	26.84(13.07)		
	Total	455(100)			
Do you drink alcoholic drinks?	No	444(97.6)	26(13.8)	0.134	0.446
	Occasionally	11(2.4)	19(13.17)		
	Daily	0			
	Total	455(100)			
What is the type of your university?	Government	359(78.9)	25.5(14)	0.439	0.207
	Foundation / Private	96(21.1)	26.72(13.4)		
	Total	455(100)			
Which subject are you currently studying?	Medical Field	209(45.9)	26.7(14)	0.104	0.164
	Engineering	165(36.3)	26.34(13.3)		
	Social Sciences	72(15.8)	25.46(14.24)		
	Natural Sciences	8(1.8)	22.45(14.5)		
	Total	454(99.8)			
	Missing data	1(0.2)			
What is your current academic year in school?	Preschool	10(2.2)	24.9 (17)	0.594	0.119
	Year1	97(21.3)	25.3 (15)		
	Year2	127(27.9)	25.3(14)		
	Year3	98(21.5)	28.4(13.6)		
	Year4	96(21.1)	24.5(11.7)		
	Year5	26(5.7)	25.6(15.3)		
	Year6	1(0.2)			
	Total	455(100)			

What is your most recent GPA?	Less than 2.5	118(25.9)	26.8(13.9)	0.399	0.374
	2.5-2.9	170(37.4)	26 (13.3)		
	3.0-3.4	118(25.9)	25.5 (14.24)		
	3.5-4.0	49(10.8)	22.45 (14.5)		
	Total	455(100)			
Have you previously failed a year in college?	No, I have not failed a year.	373(82)	25.38(14.2)	0.103	0.013*
	Yes, I have failed a year/ years.	82(18)	27.7(12)		
	Total	455(100)			
Do you have Family history of mental issues?	No, we don't.	380(83.5)	24.8(13.5)	0.001*	0.009*
	Yes, we have.	75(16.5)	30.8(14.4)		
	Total	455(100)			
Do you use psychiatric medication?	No, I don't.	415(91.2)	25.5 (13.4)	0.213	0.124
	Yes, I do.	40(8.8)	28.8 (17.4)		
	Total	455(100)			
What do you think is your level of Turkish language?	Poor	29(6.4)	25.9(14)	0.0524	0.260
	Fair	79(17.4)	27.5(15.2)		
	Good	175(38.5)	27.4(13.4)		
	Very good	172(37.8)	23.4(13.4)		
	Total	455(100)			
At present who do you live with?	Alone	43(9.5)	24 (14.8)	0.881	0.909
	With Parents	180(39.6)	26.2 (13.4)		
	With housemates	90(19.8)	25.6 (13.7)		
	With relatives	11(2.4)	24.3 (13.7)		
	In a dormitory	131(28.8)	26.1 (14.4)		
	Total	455(100)			
Do you exercise?	No	144(31.6)	30.5(13.8)	<0.0001*	<0.0001*
	Yes, occasionally.	190(41.8)	25.1(13.3)		
	Yes, less than 90 minutes per week.	27(5.9)	24.74(12.9)		
	Yes, more than 90 minutes per week.	94(20.7)	20.4(13.1)		
	Total	455(100)			
How would you define the climate between you and your classmates?	Friendly and relaxed.	256(56.3)	23.2(13.32)	<0.0001*	<0.0001*
	Competitive and stimulating	32(7.0)	23.44(10.2)		
	Competitive and hostile	29(6.4)	29.2(15.92)		
	I don't have an opinion yet.	124(27.3)	30.7(13.9)		
	I don't have friends.	13(2.9)	29.8(12.5)		
	Total	454(99.8)			
Missing	1(0.2)				

*Indicates significance at p-value < 0.05.

232 Predictors of social anxiety disorder (SAD)

233 The multiple regression analysis aimed to predict Social Phobia Inventory (SPIN) scores
234 based on six independent variables: gender, smoking status, previous failure in university,
235 family history of mental health issues, exercise frequency, and type of relationships with
236 classmates. The model significantly predicted SPIN scores ($F(6, 447) = 13.165, p <$
237 0.001), explaining 15% of the variance with an adjusted R^2 of 14%. Gender, previous
238 failures, family history, exercise, and relationships with classmates were all significant
239 predictors, while smoking status was not. This indicates that gender, family history,
240 exercise, and social relationships play important roles in the severity of social anxiety
241 symptoms, whereas smoking does not have a statistically significant impact. (Table 3).

242 **Table 3, Predictors of Social anxiety disorder (SAD)**

SPIN-Score	B	95% CI for B		SE B	Beta	R2	P
Model		Lower	Upper			.139	
Constant	26.975	24.531	29.420	1.244			.000*
Gender	-3.773	-6.406	-1.141	1.339	-.136		.005*
Smoking	-.032	-2.306	2.241	1.157	-.001		.978
Previously Failed	3.067	-.050	6.185	1.586	.085		.054
Family History	4.912	1.704	8.120	1.632	.132		.003*
Exercise	-2.384	-3.534	-1.233	.585	-.187		.000*
Classmates	1.942	1.081	2.804	.438	.197		.000*

243 Table 3: The multiple regression model. B= un-standardized regression coefficient, CI= confidence interval, SE B=
244 Standard error of coefficient, Beta = Standardized coefficient, R2 = coefficient of determination

245 *Indicates significance at p-value < 0.05

246

247

248 Factors relating to the presence/absence of social anxiety

249 disorder (SAD)

250 **Table 4** shows the results of the predictors for SAD according to the logistic regression
251 model. The model shows females are found to have a higher probability of SAD
252 compared to males (OR = 1.976, 95% CI = 1.209 – 3.228, $p = 0.007$), individuals

253 engaging in more than 90 minutes of exercise per week are less likely to have SAD
 254 compared to those who do not exercise (OR = 0.383, 95% CI = 0.207 - 0.708, p = 0.002).
 255 Occasional smokers are shown to have a lower risk of SAD compared to non-smokers
 256 (OR = 0.422, 95% CI = 0.183 - 0.973, p = 0.043). Furthermore, individuals who are
 257 undecided about their relationships with classmates are at a higher risk of SAD compared
 258 to those who perceive their relationships as friendly and relaxed (OR = 2.298, 95% CI =
 259 1.367 - 3.865, p = 0.002).
 260 The model was highly significant (χ^2 (29) = 69.934, p < 0.001) with good fit (Hosmer-
 261 Lemeshow p = 0.800), explaining 19.4% (Nagelkerke R²) of variance in social phobia
 262 and accurately classifying 68.4% of cases.

263 **Table 4, Factors relating to the presence/absence of social anxiety disorder (SAD)**

Variable	OR	95% CI For B		P-value
		Lower	Upper	
Gender				
Male	1(ref)			
Female	1.976	1.209	3.228	.007*
Monthly budget				
Very low	1(ref)			.646
Low	.826	.396	1.722	.610
Average	.781	.390	1.562	.484
Above average	.583	.231	1.470	.253
High	.200	.017	2.401	.204
Do you smoke?				
No	1(ref)			.101
Occasionally	.422	.183	.973	.043*
Daily	1.188	.506	2.790	.692
Studying field				
Medical Field	1(ref)			.517
Engineering	1.435	.876	2.352	.152
Social Sciences	1.053	.566	1.959	.871
Natural Sciences	1.476	.269	8.092	.654
GPA score				
Less than 2.5	1(ref)			.532
2.5-2.9	1.008	.576	1.763	.979
3.0-3.4	.819	.453	1.478	.506
3.5-4.0	.623	.290	1.336	.224
Family history				
No, we don't.	1(ref)			
Yes, we have.	1.488	.803	2.757	.207
Psychiatric medication				
No	1(ref)			
Yes	1.019	.458	2.267	.964
Turkish language proficiency				
Poor	1(ref)			.399
Fair	1.518	.571	4.031	.403
Good	1.233	.501	3.035	.649
Very good	.920	.368	2.297	.858
Whom do you live with?				
Alone	1(ref)			.537
With Parents	1.454	.647	3.267	.365

With housemates	1.544	.655	3.639	.321
With relatives	.555	.123	2.497	.443
In a dormitory	1.525	.675	3.448	.311
Do you exercise?				
No	1(ref)			.013*
Yes, occasionally.	.535	.321	.891	.016*
Yes, less than 90 minutes per week.	.758	.282	2.039	.582
Yes, more than 90 minutes per week.	.383	.207	.708	.002*
Classmates' relationship				
Friendly and relaxed.	1(ref)			.009*
Competitive and stimulating	2.087	.898	4.853	.087
Competitive and hostile	2.338	.927	5.897	.072
I don't have an opinion yet.	2.298	1.367	3.865	.002*
I don't have friends	2.630	.643	10.755	.178

264 Table 4: Logistic regression. OR = Odds ratio, 95% CI For B = 95% Confidence Interval for the coefficient (B)
 265 *Indicates significance at p-value < 0.05

266 Discussion

267 As Türkiye continues to attract foreign students from around the globe to pursue higher
 268 education at its universities, a significant portion of this diverse group aims to engage
 269 with the Turkish community. The country's geographical location and its system of
 270 university entrance exams, along with its population diversity, make it a chosen
 271 destination for foreign students. This study offers insights into this population of
 272 undergraduate foreign students, assessing SAD among them and identifying several
 273 factors that may be associated with the presence of SAD. As far as we know, no study
 274 examined the incidence of SAD in this population. Given their immersion in a new
 275 cultural environment, they encounter challenges in adaptation and language acquisition
 276 that can affect their psychological and social health.

277
 278 Our study demonstrates that approximately 60.8% of students experienced symptoms of
 279 SAD. Among them, 4.2% experienced very severe symptoms, while about one-fifth
 280 (11.6%) suffered from severe symptoms. Additionally, about two-thirds of students had
 281 mild to moderate SAD symptoms (23.7% and 21.3%), respectively. In agreement with
 282 our result, a study reported by Wejdan Al-Johani et al., which included medical students
 283 in Saudi Arabia, found that the prevalence of SAD was almost 51%. Moreover, 8.21%
 284 and 4.21% reported severe and very severe SAD, respectively [14]. Another study in
 285 Saudi Arabia, conducted in 2020 by Al-Hazmi et al., had results that aligned with the
 286 outcomes of our study. Using the same SPIN questionnaire we used, they found that

287 among 504 medical students, 13.5% of the participants experienced severe to very severe
288 SAD symptoms [32].

289

290 Our study's current results indicate a higher prevalence of SAD compared to a previous
291 study conducted in 2020 by Yared Reta et al. in Ethiopia. Their study found that the
292 prevalence of SAD among 293 medicine and health science undergraduate students was
293 32.8% [33]. Another study that included 523 university students in Sweden investigated
294 the prevalence of SAD. The point prevalence of SAD among the participants was 16.1%
295 [34]. In the United Kingdom, Graham Russell and Steve Shaw conducted research on
296 students studying in higher education; the findings indicated that approximately 10% of
297 students reported marked to severe SAD [35].

298

299 However, comparing our findings with those of the mentioned studies is challenging due
300 to the diverse backgrounds of our participants. Our study focused on assessing the
301 prevalence of SAD among foreign students from various faculties, representing different
302 social factors and cultures. The study included participants from over 25 countries
303 encompassing a wide range of university branches, unlike previous studies that were
304 often limited to specific populations, such as medical students. Syrian students accounted
305 for the majority of responses (59.6%). This outcome was expected as it represents the
306 reality shaped by the civil war in Syria and its proximity to Türkiye, driving many Syrian
307 individuals, particularly students, to pursue education in Türkiye. According to the data
308 of the Council of Higher Education (YÖK) in Türkiye, in 2020, the total number of
309 Syrian refugee students reached 37,236 [36]. It is crucial to consider that Syrian refugees
310 in Türkiye experience symptoms of mental disorders such as anxiety, depression, and
311 PTSD [37].

312 Moreover, our study revealed a significant association between tense relationships with
313 classmates, the hostile social environment within the university, and the development of
314 SAD. This observation underscores the challenges students face in integrating into
315 Turkish society and forming supportive relationships to create an encouraging study
316 environment. This is similar to the findings of Malecha et al., who examined the literature
317 on foreign-born nursing students in the United States; their review reported that foreign-

318 born students face many difficulties in language and communication; barriers such as
319 isolation, discrimination, and cultural differences also affect the educational experience
320 [22]. According to a study by Rosellini and colleagues (2013), 78.8% of individuals with
321 SAD were able to identify the presence of an acute or chronic stressor when their SAD
322 first manifested; this finding aligns with our research, suggesting that exposure to acute
323 stressors, such as a stressful university environment, may influence the development of
324 SAD [38]. In cross-sectional studies, there is a consistent positive correlation between
325 social anxiety and SAD diagnosis with increased levels of relational victimization and
326 social exclusion. Additionally, these conditions are linked to having fewer friends and
327 experiencing challenges in establishing new friendships, lower peer acceptance, more
328 negative interactions within close friendships, as well as greater tendencies toward social
329 avoidance and withdrawal [39-42].

330

331 The prevalence of SAD was higher in female participants compared to their male
332 counterparts. This finding aligns with the DSM-5 statement that indicates that SAD is
333 more frequent in women [43]. Consistent patterns emerged in a comprehensive study
334 conducted by Dan J. Stein et al. wherein the analysis of data from 28 community surveys
335 within the World Mental Health Survey Initiative similarly underscored a heightened
336 prevalence of SAD among women [44]. Furthermore, other studies from diverse regions
337 across the globe consistently reveal comparable results [14,45-47]. In a 2023 study
338 conducted in Hong Kong, however, males showed higher rates of SAD than females; this
339 contradicts results from other studies and can be explained by the acknowledged
340 sampling size bias within the dataset, as the female sample was significantly larger than
341 the male sample [48]. We can understand the higher frequency of SAD in women most
342 effectively through a vulnerability-stress perspective. The notion revolves around the idea
343 that varying exposure to psychosocial stressors, combined with increased susceptibility to
344 anxiety in women, is put forward as a contributing factor to the identified sex differences
345 in anxiety disorders [49].

346

347 In addition, our research revealed a favorable correlation between engaging in regular
348 exercise and experiencing fewer symptoms of SAD. This result is in line with public

349 consensus and established scientific evidence supporting the positive influence of
350 exercise on overall mental health, as well as its role in preventing and treating mental
351 disorders [50-54]. The positive influence of exercise may be attributed to different
352 mechanisms. Firstly, exercise could function as a diversion from negative thoughts.
353 Additionally, repeated social interaction may play a role in this process. From a
354 physiological perspective, engaging in physical activity can cause a reduction in cortisol,
355 the stress hormone, potentially leading to an improvement in mood [55]. Also, studies
356 propose that exercise not only fosters the growth of new nerve cells but also releases
357 proteins like brain-derived neurotrophic factor, supporting the survival of nerve cells [56,
358 57].

359

360 Another significant finding was the association between the prevalence of SAD and a
361 family history of mental illness, with a one-half fold increase in the risk of SAD in people
362 with positive family history. This finding aligns with a study conducted in Sweden by
363 Maria Tillfors et al.; where they found that having an affected family member resulted in
364 a two- to three-fold increase in the risk of SAD [58]. Furthermore, heredity has been
365 identified as a predisposing factor for SAD, and genetic studies have provided strong
366 evidence that at least some genetic component contributes to the development of the
367 disorder [43, 59].

368

369 As shown in the results section, the occasional smokers had a mean SPIN score of 19.48,
370 which is considered to have non to very mild social anxiety. This unexpected finding of the
371 current study also is shown as a statistically significant inverse relationship between
372 occasional smoking and social anxiety rates seen in our logistic regression model, with an
373 odds ratio of 0.422. Several reasons could be suggested to explain this paradoxical
374 discovery. To begin, it is possible that smoking on occasion acts as a coping tool for people
375 who suffer from milder forms of social anxiety. In addition, the social situation in which
376 occasional smoking happens may have an impact on social anxiety levels. Previous studies
377 have revealed that smokers frequently smoke in social contexts to satisfy their desire for
378 greater social engagement. This is a frequent condition in smokers with low nicotine
379 dependence [60, 61]. Occasional smoking is frequently associated with social gatherings,

380 where people may feel more comfortable and at ease in the presence of others [62].
381 However, both daily smokers and non-smokers experience mild social anxiety symptoms,
382 with mean SPIN scores of 26.84 and 26.25, respectively. Yet, there is no statistically
383 significant relation in our logistic regression model. Additional studies on this topic must
384 be conducted to understand whether social anxiety is one of the reasons people start
385 smoking.

386

387 As an expected finding, failing a year in college was found to be associated with mild social
388 anxiety symptoms among international students. This finding highlights the specific
389 obstacles that international students may encounter when navigating academic
390 surroundings and transitioning to new social contexts, which leads to an increased
391 vulnerability to social anxiety symptoms following academic losses.

392

393 **Limitations**

394 Our study, being a cross-sectional study, may not have the ability to establish strong
395 conclusions regarding causation. Therefore, longitudinal studies with larger sample sizes
396 are needed to determine other possible factors associated with SAD. Attempting to contact
397 students through social media platforms restricts access to a subset of students,
398 necessitating a comprehensive approach that ensures outreach for most international
399 students. Additionally, the decision to design the questionnaire in Arabic and English
400 excludes students who are not proficient in either language. Furthermore, our study's
401 selection of explanatory variables was not all theoretically supported. Another limitation is
402 the stigma surrounding mental illness among the students, which may prevent individuals
403 from disclosing their condition.

404

405 **Conclusion and Recommendations**

406 Our study revealed a significantly high prevalence of SAD among foreign-born
407 undergraduate students in Türkiye. Those with noticeable symptoms of SAD tended to be

408 female and have a family history of mental illnesses. Additionally, studying in a stressed
409 atmosphere and experiencing failure were associated with a higher prevalence of SAD.
410 Interestingly, those who smoked occasionally exhibited better symptoms of SAD. These
411 findings necessitate more extra efforts in recognizing and treating SAD among this
412 population and minimizing the negative consequences of adjusting to new life
413 circumstances. This is crucial to protect the mental health of these students. Further research
414 could explore, for instance, the prevalence of SAD among non-foreign-born and foreign-
415 born students in Türkiye. Moreover, future studies could investigate the relationship
416 between SAD and various variables, including the potential mediating effects of other
417 factors on the experience of SAD.

418 **Acknowledgments**

419 We express our gratitude to all respondents who completed the questionnaire, as well as
420 to the administrators of the international students' online groups and the international
421 student associations, for their invaluable assistance in reaching our population.

422 **Supporting information**

423 **S1 Tool. Study questionnaire.**

424 (DOCX)

425 **References**

- 426
- 427 1. Penninx BW, Pine DS, Holmes EA, Reif A. Anxiety disorders. *Lancet* (London,
428 England). 2021;397(10277):914–927. doi:10.1016/S0140-6736(21)00359-7.
 - 429 2. GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional,
430 and national incidence, prevalence, and years lived with disability for 328 diseases and
431 injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of
432 Disease Study 2016. *Lancet* (London, England). 2017;390(10100):1211–1259.
433 doi:10.1016/S0140-6736(17)32154-2.
 - 434 3. Substance Abuse and Mental Health Services Administration. DSM-5 Changes:
435 Implications for Child Serious Emotional Disturbance [Internet]. Rockville (MD):
436 Substance Abuse and Mental Health Services Administration (US); 2016 Jun. PMID:
30199184.


- 437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
4. Stein MB, Stein DJ. Social anxiety disorder. *Lancet* (London, England). 2008;371(9618):1115–1125. doi:10.1016/S0140-6736(08)60488-2.
 5. Steinert C, Hofmann M, Leichsenring F, Kruse J. What do we know today about the prospective long-term course of social anxiety disorder? A systematic literature review. *Journal of anxiety disorders*. 2013;27(7):692–702. doi:10.1016/j.janxdis.2013.08.002.
 6. Cox BJ, Fleet C, Stein MB. Self-criticism and social phobia in the US national comorbidity survey. *Journal of affective disorders*. 2004;82(2):227–234. doi:10.1016/j.jad.2003.12.012.
 7. Koban L, Schneider R, Ashar YK, Andrews-Hanna JR, Landy L, Moscovitch DA, et al. Social anxiety is characterized by biased learning about performance and the self. *Emotion* (Washington, D.C.). 2017;17(8):1144–1155. doi:10.1037/emo0000296.
 8. Acarturk C, de Graaf R, van Straten A, Have MT, Cuijpers P. Social phobia and number of social fears, and their association with comorbidity, health-related quality of life and help seeking: a population-based study. *Social psychiatry and psychiatric epidemiology*. 2008;43(4):273–279. doi:10.1007/s00127-008-0309-1.
 9. Leigh E, Chiu K, Clark DM. Is concentration an indirect link between social anxiety and educational achievement in adolescents? *PloS one*. 2021;16(5):e0249952. doi:10.1371/journal.pone.0249952.
 10. Hajure M, Abdu Z. Social Phobia and Its Impact on Quality of Life Among Regular Undergraduate Students of Mettu University, Mettu, Ethiopia. *Adolesc Health Med Ther*. 2020 Jun 18;11:79-87. doi: 10.2147/AHMT.S254002.
 11. Desalegn GT, Getinet W, Tadie G. The prevalence and correlates of social phobia among undergraduate health science students in Gondar, Gondar Ethiopia. *BMC Res Notes*. 2019;12:438. doi:10.1186/s13104-019-4482-y
 12. Waghachavare VB, Quraishi SR, Dhumale GB, Gore AD. A Cross-sectional Study of Correlation of Body Image Anxiety with Social Phobia and Their Association with Depression in the Adolescents from a Rural Area of Sangli District in India. *Int J Prev Med*. 2014 Dec;5(12):1626-9.
 13. Alsibani F, Al Hashmi I, Al-Anqoodi R, Al-Badi M. Social Phobia in Undergraduate University Students in Oman. *Hunan Daxue Xuebao/Journal of Hunan University Natural Sciences*. 2022;49. doi:10.55463/issn.1674-2974.49.3.24.
 14. Al-Johani WM, AlShamlan NA, AlAmer NA, et al. Social anxiety disorder and its associated factors: a cross-sectional study among medical students, Saudi Arabia. *BMC Psychiatry*. 2022;22:505. doi:10.1186/s12888-022-04147-z
 15. Hutchesson MJ, Duncan MJ, Oftedal S, Ashton LM, Oldmeadow C, Kay-Lambkin F, Whatnall MC. Latent Class Analysis of Multiple Health Risk Behaviors among Australian University Students and Associations with Psychological Distress. *Nutrients*. 2021;13(2):425. doi:10.3390/nu13020425
 16. Yeh C, Inose M. International students reported English fluency, social support satisfaction, and social connectedness as predictors of acculturative stress. *Counseling Psychology Quarterly*. 2003;16:15–28. doi:10.1080/0951507031000114058.
 17. McCabe L. Mental health and study abroad: Responding to the concern. Washington: Association of International Educators; c2005 [cited Feb 27]. Available from:<https://www.nafsa.org/sites/default/files/ektron/files/underscore/InternationalEducator/EducationAbroadNovDec05.pdf>
 18. Mikulas J, Jitka S. Statistical analysis of study abroad experiences of international students in five major host countries of Europe. *Journal of International Students*. 2019;9:1–18. doi:10.32674/jis.v9i1.262
 19. Wang C, Mallinckrodt B. Acculturation, attachment, and psychosocial adjustment of Chinese/Taiwanese international students. *Journal of Counseling Psychology*. 2006;53:422–433. doi:10.1037/0022-0167.53.4.422
 20. Lucas J. Over-stressed, overwhelmed, and over here: Resident directors and the challenges of student mental health abroad. *Frontiers: The Interdisciplinary Journal of Study Abroad*. 2009;18:187–216.
 21. Pan JY, Wong DF, Joubert L, Chan CL. The protective function of meaning of life on life satisfaction among Chinese students in Australia and Hong Kong: a cross-cultural

- 493 comparative study. *Journal of American college health: J of ACH*. 2008;57(2):221–231.
494 doi:10.3200/JACH.57.2.221-232
- 495 22. Malecha A, Tart K, Junious DL. Foreign-born nursing students in the United States: a
496 literature review. *J Prof Nurs*. 2012 Sep-Oct;28(5):297-305.
497 doi:10.1016/j.profnurs.2012.03.001. PMID: 23006652.
- 498 23. Yükseköğretim Kurulu(YÖK). ÜNİVERSİTE İZLEME VE DEĞERLENDİRME
499 GENEL RAPORU – 2023. [cited 2023 September 11]. Available from:
500 <https://www.yok.gov.tr/yayinlar/yayinlarimiz>
- 501 24. Gültekin BK, Dereboy IF. The prevalence of social phobia, and its impact on quality of
502 life, academic achievement, and identity formation in university students. *Turk Psikiyatri*
503 *Derg*. 2011 Fall;22(3):150-8.
- 504 25. Izzıç F, Akyüz G, Doğan O, Kuğu N. Social phobia among university students and its
505 relation to self-esteem and body image. *Can J Psychiatry*. 2004 Sep;49(9):630-4. doi:
506 10.1177/070674370404900910.
- 507 26. Köksal B, Topkaya N. Üniversite öğrencilerinin yaşadığı sorunlar ve sorun alanlarının
508 depresyon, anksiyete ve stresle ilişkisi. *TEBD*. 2021;19(2):926-949.
509 doi:10.37217/tebd.933931.
- 510 27. Snoubar Y. Türkiye’de Öğrenim Gören Uluslararası Öğrencilerin Sorunları ve Sosyal
511 Hizmet Gereksinimleri. [Doctoral Thesis]. Hacettepe Üniversitesi Sosyal Bilimler
512 Enstitüsü; 2015.
- 513 28. udef.org.tr [Internet]. Federation of International Students Associations; [cited 2024 Feb
514 26]. Available from:<https://udef.org.tr/uluslararası-ogrencilik/>
- 515 29. Connor KM, Davidson JR, Churchill LE, Sherwood A, Foa E, Weisler RH:
516 Psychometric properties of the Social Phobia Inventory (SPIN). New self-rating scale.
517 *Br J Psychiatry* 2000, 176:379-386.
- 518 30. Sfeir M, Saliba G, Akel M, Hallit S, Obeid S: Association between perfectionism and
519 life satisfaction among a sample of the Lebanese population: The indirect role of social
520 phobia and validation of the Arabic version of the Social Phobia Inventory. *Perspect*
521 *Psychiatr Care* 2022, 58(4):2513-2523.
- 522 31. T.C. Sağlık Bilimleri Üniversitesi. (n.d.). Hamidiye Bilimsel Araştırmalar Etik Kurulu.
523 [Cited 2024, Feb 11] Available from: [https://www.sbu.edu.tr/tr/arastirma/etik-](https://www.sbu.edu.tr/tr/arastirma/etik-kurullar/hamidiye-bilimsel-arastirmalar-etik-kurulu)
524 [kurullar/hamidiye-bilimsel-arastirmalar-etik-kurulu](https://www.sbu.edu.tr/tr/arastirma/etik-kurullar/hamidiye-bilimsel-arastirmalar-etik-kurulu)
- 525 32. Al-Hazmi BH, Sabur SS, Al-Hazmi RH. Social anxiety disorder in medical students at
526 Taibah University, Saudi Arabia. *J Family Med Prim Care*. 2020;9(8):4329
- 527 33. Reta Y, Ayalew M, Yeneabat T, Bedaso A. Social Anxiety Disorder Among
528 Undergraduate Students of Hawassa University, College of Medicine and Health
529 Sciences, Ethiopia. *Neuropsychiatr Dis Treat*. 2020 Feb 26;16:571-577. doi:
530 10.2147/NDT.S235416. PMID: 32161462; PMCID: PMC7049747.
- 531 34. Tillfors M, Furmark T. Social phobia in Swedish university students: prevalence,
532 subgroups and avoidant behavior. *Soc Psychiatry Psychiatr Epidemiol*. 2007
533 Jan;42(1):79-86. doi: 10.1007/s00127-006-0143-2. Epub 2006 Dec 11. PMID: 17160591.
- 534 35. Russell G, Shaw S. A study to investigate the prevalence of social anxiety in a sample of
535 higher education students in the United Kingdom. *J Ment Health*. 2009;18(3):198–206.
536 doi:10.1080/09638230802522494
- 537 36. Acarturk C, McGrath M, Roberts B, et al. Prevalence and predictors of common mental
538 disorders among Syrian refugees in Istanbul, Turkey: a cross-sectional study. *Soc*
539 *Psychiatry Psychiatr Epidemiol*. 2021;56:475-484. doi:10.1007/s00127-020-01941-6.
- 540 37. Metin S, Ekinci F, Taş TA, Kutlu TR. The Living Conditions and Psychosocial
541 Circumstances of Syrian Refugees Who Migrated from Their Country After War: The
542 Example of Bursa City. *Karya Journal of Health Science*. 2022;3(2):140-144. doi:
543 10.52831/kjhs.1142534.
- 544 38. Rosellini A, Rutter L, Bourgeois M, Emmert-Aronson B, Brown T. The relevance of age
545 of onset to the psychopathology of social phobia. *Journal of Psychopathology and*
546 *Behavioural Assessment*. 2013;35:356–365.

- 547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
39. Beidel DC, Turner SM. Shy children, phobic adults: The nature and treatment of social anxiety disorder (2nd ed.). Washington, DC: American Psychological Association; 2007.
 40. Boulton M. Associations between adults' recalled childhood bullying victimization, current social anxiety, coping, and self-blame: Evidence for moderation and indirect effects. *Anxiety, Stress, & Coping*. 2013;26:270-292.
 41. Dempsey AG, Storch EA. Relational victimization: The association between recalled adolescent social experiences and emotional adjustment in early adulthood. *Psychology in the Schools*. 2008;45(4):310-322. doi:10.1002/pits.20298.
 42. Erath SA, Flanagan K, Bierman K. Social anxiety and peer relations in early adolescence: Behavioral and cognitive factors. *Journal of Abnormal Child Psychology*. 2007;35:405-416.
 43. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association, 2013.
 44. Stein DJ, Lim CCW, Roest AM, et al. The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative. *BMC Med*. 2017;15:143. doi:10.1186/s12916-017-0889-2.
 45. MacKenzie MB, Fowler KF. Social anxiety disorder in the Canadian population: Exploring gender differences in sociodemographic profile. *Journal of Anxiety Disorders*. 2013;27(4):427-434. doi:10.1016/j.janxdis.2013.05.006.
 46. Asher M, Aderka IM. Gender differences in social anxiety disorder. *Journal of Clinical Psychology*. 2018;74(10):1730-1741. doi:10.1002/jclp.22624.
 47. Kupper N, Denollet J. Social anxiety in the general population: introducing abbreviated versions of SIAS and SPS. *Journal of affective disorders*. 2012;136(1-2):90-98. doi:10.1016/j.jad.2011.08.014.
 48. Ganotice FA Jr, Shen X, Yuen JKY, Chow YMA, Wong AMY, Chan KMK, et al. Students' interaction anxiety and social phobia in interprofessional education in Hong Kong: mapping a new research direction. *Annals of medicine*. 2023;55(1):2210842. doi:10.1080/07853890.2023.2210842.
 49. Hallers-Haalboom ET, Maas J, Kunst LE, Bekker MH. The role of sex and gender in anxiety disorders: Being scared "like a girl"? *Handb Clin Neurol*. 2020;1(175):359-68.
 50. Aylett E, Small N, Bower P. Exercise in the treatment of clinical anxiety in general practice - a systematic review and meta-analysis. *BMC health services research*. 2018;18(1):559. doi:10.1186/s12913-018-3313-5.
 51. Silva LAD, Tortelli L, Motta J, Menguer L, Mariano S, Tasca G, et al. Effects of aquatic exercise on mental health, functional autonomy and oxidative stress in depressed elderly individuals: A randomized clinical trial. *Clinics (Sao Paulo, Brazil)*. 2019;74:e322. doi:10.6061/clinics/2019/e322.
 52. López-Torres Hidalgo J, DEP-EXERCISE Group. Effectiveness of physical exercise in the treatment of depression in older adults as an alternative to antidepressant drugs in primary care. *BMC psychiatry*. 2019;19(1):21. doi:10.1186/s12888-018-1982-6.
 53. Smith PJ, Merwin RM. The Role of Exercise in Management of Mental Health Disorders: An Integrative Review. *Annual review of medicine*. 2021;72:45-62. doi:10.1146/annurev-med-060619-022943.
 54. Gordon BR, McDowell CP, Lyons M, Herring MP. The Effects of Resistance Exercise Training on Anxiety: A Meta-Analysis and Meta-Regression Analysis of Randomized Controlled Trials. *Sports medicine (Auckland, N.Z.)*. 2017;47(12):2521-2532. doi:10.1007/s40279-017-0769-0.
 55. Duclos M, Gouarne C, Bonnemaïson D. Acute and chronic effects of exercise on tissue sensitivity to glucocorticoids. *Journal of applied physiology*. 2003;94(3):869-875.
 56. Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends in neurosciences*. 2002;25(6):295-301.
 57. Ernst C, Olson AK, Pintel JP, Lam RW, Christie BR. Antidepressant effects of exercise: evidence for an adult-neurogenesis hypothesis?. *Journal of Psychiatry and Neuroscience*. 2006;31(2):84-92.

- 602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
58. Tillfors M, Furmark T, Ekselius L, Fredrikson M. Social phobia and avoidant personality disorder as related to parental history of social anxiety: a general population study. *Behaviour Research and Therapy*. 2001;39(3):781-790.
 59. Sadock BJ, Sadock VA, Ruiz P. *Synopsis of Psychiatry: Behavioral Sciences, Clinical Psychiatry*. Wolters Kluwer; 2015. 2.
 60. Bondy SJ, Victor JC, Diemert LM, Mecredy GC, Chaiton M, Brown KS, Cohen JE, McDonald PW, Ferrence R, Garcia JM, et al. Transitions in smoking status over time in a population-based panel study of smokers. *Nicotine Tob Res*. 2013;15:1201-1210. doi: 10.1093/ntr/nts259.
 61. Lisha NE, Delucchi KL, Ling PM, Ramo DE. Prevalence and correlates of social smoking in young adults: Comparisons of behavioral and self-identified definitions. *Nicotine Tob Res*. 2015;17:1076-1084. doi: 10.1093/ntr/ntu242.
 62. Schane RE, Glantz SA, Ling PM. Nondaily and social smoking: An increasingly prevalent pattern. *Arch Intern Med*. 2009;169:1742-1744. doi: 10.1001/archinternmed.2009.315.

Distribution of Participants by Social Anxiety Disorder (SAD) Categories



■ none or very mild (less than 20) ■ Mild (21-30) ■ Moderate (31-40) ■ Severe (41-50) ■ Very severe (51 or more)

Fig 1. Prevalence of SPIN categories.