Getting seniors back on their bicycle; a pretest-posttest case-control study on the improvement of bicycle balance control

Eric Maris*, PhD

*Donders Institute for Brain, Cognition, and Behaviour, Radboud University, P.O. Box 9104 HE Nijmegen

Corresponding Author:

Eric Maris

e-mail: eric.maris@donders.ru.nl

tel: +31243612651

25   Abstract

26   Objective

27   Quantify the effectiveness and generalizability of an intervention that aims at
28   restoring bicycle balance control skills in seniors that have quit cycling.

30   Methods

The intervention was designed such that the difficulty of the bicycle balance
control task could be increased in a stepwise fashion, gradually approaching
the difficulty of bicycle balance control on the public roads. The intervention
lasted 11 weeks and involved three components: (1) training on an exercise
bicycle, (2) balance control training on a bicycle simulator, and (3) cycling on
the public roads with a safe start-and-stop technique that was practiced on
the bicycle simulator. The intervention was administered to a group of
community-dwelling senior citizens (N=23) that had quit cycling (N=19) or
that were on the verge of doing so (N=4). The effectiveness of the
intervention was evaluated by comparing balance control skill and
confidence between a post- and a pre-intervention measurement. As a check
for possible spontaneous recovery, the same comparison was also performed
in a matched control group that did not participate in the intervention.

45   Results

46   This intervention produced a very large improvement (Cohen's $d = 1.8$,
47    $t(16)=7.5$, $p<.001$) in balance control skills and confidence on the public
48   roads.

50   Conclusion

Bicycle balance control skills and confidence can be restored over a short
period of time. However, this does not rule out the importance of the slower
process of acquiring a sufficient lower body strength.

Impact Statement

Increasing balance control skills and confidence is a realistic short-term
target for rehabilitation interventions.

58
59
60

62   <u>Keywords</u>
63   *Balance control, bicycle balance, motor skill training, bicycle simulator,*
64   *senior citizens, elderly, mobility, intervention study, case-control.*
65
66
67
68

With age, there is a sharp decrease in the percentage of citizens that

continues to ride their bicycle. This is well documented in the Netherlands,

where everyone learns to cycle at a young age, and cycling is the default

means of transportation for short distances. In the Netherlands, the

percentage of inhabitants that cycles at least several times per month

fluctuates around 77% between age 30 and 70 [1]. However, over an average

period of 15 years, this percentage drops to 39% [1]. Thus, a substantial

fraction of the senior citizens stops cycling before the end of their life, which

is on average 16 years after one has reached 70 years [2].

The physical health benefits of cycling are well documented [3,4]. For instance,

there is a strong inverse relationship between commuter cycling and all-

cause mortality [4]. Because pedaling force can be easily controlled (e.g., by

cycling uphill), cycling is an excellent training stimulus for the improvement

of lower body strength and power. In addition, it has a very low impact on the

joints, and it offers increased mobility options that are necessary for a

meaningful social life.

In this paper, I describe, motivate, and evaluate a rehabilitation intervention

focused on balance control (BC) skills that aims at getting senior citizens

back on their bicycle. Bicycle BC pertains to two centers of gravity (CoGs)

that both must be kept above their respective area of support (AoS): (1) the

CoG of the rider-bicycle combination must be kept above the line that

connects the two wheel-road contact points, and (2) the CoG of the rider's

upper body must be kept above the saddle [5]. To achieve these two objectives,

steering (turning the handlebars) is the most important control action [5,6].

BC can be understood from the perspective of optimal feedback control

(OFC) [5,7-13], which is the dominant theory in the neural control of movement

(see Fig. 1). OFC distinguishes between two determinants of BC skill: (1) the

sensory-motor control algorithms implemented in the neuronal networks of

our central nervous system (CNS), and (2) strength, power and joint range-

of-motion as implemented in the muscles, tendons, and joints (the human

part of the mechanical system; see Fig. 1). Our nervous system also has a

somatic nervous system (SNS) which sends out motor commands and

receives sensory feedback (see Fig. 1).

Fig. 1: A closed-loop feedback control model for bicycle balance. The model

has four components: (1) a sensory system that represents the sensory

division of the somatic nervous system (SNS), consisting of the eyes, the

inner ear (labyrinth, vestibular organ), the proprioceptive sensory organs

(muscle spindles and Golgi tendon organs), and their afferent axons, (2) a

computational system that represents the central NS (CNS), (3) a motor

output system that represents the sensory division of the SNS, consisting of

the efferent axons of the spinal motor neurons, and (4) a mechanical system

that represents the non-neuronal part of the human body (extrafusal muscle

fibers, skeleton, …) and the bicycle.

OFC is consistent with the fact that the effects of different types of balance

training are specific to the balance task that is trained [14,15]. In fact, OFC

postulates that, separately for every balance task, the CNS optimizes BC

under the constraints/limitations of the mechanical and the SNS (e.g.,

precision of the sensory feedback, torque and range limits of the mechanical

system, motor noise); there is no set of component BC skills that is assembled

in different ways according to different sets of task requirements. For a

bicycle BC training, this task-specificity would imply that a training effect

would not generalize to cycling on the public roads.

For a lost motor skill that must be restored, it is important to know to what

extent it depends on (1) plasticity in the CNS (responsible for the mapping of

sensory feedback onto control actions), and (2) plasticity in the organs that

are responsible for registering feedback and executing actions. Plasticity in

the CNS depends on the signaling properties of neurons which can change

over a much shorter timescale (hours, days) than the organs that are

responsible for sensory registration and motor execution [16-18]. To the best of

my knowledge, plasticity in the sensory organs has not yet been

demonstrated, and plasticity in the muscles with functional consequences

(strength/power increase) operates at a timescale of at least weeks.

I designed a bicycle BC training with a double objective: (1) realizing training

effects that generalize to bicycle BC on the public roads, (2) dissociating

central (CNS-dependent) and muscular contributions to a possible

improvement in BC skill. To realize the first objective, the difficulty of the

bicycle BC task was increased in a stepwise fashion, gradually approaching

the difficulty of bicycle BC on the public roads. And to realize the second

objective, the training included sessions on exercise (spinning) bicycles on

which the functional threshold power (FTP) could be monitored.

To realize a stepwise increase in the difficulty of the bicycle BC task, it is

useful to consider BC skill as a continuum defined by BC challenges, which

are perturbations that distort the balance. An important BC challenge are the

upward reaction forces that result from pedaling, deform the upper body and

perturb the steering. The impact of a BC challenge is modulated by the

cycling speed, which is reflected in the fact that it is much easier to balance a

bicycle when we are up to speed, as compared to when we are close to a

standstill. The modulating role of speed follows from the mechanical

properties of the bicycle, which are responsible for self-stabilizing forces that

increase with speed [6].

In the bicycle BC training, I make use of the fact that the BC challenge level

increases with pedaling force and decreases with speed. To control the BC

challenge level, I have developed a bicycle simulator that requires the same

type of BC as on the public roads (see Methods). On this simulator, also a safe

start-and-stop technique was practiced which allowed the rider to continue

with his BC exercises while cycling safely outdoors.

Study setup/Methods

Participants

Participants were recruited via an article in a local newspaper (De

Gelderlander) about the goal of the study (March 31, 2023). The article

invited participants that had stopped cycling at least six months ago, and

wanted to get back on their bicycle. A total of 105 candidates replied, which

were all invited for an information session. Based on their responses to a

questionnaire, I excluded candidates that still cycled occasionally or suffered

from balance-relevant sensory disorders (peripheral neuropathy,

labyrinthitis, vestibular neuritis, ...). I then selected the 15 participants that

lived the closest to the Radboud University campus, the location of the

intervention.

All subjects gave their informed consent for inclusion before they

participated in the study. The study was conducted in accordance with the

Declaration of Helsinki, and the protocol was approved by the Ethics

Committee of the faculty of Social Sciences (ECSS) of the Radboud University

(ECSS-2020-132).

The intervention started with cycling on a stationary exercise bicycle

(spinning) in the second week of June 2023. This part of the intervention was

in the Radboud University Sports Center. On June 15, a second newspaper

article was published in which the location of the intervention was revealed.

After this publication, 8 additional candidates spontaneously joined the

spinning sessions, and requested to participate in the study, to which I

agreed. Four of these additional participants still cycled on rare occasions but

considered quitting because of fear of falling.

A total of 23 participants started the BC training on the bicycle simulator, and

17 participants completed it (mean age 75,3 years, age range 66-88 years).

From the 6 participants that dropped out, 2 did so for a reason that was

related to the BC training: one participant's legs were so short that she could

not touch the ground while sitting on the saddle, and one participant lacked a

sufficient force to keep herself balanced on top of a stationary bicycle. Four

participants dropped out for reasons that were unrelated to the BC training:

a fracture due to a fall at home (two participants), bursitis, and an urgent eye

surgery.

The 17 participants that completed the BC training were matched to 17

candidates (the control group) that were not selected for the study because

they were living farther away from the location of the intervention as

compared to the experimental group. The matching criteria were gender and

age.

Bicycle simulator

The bicycle simulator [19] is shown in Fig. 1. The core elements of the simulator

are three rollers with a width equal to the width of an average Dutch bicycle

path (1.75 m). The two front rollers are covered with a conveyer belt that is

supported from underneath by a low-friction platform (not visible on Fig. 1).

The bicycle's front wheel rests on the conveyer belt and the rear wheel rests

on the rear roller. The rollers' rotational inertia is close to the linear inertia of

the average Dutch rider-bicycle combination, and this ensures that the

acceleration response to a pedal stroke feels natural. A torque motor is

attached to the axis of the rear roller, and this motor is essential to assist

weak participants in the first stages of the training. Finally, the bicycle

simulator has a virtual-reality (VR) component that projects an animation of

a bicycle lane, which is controlled by the cycling speed and the cyclist's lateral

position on the left-right axis of the simulator (as measured using a motion

capture system).

226
227 **Fig. 2: Participant riding on the bicycle simulator.**
228
229 In between the front and the rear rollers, there is a platform on which the
230 rider can place his feet and the trainers can stand. The rider has one trainer
231 on every side, supporting him by means of gentle pushes on his shoulders. As
232 the rider's BC skills improve, the trainers increase their distance from the
233 rider.
234
235 The rider wears a safety harness with which he is connected to a support
236 construction above the rider's head (see Fig. 2). The safety harness does not
237 prevent the rider from falling, but in case he does, his knees will not touch the
238 floor before the suspension becomes active.
239

Intervention

The intervention was over a period of 11 weeks. In that period, the
participants engaged in (1) spinning, (2) BC training on the bicycle simulator,
and (3) cycling on the public roads with a safe start-and-stop technique that
was practiced on the bicycle simulator. The three components of the
intervention were also administered in this order, with spinning and BC
training overlapping in time for the last 8 weeks.

Spinning

The first three weeks of the intervention only involved a group spinning

training, which was given by the first author and two professional spinning

coaches. Participants trained at an individually determined power level,

controlled by the one-minute functional threshold power (1minFTP).

One spinning session lasted 45 min., excluding warming-up and cooling-

down, but including a 5 min. break in the middle. Every session started with a

3-minute block in which the 1minFTP was assessed. The remainder of the

session had the structure of an interval training. The high intensity blocks

lasted at most 1 min. and were alternated with low intensity blocks at 70

percent of 1minFTP that lasted at most 2,5 min. Increasing intensity was

achieved either by increasing the resistance (and keeping the RPM fixed) or

by increasing the RPM (and keeping the resistance fixed).

Assessing the 1minFTP

The 1minFTP was used for a double purpose: (1) as a measure of leg power,

to investigate its relationship with bicycle BC skill, and (2) to train at an

individually determined power level. The 1minFTP is the highest power a

participant can maintain for one minute while cycling at 60 revolutions per

minute (RPM). The 1minFTP was assessed by having participants monitor

their breath after a one-minute effort at some power level: if the participant

could say only a few words at a time (a proxy for reaching the ventilatory

threshold), the 1minFTP was reached.

Assessing the 1minFTP required keeping a constant high-power output, and

this turned out to be very difficult for our participants. First, assessing the

1minFTP required that the participants could operate the bicycle computer

that was used to provide feedback about their power output. It took

participants two weeks to learn to operate this computer. Second, it took

several participants substantially more time to learn to produce a constant

279 power output; initially, all participants lowered their RPM when the

280 resistance was increased above a certain level, thereby preventing a higher

281 power output. Third and last, participants' performance was highly variable

282 with respect to the maximum power output they endured, and this was both

283 within sessions (no longer following the trainer's instructions with respect to

284 power output) and between sessions (lowering the 1minFTP at the

285 beginning if one was feeling tired).

287 *Bicycle simulator BC training*

288 BC training started in the fourth week of the intervention and was in one of

289 the research labs of the Radboud University (Faculty of Social Sciences). The

290 BC training was individual and was given by the first author and two

291 research assistants. A BC training session lasted approximately 30 min. At the

292 start, participants put on a safety harness and were connected to the support

293 construction. The saddle height was adjusted such that the participant's feet

294 touched the platform in between the front and the rear rollers. When the

295 participant showed signs of fear, the trainers gently pushed their hands

296 against the participant's shoulders and gradually released the pressure as the

297 participant gained confidence.

299 The guiding principle in the BC training was increasing the difficulty in small

300 steps and following the participant's progressing skill and confidence. Every

301 session started with a BC challenge level at which the participant was

302 comfortable, and that level was slowly increased if the participant agreed. In

303 the first session, the bicycle was brought up to a speed of approximately 12

304 km./h. by means of a torque motor. In this condition, the participant's only

305 task was turning the handlebars for balance control. The BC challenge level

306 was increased in four ways: (1) by decreasing the speed, (2) by decreasing

307 the assistive force at the rear roller, (3) by asking participants to accelerate

308 from standstill and to stop at any desired moment, and (4) by asking the

309 participant to tap the support surface with both feet, place them back on the

310 pedals, and to continue pedaling (feet tapping exercise).

Cycling outdoors

The participant was instructed to start cycling outdoors on his own bicycle

when he could do 10 feet taps in a row without stopping in between, and at a

resistance level slightly above the resistance on the public roads. The

participant was instructed to use the safe start-and-stop technique (feet

touching the road surface) that was practiced on the bicycle simulator, to

start cycling in an environment without other road users (a park, a parking

lot), and to gradually move to more busy environments.

Causal effect estimation in a pretest-posttest design

A randomized experiment is often considered the only possible way to obtain

an unbiased estimate of a causal effect. However, according to statistical

theory this is not the only way [20]. In fact, if spontaneous recovery can be

ruled out (also denoted as the assumption of temporal stability [20]), a pretest-

posttest design also allows for such an unbiased estimate. The argument

involves that if, in the population of interest, the average of some outcome

variable (e.g., bicycle BC skill) does not change spontaneously, then the

average post- minus pre-intervention change in this outcome variable is an

unbiased estimate of the treatment's causal effect. For bicycle BC skill in our

population of interest (senior citizens), it is unlikely that this skill

spontaneously returns (i.e., without a deliberate intervention) after it has

been lost. Of course, numerous other outcome variables do change

spontaneously on average (body temperature in a population that has the flu,

feelings of depression in a population that visits a psychotherapist, walking

BC skills in healthy young adults that had bed rest for a week, …). To remove

all doubts about a possible spontaneous recovery, I included a control group

that did not receive the intervention, but in which the change in the outcome

variable was assessed over the same time interval as in the experimental

group (see Participants).

The effect of the intervention was then evaluated statistically using a

dependent samples t-test that compared the change in the outcome variable

between the matched experimental and the control group participants.

Outcome variable

The outcome variable was obtained from a questionnaire that evaluated

participants' bicycle BC skills and confidence while riding on the public roads.

This questionnaire involved three multiple-choice questions with three

answer options each. The introduction to the questionnaire was the

following:

Three environments will be described, and you must indicate whether

you rode your bicycle in that environment in the past two weeks. By

"riding your bicycle", we mean starting from a standstill and stopping at a

desired location. For every environment, you must indicate what applies

to you: (1) I did not ride my bicycle here, (2) I rode my bicycle here, but I

try to avoid it, and (3) I rode my bicycle here without problems. The three

environments are the following: (1) without traffic (e.g., a park, a parking

lot), (2) with little traffic (e.g., small roads in an outside area, a bicycle

lane on a quiet moment), and (3) in busy traffic (e.g., in the city center

during traffic hours, around a busy marketplace).

The answers to every question/environment were converted into a score by

assigning 0, 1, and 2 points to the respective options. Summing these scores

over the three questions/environments results in a number between 0 and 6,

which is the outcome variable.

In the experimental group, this questionnaire was administered twice, the

first time in the beginning of the intervention and the second time six weeks

after the end of the intervention. The questionnaire was sent as an

attachment to an email, and participants were called on their phone if they

had not replied after one week. For some participants, it was necessary to

clarify some aspects of the questions. In the control group, the questionnaire

was administered only once and at around the same time that the second

questionnaire was sent to the experimental group. None of the control group

participants was riding their bicycle at the start of the intervention and they

were asked whether and how this had changed in the meantime.

378   Results

I conducted a pretest-posttest case-control study to evaluate the

effectiveness of an intervention aimed at restoring bicycle BC skill. The

experimental group consisted of 23 participants, of which 17 completed the

intervention (see Methods).

The intervention lasted 11 weeks and involved three components: (1) cycling

on a stationary exercise bicycle (spinning), (2) BC training on a bicycle

simulator, and (3) cycling on the public roads with a safe start-and-stop

technique that was practiced on the bicycle simulator (see Methods). The

effectiveness of the intervention was evaluated in terms of the participant's

skill/confidence on his own bicycle when cycling outdoors (see Methods). In

the experimental group, this outcome measure was assessed twice, first in

the beginning of the intervention, and then six weeks after the intervention.

To correct for a possible spontaneous recovery of cycling skill/confidence, I

used a matched control group that did not receive the intervention, but in

which the change in cycling skill/confidence over the same time interval was

assessed (see Methods).

The improvement in cycling skill/confidence was significantly larger in the

experimental group (M=3.1, SD=2.1) compared to a matched control group

(M=0, SD=0), t(16)=6.0, p<.001, 95% CI [2.0 4.1]. Note that none of the

control group participants had a spontaneous recovery of their cycling

skill/confidence. The raw outcome measures of the experimental group are

shown in Fig. 3. The effect size (Cohen's d) was 1.5, which is between a very

large and a huge effect [21]. This effect was obtained with an average of 4.8

simulator BC training sessions of 30 min. each.

Fig. 3: Raw outcome measures of the experimental group (a small random jitter was added to see overlapping lines).

It was my original plan to dissociate central (CNS-dependent) and muscular

contributions to a possible improvement in BC skill. For that, it was

necessary to quantify the increase in maximum power output on the spinning

bicycle, for which I used the one-minute functional threshold power

(1minFTP), which was assessed at the beginning of every spinning session.

However, keeping a constant high-power output (a requirement for a reliable

FTP assessment) turned out to be very difficult for our participants and

required several weeks of training (see Methods). I conclude that I failed in

obtaining a reliable quantification of the increase in maximum leg power

increase in this group of participants.

However, it is useful to report the mean 1minFTP on which participants

converged after learning to produce a constant force output. This mean

1minFTP was 93.6 Watt (sd=17.2). FTP is usually expressed relative to

bodyweight, and in this participant group the average 1 minute power-to-

weight ratio was 1.2 Watt/kg (sd=0.3). For reference, it is useful to compare

this value with published 1 minute power-to-weight ratios for untrained

amateur cyclists [22] (defined as the 5-th percentile of the population of

amateur cyclists): 3.9 for males and 3.3 for females. Thus, the participants in

this study had an average 1 minute power-to-weight ratio that is

approximately three times smaller than that of an untrained cyclist.

Discussion and conclusions

A very large improvement in senior cyclists' BC skills and confidence can be

realized by means of a short intervention (11 weeks), even if that person has

quit cycling for a long time. This is remarkable given the task-specificity of

the effects of different types of balance training [14,15], and is most likely due to

the high similarity between the BC training on the simulator and cycling

outdoors.

Limitations

This study did not expose participants to some important BC challenges (e.g.,

curbstone, pothole, sharp turn) and it is unknown whether they can deal with

them successfully. It is very likely that several BC challenges require a

sufficient lower body strength to prevent perturbations from affecting the

steering movements. For example, (1) standing on the pedals while covering

an obstacle, and (2) core muscle co-contraction in a sharp turn at high speed.

Cycling safely not only involves BC (staying upright) but also navigation

(avoiding obstacles and/or following a track), which requires additional

skills such as selective attention, anticipation, and planning. These have not

been investigated.

From the perspective of OFC, it is tempting to conclude that the improvement

is mainly due to the fast plasticity in the CNS, and not to the comparatively

slow plasticity in the muscles. However, to make a strong case for this claim,

it is essential to have a reliable measure of maximum power output, and this

turned out to be very difficult. The existing literature on the relation between

(1) muscle strength/power, and (2) the incidence of falls and balance task

performance is inconsistent, with one systematic review (focusing on falls) [23]

arguing for a strong relation and another (focusing on balance task

performance) [24] for a very weak one. This is probably related to the fact that,

in the population of interest, falls cannot be predicted from the performance

on balance tasks [25].

For several reasons, the relation between muscle strength/power, balance

control, and falls, is a complicated one. First, many studies have been

conducted in homogeneous groups (institutionalized participants, small age

range, …) and this restriction-of-range results in an underestimation of the

correlation in the heterogeneous/unselected population. Second, most

people are aware of the range of BC challenges that their BC skill allows, and

they try to stay within this range. This restriction-of-range at the individual

level reduces the correlation between BC skills and the incidence of falls.

Third, there may be large differences between muscle groups and their

associated joint ranges-of-motion with respect to their relevance for BC. In

most studies, only leg strength/power was measured, but the core muscles in

the trunk and hip may very well be more relevant for BC. Fourth, BC not only

depends on the muscles for bringing the AoS under the CoG (the muscles'

motor role) but also for informing the CNS about the position of CoG relative

to the AoS, which depends on the muscle spindles (the muscles' sensory

feedback role) [13,26]. To be useful, muscle spindle output must reflect the

tension in the force-producing fibers, but its precise contribution to the loss

of BC is unknown.

Conclusion

I have demonstrated that it is possible to realize a very large improvement in

a senior cyclist's BC skills and confidence over a short period (11 weeks).

This is most likely due to the high similarity between the BC training and

cycling outdoors, but it does not rule out the importance of the slower

process of building (and loosing) muscular strength/power.

Author's contributions

All authors contributed equally to the manuscript and read and approved the

final version of the manuscript.

Acknowledgments

The author would like to thank Maartje Floris and Loes van Leeuwen for their

assistance in the training and the data collection.

Funding

This research was funded by the Radboud Center Social Sciences. The

authors report no involvement in the research by the sponsor that could have

influenced the outcome of this work.

Conflict of Interest

None declared.

References

1.   Research facts and numbers 'Cycling for everyone' [Onderzoek feiten en

cijfers 'Fietsen voor Iedereen'] (Mobycon, commissioned by the Ministery of

Infrastructure and Water Management.) (2022).

2.   VZinfo.nl. RIVM. Accessed 25 January, 2024.

https://www.vzinfo.nl/levensverwachting/leeftijd-en-geslacht#Levensverwachting

3.   De Hartog JJ, Boogaard H, Nijland H, Hoek G. Do the health benefits of

cycling outweigh the risks? *Environmental health perspectives.*

2010;118(8):1109-1116.

4.   Oja P, Titze S, Bauman A, et al. Health benefits of cycling: a systematic

review. *Scandinavian journal of medicine & science in sports.* 2011;21(4):496-

509.

5.   Maris E. A bicycle can be balanced by stochastic optimal feedback control

but only with accurate speed estimates. *PLoS one.* 2023;18(2):e0278961.

6.   Meijaard JP, Papadopoulos JM, Ruina A, Schwab AL. Linearized

dynamics equations for the balance and steer of a bicycle: a benchmark and

review. *Proceedings of the Royal Society of London A: Mathematical, Physical*

and Engineering Sciences. 2007;463(2084):1955-1982.

7.   Åström KJ. *Introduction to stochastic control theory.* Courier Corporation;

2012.

8.   Todorov E. Optimality principles in sensorimotor control. *Nature*

neuroscience. 2004;7(9):907-915.

9.   Todorov E, Jordan MI. Optimal feedback control as a theory of motor

coordination. *Nature neuroscience.* 2002;5(11):1226-1235.

10.   Wolpert DM. Computational approaches to motor control. *Trends in*

cognitive sciences. 1997;1(6):209-216.

11.   Scott SH. Optimal feedback control and the neural basis of volitional

motor control. *Nature Reviews Neuroscience.* 2004;5(7):532-545.

12.   Wolpert DM, Ghahramani Z, Jordan MI. An internal model for

sensorimotor integration. *Science.* 1995;269(5232):1880-1882.

13.   Maris E. An internal sensory model allows for balance control based on

non-actionable proprioceptive feedback. *arXiv.*

2024;doi:10.48550/arXiv.2403.00951

14.   Giboin L-S, Gruber M, Kramer A. Task-specificity of balance training.

Human movement science. 2015;44:22-31.

15.   Kümmel J, Kramer A, Giboin L-S, Gruber M. Specificity of balance

training in healthy individuals: a systematic review and meta-analysis. *Sports*

Medicine. 2016;46:1261-1271.

16.   Debanne D, Inglebert Y, Russier M. Plasticity of intrinsic neuronal

excitability. *Current opinion in neurobiology*. 2019;54:73-82.

17.   Magee JC, Grienberger C. Synaptic plasticity forms and functions. *Annual*

review of neuroscience. 2020;43:95-117.

18.   Mozzachiodi R, Byrne JH. More than synaptic plasticity: role of

nonsynaptic plasticity in learning and memory. *Trends in neurosciences*.

2010;33(1):17-26.

19.   Maris E, inventor; Bicycle simulator [Patent]. The Netherlands patent

application PCT/NL2022/050304. 2022.

20.   Holland PW. Statistics and causal inference. *Journal of the American*

statistical Association. 1986;81(396):945-960.

21.   Sawilowsky SS. New effect size rules of thumb. *Journal of modern applied*

statistical methods. 2009;8(2):26.

22.   Johnstone D. How does your cycling power output compare? Accessed 29-

2, 2024. https://www.cyclinganalytics.com/blog/2018/06/how-does-your-cycling-

power-output-compare

23.   Moreland JD, Richardson JA, Goldsmith CH, Clase CM. Muscle weakness

and falls in older adults: a systematic review and meta-analysis. *Journal of the*

American Geriatrics Society. 2004;52(7):1121-1129.

24.   Muehlbauer T, Gollhofer A, Granacher U. Associations between measures

of balance and lower-extremity muscle strength/power in healthy individuals

across the lifespan: a systematic review and meta-analysis. *Sports medicine*.

2015;45:1671-1692.

25.   Boulgarides LK, McGinty SM, Willett JA, Barnes CW. Use of clinical and

impairment-based tests to predict falls by community-dwelling older adults.

Physical therapy. 2003;83(4):328-339.

26.   Proske U, Gandevia SC. The proprioceptive senses: their roles in signaling

body shape, body position and movement, and muscle force. *Physiological*

reviews. 2012;