A program for real-time surveillance of SARS-CoV-2 genetics

Hayden N. Brochu[1], Kuncheng Song[1], Qimin Zhang[1], Qiandong Zeng[1], Adib Shafi[1], Matthew Robinson[1], Jake Humphrey[1], Bobbi Croy[2], Lydia Peavy[3], Minoli Perera[3], Scott Parker[3], John Pruitt[3], Jason Munroe[4], Rama Ghatti[5], Thomas J. Urban[3], Ayla B. Harris[3], David Alfego[1], Brian Norvell[3], Jonathan D. Williams[3], Deborah Boles[3], Melinda B. Nye[6], Suzanne E. Dale[6], Michael Sapeta[6], Christos J. Petropoulos[7], Jonathan Meltzer[6], Marcia Eisenberg[3], Oren Cohen[8], Stanley Letovsky[1], & Lakshmanan K. Iyer[1],*

[1]Labcorp Center for Excellence in Data Science, AI and Bioinformatics, Burlington, NC 27215, USA; [2]Labcorp Information Technology, Burlington, NC 27215, USA; [3]Labcorp Research and Development, Burlington, NC 27215, USA; [4]Labcorp Consumer Genetics Department, Burlington, NC 27215, USA; [5]Labcorp-Sequenom, San Diego, CA 92121, USA; [6]Labcorp Center for Esoteric Testing, Burlington, NC 27215, USA; [7]Labcorp-Monogram Biosciences, South San Francisco, CA 94080, USA; [8]Fortrea Inc., Durham, NC 27703, USA; *Corresponding author

Please direct all correspondence to Lakshmanan K. Iyer (iyerl@labcorp.com)

A program for real-time surveillance of SARS-CoV-2 genetics

Abstract

The COVID-19 pandemic brought forth an urgent need for widespread genomic surveillance for rapid detection and monitoring of emerging SARS-CoV-2 variants. It necessitated design, development, and deployment of a nationwide infrastructure designed for sequestration, consolidation, and characterization of patient samples that disseminates de-identified information to public authorities in tight turnaround times. Here, we describe our development of such an infrastructure, which sequenced 594,832 high coverage SARS-CoV-2 genomes from isolates we collected in the U.S. from March 13th 2020 to July 3rd 2023. Our sequencing protocol ('Virseq') generates mutation-resistant sequencing of the entire SARS-CoV-2 genome, capturing all major lineages. We also characterize 379 clinically relevant SARS-CoV-2 multi-strain co-infections and ensure robust detection of emerging lineages via simulation. The modular infrastructure, sequencing, and analysis capabilities we describe support the U.S. Centers for Disease Control national surveillance program and serve as a model for rapid response to emerging pandemics at a national scale.

Introduction

The rapid emergence of COVID-19 and looming burden on global healthcare systems warranted swift responses from the international community. The causal virus, SARS-CoV-2, was first identified by metagenomic RNA sequencing[1] as well as Sanger- and PCR-based detection methods[2,3]. Very early in the pandemic response, we prioritized the development of SARS-CoV-2 diagnostic assays to meet the demand for detection methods, offering one of the first PCR-based tests and performing as many as 275,000 tests daily[4]. This immense scale of PCR testing enabled

us to assess the dynamics of COVID-19 infection as it pertains to PCR positivity[5] and also provide

population-based analysis on the maintenance of antibody titers[6,7].

Similar to other betacoronaviruses, the SARS-CoV-2 genome mutated as it infected and

spread across the population, with a mutation rate of approximately 1-2 x 10^{-6} mutations per

nucleotide per replication cycle[8]. Such genetic changes are known to impact the severity and

transmissibility of infection as well as vaccine efficacy[9], thus requiring close to real-time

surveillance using next generation sequencing (NGS)-based methods to inform public health

policies. Multiple whole genome sequencing approaches have been applied to support this need,

namely the ARTIC SARS-CoV-2 amplicon-based protocol for whole genome sequencing[10], direct

RNA sequencing[11], and sequence hybridization[12]. Shotgun metagenomic sequencing of

wastewater has also been an effective surveillance strategy for approximating variant abundances[13]

and identifying so-called 'cryptic lineages'[14,15], as it became apparent that COVID-19 patients

exhibit fecal viral shedding[16,17].

Ultimately, a greater need for high-throughput, real-time genomic sequencing of SARS-

CoV-2 emerged in the United States through the SARS-CoV-2 Sequencing for Public Health

Emergency Response, Epidemiology and Surveillance (SPHERES) spearheaded by the Centers

for Disease Control (CDC). To address this public health need and in collaboration with CDC

SPHERES, we rapidly developed a national surveillance apparatus using our purposely designed

infrastructure for flexible sampling, a unique approach for SARS-CoV-2 whole genome

sequencing, and tailored analytical methodologies that ensure continuously robust SARS-CoV-2

lineage determination using the Phylogenetic Assignment of Named Global Outbreak (PANGO)

nomenclature[18]. Our assay, which we call 'Virseq', is distinguished from other NGS-based SARS-

CoV-2 whole genome sequencing approaches through its use of probe-based tiling[19] and long

reads, which provide versatile, mutation-resistant capabilities. As of July 3rd, 2023, we have

sequenced 594,832 genomes (10X median depth of coverage) and have provided 524,498 high-

quality SARS-CoV-2 genomes (10X median depth of coverage, >90% genome coverage, complete

S gene coverage) and patient demographic data to the CDC using our Virseq assay, representing

continuous snapshots of SARS-CoV-2 viral evolution.

In this study, we provide a retrospective analysis of our Virseq assay using our vast

repertoire of high-quality genomes, showcasing our surveillance capabilities and the modular

resources that support its continued use. We show that Virseq has generated uninterrupted

surveillance reflecting the nationwide prevalence of SARS-CoV-2 consistent with our RT-PCR

sample collection, and we further demonstrate the robustness of SARS-CoV-2 lineage

determination through our in-house analytical capabilities. We also address the analytical

challenges posed by rare SARS-CoV-2 co-infections by developing a custom workflow that yields

haplotype-resolved consensus genomes. Finally, we pose this suite of resources as a model for

mounting rapid and robust large-scale surveillance networks.

Results

An infrastructure for nationwide COVID-19 surveillance

The rapid emergence and continuous evolution of SARS-CoV-2 necessitated setting up a national

surveillance program that could provide real-time epidemiological snapshots across the United

States. In response to the CDC's basal surveillance program, we organized and implemented a

nationwide infrastructure to identify SARS-CoV-2 positive patient samples across the United

States, consolidate these samples and their associated demographic data, and sequence their

genomes to determine their SARS-CoV-2 PANGO lineages. Our surveillance system also includes

mechanisms for monitoring emerging lineages and their potential impacts on qPCR and

sequencing performance. Further, this setup is flexible and modular, enabling rapid responses and

developments to our pathogen surveillance.

A schematic of this modular infrastructure and its associated sequencing protocols is shown

in **Figure 1**. In **Figure 1a**, we show a high-level view of our surveillance pipeline that begins with

sample accessioning and resulting in whole genome sequencing reports with de-identified sample

metadata. Upper respiratory samples (nasopharyngeal (NP) or nasal swabs) collected in 0.9%

saline or viral transport media were collected and analyzed through our COVID-19 RT-PCR Test

at various Labcorp® service centers and laboratories, and the resulting PCR extraction plates

containing SARS-CoV-2-positive samples were then shipped to Labcorp® central locations and

consolidated into plates with high viral titer samples (i.e. N1 Ct < 31) using our custom developed

plate selector app (**Figure 1b**, Methods). The selection of plates using this app is critical, as it de-

identifies data thereby ensuring HIPPA compliance and allows flexibility in choosing samples. For

example, we can focus on certain geographic regions where outbreaks are underway, put

limitations on the amount of data from a state to ensure uniform surveillance, and threshold on the

PCR amplification N1 Ct value to ensure sufficient genetic material is available for sequencing in

each sample. The resulting consolidated plates of high viral titer samples were then shipped to our

sequencing labs where samples were processed through custom tiled Molecular Loop® probe

amplification, followed by library preparation and sequencing on the PacBio® Sequel II™ platform

in a highly multiplexed fashion (**Figure 1c-d**). PacBio® raw data was then processed to generate

Circular Consensus Sequencing (CCS) reads which were then analyzed using our custom

bioinformatics workflow to generate consensus genomes for each sample (**Figure 1d**). Stringent

sequence coverage quality control was then applied followed by PANGO lineage determination

for each sample (Methods), and results were then merged with patient demographic data and de-

identified with a new custom ID generated for each sample (**Figure 1d**). Consensus genomes,

summary reports, raw CCS reads, and alignment variant calls were then provided to CDC who in

turn processed our submission and deposited the relevant data to public repositories, namely

GISAID[20] and NCBI[21].

High-throughput, high-fidelity SARS-CoV-2 genome surveillance pipeline

The rapid pathogen evolution during a pandemic and the possibility of sporadic outbreaks

necessitates a highly robust genomic surveillance pipeline. From January 2021 to July 2023, we

have used our Virseq pipeline (**Figure 1**) to report 524,498 high-quality SARS-CoV-2 genomes

(10X median depth of coverage, >90% genome coverage, complete S gene coverage) and patient

demographic data to the CDC. These sequences captured all major lineages that have emerged

throughout the COVID-19 pandemic since the inception of this surveillance effort, including

Alpha, Delta, Omicron, and the many Omicron subvariants (**Figure 2**, Methods). When overlaying

the positivity rate of our diagnostic PCR assays used for sample picking, we observed multiple

fluctuations matching variant emerges, such as BA.1, BA.4/BA.5, and XBB.1.5 (**Figure 2b**).

Stratifying genomes by U.S. Health and Human Services (HHS) region (HHS regions 1-10;

Methods), we found that the HHS regions 1 and 2 (corresponding to the U.S. Northeast) often

served as a harbinger to predict variant emergence for all other regions (**Figure S1**). For example,

the initial Omicron variant (BA.1) and the more recent XBB.1.5 variant reached ~50% prevalence

in HHS regions 1 and 2 approximately one week and four weeks prior to all other regions,

respectively (**Figures S1 and S2**).

After the BA.1 wave, reports indicated that SARS-CoV-2 infections by Omicron variants

exhibited lower viral loads[22,23], prompting us to investigate the diagnostic PCR N1 Ct values of

our samples, which are a useful proxy for viral load. As expected, lower sample Ct values were correlated with both increased average depth of coverage and higher consensus genome coverage (**Figure S3a**). We also observed shifts in sample diagnostic N1 Ct values throughout the pandemic that ranged from 20-21 in 2021 and exceeded 24 during the BA.1 wave in the 2021-2022 winter season, prior to reaching a steady state between 22-23. (**Figure S3b**). In our later analysis of SARS-CoV-2 co-infections, we found that co-infected sample N1 Ct values followed the same trends as those of samples where only a single SARS-CoV-2 lineage was detected. Together, these results indicate that our surveillance captures the overall kinetics and patterns of circulating variants.

This collection of high-quality genomes also uniquely captured demographic trends from all 50 states in the U.S. and the District of Columbia. States with the highest representation include California (n>62,000) and New Jersey (n>48,000), while other key states from HHS regions 9 and 10 (WA, n >19,000; AZ, n>17,000), HHS region 5 (IL, n>23,000; OH, n>10,000), and HHS regions 4 and 6 (NC, n>45,000; FL, n>36,000; TX, n>14,000) also had strong sampling (**Figure 3a**, **Table S1**). We also observed that the number of Virseq-generated genomes has represented a consistent proportion of total SARS-CoV-2-positive samples collected in each HHS region with minor fluctuations (**Figure 3b**). These fluctuations are expected as the timing of variant outbreaks (e.g. BA.1) can vary across HHS regions along with concomitant surges in PCR positivity rate (**Figure 2b**). Despite these changes, we still observed sustained census normalized sampling across both our diagnostic PCR assays and Virseq (**Figure S4a**) even with an end to the COVID-19 emergency response and the reduction in our surveillance volume in 2023 (**Figure S4b**). Notably, we observed a reduction in Virseq surveillance in HHS regions 6-8 at the onset of the BA.2 wave, as BA.2 prevalence spiked elsewhere in the U.S. before affecting these regions (February 2022; **Figures 2b, 3b**, **S1, S2, and S4**).

We also found that the age distribution of samples collected in each geographic region

shifted throughout the course of the pandemic, with a plurality from pediatrics in 2021, shifting to

a more even distribution of ages in 2022, and finally shifting to a plurality from older segments of

the population in 2023 (**Figures S5 and S6**). Interestingly, there was also a modest trend in patient-

reported gender, as the proportion of samples from female patients appeared to increase from 2021

to 2022 and once again in 2023 (**Figures S5 and S6**), despite there not being any difference in

PCR positivity between males and females (**Figure S7a**). We also observed a bifurcation in PCR

positivity across age groups in March 2022, as pediatric PCR positivity lowered to approximately

half of the positivity in most other age groups and this relative difference has not changed since

(**Figure S7b**). Intriguingly, two months prior (January 2022), the CDC and FDA announced

multiple expansions of pediatric COVID-19 vaccination availability[24–26] (**Figure S7b**). We also

observed that in June 2022, 18–19-year-olds had a reduction in PCR positivity relative to older

age groups as well, and most recently in June 2023 we observed an increase in geriatric (80+ year

olds) PCR positivity (**Figure S7b**). These demographic- and region-specific trends in variant

prevalences and PCR positivity rates require a surveillance apparatus like ours that is flexible and

robust to the rapid evolution of the SARS-CoV-2 genome.

Robust, mutation-resistant S gene sequencing using a probe-based long-read strategy

To maintain nationwide surveillance of pathogen genome evolution, the selected whole-genome

sequencing approach must be able to withstand sudden changes in genetic diversity. Our

surveillance apparatus uniquely employs a probe-based long-read sequencing approach that is

mutation-resistant by design due to its ~22X genome tiling of >99% of the SARS-CoV-2 genome,

except for a few hundred base pairs of the 5'- and 3' peripheral genomic regions (**Figure 1c**). At

the end of 2021, the Omicron variant (BA.1) emerged and swept through the U.S. in a matter of

weeks (**Figures 2**, **S1, and S2**), dramatically shifting the diversity of the circulating lineages across

the U.S. population (**Figure 4a**). Unlike earlier variants like Delta which were predominantly

mutated in the ORF1a genic region, the original Omicron variant (BA.1) introduced a surge of

novel S gene mutations (27 SNPs and three deletions compared to Delta) (**Figure 4b**), raising

concern regarding the ability of PCR- and amplicon-based assays to detect BA.1. In fact, the Spike

gene target failure (SGTF) genomic signature was so common that it became a useful proxy for

the PCR detection of emerging variants, such as Alpha and Omicron[27]. While interruptions in our

surveillance were not observed (**Figure 2**), we verified the fidelity of our sequencing of BA.1*

using an *in silico* approach to check for probe dropout caused by lineage defining mutations

(Methods). We found that our assay retained ~20X *in-silico* tiling of the S gene during the initial

Omicron wave (**Figure 4c**), and in the worst-case scenario where we allowed zero SNP tolerance

in probe binding regions, we retained a minimum of ~10X probe tiling (**Figure S8**). Furthermore,

as the total number of unique mutations and the concurrent prevalence of multiple circulating

lineages measured in terms of their entropy continues to increase with more recent XBB

subvariants, we continue to observe profoundly stable genome-wide probe tiling *in silico* (**Figure**

4c). This robust probe tiling is especially important as chronic infections and widespread

vaccination have altered the evolutionary trajectory of the SARS-CoV-2 genome[28] and the receptor

binding domain of the S gene remains under considerable selective pressure in the Omicron era[8].

We also confirmed the robustness of our sequencing strategy by analyzing the genome-

wide per-base coverage of Variants of Concern (VOCs) that have emerged throughout the

pandemic (**Figure S9**, starting with B.1.1.7 or 'Alpha' up until XBB.1.5). We found that overall

per-base coverage has remained stable and well above our minimum per-base coverage required

for base calling in our consensus genomes, despite the heavily mutated S gene of Omicron and

subvariants thereof (**Figure S9**). Together, these results show that the Virseq assay is a stable and

effective sequencing strategy and is a critical component of our surveillance apparatus. However,

while we are confident in our response to variant outbreaks thus far, it is imperative that proactive

measures are taken to preclude future surveillance interruptions.

Modeling the performance of the Virseq assay by simulation

One challenge of sustaining continuous whole genome surveillance is the need to predict changes

in sequencing performance that may occur and its effect on the characterization of the pathogen

variants. Many SARS-CoV-2 lineages emerge in regions outside of our surveillance network (i.e.

in countries other than U.S.) and may be too rare for detection once they initially spread to our

surveilled regions. To address this challenge, we developed a Virseq performance simulator that

models our entire production process from raw reads to PANGO lineage determination, capturing

the sequencing and other systematic errors that might propagate into consensus genomes. This

simulator was constructed using a representative batch of samples and incorporates the per-base

coverage and minor allele fractions commonly observed at each genomic position (Methods,

Figure S10). Application of this simulator begins with an input sequence that is then mutated to

reflect any errors introduced by our production process, which can then be compared with the

original sequence via concordance analysis of their PANGO lineage determinations.

We routinely use this simulator to monitor newly designated lineages in the PANGO

nomenclature (largely VOCs) as well as randomly selected sequences from previous months of

surveillance, leveraging the GISAID database[20]. This routine monitoring mechanism is a crucial

component of our FDA EUA and could be used to help maintain future pandemic surveillance

networks. In this study, we expanded this analysis to include up to 100 sequences each from 1,899

VOCs (97,421 total sequences, 'VOC experiment') and descendant lineages thereof, current and

former, and 10,000 randomly selected sequences from each month spanning from January 2021

until July 2023 (310,000 total sequences, 'retrospective experiment') (Methods). As expected, we

observed similar coverage profiles between the two experiments and the simulator model,

indicating that the simulated genomes accurately reflected Virseq-generated sequences (**Figure**

S10b). When assessing the PANGO lineage concordance between the simulated and original

genomes, we first checked for exact matches then also checked for parent/child relationships

between the lineages compared (e.g. BA.5 is a parent of the child BA.5.1), deeming these parent

matches (Methods).

Overall, we observed strong concordance in both the retrospective (99.55% exact, 99.97%

parent) and VOC experiments (99.06% exact, 99.84% parent) (**Figure 5**). In the retrospective

experiment we observed strong concordance across all 31 months analyzed with some month-to-

month fluctuations (>98.95% exact, >99.9% parent) (**Figure 5a**). We also observed that some

fluctuations coincided with shifts in circulating lineage diversity and the timing of VOC

emergences (**Figure 4**, **Figure 5a**). Intriguingly, while some VOC emergences resulted in slight

reductions of concordance (BA.1, BA.4/BA.5), others counterintuitively coincided with improved

concordance (BA.2, BQ, XBB.1.9/XBB.1.16) (**Figure 4**, **Figure 5a**).

When assessing the concordance of individual VOCs, we found that 98.21% (1,865) and

99.63% (1,892) of VOCs had >90% exact and parent lineage concordance, respectively (**Figure**

5b). In total, we observed seven VOCs with a small number of discordant calls, and four of these

(BA.2.2.1, BA.5.10, BQ.1.19, and BY.1) had UShER tree placement conflicts with their designated

hashes, i.e. these sequences were representative of those lineages but could not be placed

accordingly in the UShER tree (**Table 1**). This indicated that the original sequences of these VOCs

had unstable PANGO lineage designations. For example, one of four simulated BA.2.2.1

sequences was called a BA.1, and we later found that the original sequence was hashed as BA.2.2.1

but placed in the UShER tree as BA.1 (**Table 1**). Inspection of the other three VOCs revealed that

they were recombinants with discordant calls corresponding to one of the recombined lineages

(**Table 1**), indicating that the simulated genomes had a loss of resolution.

We then investigated features of the simulated genomes, including key drivers of

discordant lineage calls. As expected, we found that genome coverage was significantly lower

among genomes that had discordant lineage calls or that were only parent concordant compared to

those with exact concordance in both experiments ($p<0.001$ in all comparisons, Wilcoxon rank-

sum test, **Figure S11a-b**). Both experiments yielded similar genomic positional dependence of

consensus genome errors (**Figure S11c**), and these errors were often found in regions modeled

with poor coverage (5'/3' peripheral genomic regions) instead of positions modeled with higher

base calling errors (**Figure S11d-e**). These simulation experiments collectively show that the

Virseq assay generates consensus genomes with accurate PANGO lineage designations and that

accuracy is predominantly driven by genomic coverage, which is expected behavior for the

pangolin software in general[29]. This Virseq simulator is a crucial component of our surveillance

apparatus, ensuring that we anticipate potential sequencing interruptions and serving as a model

for viral sequencing simulators in general.

Detection and haplotype phasing of SARS-CoV-2 mixtures

The concurrent circulation of multiple lineages of the same virus during a pandemic may result in

co-infections of different viral lineages, which in some cases result in more severe clinical

outcomes in COVID-19 patients[30]. Thus, another requirement for effective surveillance machinery

is the ability to distinguish and characterize co-infections, which may complicate consensus

genome generation and PANGO lineage determination. After an initial finding of within-host

SARS-CoV-2 diversity[31], reports emerged describing patients likely co-infected with co-

circulating SARS-CoV-2 lineages[32–37]. Since our SARS-CoV-2 whole genome sequencing dataset

robustly captures these pandemic-wide trends in circulating lineages (**Figure 2**) and is highly

stable (**Figure 4**, **Figure 5**), we posited that recovery of multiple SARS-CoV-2 lineage detections

from the same sample would be possible.

To identify these potential mixtures (i.e. co-infections), we developed a custom workflow

utilizing freyja[14], an off-the-shelf mixture deconvolution algorithm (Methods). We found that

deeply sequenced samples (20.7% or 123,373 of 594,832) produced stable lineage mixture results

(>99% genome coverage, 100% S gene coverage, and average depth of coverage > 200X) (**Figure

S12a**). We then imposed three criteria for a sample to be classified as a mixture. The first two

require the most and second most abundant lineages to have relative abundances no greater than

0.8 and no less than 0.2, respectively, which only 571 (0.46% of 123,373) samples satisfied (**Figure

S12b-c**). Thirdly, we required the mixed lineages to differ by at least three defining SNPs

(heretofore defined as 'discriminating SNPs'), since such mixtures were found to have stronger

concordance between lineage relative abundances and discriminating SNP allele fractions **Figure

S12d-f**). This process yielded a final confident set of 379 mixtures (**Supplementary Data**) likely

of similar quality to their non-mixture counterparts, as they were found to have similar median

depth of coverage (mixtures: 338.4, non-mixtures: 335.6, p=0.46, Wilcoxon rank-sum test)

(**Figure S12g**). These mixtures had similar Ct values as non-mixtures, with the same differences

observed between the BA.1 wave samples and those before and afterwards (**Figure S3c**). We also

found that these mixtures had lineage compositions concordant with the original lineage called by

pangolin[29], since in most cases the pangolin-derived lineage either closely matched the majority

mixture lineage (217 or 57.3%) or was a parent of both majority and minority mixture lineages

(142 or 37.5%) (**Figure S13**).

Not surprisingly, we detected a plurality of mixtures during the BA.1 wave (159 or 42%),

which represents the largest portion of the dataset analyzed (18% or 22,169 samples). We also

observed the highest prevalence of mixtures during the BA.1 wave, peaking at 1.4% the last week

of December 2021 (**Figure 6a**). When categorizing the lineages comprised by these mixtures using

their lineage groups, we found that most mixtures were of lineages from the same group (347 or

91.6%), e.g. BA.1* mixed with BA.1* (**Figure 6b**, **Table S2**). This is likely due to variants

emerging in blocks (**Figure 2**), though we did observe some mixtures of different lineage groups

collected during the transitions between these blocks, e.g. one Delta*-Mu* and three BA.1*-Delta*

mixtures (**Figure 6b**, **Table S2**).

Since these mixtures have robust sequencing depth (**Figure S12g**) and comprise lineages

with as many as 77 discriminating SNPs (**Supplementary Data**), we hypothesized that it would

be feasible to resolve lineage haplotypes. Using a standard haplotype phasing tool for long reads

(Methods), we were able to produce haplotype blocks in most mixtures (277 or 73.1%), favoring

mixtures harboring discriminating SNPs greater in number and closer together (**Figure 6c**). We

also developed a custom approach that employs a greedy strategy to merge haplotype blocks

together based on the alleles of the discriminating SNPs in each haplotype block (Methods). This

greatly increased haplotype block resolution, on average increasing the number of SNPs in the

largest (merged) haplotype block by ~100% and increasing the size of haplotype blocks to as long

as 15.8kbp (**Figure 6d-e**). These exceptionally large haplotype blocks were found among the

BA.1*/Delta* mixtures, which have the largest number of discriminating SNPs among the

mixtures identified (**Supplementary Data**). One example is LC0471172, which is a mixture of

BA.1.1.18 and AY.39 that had a final merged haplotype block of length 15.8kbp harboring 61 SNPs

and spanning most of ORFs 1a and 1b as well as the entire S gene and 3' end of the genome (**Figure**

S14). These rare, finely resolved mixture haplotypes are evidence that combining haplotype

reconstruction with mixture analysis has potential to unveil unique sample characteristics. This

mixture analysis workflow thus provides our surveillance apparatus with the essential ability to

detect co-infections and ensures that consensus genomes are correctly reported in such cases.

Discussion

Genomic surveillance, globally and through our contribution to CDC SPHERES, proved to be

critical for monitoring the emergence of highly mutated SARS-CoV-2 variants and their potential

influence on disease severity[38] and the hundreds of vaccine development efforts worldwide (183

in clinical development as of March 30, 2023[39]). In this report we showcased our robust,

comprehensive U.S.-based SARS-CoV-2 surveillance network enabled through our infrastructure

and sequencing capabilities. Our probe-based tiling of the genome precluded surveillance

interruptions, while other amplicon-based assays have required multiple updates[10]. However,

while our tiled approach has ensured robust lineage detection so far, there is always a possibility

that an emerging novel lineage may introduce mutations that could potentially affect our ability to

detect it. To address this uncertainty we routinely assess emerging SARS-CoV-2 lineages using

our Virseq assay simulator before they are widely circulating among the U.S. population. These

resources ensure that we are prepared to swiftly respond to any sudden and/or large mutations in

the SARS-CoV-2 genome.

One key feature of our surveillance network is the rapid sequestration and consolidation of

high viral titer samples before sequencing. This strategy could in theory be applied to any

infectious agent for which we or others have robust diagnostic assays. In the case of SARS-CoV-

2 in this study we apply an N1 Ct value upper limit for sample inclusion to guarantee sufficient

genetic material for whole genome sequencing. While necessary for sequencing feasibility, this

thresholding may introduce bias in the sample selection and comprehensive lineage coverage.

Another important aspect of our surveillance is the dense network of various Labcorp® testing

centers throughout the U.S. In this study we that show our SARS-CoV-2 diagnostic samples come

from all HHS regions and are sent for whole genome sequencing through our Virseq assay with

limited bias. Importantly, this comprehensive demographic coverage is contingent on the

availability of samples from our testing centers, which could potentially change due to myriad

factors such as local mandates and/or health care coverage.

Our targeted long-read sequencing approach is also equipped for recovery of haplotype-

resolved viral genomes. To our knowledge, our study is the first to provide a pandemic-wide, high-

resolution evaluation of co-infections at this scale, though there have been other systematic efforts

that are smaller[40] or target specific types of co-infections, e.g. Omicron/Delta[37]. A crucial step in

confirming co-infections is the haplotype phasing of observed heterozygous mutations, which is

generally limited to long-read sequencing approaches described here with PacBio® sequencing and

by others using Oxford Nanopore Technologies®[41]. In our study we observe most co-infections

from the same lineage group (e.g. BA.1*/BA.1*), likely representing individuals who were

exposed to unique variants in rapid succession rather than those who are chronically ill from a

previous infection. These co-infections may not only have clinical relevance[30], but also represent

potential recombination events. For example, the BA.1 wave showed the highest prevalence of co-

infections in our dataset and incidentally introduced numerous recombinants, including those

formed from Delta and BA.1[42–44]. Ultimately, the use of long reads is uniquely suited for

distinguishing co-infections from these recombinant cases as well as other sources of intra-host

variation of the virus that have been described[31].

The surveillance apparatus we describe is not only robust to the pandemic undulations but

is also flexible and modular. For example, we recently adapted this workflow to employ an Oxford

Nanopore Technologies®-based ClearLabs® sequencing approach in lieu of our probe-based long-

read (PacBio®-based) sequencing approach. This alternative EUA approved pipeline enables rapid

turnaround time (~1 day) and retains the same suite of analytical tools as our primary surveillance

apparatus. We have also leveraged our infrastructure to provide additional features beyond

monitoring SARS-CoV-2 genetic evolution. Early in the COVID-19 pandemic it became clear that

convalescent sera from COVID-19 survivors would be essential for development of antibody

therapies[45]. We and others analyzed sera specimens collected from over 3,000 unvaccinated

individuals and found that most did not exceed antibody concentrations associated with 90%

vaccine efficacy, indicating that vaccination is necessary for maximum protection against SARS-

CoV-2 infection[46]. We also use our infrastructure to systematically obtain sera from individuals

recently infected with SARS-CoV-2 VOCs as part of multiple efforts to investigate antibody cross-

reactivity, which is essential for development of COVID-19 vaccine boosters.

As the threat of COVID-19 wanes and global SARS-CoV-2 surveillance networks scale

back, there is a strong need for continued development of rapid response tools. Diagnostics that

target multiple pathogens, such as our seasonal respiratory panel[47], are increasingly useful to this

end. These diagnostics may serve as outbreak detection tools when their negativity rate spikes,

indicating the emergence of a novel pathogen. Agnostic surveillance techniques, such as those

monitoring wastewater via shotgun sequencing[13–15,17], have already shown promise by detecting

Poliovirus Type 2 in New York wastewater[48]. If an outbreak does not immediately attenuate, then

our surveillance apparatus described in this study could serve as a model for sustained monitoring

of whole genome variations that could impact disease severity, outbreak dynamics, and the

efficiency of targeted diagnostic assays.

In this retrospective study, we showcase our unique positioning for rapid development and

maintenance of robust pathogen surveillance. Our nationwide surveillance network and its suite of

analytical and sequencing components collectively serve as a model for future large-scale

surveillance efforts. Looking to the future, it is our mission to stay vigilant and continue refining

this model to combat the many emergent infectious diseases posing imminent threats to public

health.

Methods

Ethical Statement

The use of residual de-identified samples for this study was determined as not a human subject

research requiring IRB review.

SARS-CoV-2 surveillance and whole genome sequencing

Extracted total nucleic acid from positive specimens identified through the Labcorp® FDA EUA

approved COVID-19 RT-PCR Test or SARS-CoV-2 & Influenza A/B Assay Test were sequestered

and consolidated using a Hamilton Microlab® STAR™ instrument and plate selector app, retaining

only positive samples with N1 Ct values less than 31. Sample RNA was reverse transcribed to

cDNA and a specially designed SARS-CoV-2 probe set containing ~1000 tiled Molecular Loop®

Loopcap™ Molecular Inversion Probes (MIPS) was used to amplify the cDNA from 99.6% of the

SARS-CoV-2 genome with most bases covered by 22 MIPs[19]. The product synthesized in-between

the MIPs was enriched and had sample specific molecular barcodes added via amplification for

long-read sequencing on a Pacific Biosciences® Sequel II™[49].

Sequence quality control and post-processing

After sequencing, circular consensus sequence (CCS) bam files were generated using the PacBio®

SMRT LINK™ software v9.0 ccs program[50] and subsequently demultiplexed using lima with the

following parameters: "--min-score-lead -1", "--min-score 80", "--window-size-multi 1.1", "--

neighbors". Molecular Loop® barcodes were then trimmed by aligning sequences to barcodes

using pbmm2 with parameters "--sort" and "--preset HIFI" and custom processing scripts. Final

sample fastq files were generated by converting the resulting bam files using BamTools[51].

Sequence fastq files were analyzed using a genome analysis pipeline implemented in the

CLC Genomics Server version 9.1.1[52]. This workflow starts with a sample-level fastq file and uses

Minimap2[53] to align reads to the SARS-CoV-2 reference genome (NCBI GenBank reference

NC_045512.2) to generate a bam file of the alignment as well as a VCF file containing the variants

called using a custom variant caller in CLC. A consensus sequence for each sample was then

generated using VCFCons v8.5.0[54]. When VCFCons calls a nucleotide sequence for genome

construction it was required to have least 4 CCS reads covering that base pair and an alternate

allele frequency compared to the reference of at least 80%. If the alternate allele frequency was

between 20% and 80%, then the appropriate ambiguous IUPAC nucleotide was called. If a

nucleotide was covered by less than 4 CCS reads it was reported as ambiguous (N) in the consensus

sequence.

Three different coverage quality control metrics were used to ensure high accuracy of

resulting consensus genome sequences. Firstly, the median CCS read coverage was calculated

separately for 29 ~1kb genomic regions and each sample was required to have at minimum 10X

mean of median amplicon coverage (depth of coverage). For samples to be kept for downstream

Phylogenetic Assignment of Named Global Outbreak (PANGO) lineage determination[18] and

sequence analysis, a minimum genome coverage of 50% was required. For sample genome

sequences to be reported to the CDC (and later deposited to GISAID[20]), a more stringent genome

coverage threshold of 90% was applied along with a third coverage filter that required no more

than 1 ambiguous base call in each 6bp sliding window of the S gene. Furthermore, samples were

reported within 21 days of collection.

Sample consensus genome sequences with at least 10X depth of coverage and genome

coverage of at least 50% were further analyzed using pangolin software[29] (v4.3.1 with pangolin

data v1.22) and the UShER algorithm[55] with default parameters to determine SARS-CoV-2

PANGO lineages. Sequences and their mutations were also characterized using Nextclade

v2.14.0[56] and the Nextclade SARS-CoV-2 dataset compiled on August 9th, 2023. In all subsequent

analyses, PANGO lineages were assigned groups by a pre-determined set of parent lineages,

representing key variants of concern (VOCs) throughout the pandemic. A lineage and its

descendants are indicated by appending "*". When a lineage is a descendant of two of the

following parent lineage groups, the closest parent was selected. If a lineage was not a descendant

of any of the parents, then it was placed in the "other" group. Parent lineage groups are as follows:

Alpha (B.1.1.7*), Beta (B.1.351*), Gamma (P.1*), Epsilon (B.1.427* and B.1.429*), Eta

(B.1.525*), Iota (B.1.526*), Kappa (B.1.617.1*), Mu (B.1.621*), Zeta (P.2*), Delta (B.1.617.2*),

BA.1*, BA.2*, BA.4*, BA.5*, BQ*, XBB*, XBB.1.5*, XBB.1.9*, and XBB.1.16*.

Demographic and phylogenetic analysis

Time-resolved phylogenetic analysis of SARS-CoV-2 consensus genome sequences was

performed using augur v.22.2.0[57] and auspice.us v0.12.0 (Auspice 2.49.0) within the Nextstrain

framework[58]. Consensus genomes were restricted to those with 100% S gene coverage and at least

99% genome coverage sequenced between January 2021 and June 2023. The augur filter utility

was used to limit the dataset to a maximum of 1,000 sequences per month each with metadata

including sample collection date and PANGO lineage. Next, the augur refine utility was used to

create a time-resolved phylogenetic tree using the TreeTime algorithm[59]. Finally, the resulting tree

was annotated with lineage information using the augur export utility, and the final tree was

displayed on auspice.us.

Age and gender distributions of all samples in this study with consensus genomes passing

the minimal coverage criteria for PANGO lineage determination were analyzed, restricting to those

collected between January 2021 and June 2023 with weekly sample counts of at least 100.

Distributions were also stratified by U.S. HHS regions[60]. For each region, annual gender

proportions were calculated, and age distributions were determined by aggregating ages into

different groups as follows: 0-17 (pediatrics), 18-19, 20-29, 30-39, 40-49, 50-59, 60-69, 70-79,

and 80 or older.

RT-PCR sampling, positivity, and N1 Ct value analysis

Demographic data and positive/negative results of RT-PCR samples collected through the

surveillance network described in this study were aggregated for multiple analyses. First, the RT-

PCR sampling was compared with Virseq assay sampling across each HHS region and month

where at least 100 Virseq samples were sequenced. The geographic distribution of RT-PCR and

Virseq sampling were also independently assessed by normalizing with U.S. state census data from

2020-2022[61]. Census data from 2023 was estimated by averaging that from 2020-2022. Overall

RT-PCR positivity was analyzed weekly and was also stratified by gender and age groups

(described in Methods section 'Demographic and phylogenetic analysis'). N1 Ct values of Virseq

samples were analyzed weekly and stratified based on whether samples were collected before or

after the Omicron wave (November 2021 to February 2022), denoted as 'pre-wave', 'Omicron

wave', and 'post-wave'. N1 Ct values were also compared based on co-infection status using only

deeply sequenced samples from the co-infection analysis described later in Methods section

'Workflow for detection and characterization of SARS-CoV-2 mixtures'.

Whole genome coverage and mutation frequency analysis

Genome-wide assessment of SARS-CoV-2 sequence mutation frequencies was performed using

all results obtained from Nextclade v2.14.0[56] for samples passing minimal coverage quality

control. Mutations were considered as "lineage-defining" if they appeared in at least 70% of the

genome sequences assigned to that lineage. The number of mutations with at least 5% prevalence

across the entire genome and within specific SARS-CoV-2 genic regions was calculated for each

sample collection week. Circulating lineage diversity was calculated in each week using unique

lineage counts and the Shannon entropy implementation in the vegan R package[62]. In all analyses

that indicate dominant lineage groups during specific time periods, this was calculated using the

first week where at least 5% of the lineages were assigned to that lineage group. In the case of

XBB.1.9 and XBB.1.16, these were combined, and the sum of their occurrences was used.

Per-base CCS coverage analysis was performed using selected VOCs (B.1.1.7, B.1.617.2,

BA.1, BA.2, BA.4, BA.5, BQ.1, and XBB.1.5), where the sequences chosen were required to have

depth of coverages of 100 +/- 10. 50 sequences were randomly chosen for each VOC that met the

coverage constraint and had the exact PANGO lineage determination of the VOC. Per-base

coverage was determined using Samtools[63] depth with parameters "-q 0 -Q 0" applied to the sample

alignment bam files. Median per-base coverage was then calculated for each VOC and smoothed

using a 30bp sliding window. Lineage defining mutation density was also determined for each

VOC by enumerating mutations at each genomic position and smoothing over a 1kbp sliding

window.

Virseq performance simulator

The Virseq pipeline performance was assessed by constructing a process whereby an input whole

genome sequence is mutated in a manner that simulates the coverage and errors introduced by

sequencing and post-processing analysis. This simulator was constructed using coverage and error

models of a representative sequencing batch selected from February 2022 containing 520 samples

reported to CDC (**Supplementary Data**). The coverage model was designed using a min max

normalization strategy, where two components were stored for later application of the model: 1) a

list of each sample's maximum per-base coverage and 2) the mean min max normalized coverage

at each position of the genome. In practice, since the peripheral 5' and 3' ends of the genome do

not have tiled coverage by design, a sample's min max normalized coverages equate to positional

coverage divided by the maximum per-base coverage. The error model was generated under the

assumption that the consensus at each base position is the correct base call. Thus, the probability

of an error was calculated using the maximum minor allele frequency. At each genomic position

the number of each base call was enumerated in R using Rsamtools[64] to identify the error rates for

each sample, and the mean error rate at each position was recorded. If the median coverage at a

position was less than 3, then the global median error rate was used. If the error rate was zero (i.e.

there were not any alternate base calls at a position), then half of the global minimum non-zero

error rate was used instead.

These two models were collectively applied to an input SARS-CoV-2 whole genome

sequence first by identifying positional coverages. A maximum coverage was randomly selected

from the list obtained from the representative batch and the expected mean coverage was computed

at each position by multiplying this maximum coverage by each pre-computed mean positional

fractional coverage. Next, the coverage at a position was sampled from a Poisson distribution using

the mean obtained in the prior step. If the sampled coverage was less than the minimum per-base

coverage threshold of 4, then an ambiguous base call was simulated at that position (see Methods

section 'Sequence quality control and post-processing'). Next, for all remaining unambiguous

positions a consensus base call was simulated using the sampled coverage and the pre-computed

average error at each position. Bases were sampled using a cumulative binomial using the average

error and coverage, and if the number of errors exceeded half of the base calls a random base was

called at the position; otherwise, the reference base was used. Finally, the PANGO lineages were

determined for both the original and simulated sequence using pangolin software[29] (v4.3.1 with

pangolin data v1.22) and checked for concordance. If the lineages were identical, the result was

considered an 'exact match'. If one lineage was a descendant of the other, then the result was

categorized as a 'parent match'. All other cases were considered 'discordant'.

Two simulation experiments were used to assess the performance of the Virseq pipeline.

Sequences were retrieved from GISAID[20] (accessed August 25th, 2023) and those used in the

experiments were restricted to those with at least 99% genome coverage, ensuring high quality

lineage calls. The first experiment assessed up to 100 sequences from each VOC and descendant

lineages thereof, current and former, designated in pangolin data v1.22. The second experiment

used a random selection of 10,000 sequences from each month ranging from January 2021 to July

2023.

Workflow for detection and characterization of SARS-CoV-2 mixtures

A custom workflow was developed to detect and characterize samples containing more than one

unique PANGO lineage, i.e. mixtures or co-infections. All bam files of samples passing minimum

coverage metrics were processed through the recommended freyja[14] processing workflow, first

calling variants using iVar[65] and subsequently using the freyja demixing algorithm[14] to determine

the lineage abundances in each sample. This algorithm attempts to identify a parsimonious set of

lineages best explaining the UShER[55] defining single nucleotide polymorphisms (SNPs) detected

in the sample.

Lineage abundances were then processed, greedily aggregating abundances of lineages

with parent/descendant relationships (e.g. BA.1 and BA.1.1) starting with the most abundant

lineage. This was performed since one lineage would have UShER[55] defining SNPs forming a

subset or superset of the other lineage, and it was assumed that this splitting of highly similar

lineages was due to sequencing noise. Samples were then filtered, requiring an empirically

determined minimum depth of coverage where the rate of mixture detection was stable and low

(**Supplementary Figure 12a**). Mixtures were then required to satisfy three criteria: 1) Top lineage

relative abundance no greater than 0.8, 2) second lineage relative abundance no less than 0.2, and

3) minimum of 3 UShER[55] defining SNPs discriminating the two lineages comprising the mixture

(**Supplementary Figure 12b-f**). Each resulting mixture sample was then categorized using the

parent lineage groups of the two mixed lineages (see Methods section 'Sequence quality control

and post-processing').

Mixture samples were then processed using WhatsHap[66], a standard haplotype assembly

tool suitable for long sequencing reads. If a sample did not yield any haplotype blocks (i.e. no two

SNPs were phased), then analysis was halted. If only a single haplotype block was obtained, then

the block length and phased mutations therein were recorded. If a sample had at least two haplotype

blocks, then a greedy algorithm was applied to merge these blocks while leveraging *a priori*

mixture knowledge yielded by freyja[14]. The number of UShER[55] defining SNPs unique to each

mixture lineage (i.e. lineage-discriminating SNPs) was recorded for each block, and blocks were

iteratively merged in order of descending number of lineage-discriminating SNPs (ties broken by

using the larger block length). Blocks were merged to maximize the number of correctly phased

lineage-discriminating SNPs. If the addition of a block to this greedily merged block didn't

improve this optimization criterion, then it was skipped, and the next block was assessed. This

process continued until no further blocks with lineage-discriminating SNPs remained. A custom

script was used to modify the haplotagged bam output by WhatsHap[66] for visualization of merged

haplotype blocks in Integrative Genomics Viewer v2.16.2[67].

Author Contributions

Development and maintenance of Virseq surveillance apparatus – all authors. Manuscript

supervision and administration – L.K.I. Manuscript conceptualization and methodology – H.N.B.,

K.S., Q.Zh., J.D.W., S.L., and L.K.I. Formal analysis and data curation – H.N.B. and K.S. Formal

analysis review – H.N.B., K.S., Q.Zh., S.L., and L.K.I. Preparation of manuscript figures and

tables – H.N.B. and K.S. Original manuscript draft preparation – H.N.B. Manuscript review and

editing – H.N.B., K.S., Q.Zh., Q.Ze., J.D.W., M.B.N., S.E.D., J.Me., M.E., S.L., and L.K.I.

Approval of final manuscript – all authors.

Funding Statement

564   No grants or financial support were used for this study.

Acknowledgements

566   We would like to acknowledge the dedicated members of our IT infrastructure team who continue

567   to be instrumental in maintaining the resources needed for this surveillance effort. We would also

like to thank the hundreds of technicians and technologists who processed SARS-CoV-2 PCR testing at Labcorp during the pandemic. Without the hard work and dedication of the broader Labcorp enterprise, this surveillance effort would not have been possible. We would also like to extend our gratitude to the dedicated teams at PacBio and Molecular Loop who were instrumental in helping us set up their respective resources for use in the Virseq pipeline. We would especially like to thank Elizabeth Tseng at PacBio for her significant contribution towards the development of the VCFCons software used in the Virseq pipeline. Lastly, we would like to thank the U.S. Centers for Disease Control (CDC) for their guidance throughout this surveillance effort.

Competing Interests

Data Availability

Figures S1-13 and **Table S1** may be found in the **Supplementary Materials**. A list of de-identified LCIDs and their corresponding GISAID EPI_ISL IDs from samples analyzed in this study that were reported to CDC may be found in the **Supplementary Data**. A list of LCIDs used to construct the empirical simulation model are included in the **Supplementary Data**. Additionally, all processed data used to generate **Figures 2b**, **3b**, **4-6**, and **S1-13** are provided in the **Supplementary Data**. Data used to generate **Figure 3a** are provided in **Table S1**.

References

1.   Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265–269 (2020).

2.   Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus:

implications for virus origins and receptor binding. *The Lancet* **395**, 565–574 (2020).

3.   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat

origin. *Nature* **579**, 270–273 (2020).

4.   Cohen, O. *et al.* Labcorp's perspective: Responding to SARS-CoV-2 and the next pandemic.

Nat. Portf. (2022).

5.   Sullivan, A. *et al.* Follow-Up SARS-CoV-2 PCR Testing Outcomes From a Large Reference

Lab in the US. *Front. Public Health* **9**, 679012 (2021).

6.   Sullivan, A. *et al.* Antibody titer levels and the effect on subsequent SARS-CoV-2 infection

in a large US-based cohort. *Heliyon* **9**, e13103 (2023).

7.   Alfego, D. *et al.* A population-based analysis of the longevity of SARS-CoV-2 antibody

seropositivity in the United States. *EClinicalMedicine* **36**, 100902 (2021).

8.   Markov, P. V. *et al.* The evolution of SARS-CoV-2. *Nat. Rev. Microbiol.* **21**, 361–379 (2023).

9.   Tao, K. *et al.* The biological and clinical significance of emerging SARS-CoV-2 variants.

Nat. Rev. Genet. **22**, 757–773 (2021).

10. Itokawa, K., Sekizuka, T., Hashino, M., Tanaka, R. & Kuroda, M. Disentangling primer

interactions improves SARS-CoV-2 genome sequencing by multiplex tiling PCR. *PLOS*

ONE **15**, e0239403 (2020).

11. Vacca, D. *et al.* Direct RNA Nanopore Sequencing of SARS-CoV-2 Extracted from Critical

Material from Swabs. *Life Basel Switz.* **12**, 69 (2022).

12. Rehn, A. *et al.* Catching SARS-CoV-2 by Sequence Hybridization: a Comparative Analysis.

mSystems **6**, e0039221 (2021).

13.   Baaijens, J. A. *et al.* Lineage abundance estimation for SARS-CoV-2 in wastewater using

612   transcriptome quantification techniques. *Genome Biol.* **23**, 236 (2022).

613   14.   Karthikeyan, S. *et al.* Wastewater sequencing reveals early cryptic SARS-CoV-2 variant

614   transmission. *Nature* **609**, 101–108 (2022).

615   15.   Smyth, D. S. *et al.* Tracking cryptic SARS-CoV-2 lineages detected in NYC wastewater. *Nat.*

616   *Commun.* **13**, 635 (2022).

617   16.   Wang, X. *et al.* Fecal viral shedding in COVID-19 patients: Clinical significance, viral load

618   dynamics and survival analysis. *Virus Res.* **289**, 198147 (2020).

619   17.   Kitajima, M. *et al.* SARS-CoV-2 in wastewater: State of the knowledge and research needs.

620   *Sci. Total Environ.* **739**, 139076 (2020).

621   18.   O'Toole, Á., Pybus, O. G., Abram, M. E., Kelly, E. J. & Rambaut, A. Pango lineage

622   designation and assignment using SARS-CoV-2 spike gene nucleotide sequences. *BMC*

623   *Genomics* **23**, 121 (2022).

624   19.   LoopCap^{TM} Technology Elegant, High-Performance Targeted NGS.

625   https://molecularloop.com/wp-content/uploads/2023/04/Molecular-Loop-Tech-

626   Note_0423_Final.pdf (2023).

627   20.   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative

628   contribution to global health. *Glob. Chall. Hoboken NJ* **1**, 33–46 (2017).

629   21.   Sayers, E. W. *et al.* Database resources of the national center for biotechnology information.

630   *Nucleic Acids Res.* **50**, D20–D26 (2022).

631   22.   Puhach, O., Meyer, B. & Eckerle, I. SARS-CoV-2 viral load and shedding kinetics. *Nat. Rev.*

632   *Microbiol.* **21**, 147–161 (2023).

23. Sentis, C. *et al.* SARS-CoV-2 Omicron Variant, Lineage BA.1, Is Associated with Lower
Viral Load in Nasopharyngeal Samples Compared to Delta Variant. *Viruses* **14**, 919 (2022).

24. CDC Expands Booster Shot Eligibility and Strengthens Recommendations for 12-17 Year
Olds. https://www.cdc.gov/media/releases/2022/s0105-Booster-Shot.html (2022).

25. CDC Recommends Pfizer Booster at 5 Months, Additional Primary Dose for Certain
Immunocompromised Children. https://www.cdc.gov/media/releases/2022/s0104-Pfizer-
Booster.html (2022).

26. Coronavirus (COVID-19) Update: FDA Expands Eligibility for Pfizer-BioNTech COVID-
19 Vaccine Booster Dose to Children 5 through 11 Years. https://www.fda.gov/news-
events/press-announcements/coronavirus-covid-19-update-fda-expands-eligibility-pfizer-
biontech-covid-19-vaccine-booster-dose (2022).

27. McMillen, T., Jani, K., Robilotti, E. V., Kamboj, M. & Babady, N. E. The spike gene target
failure (SGTF) genomic signature is highly accurate for the identification of Alpha and
Omicron SARS-CoV-2 variants. *Sci. Rep.* **12**, 18968 (2022).

28. Gayvert, K. *et al.* Evolutionary trajectory of SARS-CoV-2 genome shifts during widespread
vaccination and emergence of Omicron variant. *Npj Viruses* **1**, 5 (2023).

29. O'Toole, Á. *et al.* Assignment of epidemiological lineages in an emerging pandemic using
the pangolin tool. *Virus Evol.* **7**, veab064 (2021).

30. Pedro, N. *et al.* Dynamics of a Dual SARS-CoV-2 Lineage Co-Infection on a Prolonged Viral
Shedding COVID-19 Case: Insights into Clinical Severity and Disease Duration.
Microorganisms **9**, 300 (2021).

31. Lythgoe, K. A. *et al.* SARS-CoV-2 within-host diversity and transmission. *Science* **372**,
eabg0821 (2021).

32.   Wawina-Bokalanga, T. *et al.* Genomic evidence of co-identification with Omicron and Delta

657   SARS-CoV-2 variants: a report of two cases. *Int. J. Infect. Dis.* **122**, 212–214 (2022).

658   33.   Combes, P. *et al.* Evidence of co-infections during Delta and Omicron SARS-CoV-2 variants

659   co-circulation through prospective screening and sequencing. *Clin. Microbiol. Infect.* **28**,

660   1503.e5-1503.e8 (2022).

661   34.   Hosch, S. *et al.* Genomic Surveillance Enables the Identification of Co-infections With

662   Multiple SARS-CoV-2 Lineages in Equatorial Guinea. *Front. Public Health* **9**, 818401

663   (2022).

664   35.   Zhou, H.-Y. *et al.* Genomic evidence for divergent co-infections of co-circulating SARS-

665   CoV-2 lineages. *Comput. Struct. Biotechnol. J.* **20**, 4015–4024 (2022).

666   36.   Francisco Jr, R. D. S. *et al.* Pervasive transmission of E484K and emergence of VUI-NP13L

667   with evidence of SARS-CoV-2 co-infection events by two different lineages in Rio Grande

668   do Sul, Brazil. *Virus Res.* **296**, 198345 (2021).

669   37.   Rockett, R. J. *et al.* Co-infection with SARS-CoV-2 Omicron and Delta variants revealed by

670   genomic surveillance. *Nat. Commun.* **13**, 2745 (2022).

671   38.   Nguyen, N. N., Nguyen, Y. N., Hoang, V. T., Million, M. & Gautret, P. SARS-CoV-2

672   Reinfection and Severity of the Disease: A Systematic Review and Meta-Analysis. *Viruses*

673   **15**, 967 (2023).

674   39.   WHO   COVID-19   vaccine   tracker   and   landscape.

675   https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccines

676   (2023).

677   40.   Peñas-Utrilla, D. *et al.* Systematic genomic analysis of SARS-CoV-2 co-infections

678   throughout the pandemic and segregation of the strains involved. *Genome Med.* **15**, 57

679   (2023).

680   41.   Li, Y. *et al.* Both simulation and sequencing data reveal coinfections with multiple SARS-

681   CoV-2 variants in the COVID-19 pandemic. *Comput. Struct. Biotechnol. J.* **20**, 1389–1401

682   (2022).

683   42.   Arora, P. *et al.* The SARS-CoV-2 Delta-Omicron Recombinant Lineage (XD) Exhibits

684   Immune-Escape Properties Similar to the Omicron (BA.1) Variant. *Int. J. Mol. Sci.* **23**, 14057

685   (2022).

686   43.   Lacek, K. A. *et al.* SARS-CoV-2 Delta–Omicron Recombinant Viruses, United States.

687   *Emerg. Infect. Dis.* **28**, 1442–1445 (2022).

688   44.   Mohapatra, R. K., Kandi, V., Tuli, H. S., Chakraborty, C. & Dhama, K. The recombinant

689   variants of SARS-CoV-2: Concerns continues amid COVID-19 pandemic. *J. Med. Virol.* **94**,

690   3506–3508 (2022).

691   45.   Casadevall, A. & Pirofski, L. The convalescent sera option for containing COVID-19. *J. Clin.*

692   *Invest.* **130**, 1545–1548 (2020).

693   46.   Schuh, A. J. *et al.* SARS-CoV-2 Convalescent Sera Binding and Neutralizing Antibody

694   Concentrations Compared with COVID-19 Vaccine Efficacy Estimates against Symptomatic

695   Infection. *Microbiol. Spectr.* **10**, e01247-22 (2022).

696   47.   Larkin, H. D. First Nonprescription COVID-19 Test That Also Detects Flu and RSV. *JAMA*

697   **328**, 11 (2022).

48.   Ryerson, A. B. *et al.* Wastewater Testing and Detection of Poliovirus Type 2 Genetically

Linked to Virus Isolated from a Paralytic Polio Case — New York, March 9–October 11,

2022. *MMWR Morb. Mortal. Wkly. Rep.* **71**, 1418–1424 (2022).

49.   Rhoads, A. & Au, K. F. PacBio Sequencing and Its Applications. *Genomics Proteomics*

Bioinformatics **13**, 278–289 (2015).

50.   Wenger, A. M. *et al.* Accurate circular consensus long-read sequencing improves variant

detection and assembly of a human genome. *Nat. Biotechnol.* **37**, 1155–1162 (2019).

51.   Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P. & Marth, G. T. BamTools:

a C++ API and toolkit for analyzing and managing BAM files. *Bioinforma. Oxf. Engl.* **27**,

1691–1692 (2011).

52.   Liu, C.-H. & Di, Y. P. Analysis of RNA Sequencing Data Using CLC Genomics Workbench.

Methods Mol. Biol. Clifton NJ **2102**, 61–113 (2020).

53.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

54.   Tseng, E., Zeng, Q. & Iyer, L. VCFCons: a versatile VCF-based consensus sequence

generator for small genomes. *bioRxiv* (2021) doi:10.1101/2021.02.26.433111.

55.   Turakhia, Y. *et al.* Ultrafast Sample placement on Existing tRees (UShER) enables real-time

phylogenetics for the SARS-CoV-2 pandemic. *Nat. Genet.* **53**, 809–816 (2021).

56.   Aksamentov, I., Roemer, C., Hodcroft, E. & Neher, R. Nextclade: clade assignment, mutation

calling and quality control for viral genomes. *J. Open Source Softw.* **6**, 3773 (2021).

57.   Huddleston, J. *et al.* Augur: a bioinformatics toolkit for phylogenetic analyses of human

pathogens. *J. Open Source Softw.* **6**, 2906 (2021).

58. Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution. *Bioinforma. Oxf.*
Engl. **34**, 4121–4123 (2018).

59. Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood phylodynamic
analysis. *Virus Evol.* **4**, (2018).

60. HHS Regional Offices. https://www.hhs.gov/about/agencies/iea/regional-offices/index.html.
(2021).

61. U.S.   Census   Bureau.   Annual   Population   Estimates,   2020-2022.
https://www2.census.gov/programs-surveys/popest/datasets/2020-2022/state/totals/NST-
EST2022-ALLDATA.csv.

62. Oksanen,   J.   *et   al.*   vegan: Community   Ecology   Package.   https://CRAN.R-
project.org/package=vegan. (2022).

63. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).

64. Morgan, M., Pagès, H., Obenchain, V. & Hayden, N. Rsamtools: Binary alignment (BAM),
FASTA,   variant   call   (BCF),   and   tabix   file   import.
https://bioconductor.org/packages/Rsamtools. (2021).

65. Grubaugh, N. D. *et al.* An amplicon-based sequencing framework for accurately measuring
intrahost virus diversity using PrimalSeq and iVar. *Genome Biol.* **20**, 8 (2019).

66. Patterson, M. *et al.* WhatsHap: Weighted Haplotype Assembly for Future-Generation
Sequencing Reads. *J. Comput. Biol. J. Comput. Mol. Cell Biol.* **22**, 498–509 (2015).

67. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

Figure Legends

Figure 1. Journey of a sample from raw nasopharyngeal and nasal swabs to CDC reporting.
a) Overview of end-to-end genomic surveillance setup. **b)** Consolidation of positive high viral titer samples. **c)** Probe amplification protocol. **d)** Long read sequencing analysis and bioinformatics workflow to prepare high-quality SARS-CoV-2 genomic sequences with demographic metadata for CDC.

Figure 2. SARS-CoV-2 PANGO lineage analysis of 594,832 high-quality genomes from USA samples collected between January 2021 and July 2023. In each plot samples are grouped and colored by the closest parent listed on the top left. **a)** Time-resolved phylogeny of a subset of samples (n=28,069) clustered based on the oldest collected sample with at most 1,000 samples per month. **b)** SARS-CoV-2 lineage proportions across weeks with at least 100 samples (left y-axis) and the PCR positivity rate (%) indicated by the black line (right y-axis).

Figure 3. U.S. nationwide SARS-CoV-2 genomic surveillance consisting of 594,832 high-quality genomes from samples collected between January 2021 and July 2023. In both panels, the U.S. is divided into 10 Health and Human Services (HHS) regions, each in separate colors. **a)** Each state is notated by its two-letter abbreviation and the shade indicates the number of samples collected. Each region uses a shared color gradient, shown on the right in grayscale with maximum and minimum values of 19,000 and 1,000, respectively. **b)** Line plots showing the ratio of Virseq-generated genomes versus the total SARS-CoV-2-positive RT-PCR samples collected on the log2 scale over each sample collection month. January 2021 and July 2023 were excluded, as they each have fewer than 100 Virseq-generated genomes in the dataset analyzed. Vertical black lines denote the start of years 2022 and 2023.

Figure 4. Pandemic-wide trends in SARS-CoV-2 genomic mutations and the robustness of

whole genome probe-based sequencing. a) Number of mutations with at least 5% prevalence in

the lineage population (black) and circulating lineage shannon diversity (red). **b)** Number of

mutations with at least 5% prevalence in the lineage population separated by genic origin. **c)**

Heatmap showing the genome-wide probe coverage of the most common lineage in circulation for

each collection week with genomic positions shown 5' (bottom) to 3' (top). Probes were considered

failures if a deletion, insertion, or >3 SNPs were detected in either probe arm. Large genic regions

(ORF1a, ORF1b, S) are indicated by horizontal lines and are labeled on the right. In all panels,

results are stratified by sample collection week with vertical bars separating the major waves of

the pandemic with the causal variant shown above. Waves are demarcated using the collection

week when the causal variant first reached 5% prevalence.

Figure 5. Virseq simulation results. Exact and parent concordance for the retrospective (**a**) and

VOC (**b**) simulation experiments. Retrospective results are shown with exact (red) and parent

(blue) concordance as separate lines over the sample collection months analyzed, with years

demarcated by vertical lines. VOC results are shown with exact and parent concordance results for

each VOC data point with 90% thresholds marked by dashed red lines. VOCs colored black have

>90% exact concordance, those colored yellow only have >90% parent concordance, and those

colored red are below both concordance thresholds. Vertical gray bars separate the major waves of

the pandemic with the causal variant shown above. Waves are demarcated using the collection

week when the causal variant first reached 5% prevalence.

Figure 6. Haplotype phasing of SARS-CoV-2 mixture samples (i.e. coinfections). a-b)

Prevalence and number of mixtures in each sample collection week with major pandemic waves

demarcated as they were in **Figure 3**. In **(a)**, the weekly average (~0.2%) is shown as a horizontal

dashed red line. In **(b)**, mixtures are colored based on the lineage family of the major and minor

mixture components, e.g. BA.1*-Delta indicates a mixture of BA.1 and Delta sublineages with

majority BA.1. **c)** Sample-level comparison of the number of and minimum distance between

UShER defining mutations that discriminate the two mixture lineages. Samples are colored based

on their number of defining mutations phased: >1 phased (black, n=149), 1 phased only with non-

defining mutation(s) (yellow, n=128), and 0 phased (red, n=102). **d)** Comparison of sample

defining mutation phasing among largest resolved haplotype blocks. Results are compared

between original whatshap haplotype blocks and those same haplotype blocks merged using freyja

mixture results. **e)** The number of defining mutations phased and the lengths of sample merged

haplotype blocks.

TABLES

Table 1. Summary of VOCs with more frequent discordant PANGO lineage calls. Each row shows

a lineage with < 90% parent concordance and the most common discordant lineage observed

among the simulated sequences.

PANGO lineage	Concordance (%)	Total sequences analyzed	Most common discordant lineage
BA.2.2.1	75.00%	4	BA.1
BA.5.10	60.00%	10	BA.5.2
BQ.1.19	75.00%	4	BQ.1.2
BY.1*	86.67%	15	BA.2.75.7
XB*	55.00%	100	B.1
XBV*	78.57%	14	XBB.1
XP*	85.71%	7	BA.1.1

801   * BY is the alias of BA.2.75.6. XB, XBV, and XP are the following recombinants, respectively:

802   B.1.634/B.1.631, CR.1/XBB.1, and BA.1.1/BA.2.

803   **Figure 1**

805   **Figure 2**

Figure 3

809   **Figure 4**

811 **Figure 5**

813   **Figure 6**