**Title:** Demographic analysis of mortality changes associated with the COVID-19

pandemic in Japan, 2020–2022

**Running title:** Mortality change in Japan during COVID-19 pandemic

**Article type:** Original Article

Authors:

Yuta Okada, M.D., Ph.D.[a], Hiroshi Nishiura, M.D., Ph.D.[a]

[a] Graduate School of Medicine, Kyoto University, Kyoto, Japan

Correspondence to: Hiroshi Nishiura

Address: Kyoto University, Yoshidakonoecho, Sakyoku, Kyoto city, Kyoto 6068503,

Japan

Tel: +81-75-753-4456; Fax: +81-75-753-4458

Email: nishiura.hiroshi.5r@kyoto-u.ac.jp

**Declarations of interest**

None.

**Abstract**

**Objectives:** To evaluate the mortality impact of the COVID-19 pandemic by age and

prefecture in Japan.

**Study design:** Retrospective demographic analysis.

**Methods:** A hierarchical time-varying regression model was applied to prefectural life

table data in Japan during 2000–2019. Pre-pandemic baseline mortality rates and death

counts during 2020–2022 were projected by this model to evaluate the mortality impact

of the COVID-19 pandemic on mortality during 2020–2022.

**Results:** Nationally, the observed mortality rates were higher than projected rates

among those aged 10–29 years and lower for ages 1–9 and >80 years in 2020. During

2021 and 2022, mortality rates consistently increased for most age groups, with

substantial regional heterogeneity in those aged <14 years. Consistently, the neutral

gaps between the observed and projected life expectancy across prefectures in 2020

turned negative in most prefectures during 2022. In 2020, the negative gap in life

expectancy ranged from a substantial shortening of −1.41 years (95% prediction interval

[PI]: −1.69, −1.12) in Okinawa to a slight shortening of −0.23 years (95% PI: −0.63,

0.17) in Tottori. Excess deaths per population also grew consistently in all prefectures

during 2020–2022; Miyazaki yielded the largest estimate at 186.2 (95% PI: 158.6,

214.0) per 100,000 population.

**Conclusions:** Our framework highlights the usefulness of life table data for evaluating

the mortality impact caused by an event such as the pandemic. Our estimates, adjusted

by pre-pandemic demographic trends, revealed the massive mortality impact of the

COVID-19 pandemic across a wide age range.

**Keywords:** Excess death; COVID-19; life table; Bayesian hierarchical model; Japan

**Introduction**

COVID-19 has caused substantial morbidity and mortality worldwide since its

emergence in Wuhan, China in November 2019. In addition to documented COVID-19

deaths, substantial mortality directly or indirectly associated with COVID-19 has

stemmed from various mechanisms, including cardiovascular complications or limited

health care access.[1–4] At the global level, an estimated 14.9–15.9 million excess deaths

occurred up to the end of 2021[5–7], with excess deaths consistently reported even after

2022. [8,9] Trends in life expectancy estimates are also in line with those suggested by the

excess mortality estimates, elucidating a substantial impact on mortality during the early

phases of the COVID-19 pandemic in many countries. [6,10–13]

Globally, evaluation of the total mortality impact owing to the COVID-19

pandemic has been carried out using excess mortality estimates. This approach is widely

accepted for quantifying the mortality impact of epidemics, such as seasonal influenza,

which is not fully observable by relying on indicator-based surveillance. Technically,

excess mortality, expressed either as an absolute number or rate per population, is

typically estimated as the difference between the observed and predicted mortality, or

the "baseline" mortality projected by a statistical model fitted to the historical

baseline.[5,6,14–20] The definition of "baseline" mortality is a practical issue in its

application to COVID-19. The baseline is updated according to the latest available data

in widely used approaches for the calculation of excess mortality; thus, in the case of

COVID-19, the baseline spans an extended period. This leads to difficulty in

interpreting the estimated excess deaths as the genuine impact of mortality owing to the

pandemic. One such example is the weekly excess mortality estimates in Japan provided

in the "Excess and Exiguous Deaths Dashboard in Japan" (URL: https://exdeaths-

japan.org/en/; hereinafter, ExDeath Dashboard).[9] In that data source, the excess

mortality at national and prefectural levels is calculated using the Farrington algorithm,

which updates the baseline to the latest 5-year period of available weekly data.[9,18]

As for the estimation of excess mortality, another practical issue in

conventional approaches is that these estimates do not provide results that are age-

standardized or age-stratified. This can potentially lead to inappropriate interpretation of

the mortality impact caused by external factors including COVID-19, especially when

the underlying demographic change is not negligible. One measure to circumvent this

issue is to deploy demographic approaches based on life table data to capture the

temporal change in the age-wise mortality structure as well as to forecast mortality.[21–24]

Compared with conventional excess mortality estimates on a weekly basis, an apparent

limitation of a life table-based approach is that life table data are usually only available

on a yearly basis, with a certain time delay in reporting. However, an obvious strength

of a life table-based approach is that key backbones in the age-mortality correlation and

the underlying demographic trends can be captured, which may enable technically more

sound projections of pre-pandemic trends in mortality beyond 2020, compared with

existing approaches.

Given the above background, in the present study, we aimed to establish a

novel framework to estimate post-pandemic changes in mortality by age group at

national and subnational levels using longitudinal life table data provided in the

Japanese Mortality Database.[25] Using life table data during 2000–2019, we applied a

time-varying regression model with a hierarchical structure that captured the pre-

pandemic trend in mortality schedules by age group. The mortality during 2020–2022

projected by this model revealed the changes in mortality and life expectancy

attributable to the COVID-19 pandemic during 2020–2022 at national and prefectural

levels.

**Methods**

**Epidemiological data**

We retrieved national and prefectural life table data in Japan from 2020 to 2022

from The Japanese Mortality Database. [25] We recalculated these life tables in order to

handle those over 100 in a single age group. Death counts in each age group (not

integers) were rounded to the closest integer values for subsequent analysis.

We retrieved data on mortality caused by the 2011 Tohoku earthquake from the vital

statistics of 2011 published by the Ministry of Health, Labour and Welfare. [26] For

subsequent statistical inference, we basically subtracted disaster-associated death counts

owing to the Tohoku earthquake from age groups >5 years for Iwate, Miyagi, and

Fukushima prefectures. In the Tohoku earthquake data, the youngest age group was

categorized as 0–4 years, which does not match the classification in the life tables (ages

0 and 1–4 as separate groups); we therefore accounted for the disaster-related mortality

in the statistical model.

We also retrieved estimated weekly excess or exiguous death estimates from the

ExDeath Dashboard[9] during 2020–2022 for comparison with our excess death

estimates.

**Decomposition of the national-level mortality table during 2000–2019**

We used the age-stratified mortality rates from national life tables during

2000–2019 to compose a mortality schedule matrix $\mathbf{M}$. Singular value decomposition of

$\mathbf{M}$ is a widely used method in demography to extract key characteristics in mortality

schedules. The result of decomposition can be written in the form

$$\mathbf{M} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^{\mathrm{T}}, \tag{1}$$

where $\mathbf{U}$ is a time-dependent component, $\mathbf{\Sigma}$ generates singular values, and $\mathbf{V}$ contains

column vectors that represent key structures in mortality along age. We applied singular

value decomposition to the national longitudinal mortality schedule matrix, rather than

to the mortality schedule matrix obtained by combining data from all regions, as in

Alexander et al.[24]

**Hierarchical time-varying regression model**

We built a statistical model to capture national-level mortality trends while also

allowing variability by prefecture, similar to that in Alexander et al., with some

modifications.[24] Briefly, we modeled the mortality rate in age group $x$ and in prefecture

$a$ in year $t$ as a linear combination of $Y_x^{(1)}$, $Y_x^{(2)}$, and $Y_x^{(3)}$, which are the three leading

column vectors of $\mathbf{V}$ described above:

$$\log(m_{x,a,t}) = \sum_{p=1}^{3} \beta_{a,t}^{(p)} Y_x^{(p)} + u_{x,a}, \tag{2}$$

where $\beta_{a,t}^{(p)}$ ($p = 1, 2, 3$) are the time-varying coefficients in year $t$. For details on the

shapes and reason for choosing $Y_x^{(1)}$, $Y_x^{(2)}$, and $Y_x^{(3)}$, see the Supplementary Methods 3.

The final term $u_{x,a}$ in equation (2) is the random effects that were assumed to be time-

invariant in our model. We also assumed that $\beta_{a,t}^{(p)}$ is modeled as the sum of the national

averages $\overline{\beta_t^{(p)}}$ and random effects $\psi_{\beta|a,t}^{(p)}$ in year $t$; see the Supplementary Methods 1 for

details.

A main part of the likelihoods considered in the present study was based on the

observation process of death counts in each prefecture and in each age group. We

assumed that the observed mortality rate in age group $x$, prefecture $a$, in year $t$ follows a

Poisson distribution. Additionally, the likelihood for total death counts across all age

groups in each prefecture was also taken into consideration.[27] See the Supplementary

Methods 1 on how the effect of the 2011 earthquake is accounted for with respect to

$m'_{x,a,t}$.

**Statistical inference of temporal changes in mortality**

140  The statistical model for the temporal changes in $\overline{\beta_t^{(p)}}$ was dealt with as time-

141  dependent and was assumed as a second-order random walk:

$$\overline{\beta_{t+1}^{(p)}} - \overline{\beta_t^{(p)}} \sim Normal\left(\overline{\beta_t^{(p)}} - \overline{\beta_{t-1}^{(p)}}, \sigma_{\beta,time}^{(p)}\right) \tag{3}$$

We used the Markov chain Monte Carlo method to estimate unknown parameters.

On the basis of the estimates of $\beta_{a,t}^{(p)}$ from the 2000–2019 data, we projected

$\beta_{a,t}^{(p)}$ up to 2022. Therefore, we assumed the following: (i) the change rate of $\overline{\beta_t^{(p)}}$ from

2018 to 2019 was maintained up to 2022, (ii) the prefectural random effects $\psi_{\beta|a,2022}^{(p)}$,

$\psi_{\beta|a,2021}^{(p)}$, and $\psi_{\beta|a,2020}^{(p)}$ are equal to $\psi_{\beta|a,2019}^{(p)}$; consequently, we obtained the projected

mortality rates at the national level, $\overline{m_{x,t}^{',projected}}$, and at prefectural levels, $m_{x,a,t}^{',projected}$,

up to 2022. Excess or exiguous deaths were computed as the gap between projection-

based prediction and observations of death counts during 2020–22. The gaps between

observed and projected life expectancy at birth, $\Delta e_{0|a,t} = e_{0|a,t}^{observed} - e_{0|a,t}^{projected}$, were

also evaluated by calculating $e_{0|a,t}^{projected}$ using predicted values of $m_{x,a,t}^{',projected}$ ($t =$

2020-2022). The calculation of life expectancy was based on standard methods to

calculate period life tables[28]. See the Supplementary Methods 1-2 for further technical

details.

**Statistical analysis software**

All statistical and numerical analyses in the present study were performed

using R version 4.2.2 (The R Project for Statistical Computing, Vienna, Austria) and

CmdStan version 2.33.0.[29,30]

**Results**

Changes in the national mortality rates by age group during 2019–2022 are

shown in Figure 1. The mortality curves during 2019–2022 were not substantially

different and were lower overall across all age groups, compared with 2000 or 2010

(Figure 1A). The difference in mortality curves during 2020-2022 compared with that of

2019 is shown in Figure 1B. During 2020–2022, the overall mortality rates among those

younger than 10 years old were lower than those in 2019, which contrasts with the

higher mortality rates during the same period for individuals in their 10s and 20s. In age

groups >30 years, 2022 is outstanding in that mortality rates among all age groups >30

years were higher than those in 2019.

The difference between the observed and projected mortality rates in 2020–

2022 is shown in Figure 2. The overall trend seen in the results from 2020 (panel A) is

substantial exiguous mortality in those aged 1–9 and ≥80 years and substantial excess

mortality in those aged 15–59 and 65–74 years. This substantial excess mortality rate

became greater in magnitude during 2021 and 2022, and the age range with excess

mortality rate widened up to those aged 75–79 years. The exiguous mortality rate

among individuals aged 1–9 years disappeared in 2021 and 2022 whereas that among

those aged ≥80 years was still observable among individuals aged ≥95 years.

The prefectural version of the values for the calculated gap between observed

and projected mortality rates during 2020–2022 is shown in Figure 3. The overall

tendency during 2020–2022 is a gradual increase in these gaps among those aged ≥30

years, where most age groups experienced consistent increases in mortality. Conversely,

greater heterogeneity among prefectures was observed in those younger than 30 years

old for the entire period, but the overall trend showed an increase in excess mortality,

even in those younger age groups.

A real-time update of the gap between observed and projected life expectancy

at birth is depicted in Figure 4, showing how long the life expectancy was shortened in

each pandemic year. In 2020, the life expectancy gaps were neutral to positive in all

except a few prefectures. The finding in 2021 differed in that these gaps became mostly

neutral to negative, and only the gap in Iwate was clearly in the positive range. The

negative tendency grew greater in 2022 when these gaps were clearly negative in all

prefectures except Tottori, where the gap was −0.235 (95% prediction interval [PI]:

−0.631, 0.168). The greatest negative gap was −1.41 years (95% PI: −1.69, −1.12) in

Okinawa.

Figure 5 shows the mortality gap between observed and projected deaths per

100,000 population in all prefectures, in comparison with the yearly aggregated mean

predicted gaps provided in ExDeath dashboard.[9] Our estimates showed neutral to

exiguous deaths in most prefectures in 2020 (Figure 5A). However, neutral to excess

deaths were seen in 2021, and this trend grew worse in 2022 when all prefectures

experienced substantial excess deaths. The worst prefecture in terms of excess deaths

was Miyazaki with 186.2 (95% PI: 158.6, 214.0) excess deaths per 100,000 population;

this was followed by Okinawa (146.9, 95% PI: 127.5, 166.1) and Toyama (145.5, 95%

PI: 117.2, 174.0). Comparison with the data from ExDeath Dashboard in 2022 revealed

some discrepancy with our estimates, especially in highly populated prefectures

including Tokyo, Osaka, and their neighboring prefectures.

Regarding gaps in life expectancy and death counts, results in the 5-year

baseline scenario were not substantially different from the main scenario, as presented

in the main text (see the Supplementary Figures 7-8).

**Discussion**

In the present study, we proposed a framework to evaluate excess mortality by

age and prefecture during the COVID-19 pandemic from 2020 to 2022 in Japan. By

integrating demographic and statistical frameworks developed in previous studies, we

projected the baseline mortality rates during 2000–2019 by extending the pre-pandemic

trend up to 2022 and evaluated the difference between observed mortality rates and

projected ones. Consideration for the age structures in mortality schedules in the present

framework is a key distinction from established regression-based excess mortality

estimations,[5,6,14–20] making our framework an important addition to an array of

conventional statistical and demographic approaches used in common excess mortality

estimations. Although our analysis was limited to the mortality impact of the COVID-

19 pandemic in Japan, the range of potential applications of our framework is wide

because the proposed approach only requires yearly life table data that are readily

available in many countries.

221    The key finding in our study was the substantial and steady increase in

222 observed mortality rates and the loss of life expectancy at birth in 2021 and 2022, as

223 illustrated in Figures 2–4. These results support our previous findings[31,32] that mortality

224 rates among individuals aged >30 years were significantly elevated during 2020–2022.

225 Another important finding from our study was the importance of accounting for the

226 underlying trend in the age-wise demographic structure, as suggested by the findings in

227 Figure 5. Although our results basically align with the ExDeath Dashboard,

228 substantially higher estimates are given in the ExDeath Dashboard for Tokyo, Osaka,

229 and their neighboring prefectures, despite partial use of post-pandemic data. This

230 finding might be due to our model accounting for baseline trends in age structure, in

231 contrast to the Farrington algorithm used in the ExDeath Dashboard, which is designed

232 to capture historical seasonal patterns. One future scope of methodological development

233 might encompass the combination of age-structured models like ours with conventional

234 statistical methodologies that also capture seasonality, which would be enabled when

235 finer-scale chronological data become available.

236    Technically, the key contribution of our study is that we extended the

237 framework by Alexander et al.,[24] which combines matrix factorization of the mortality

238 schedule matrix with time-varying regression analysis to enable excess mortality

estimation. Although our primary goal was not to improve regional estimates as in

Alexander et al., we projected future mortality schedules based on time-varying

regression of the baseline period, which may likely reflect pre-pandemic demographic

trends.

The present study has two limitations. First, our framework is designed to

handle yearly life table data from Japan, which are usually reported with a time lag

longer than a year; this naturally means that results cannot be obtained in a timely

manner. However, our framework is also applicable with minor modifications when up-

to-date data with shorter reporting intervals become available. Another important

limitation is that we simply extended the pre-pandemic demographic trends to obtain

the projected "baseline" mortality, which may have changed even in the absence of the

COVID-19 pandemic. Even conventional regression-based approaches are not free of

this limitation, but potential non-pandemic trend changes might have a larger impact on

our approach because this method entirely omits post-pandemic data from the baseline

estimation. However, our future projections of pre-pandemic mortality trends

incorporated uncertainty, which should have alleviated such potential shortcomings by

considering the cumulative uncertainty of the projections up to 2022. For further

improvement regarding this issue, exploration of methodologies that are both practically

feasible and ensure interpretability is warranted. Additional discussion regarding the

length of the baseline period can be found in the Supplementary Discussion.

In conclusion, we successfully established a novel framework that combines

demographic approaches with hierarchical Bayesian time-varying regression models to

evaluate the mortality impact of the COVID-19 pandemic in Japan during 2020–2022.

The application of this framework not only is limited to Japan or COVID-19 but also

may serve as a tool for assessing excess mortality caused by unexpected external events,

such as future pandemics, where longitudinal life table data are available.

**Acknowledgment**

We thank Analisa Avila, MPH, ELS, of Edanz (https://jp.edanz.com/ac) for editing a

draft of this manuscript.

**Funding Sources**

Y.O. received funding from the SECOM Science and Technology Foundation.

H.N. received funding from Health and Labour Sciences Research Grants (grant

numbers 20CA2024, 21HB1002, 21HA2016, and 23HA2005), the Japan Agency for

Medical Research and Development (grant numbers JP23fk0108612 and

JP23fk0108685), JSPS KAKENHI (grant numbers 21H03198 and 22K19670), the

Environment Research and Technology Development Fund (grant number

JPMEERF20S11804) of the Environmental Restoration and Conservation Agency of

Japan, the Daikin GAP Fund of Kyoto University, Japan Science and Technology

Agency SICORP program (grant numbers JPMJSC20U3 and JPMJSC2105), the

CREST program (grant number JPMJCR24Q3), and RISTEX program for Science,

Technology, and Innovation Policy (grant number JPMJRS22B4). The funders had no

role in the study design, data collection and analysis, decision to publish, or preparation

of the manuscript.

**285     Conflict of interest**

286     We declare that we have no conflicts of interest.

**287     Ethical approval statement**

288     Ethical approval was not required because all data used in the present study did not

289     include any personally identifiable information.

**290     Data availability**

All data used in the present study are publicly available from the Japanese Mortality

Database (https://www.ipss.go.jp/p-toukei/JMD/index-en.asp) and from the "Excess and

Exiguous Deaths Dashboard in Japan" (URL: https://exdeaths-japan.org/en/). The

supplementary files include the summary data from the results of our analyses.

**References**

1.    Knight R, Walker V, Ip S, et al. Association of COVID-19 with major arterial and
venous thrombotic diseases: A population-wide cohort study of 48 million adults in
England and wales. *Circulation*. 2022;146(12):892-906.
doi:10.1161/CIRCULATIONAHA.122.060785

2.    Rossman H, Meir T, Somer J, et al. Hospital load and increased COVID-19 related
mortality in Israel. *Nat Commun*. 2021;12(1):1904. doi:10.1038/s41467-021-
22214-z

3.    French G, Hulse M, Nguyen D, et al. Impact of hospital strain on excess deaths
during the COVID-19 pandemic - United States, July 2020-July 2021. *MMWR*
*Morb Mortal Wkly Rep*. 2021;70(46):1613-1616. doi:10.15585/mmwr.mm7046a5

4.    Raisi-Estabragh Z, Cooper J, Salih A, et al. Cardiovascular disease and mortality
sequelae of COVID-19 in the UK Biobank. *Heart*. 2022;109(2):119-126.
doi:10.1136/heartjnl-2022-321492

5.    Msemburi W, Karlinsky A, Knutson V, Aleshin-Guendel S, Chatterji S, Wakefield
J. The WHO estimates of excess mortality associated with the COVID-19
pandemic. *Nature*. 2023;613(7942):130-137. doi:10.1038/s41586-022-05522-2

6.    GBD 2021 Demographics Collaborators. Global age-sex-specific mortality, life
expectancy, and population estimates in 204 countries and territories and 811
subnational locations, 1950-2021, and the impact of the COVID-19 pandemic: a
comprehensive demographic analysis for the Global Burden of Disease Study
2021. *Lancet*. 2024;403(10440):1989-2056. doi:10.1016/S0140-6736(24)00476-8

7.    COVID-19 Excess Mortality Collaborators. Estimating excess mortality due to the
COVID-19 pandemic: a systematic analysis of COVID-19-related mortality, 2020-
21. *Lancet*. 2022;399(10334):1513-1536. doi:10.1016/S0140-6736(21)02796-3

8.    Excess mortality within England: 2023 data - statistical commentary. Gov.uk.
October 17, 2024. Accessed October 31, 2024.
https://www.gov.uk/government/statistics/excess-mortality-within-england-post-
pandemic-method/excess-mortality-within-england-2023-data-statistical-
commentary

9.    Excess and Exiguous Deaths Dashboard in Japan. Excess and Exiguous Deaths
Dashboard in Japan. September 24, 2024. Accessed October 31, 2024.
https://exdeaths-japan.org/

10.   Schöley J, Aburto JM, Kashnitsky I, et al. Life expectancy changes since COVID-
19. *Nat Hum Behav*. 2022;6(12):1649-1659. doi:10.1038/s41562-022-01450-3

11.   Aburto JM, Schöley J, Kashnitsky I, et al. Quantifying impacts of the COVID-19
pandemic through life-expectancy losses: a population-level study of 29 countries.
*Int J Epidemiol*. 2022;51(1):63-74. doi:10.1093/ije/dyab207

12.   Huang G, Guo F, Liu L, et al. Changing impact of COVID-19 on life expectancy
2019-2023 and its decomposition: Findings from 27 countries. *SSM Popul Health*.
2024;25(101568):101568. doi:10.1016/j.ssmph.2023.101568

13.   Bilal U, Hessel P, Perez-Ferrer C, et al. Life expectancy and mortality in 363 cities
of Latin America. *Nat Med*. 2021;27(3):463-470. doi:10.1038/s41591-020-01214-4

14.   Mostert S, Hoogland M, Huibers M, Kaspers G. Excess mortality across countries
in the Western World since the COVID-19 pandemic: 'Our World in Data'
estimates of January 2020 to December 2022. *bmjph*. 2024;2(1):e000282.
doi:10.1136/bmjph-2023-000282

15.   Serfling RE. Methods for current statistical analysis of excess pneumonia-
influenza deaths. *Public Health Rep*. 1963;78(6):494-506. doi:10.2307/4591848

16.   Choi K, Thacker SB. An evaluation of influenza mortality surveillance, 1962-
1979. II. Percentage of pneumonia and influenza deaths as an indicator of influenza

activity. *Am J Epidemiol*. 1981;113(3):227-235.

doi:10.1093/oxfordjournals.aje.a113091

17.  Simonsen L, Clarke MJ, Williamson GD, Stroup DF, Arden NH, Schonberger LB.

The impact of influenza epidemics on mortality: introducing a severity index. *Am J*

*Public Health*. 1997;87(12):1944-1950. doi:10.2105/ajph.87.12.1944

18.  Farrington CP, Andrews NJ, Beale AD, Catchpole MA. A statistical algorithm for

the early detection of outbreaks of infectious disease. *J R Stat Soc Ser A Stat Soc*.

1996;159(3):547. doi:10.2307/2983331

19.  Noufaily A, Enki DG, Farrington P, Garthwaite P, Andrews N, Charlett A. An

improved algorithm for outbreak detection in multiple surveillance systems. *Stat*

*Med*. 2013;32(7):1206-1222. doi:10.1002/sim.5595

20.  Lee WE, Woo Park S, Weinberger DM, et al. Direct and indirect mortality impacts

of the COVID-19 pandemic in the United States, March 1, 2020 to January 1,

2022. *Elife*. 2023;12:e77562. doi:10.7554/eLife.77562

21.  Lee RD, Carter LR. Modeling and forecasting U.s. mortality. *J Am Stat Assoc*.

1992;87(419):659-671. doi:10.1080/01621459.1992.10475265

22.  Wilmoth JR. Variation in vital rates by age, period, and cohort. *Sociol Methodol*.

1990;20:295-335. doi:10.2307/271089

23.  Clark SJ. A general age-specific mortality model with an example indexed by child

mortality or both child and adult mortality. *Demography*. 2019;56(3):1131-1159.

doi:10.1007/s13524-019-00785-3

24.  Alexander M, Zagheni E, Barbieri M. A flexible Bayesian model for estimating

subnational mortality. *Demography*. 2017;54(6):2025-2041. doi:10.1007/s13524-

017-0618-7

25.  National Institute of Population and Social Security Research. The Japanese

Mortality Database. December 5, 2023. Accessed October 24, 2024.

https://www.ipss.go.jp/p-toukei/JMD/index-en.asp

26.  Ministry of Health, Labour and Welfare. Annual Report on the Vital Statistics of
Japan, 2011 (Confirmed Data). September 6, 2012. Accessed October 24, 2024.
https://www.mhlw.go.jp/toukei/saikin/hw/jinkou/kakutei11/

27.  Raikov D. On the decomposition of Poisson laws. *Dokl Akad Nauk SSSR*.
Published online 1937.

28.  Preston S, Heuveline P, Guillot M. *Demography*. Wiley-Blackwell; 2000.

29.  R: The R Project for Statistical Computing. 2023. Accessed September 4, 2024.
https://www.r-project.org/

30.  Gabry J, Češnovar R, Johnson A, Bronder S. R Interface to CmdStan. 2023.
Accessed September 4, 2024. https://mc-stan.org/cmdstanr/

31.  Munira MS, Okada Y, Nishiura H. Life-expectancy changes during the COVID-19
pandemic from 2019-2021: estimates from Japan, a country with low pandemic
impact. *PeerJ*. 2023;11:e15784. doi:10.7717/peerj.15784

32.  Okada Y, Nishiura H. Changes in life expectancy and life span equality during the
COVID-19 epidemic in Japan up to 2022. *medRxiv*. Published online April 19,
2024:2024.04.18.24306049. doi:10.1101/2024.04.18.24306049

**Figure 1. Changes in mortality rates by age group during 2000–2022.**

Panel (A) shows mortality schedule curves in 2020–2022 in comparison with those in 2000, 2010, and 2019. Panel (B) provides a closer look at the change in mortality rates by age group in 2020–2022 compared with that in 2019.

**Figure 2. Gaps between observed and projected mortality rates by age group, 2020–2022.**

The red points with error bars show the median and 95% prediction intervals of the gaps between observed and projected (log-scaled) mortality rates in (A) 2020, (B) 2021, and (C) 2022.

**Figure 3. Heatmaps of the gap between observed and projected mortality rates by prefecture in 2020–2022.**

The cells represent gaps between the observed and the median of projected (log-scaled) mortality rates in (A) 2020, (B) 2021, and (C) 2022 for all prefectures. Cells with colors closer to dark blue show mortality rates that were lower than expected (projected) whereas those closer to solid red indicate mortality rates that were higher than expected.

**Figure 4. Gaps between life expectancy at birth calculated from observed and projected mortality rates in 2020–2022.**

Gaps between observed and projected life expectancy at birth are shown in (A) 2020, (B) 2021, and (C) 2022 for all prefectures. In each panel and for each prefecture, red points and error bars show the median and 95% prediction interval, respectively.

**Figure 5. Gaps between observed and projected death counts per 100,000 population in 2020–2022.**

Gaps between observed and projected death counts per 100,000 population are shown in (A) 2020, (B) 2021, and (C) 2022 for all prefectures. In each panel and for each prefecture, red points and error bars show the median and 95% prediction interval, respectively. Green hollow diamonds represent the average yearly aggregated gap between observed and projected death counts provided in "Excess and Exiguous Deaths Dashboard in Japan."[9]

# (A) Mortality Rate (log-scale) by Age Groups

# (B) Change in Mortality Rate by Age Groups since 2019

**(A) Gap in Log-scaled Mortality Rates, 2020**
Observed vs Projected

**(B) Gap in Log-scaled Mortality Rates, 2021**
Observed vs Projected

**(C) Gap in Log-scaled Mortality Rates, 2022**
Observed vs Projected

(A) Gap in Life Expectancy, 2020
Observeved and Projected

(B) Gap in Life Expectancy, 2021
Observeved and Projected

(C) Gap in Life Expectancy, 2022
Observeved and Projected

**(A) Mortality Gap vs Projected, 2020**   **(B) Mortality Gap vs Projected, 2021**   **(C) Mortality Gap vs Projected, 2022**