**EVALUATING THE PREVALENCE OF HUMAN AND ANIMAL AFRICAN**

**TRYPANOSOMIASIS IN NIGERIA: A SCOPING REVIEW**

Chinwe Chukwudi[1*], Elizabeth Odebunmi[2], Chukwuemeka Ibeachu[2],

[1]Department of Veterinary Pathology and Microbiology, University of Nigeria Nsukka. Nsukka,

Enugu State, Nigeria.

[2]Department of Public Health, School of Health and Society, University of Wolverhampton,

Wolverhampton, UK

[*]Correspondence: chinwe.chukwudi@unn.edu.ng, cuchukwudi@gmail.com

**Simple Summary:** Human African Trypanosomiasis (HAT) is a zoonotic NTD targeted for

elimination in 2030. There is a scarcity of HAT epidemiological studies/data from Nigeria.

WHO records do not reflect the published data on HAT prevalence/cases in Nigeria. But the

disease remains prevalent, with a complex epidemiology and zoonotic potential. Inadequate

surveillance and diagnostics challenges limit HAT reporting from Nigeria

**ABSTRACT**

African trypanosomiasis is a protozoan disease that affects both humans and animals. Human

African Trypanosomiasis (HAT) is a Neglected Tropical Disease targeted for elimination in

2030. Although WHO has not reported HAT from Nigeria in the last decade, there are published

studies reporting seroprevalence, parasite detection/isolation, and animal carriers for HAT in

Nigeria. Interestingly, the burden of Animal African Trypanosomiasis (AAT) continues to

increase. In this study, we synthesized published reports on the prevalence of HAT and AAT in

Nigeria from 1993-2021, the trypanosome species involved, the spread of animal reservoirs, and

the variability in diagnostic methodologies employed. A scoping review was performed

following the methodological framework outlined in PRISMA-ScR checklist. Sixteen eligible

studies published between 1993 and 2021 were reviewed: 13 for AAT, 3 for HAT. Varying

prevalence rates were recorded depending on the diagnostic methods employed. The average

prevalence reported from these studies was 3.3% (HAT), and 27.3% (AAT). Diagnostic methods

employed include microscopy, PCR and Card Agglutination Test for Trypanosomiasis (CATT).

Cattle, pigs, and dogs were identified as carriers of human-infective trypanosomes. This study

highlights the scarcity of HAT epidemiological studies/data from Nigeria, the high prevalence,

complex epidemiology, inadequate surveillance and utter neglect of African Trypanosomiasis in

Nigeria. Remarkably, WHO records do not reflect the published data showing evidence of HAT

prevalence/cases in Nigeria. Unfortunately, diagnostics challenges and unrealistic disease

reporting protocols seem to limit HAT reporting from Nigeria. Therefore, adequately

coordinated epidemiological surveys and targeted intervention policies are imperative to

ascertain the true epidemiological status of HAT in Nigeria and prevent disease re-emergence

towards achieving WHO's elimination targets. The presence of animal carriers of human-

infective trypanosomes underscores the importance of a one-health approach to combat African

trypanosomiasis effectively.

Keywords: African trypanosomiasis, HAT, Trypanosoma brucei, prevalence, animal reservoir,

Neglected Tropical Disease.

## BACKGROUND

African Trypanosomiasis (AT) is one of the Neglected Tropical Diseases (NTDs) with an estimated 60 million people at risk of infection in sub-Saharan Africa, [1,2] as well as various animals such as dogs, cattle, pigs, sheep and goats. [3, 4] This vector-borne disease, caused by a protozoan parasite of the genus Trypanosoma, is transmitted by tsetse flies to both humans and animals. [5] Two sub-species of *Trypanosoma brucei* cause Human African Trypanosomiasis (HAT), also known as sleeping sickness. *Trypanosoma brucei gambiense* (Tbg), found in 24 countries in West and Central Africa, is responsible for 97% of all reported cases, and causes a chronic disease with neurologic symptoms. *Trypanosoma brucei rhodesiense* (Tbr), which is found in 13 countries in East and Southern Africa, is responsible for 3% of all reported cases and causes an acute disease. [10] Unlike Tbg, which primarily infects humans (although some animal reservoirs/carriers have been identified), Tbr is considered zoonotic, causing disease in both humans and animals. [11] Animal African Trypanosomiasis (AAT), also known as Nagana, is caused by various trypanosome species, including *T. brucei, T. congolense, T. vivax, T. equiperdum, T. evansi, T. simiae, T. suis, and T. theileri*. [6, 7] While cattle are the most affected, other animals such as goats, dogs, sheep, pigs, and wild animals are also susceptible. [8] *T. equiperdum* causes a venereal disease (Dourine) in horses and donkeys, while *T. evansi*, causes a form of trypanosomiasis known as Surra in horses, camels, buffaloes, mules, and deer. [9]

The transmission of both HAT and AAT heavily depends on tsetse fly (*Glossina sp.*), which is exclusively found in sub-Saharan Africa. [12] Nevertheless, maternal and sexual transmission has been indicated for HAT, although their exact contributions to the epidemiology of the human disease has not been fully explored. [13] The coexistence of humans, vectors, and parasites in a conducive environment in the tropical trypanosomiasis belt of Africa, coupled with the presence

of animal reservoirs/hosts, makes African trypanosomiasis a significant public health challenge,

with detrimental effects on health, the economy, poverty levels, and agricultural productivity.[14, 15]

Reported cases of HAT have substantially declined globally to fewer than 1,000 cases between

2019 and 2020.[10] This decline is attributed to HAT control strategies and inclusion in the

Neglected Tropical Diseases, with a roadmap aiming to eliminate gHAT as a public health

concern by 2020 and zero transmission to humans by 2030.[16] However, disease monitoring and

epidemiological surveillance is grossly inadequate in many endemic countries.[17] For Nigeria,

WHO records indicate that HAT was last reported from Nigeria in 2012 (except for one case that

was diagnosed in a Nigerian in the UK in 2016).[18] This has resulted in the neglect of the country

in HAT surveillance and control programs, since it is believed that the country is on course to

eliminate the disease. However, Cameroon which shares borders with the Nigerian HAT foci

continues to report cases (Fig. 1). Interestingly, there has been reports of finding the human-

infective trypanosomes in animals in Nigeria, especially in companion animals (dogs). This

raises questions about the existence and true prevalence of HAT in Nigeria, especially with

regards to the source of infection and transmission route for the animal carriers. Ignoring or

neglecting any HAT foci or infection site, no matter how small it is, has grave consequences for

the achievement of the NTD elimination targets. Such locations could serve as sources of

continuous re-infection and may become possible breeding sites for the next epidemic strain of

the parasite, leading to disease re-emergence. Therefore, adequate epidemiological information

about the co-circulation of human and animal trypanosomes is critical to inform control or

eradication policies and strategies. Hence, this study examined published evidence on the

prevalence of African trypanosomiasis in both humans and animals in Nigeria, with a view to

understanding the occurrence, host spread, parasite species involved, diagnostic techniques being

used, and the possibility of cross infection between humans and animals.

Fig 1: WHO HAT prevalence records for Nigeria and neighbouring Cameroon. Assessed from

https://www.who.int/data/gho/data/themes/topics/human-african-trypanosomiasis. Red dot

inside the map represents major known HAT foci.

**METHODS**

This study followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

(PRISMA-ScR) four-stage framework, encompassing identification, screening, eligibility

assessment, and inclusion to ensure a transparent, accurate, and high-quality reporting of the

review's findings. [19]

**Search Strategy**

The searches were performed using MEDLINE, CINAHL, and EMBASE in July 2023. The

search terms included 'Neglected Tropical Disease,' 'African trypanosomiasis,' 'human African

trypanosomiasis,' 'animal African trypanosomiasis,' 'sleeping sickness,' 'nagana,'

'trypanosoma,' 'trypanosomosis,' 'trypanosomes.' These terms were combined using Boolean

operators 'OR', as well as 'AND' to retrieve relevant articles. A manual search was also

performed to ensure that all relevant studies were captured.

**Eligibility Criteria**

The eligibility criteria for this review were informed by participants (P), phenomena of interest (I), and context (Co) or PICo, where P is humans/animals with AT, I is African trypanosomiasis infection and Co is Nigeria. Primary research studies including cross sectional, cohort, surveys, and prevalence studies that reported animal and human African trypanosomiasis infection were included. Specific inclusion criteria included studies from Nigeria, published in English language with well detailed sample size, prevalence of trypanosomiasis in sampled population, method of diagnosis and *Trypanosoma* species identified.

**Study Selection**

Study selection was performed independently by the three authors. Discrepancies between the authors were resolved through discussion until a consensus was reached. After duplicates were removed, the remaining studies were screened based on their titles and abstracts. Subsequently, non-relevant studies were excluded, and the full text of the remaining studies were assessed according to the pre-defined eligibility criteria. Finally, studies that met the eligibility criteria were included for further data extraction.

**Data Extraction**

Studies were extracted into a pre-prepared Microsoft Excel spreadsheet with the following variables: author and year of publication, study subject (human, animal), type of study design, study size, *Trypanosoma* species identified, diagnostic method used, number of positive cases, prevalence, or percentages.

**Data Synthesis**

A narrative summary was performed to synthesize findings from this review.

**RESULTS**

A total of 1,977 articles were retrieved from the search of three databases. 59% (1,159) from

MEDLINE, 36% (724) from EMBASE and 5% (94) from CINAHL. After removal of 40

duplicates, 1,937 articles were screened for titles and abstracts. A total of 1,787 non-relevant

articles were excluded in accordance with the predetermined criteria. Following full text

retrieval, another 25 studies were excluded as their full texts were not accessible. Upon review of

the full texts of the remaining 125 articles, a total of 11 were deemed eligible for inclusion in the

study. An additional 5 articles were identified from a search of reference lists. Finally, 16 articles

were included. The study selection process is represented in table I.

**Description of eligible studies**

All included studies were published between 1993 and 2021, with the number of study subjects

ranging from 19 to 7,143 and an overall sample size of 18,091. Of the 16 included studies, three

investigated human African trypanosomiasis,[20 , 21, 22] while thirteen studies reported animal

African trypanosomiasis. Five of these reported infection in cattle - bovine trypanosomiasis.[23,24,]

[25, 26,27] Two studies investigated trypanosomiasis infection in pigs – porcine trypanosomiasis. [28,]

[29] Three studies investigated infection in ruminants – sheep, goat, cattle, [30,31,32] while one study

investigated in dogs – canine trypanosomiasis.[33] The remaining two studies investigated

infection in multiple animal hosts including monkey. [34, 35] Table II provides a summary of the

included studies.

**Prevalence of HAT**

A total of 1,974 individuals underwent testing for the disease across three studies. Among them,

65 individuals were seropositive for the disease in two of the studies using the CATT kit. [20,21]

Trypanosomes were detected from the blood of 23 seropositive individuals by various methods

such as microscopic examination of blood films or buffy coat, and in vivo inoculation into

mice/rats. The detection of the parasites in the blood did not corelate with how strongly positive

the CATT test was, as parasites were detected in patients that were weakly, moderately and

strongly positive in the CATT test. One study also examined the CSF, and detected parasites in

the CSF of 4 patients. The third study, conducted in the northern part of the country, did not

detect any positive case within the sampled population.[22] The seroprevalence of HAT reported

by these studies for various study sites across the country ranged from 0% – 9.6%, with a

calculated average of 3.3% (65/1,974). The actual prevalence of HAT as confirmed by parasite

detection was as high as 4.8% in one study site (Urhouka), with a calculated average of 1.2%

(23/1974) across all studies. *Trypanosoma brucei gambiense* type 1was reported as the causative

parasite specie, as detected by the gambiense-specific TgsGP-PCR.

**Prevalence of AAT**

Thirteen included studies reported AAT with an overall sample size of 16,117. The highest

sampled animal was cattle at 12,193, while the least is monkey at 1 sample. Others are 577

sheep, 418 goat, 2,639 pigs, 31 dogs, and 318 small ruminants (unspecified). Out of the total

sampled animals, 4,404 were found positive for trypanosomiasis infection from all the 13

studies. The prevalence of AAT as reported in these studies ranged between 0.8% - 46.8% for

various animal species in various study sites, with a calculated mean prevalence of 27.3%

(4,404/16,117).

These studies detected the animal trypanosomes: *T. vivax, T. congolense, T. brucei, T. simiae, T.*

*evansi, T. theileri* in the studied samples, with *T. vivax*, *T. congolense,* and *T. brucei* report as the

most prevalent trypanosome species in animals. In addition to these trypanosomes, three of these

studies detected the human infective parasite *Trypanosoma brucei gambiense* in animals. [29, 33, 34]

**Diagnostic Techniques**

The included studies for this review utilized various diagnostic methods, including microscopy,

Card Agglutination Test for Trypanosomiasis (CATT), and Polymerase Chain Reaction (PCR) in

the detection of trypanosomes. Overall, microscopy was the most utilized diagnostic technique in

these studies, especially in animals. CATT and PCR were also utilized individually, together or

in addition to microscopy for humans.

**DISCUSSION**

The findings from this review, provides insights into the prevalence, host range, species

distribution, and diagnostics techniques of African trypanosomiasis in humans and animals in

Nigeria. According to this study's findings, the calculated mean prevalence of AAT in Nigeria is

27.3% (4,404 out of 16,117), while that of HAT is 1.2% by parasite identification and 3.3%

seroprevalence, indicating that African trypanosomiasis is still endemic in both humans and

animals in Nigeria. The presence of various trypanosome species and use of different diagnostic

methods contributes to the complexity of this disease, its epidemiological studies in Nigeria and

its public health significance. The study identified several trypanosomes, including *T. vivax, T.*

*congolense, T. brucei brucei, T. simiae, T. evansi, T. theileri* and *T. brucei gambiense* as the

causative pathogens of African trypanosomiasis in animals and humans in Nigeria. Additionally,

the most used diagnostic technique in Nigeria for detecting these trypanosomes is microscopic

examination, particularly in animals. Bearing in mind that gHAT is characterized by extremely

low parasitaemia, and hence, very difficult to detect by microscopic examination of blood

smears, the identification of *T. b gambiense* by microscopy in the few studies available indicate

that the prevalence of HAT in the country might be underestimated.

**The Prevalence of human African trypanosomiasis in Nigeria**

Despite the World Health Organization reporting only eight new HAT cases in Nigeria between

2010 and 2016,[36] the results of this analysis present a different perspective. During this period,

two of the studies included in this analysis collectively identified 65 HAT seropositive

individuals out of a total of 1,974 assessed, with the parasites isolated from 7 individuals in one

study and detected in the blood and CSF of 16 and 4 individuals respectively in another study.

[20,21] This means that in the same period when WHO data recorded 8 cases, there were 23

confirmed cases reported in research studies. This notable difference suggests that the actual

prevalence of HAT in Nigeria might be grossly under-reported. This is further buttressed by the

fact that the only reported case from 2013 was diagnosed in the UK, indicating that many clinical

cases may be escaping diagnosis in Nigeria due to lack of diagnostic capacity/resources. This

could be attributed to the challenges of clinical HAT diagnosis especially in the resource-limited

settings of the disease foci in the country, and non-inclusion of HAT into routine medical checks

in Nigeria's healthcare facilities.[21] More importantly, there is very little HAT surveillance going

on in the country, and when done, are restricted to the previously known HAT foci. The focal

nature of the disease has made the sparse surveillance efforts to be limited to the traditionally

known foci, while totally neglecting the vast majority of the country which are prone to disease

expansion due to climate change and human/animal migration. Considering that some of the

studies reported silent infections in human carriers who were apparently healthy, and that

microscopic detection of the parasites in HAT is quite challenging due to extremely low

parasitemia and difficulty in obtaining CSF, it is apparent that the true burden of HAT in Nigeria

is largely unknown. Prevalence studies are grossly inadequate in Nigeria (as only 3 eligible

studies were found within the period reviewed), despite being an endemic country. This

underscores the limited attention that HAT receives in some endemic regions, especially Nigeria.

[37]

Human African trypanosomiasis has been included in the Neglected Tropical Diseases Road

map, aiming to eliminate HAT as a public health concern by 2020 and interrupt the transmission

of *Trypanosoma brucei gambiense* (Tbg) to humans by 2030.[16] The criteria for elimination are

set at fewer than 2,000 reported cases annually and no more than 1 case per 10,000 residents in

areas at moderate or high risk.[17] Several endemic African countries, such as Benin, Côte d'Ivoire,

Equatorial Guinea, Togo, and Uganda, have successfully eliminated gambiense Human African

Trypanosomiasis (gHAT) as a public health problem.[10] However, Nigeria still faces challenges in

achieving this goal. The prevalence rates of HAT from these studies in Nigeria are much higher

when compared to 0.06% in Côte d'Ivoire and 0.88% in Uganda.[38,39] With infection rates

(seroprevalence) as high as 9.6%, parasites detected in blood and CSF in 4.8% and 1.8%

respectively, and parasites isolated from 1.5% of the population in specific study sites in the

country [20 , 21], these reviewed studies suggest that Nigeria may have a higher

incidence/prevalence rate than is being reported by WHO. The disparity in the WHO reported

data and published research data indicates that WHO reports may not necessarily reflect the

actual disease status in Nigeria, and therefore could be misleading. This is mostly due to the lack

of HAT surveillance and monitoring in several parts of the country. These findings suggest that

the disease may still pose a significant health concern in the country, but surveillance and

reporting are grossly inadequate. In view of the possibility of genetic admixture with sympatric

animal parasites that may lead to new parasites clones and disease/epidemic re-emergence, this

situation could present a huge drawback to the planned gHAT elimination targets by 2030.

**Prevalence of AAT in Nigeria**

This study has indicated a calculated mean prevalence rate of 27.3% for animal African trypanosomiasis, which is higher than the 16.1% in previous review.[40] This indicates increasing AAT burden. A notable finding from this study is the presence of the human-infective *T. brucei gambiense* in pigs, dogs, and cattle in Nigeria.[29,33,34] Similar findings have been documented in various sub-Saharan African countries.[41,42,43] This raises additional concerns regarding the potential role of animals in the epidemiology of HAT, and the possibility of HAT re-emergence with new epidemic clones of the parasite. Although the precise role of these animal hosts/carriers in gHAT transmission is not entirely clear[44], their presence poses a potential threat to the goal of eliminating HAT transmission to humans by 2030.[45] While the zoonotic potential of *T. brucei gambiense* has been a subject of significant discussion over the years, the confirmation of its presence in animal hosts highlights the possibility of transmission between humans and animals that should not be neglected. The existence of these animal reservoirs in Nigeria, could contribute to the persistence and potential spread of Human African Trypanosomiasis (HAT) beyond the known foci. One of the prerequisites for a disease to be eradicable is the absence of animal reservoirs.[46] Therefore, in addition to undetected human infections and inadequate diagnostic tools/capacity, the presence of potential animal reservoir hosts is a major challenge to the complete elimination of HAT in Nigeria and across sub-Saharan Africa.[47, 48] Without addressing the knowledge gaps regarding *T. b. gambiense* reservoirs, complete gHAT elimination may be unattainable by 2030.[45] The complex genetic and immunologic interactions in humans and animals co-hosting various parasite species/subspecies and consequent changes in disease transmission dynamics in these sites could potentially present conducive breeding grounds for new epidemic strains of the parasites. Therefore, further research to gain a

comprehensive understanding of the specific role played by these animal reservoirs in the

transmission of HAT is crucial.

**The Diagnosis of Trypanosomiasis**

The diagnosis of trypanosomiasis relies on various methods like parasitological, serological, and

molecular tests, each with different levels of accuracy, ease of use, and cost.[8] For instance, less

sensitive and less expensive diagnostic procedures such as microscopy are more likely to give

false negative results, leading to an underestimation of the true prevalence of the disease.

Conversely, more sensitive and more expensive diagnostic tests like PCR can detect more

positive cases that might be missed by less precise methods, resulting in higher reported

prevalence rates.[49,50], but may not be easily accessible/affordable. Consequently, the choice of

diagnostic methods and their accuracy can significantly affect the detection of infected

individuals or animals and the reported prevalence rates.[51]

Microscopic examination was found to be the most used method for identifying trypanosomiasis

infections in both animals and humans in Nigeria. However, this method has faced criticism for

its low sensitivity, especially in cases with low parasitemia characteristic of human infections

with TbG, which can result in the underestimation of HAT prevalence rates.[52] On the other hand,

Polymerase Chain Reaction has been acknowledged as the preferred diagnostic method for

epidemiological investigations concerning *Trypanosoma sp*. due to its high sensitivity and

capacity for processing many samples.[53] This aligns with the results of this study, where the use

of both microscopy and PCR on the same group of samples revealed that microscopic

examination detected a lower prevalence rate compared to PCR.

As observed in one of the studies, the prevalence of AAT in cattle using microscopy was 15.1%

while PCR detected a higher prevalence of 63.7%.[27] This substantial gap in prevalence rates can

be attributed to the varying sensitivities of the two methods. Similarly, when examining *T. b.*

*gambiense* infections in humans using the Card Agglutination Test for Trypanosomiasis (CATT)

and PCR, different prevalence rates were detected.[21] In one study, CATT showed an overall

prevalence rate of 1.8%, whereas PCR indicated a lower rate of 0.6% among the same sampled

population of 1200 individuals. This discrepancy in prevalence rates between the two diagnostic

techniques underscores the higher specificity of PCR and the potential for false-positive results

from CATT due to cross-reactivity with antibodies against other endemic protozoan diseases

such as malaria.[54] On the other hand, there exists a substantial likelihood of encountering false

negative results with CATT, primarily attributable to discrepancies in antigen types utilized in

the CATT assay kit especially in Nigeria and Cameroon. [55,56]

The variation in prevalence rates has also been linked to the capacity of diagnostic tools to

identify small amounts of trypanosomes in infected samples. Unlike the PCR method, which has

the capability to detect very small quantities of infecting parasites,[42] the microscopy technique is

limited in its ability to detect trypanosomes in the characteristically low parasitaemia in gHAT,

and to distinguish between different species and subspecies.[33] In this review, the polymerase

chain reaction approach was successful in detecting the human-infective parasite *T. b. gambiense*

in animals.[33,34]

In summary, the choice of diagnostic methods significantly impacts the reported prevalence rates

of trypanosomiasis. This emphasizes the importance of carefully considering the methods used

when interpreting and comparing prevalence data across different studies and locations.

However, these choices are often influenced by factors like the availability of adequately skilled

personnel, cost, ease of use, and useability of the diagnostic tools.[57] Nigeria, being a developing

nation, faces challenges related to limited resources, funding, and healthcare personnel. This may

explain why microscopy is frequently used for trypanosomiasis detection in Nigeria compared to

other methods as it requires minimal equipment and is a cost-effective way to detect the disease.[8]

**Conclusions**

This study highlights the variably high prevalence rate of African trypanosomiasis infection in

both humans and animals in Nigeria, thereby preenting evidence of the persistent threat that the

disease poses to the health of humans and animals. The identification of pigs, dogs, and cattle as

carriers for the human-infective parasite *Trypanosoma brucei gambiense* underscores the

potential for transmission between humans and animals and cross-species hybridization, thus

highlighting the zoonotic potential of *gambiense* Human African Trypanosomiasis (gHAT). This

presents a significant challenge to the 2030 objective of interrupting gHAT transmission to

human populations. To enhance the efficacy of gHAT control measures, additional research is

crucial to elucidate the precise roles played by animal reservoirs or carriers, and establish more

effective diagnostic techniques for the detection of the trypanosomes in low resource endemic

areas.

**Table I: Flow chart of study selection process**

**Table II: Summary of the characteristics of the reviewed articles**

| Author/Year | Title | Aim | City/Area | Study Design | Study Population | Sample Size | Outcome Measured | Result | Prevalence rate | Diagnostic Method | Species Identified |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Danie et al., 1993 | Prevalence of bovine trypanosomiasis in Gongola State of Northern Nigeria | To assess the prevalence of bovine trypanosomosis in Karim Lamido and Numan local government areas of Gongola State. | Adamawa: Formerly Gongola (Karim Lamido and Numan LGAs) | Cross-sectional Survey | Cattle | 1065 | Bovine trypanosomiasis | A total of 42 (3.9 %) Cattle were found to be infected with trypanosomes. Out of which 27 (64.3 %) were due to *T. vivax*, 13 (31 %) to *T. congolense* and 2 (4.8 %) to *T. brucei.* | 3.90% | Microscopy | *T. vivax* *T. congolense* *T. brucei* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Danie et al., 1994 | Prevalence of trypanosomiasis in sheep and goats in a region of northern Nigeria | To investigate the sensitivity of four techniques currently in use for the parasitologic al diagnosis of trypanosomosis. | Bauchi (Alkaleri and Gombe LGAs) | Cross-sectional Survey | Sheep Goat | 615 (258 sheep and 357 goats) | Animal African trypanosomiasis | A total of 19 (7.4%) sheep and 18 (5.0%) goats were positive giving a total infection rate of 37 (6.0%), 22 being positive with *T vivax*, 9 with *T. congolense* and 6 with *T. brucei* | 6% | Microscopy | *T. vivax* *T. congolense* *T. brucei* |
| Enwezor et al., 2009 | Survey of bovine trypanosomosis in the | To assess the prevalence of trypanosomes in cattle at | Kaduna | Cross-sectional Survey | Cattle | 1293 | Bovine trypanosomiasis | A total of 109 cattle were infected with trypanosomes giving an overall prevalence | 8.40% | Microscopy | *T. vivax* *T. congolense* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Kachia Grazing Reserve, Kaduna State, Nigeria | the Kachia Grazing Reserve (KGR) | | | | | | of 8.4%. Out of these, 105 (96.3%) was due to *T. vivax*, 2 (1.9%) to *T. congolense*, 1 (0.9%) to *T. b. brucei* and a mixed infection of *T. congolense* and *T. vivax* 1 (0.9%). | | | *T. brucei brucei* |
| Enwezor et al., 2019 | Investigation of livestock for presence of Trypanosoma brucei gambiense | Aimed at screening livestock for possible presence of T. b. gambiense within Tafa | Niger | Cross-sectional Survey | Cattle Sheep Goats Local dogs Monkey | 460 animals (177 cattle, 209 sheep, 61 goats, | Animal African trypanosomiasis | A total of 10 animals were positive for trypanosomes with an overall prevalence of 2.17 %. The morphological identification indicated the presence | 2.17% | Microscopy | *T. brucei* *T. congolense* *T. vivax* |

| | In Tafa Local Government Area of Niger State, Nigeria | Local Government Area (LGA) of Niger State, Nigeria. | | | | | 12 local dogs and 1 monkey) | | of *T. brucei, T. vivax* and a mixed infection of *T. brucei* and *T. congolense*. Details by animal species showed 3 cattle (1.69 %), 6 sheep (3.02 %) and 1 goat (1.64 %) infected with trypanosomes. Neither the dogs nor the monkey was positive for trypanosomes. | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Habee ) et al., | Molecular identification and | To investigate the | Kwara | Cross-section al | Cattle | 398 | Bovine trypanos omiasis | A total of 3 Cattle were positive by microscopy, | Microsco py: 0.8% PCR: | PCR Microsco py | *T. theileri*□ *T. evansi*□ |

| 2021 | prevalence of trypanosomes in cattle distributed within the Jebba axis of the River Niger, Kwara state, Nigeria | prevalence of trypanosome infection in cattle, and molecularly identified the species of trypanosomes in infected cattle and the spatial distribution of trypanosome-infected herds along | | Survey | | | | representing 0.8% prevalence, while 12 samples representing 3.0% tested positive by nested PCR. With *T. congolense* more prevalent (50.0%). | 3.0% | | *T. simiae* *T. congolense* *T. brucei* *T. vivax* |

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | the Jebba axis of the River Niger. | | | | | | | | |
| Kalu et al., 996 | Observations on the epidemiology of ruminant trypanosomosis in Kano State, Nigeria | To evaluate the prevalence of trypanosomosis among ruminants in Kano State and to elucidate aspects of the disease transmission in the area | Kano | Cross-sectional Survey | Ruminants (Cattle, sheep, goat) | 1424 (1106 cattle and 318 small ruminants) | Animal African trypanosomiasis | 59 cattle were infected with trypanosomes giving a prevalence of 5.3% with a confidence interval of ±1.3. Three out of 318 small ruminants sampled were infected (prevalence 0.9% ± 1.0). | Cattle: 5.3% (95% CI: 4.0–6.6%)Small ruminants (95% CI: -0.1 – 1.9%) | Microscopy | *T. vivaxT. congolense T. brucei* |

| | | To conduct an active screening of T. b. gambiense in humans in the old Gboko sleeping sickness focus in Nigeria and characterized isolates using TgsGP-polymerase | | | | | | Out of sampled people, a total of 28 were found positive. The CATT revealed an overall infection rate of 1.8% of the 1200 samples studied. PCR revealed an overall infection rate of 0.6% of the 1200 samples analysed. Tryypanosmes (TbG) was isolated from 7 of the samples | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Karshma et al., 2016 | Silent Human Trypanosoma brucei gambiense Infections around the Old Gboko Sleeping Sickness Focus in Nigeria | | Gboko | Cross-sectional Survey | Humans | 1200 | Human African trypanosomiasis | | CATT: 1.8% Parasite isolation/ PCR: 0.6% | CATT PCR | *T. b. gambiense* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | chain reaction. | | | | | | | | | |
| Karshima et al., 2016 | Animal reservoirs of Trypanosoma brucei gambiense around the old Gboko sleeping sickness focus in Nigeria | To ascertain the possible role of animal reservoirs in the epidemiology of the parasite in the old Gboko sleeping sickness focus in | Gboko | Cross-sectional Survey | Cattle Pigs | 1200 (600 Cattle, 600 Pigs) | Animal African trypanosomiasis | A total of 118 animals were positive (46 cattle, 72 pigs). The overall infection rates for the CATT and TgsGP-PCR were 8.9 and 0.9%, respectively. Trypanosomes of animal origin identified by ITS 1 PCR were *T. brucei* (4.2%), *T. congolense* forest (3.2%), *T.* | CATT: 8.9% PCR: 0.9% | CATT PCR | *T. brucei (T. b. gambiense ) T. congolense forest T. congolense savannah T. vivax* |

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | Nigeria and characterized isolates using TgsGP polymerase chain reaction. | | | | | | *congolense* savannah (2.0%), *T. vivax* (2.2%) and mixed infections (1.5%) in cattle as well as *T. brucei* (4.8%), *T. congolense* forest (1.8%), *T. congolense* savannah (1.0%) and mixed infections (1.2%) in pigs. *T. brucei gambiense* and other animal trypanosomes were identified among animals in the focus, | | | |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | indicating the existence of animal reservoirs of human infective T. b. gambiense. | | | |
| Majekodunni et al., 2013 | A longitudinal survey of African animal trypanosomiasis in domestic cattle on the Jos Plateau, Nigeria: | To determine seasonal variations in the prevalence of AAT across the Jos Plateau | Jos | Cross-sectional Survey | Cattle | 7143 | Bovine trypanosomiasis | 3, 342 cattle were found positive. The prevalence of bovine trypanosomiasis was found to be high at 46.8% (39.0 – 54.5%) and significant, seasonal variation was observed between the dry and the end of the wet season. T. b. | 46.8% (39.0 – 54.5%) | PCR | *T. congolense* *T. vivax* *T. b. brucei* |

| Author | Title | Objective | Location | Study Design | Host | Sample Size | Disease | Results | | Method | Species |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Prevalence, distribution, and risk factors | | | | | | | *brucei* was observed at a prevalence of 3.2% (1% – 5.5%); *T. congolense* at 27.7% (21.8% - 33.6%) and *T. vivax* at 26.7% (18.2% - 35.3%). | | | |
| Nmori et al., 2010 | Human African trypanosomiasis in endemic focus of Abraka, Nigeria | To investigate the prevalence of human African trypanosomiasis (HAT), caused by | Abraka, Delta | Cross-sectional Survey | Humans | 474 | Human African trypanosomiasis | Of the 474 screened, 44(9.3%) were seropositive with seroprevalence of 22(9.6%) in Urhouka, 14(9.5%) in Umeghe and 8(7.9%) for Ugonu. | CATT: 9.30% Microscopy (blood): 3.4% Microscopy (CSF): | Microscopy CATT | *T.b.gambiense* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Trypanosoma brucei gambiense in an endemic focus of Nigeria, as it relates to age, sex and occupational differences. | | | | | | | 0.8% | | |
| Omek ;, .994 | Pig trypanosom osis: prevalence and significance | To determine the prevalence of trypanosome species pathogenic to | Anambra and Benue states | Cross-section al Survey | Pigs | 1954 | Porcine trypanos omiasis | 524 (26.8 %) were positive for trypanosome infections, 348 (66.5 %) of which had a mixed *T. brucei* and *T.* | 26.80% | Microsco py | *T. congolens e* *T. brucei* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | in the endemic Middle Belt zone of southern Nigeria | pigs and the significance of pig trypanosomosis in the Middle Belt zone of Southern Nigeria. | | | | | | *congolense* burden, while 125 (23.9 %) and 43 (8.2 %) others had single *T. brucei* and *T. congolense* infections respectively. | | | |
| Onah and Ebene 2004 | Isolation of a human serum-resistant Trypanosoma brucei from a | To re-evaluate the role of domestic pig as a reservoir for *T. b. gambiense* | Enugu | Cross-sectional Survey | Pigs | 85 | Porcine trypanosomiasis | 19 positive cases were identified from the 85 sampled pigs. 15 (78.96%) were identified as single infection caused by *T. brucei*. While the | 22.35% | Microscopy | *T. brucei T. congolense T. brucei gambiense* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | naturally infected pig in the Nsukka area of Enugu State | | | | | | | remaining 4 (21.05%) were due to mixed infections of *T. brucei* and *T. congolense*. | | | |
| Takee *et al.,* 2013 | Molecular survey of pathogenic trypanosomes in naturally infected Nigerian cattle | To determine the prevalence and characteristics of trypanosome species and strains in Nigerian | Ogun and Kaduna states | Cross-sectional Survey | Cattle | 411 | Bovine trypanosomiasis | Parasite detection by microscopy observation showed 62 samples infected by one or more species of Trypanosomes, for a prevalence of 15.1% (95% CI, 12–18%). PCR detection showed | Microscopy: 15.1% (95% CI, 12–18%). PCR: 63.7% (95% CI, 59.4– | Microscopy PCR | *T. brucei* *T. vivax* *T. congolense* |

| | | cattle using PCR for the first time. | | | | | | 262 samples infected by one or more species of Trypanosoma, for an overall prevalence of 63.7% (95% CI, 59.4–68.8%) and *T congolense* was the most prevalent 48.7% (95% CI, 4.2–54.3), followed by *T. vivax* 26.0% (95% CI, 21.8–31.1%) and *T. brucei* 4.4% (95% CI, 3.3–7.1%). Prevalence of mixed | 68.8%) | | |

| | | | | | | | | infections was 13.9% (95% CI, 10.6–17.4%) being co-infection by *T. congolense* and *T. vivax.* | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Uba et al., 2016 | Knowledge and prevalence of human african trypanosomiasis among residents of kachia grazing reserve, | To determine the knowledge, practices and prevalence of HAT among residents of the grazing reserve. | Kaduna | Cross-sectional Survey | Humans | 300 | Human African Trypanosomiasis | Of the 300 respondents that were examined and screened for HAT, none had palpable cervical lymph nodes enlargement, and none tested positive on CATT; hence HAT prevalence was zero. | 0.00% | CATT | |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Kachia local government area, Kaduna state, Nigeria, 2012 | | | | | | | | | | |
| Umea kuana et al., 2019 | Identificatio n of Trypanoso ma brucei gambiense in naturally infected dogs in | To determine which trypanosome species, cause canine trypanosomo sis in the Nsukka area | Nsukka | Cross-section al Survey | Dogs | 19 | Canine African Trypanos omiasis (CAT) | All 19 dogs sampled had canine trypanosomosis caused by trypanosomes of the *T. brucei* group. Two of the dogs were shown to be infected | *Tbg:* 10.52% | PCR | *T. b. gambiense T. brucei T. congolens e* |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Nigeria | of Nigeria and whether any dogs harbor the human-infective trypanosome, Tbg1. | | | | | | with the human pathogen *T. b. gambiense* Group 1 (Tbg1). | | | |
| Wayo 2017 | Prevalence of trypanosomiasis in sheep in the Kachia grazing reserve, | To update information on the prevalence of trypanosomiasis in small ruminants in the area, to | Kaduna | Cross-sectional Survey | Sheep | 110 | Animal African trypanosomiasis | A total of 45 (40.9%) animals were found positive. The trypanosomes observed were *T. congolense* (40.0%), *T. Brucei* (28.8%), *T. vivax* (17.7%) and | 40.9 | Microscopy | *T. brucei* *T. vivax* *T. congolense* |

| Kachia, Kaduna State, Nigeria | allow for proper planning of control activities and serve as valuable scientific data. | | | | | | mixed infections (13.3%) | | | |

## DECLARATIONS

**Funding:** The authors received no specific funding for this work

**Conflicts of interest:** The authors declare no conflict of interest.

**Ethics approval:** Not applicable

**Availability of data:** The dataset supporting the conclusions of this article is included within the article**.**

**Consent for publication:** Not applicable.

**Authors' contributions:** Conception/design: CC,CI; Data acquisition/analysis: EO, CI, data interpretation: CC, CI; Initial manuscript draft: EO, Substantive manuscript revision: CC; Manuscript editing and approval: EO, CI, CC.

## REFERENCES

1. Food and Agriculture Organization. The disease | Programme Against African Trypanosomosis (PAAT) | Food and Agriculture Organization of the United Nations, https://www.fao.org/paat/the-programme/the-disease/en/ (2024, accessed Apr 9, 2024).

2. Papagni R, Novara R, Minardi ML, et al. Human African Trypanosomiasis (sleeping sickness): Current knowledge and future challenges. Frontiers in Tropical Diseases 2023.

3. Balasegaram M, Balasegaram S, Malvy D, et al. Neglected diseases in the news: a content analysis of recent international media coverage focussing on leishmaniasis and trypanosomiasis. PLoS Negl Trop Dis 2008. DOI: 10.1371/journal.pntd.0000234.

4. Simarro PP, Diarra A, Ruiz Postigo JA, et al. The Human African Trypanosomiasis Control and Surveillance Programme of the World Health Organization 2000–2009: The Way Forward. PLoS Negl Trop Dis 2011. DOI: 10.1371/journal.pntd.0001007.

5. CDC. CDC - African Trypanosomiasis - Disease, https://www.cdc.gov/parasites/sleepingsickness/disease.html (2020, accessed Aug 15, 2023).

6. Stevens JR and Brisse S. Systematics of trypanosomes of medical and veterinary importance. CABI 2004. DOI: 10.1079/9780851994758.0001.

7. Steverding D. The history of African trypanosomiasis. Parasites & vectors 2008.

8. Desquesnes M, Gonzatti M, Sazmand A, et al. A review on the diagnosis of animal trypanosomoses. Parasit Vectors 2022. DOI: 10.1186/s13071-022-05190-1.

9. Ponte-Sucre A. An Overview of Trypanosoma brucei Infections: An Intense Host–Parasite Interaction. Front Microbiol 2016. DOI: 10.3389/fmicb.2016.02126.

10. World Health Organization. Trypanosomiasis, human African sleeping sickness. , : https://www.who.int/news-room/fact-sheets/detail/trypanosomiasis-human-african-(sleeping-sickness) (2023, accessed 13 November 2023).

11. Franco JR, Simarro PP, Diarra A, et al. Epidemiology of human African trypanosomiasis. Clin Epidemiol 2014. DOI: 10.2147/CLEP.S39728.

12. Brun R, Prof, Blum J, MD, Chappuis F, MD, et al. Human African trypanosomiasis. Lancet 2010. DOI: 10.1016/S0140-6736(09)60829-1.

13. Lindner AK and Priotto G. The Unknown Risk of Vertical Transmission in Sleeping Sickness—A Literature Review. PLOS Neglected Tropical Diseases 2010. DOI: 10.1371/journal.pntd.0000783.

14. Fèvre EM, Wissmann Bv, Welburn SC, et al. The Burden of Human African Trypanosomiasis. PLOS Neglected Tropical Diseases 2008. DOI: 10.1371/journal.pntd.0000333.

15. Ilemobade AA. Tsetse and trypanosomosis in Africa: the challenges, the opportunities. Onderstepoort J Vet Res 2009. DOI: 10.4102/ojvr.v76i1.59.

16. World Health Organization. Accelerating work to overcome the global impact of neglected tropical diseases: a roadmap for implementation, https://iris.who.int/handle/10665/338712 (2012, accessed 9 April 2024).

17. Franco JR, Cecchi G, Priotto G, et al. Monitoring the elimination of human African trypanosomiasis at continental and country level: Update to 2018. PLOS Neglected Tropical Diseases 2020. DOI: 10.1371/journal.pntd.0008261.

18. Luintel A, Lowe P, Cooper A, et al. Case of Nigeria-Acquired Human African Trypanosomiasis in United Kingdom, 2016. Emerging Infectious Disease journal 2017. DOI: 10.3201/eid2307.170695.

19. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Systematic Reviews 2015. DOI: 10.1186/2046-4053-4-1.

20. Nmorsi OPG, Isaac C, Igbinosa IB, et al. Human African trypanosomiasis in endemic focus of Abraka, Nigeria. Asian Pacific Journal of Tropical Medicine 2010. DOI: 10.1016/S1995-7645(10)60107-1.

21. Solomon Ngutor K, Idris LA and Oluseyi Oluyinka O. Silent Human Trypanosoma brucei gambiense Infections around the Old Gboko Sleeping Sickness Focus in Nigeria. Journal of parasitology research; J Parasitol Res 2016. DOI: 10.1155/2016/2656121.

22. Uba BV, Aliyu A, Abubakar A, et al. Knowledge and prevalence of human african trypanosomiasis among residents of kachia grazing reserve, Kachia local government area, Kaduna state, Nigeria, 2012. Pan African Medical Journal 2016. DOI: 10.11604/pamj.2016.23.89.7999.

23. Daniel AD, Dadah AJ, Kalejaiye JO, et al. Prevalence of bovine trypanosomiasis in Gongola State of Northern Nigeria. Rev Elev Med Vet Pays Trop 1993.

24. Enwezor FNC, Umoh JU, Esievo KAN, et al. Survey of bovine trypanosomosis in the Kachia Grazing Reserve, Kaduna State, Nigeria. Vet Parasitol 2009. DOI: 10.1016/j.vetpar.2008.10.032.

25. Habeeb IF, Chechet GD and Kwaga JKP. Molecular identification and prevalence of trypanosomes in cattle distributed within the Jebba axis of the River Niger, Kwara state, Nigeria. Parasites & vectors 2021. DOI: 10.1186/s13071-021-05054-0.

26. Majekodunmi AO, Fajinmi A, Dongkum C, et al. A longitudinal survey of African animal trypanosomiasis in domestic cattle on the Jos Plateau, Nigeria: Prevalence, distribution and risk factors. Parasites and Vectors 2013. DOI: 10.1186/1756-3305-6-239.

27. Takeet MI, Fagbemi BO, De Donato M, et al. Molecular survey of pathogenic trypanosomes in naturally infected Nigerian cattle. Res Vet Sci 2013. DOI: 10.1016/j.rvsc.2012.10.018.

28. Omeke BC. Pig trypanosomosis: prevalence and significance in the endemic Middle Belt zone of southern Nigeria. Rev Elev Med Vet Pays Trop 1994.

29. Onah D and Ebenebe O. Isolation of a human serum-resistant Trypanosoma brucei from a naturally infected pig in the Nsukka area of Enugu State. Nigerian Veterinary Journal 2004. DOI: 10.4314/nvj.v24i1.3435.

30. Daniel AD, Joshua RA, Kalejaiye JO, et al. Prevalence of trypanosomiasis in sheep and goats in a region of northern Nigeria. Rev Elev Med Vet Pays Trop 1994.

31. Kalu AU and Lawani FA. Observations on the epidemiology of ruminant trypanosomosis in Kano State, Nigeria. Rev Elev Med Vet Pays Trop 1996.

32. Wayo B, Samdi SM, Fajinmi AO, et al. Prevalence of trypanosomiasis in sheep in the Kachia grazing reserve, Kachia, Kaduna State, Nigeria. African Journal of Clinical and Experimental Microbiology 2017. DOI: 10.4314/ajcem.v18i2.10.

33. Umeakuana PU, Gibson W, Ezeokonkwo RC, et al. Identification of Trypanosoma brucei gambiense in naturally infected dogs in Nigeria. Parasit Vectors 2019. DOI: 10.1186/s13071-019-3680-8.

34. Solomon NK, Lawal I, Shalangwa I, et al. Animal reservoirs of Trypanosoma brucei gambiense around the old Gboko sleeping sickness focus in Nigeria 2016. DOI: 10.5897/JPVB2015.0228.

35. Enwezor F, Emmanuel R, Olanrewaju T, et al. Investigation of livestock for presence of Trypanosoma brucei gambiense in Tafa Local Government Area of Niger State, Nigeria INVESTIGATION OF LIVESTOCK FOR PRESENCE OF TRYPANOSOMA BRUCEI GAMBIENSE IN TAFA LOCAL GOVERNMENT AREA OF NIGER STATE, NIGERIA, January 1,.

36. WHO. GHO | By category | Number of new reported cases (T.b. gambiense) - Data by country, https://apps.who.int/gho/data/node.main.A1636?lang=en (2023, accessed Aug 21, 2023).

37. Mwiinde AM, Simuunza M, Namangala B, et al. Healthcare Management of Human African Trypanosomiasis Cases in the Eastern, Muchinga and Lusaka Provinces of Zambia. Trop Med Infect Dis 2022. DOI: 10.3390/tropicalmed7100270.

38. Selby R, Wamboga C, Erphas O, et al. Gambian human African trypanosomiasis in North West Uganda. Are we on course for the 2020 target?. PLoS Negl Trop Dis 2019. DOI: 10.1371/journal.pntd.0007550.

39. Koné M, Kaba D, Kaboré J, et al. Passive surveillance of human African trypanosomiasis in Côte d'Ivoire: Understanding prevalence, clinical symptoms and signs, and diagnostic test characteristics. PLOS Neglected Tropical Diseases 2021. DOI: 10.1371/journal.pntd.0009656.

40. Odeniran PO and Ademola IO. A meta-analysis of the prevalence of African animal trypanosomiasis in Nigeria from 1960 to 2017 2018. DOI: 10.1186/s13071-018-2801-0.

41. Boundenga L, Mombo IM, Augustin M, et al. Molecular Identification of Trypanosome Diversity in Domestic Animals Reveals the Presence of Trypanosoma brucei gambiense in Historical Foci of Human African Trypanosomiasis in Gabon. Pathogens 2022. DOI: 10.3390/pathogens11090992.

42. Simo G, Asonganyi T, Nkinin SW, et al. High prevalence of Trypanosoma brucei gambiense group 1 in pigs from the Fontem sleeping sickness focus in Cameroon. Vet.Parasitol. 2006. DOI: 10.1016/j.vetpar.2006.02.026.

43. Njiokou F, Nimpaye H, Simo G, et al. Domestic animals as potential reservoir hosts of Trypanosoma brucei gambiense in sleeping sickness foci in Cameroon. Parasite 2010. DOI: 10.1051/parasite/2010171061.

44. Franco JR, Simarro PP, Diarra A, et al. The journey towards elimination of gambiense human African trypanosomiasis: not far, nor easy 2014.

45. Mehlitz D and Molyneux DH. The elimination of Trypanosoma brucei gambiense? Challenges of reservoir hosts and transmission cycles: Expect the unexpected 2019.

46. Molyneux DH, Hopkins DR and Zagaria N. Disease eradication, elimination and control: the need for accurate and consistent usage. Trends in Parasitology 2004. DOI: 10.1016/j.pt.2004.06.004.

47. Büscher P, Bart J, Boelaert M, et al. Do Cryptic Reservoirs Threaten Gambiense-Sleeping Sickness Elimination?. Trends Parasitol 2018. DOI: 10.1016/j.pt.2017.11.008.

48. Geerts M, Chen Z, Bebronne N, et al. Deep kinetoplast genome analyses result in a novel molecular assay for detecting Trypanosoma brucei gambiense-specific minicircles. NAR Genom Bioinform 2022. DOI: 10.1093/nargab/lqac081.

49. Hall MK, Kea B and Wang R. Recognising Bias in Studies of Diagnostic Tests Part 1: Patient Selection. Emerg Med J 2019. DOI: 10.1136/emermed-2019-208446.

50. Shreffler J and Huecker MR. Diagnostic Testing Accuracy: Sensitivity, Specificity, Predictive Values and Likelihood Ratios. In: Anonymous *StatPearls*. Treasure Island (FL): StatPearls Publishing, 2023.

51. Okello I, Mafie E, Eastwood G, et al. African animal trypanosomiasis: A systematic review on prevalence, risk factors and drug resistance in sub-Saharan Africa. J.Med.Entomol. 2022.

52. Picozzi K, Tilley A, Fevre EM, et al. The diagnosis of trypanosome infections: applications of novel technology for reducing disease risk 2002.

53. Contreras Garcia M, Walshe E, Steketee PC, et al. Comparative Sensitivity and Specificity of the 7SL sRNA Diagnostic Test for Animal Trypanosomiasis 2022.

54. Inojosa WO, Augusto I, Bisoffi Z, et al. Diagnosing human African trypanosomiasis in Angola using a card agglutination test: observational study of active and passive case finding strategies. BMJ 2006. DOI: 10.1136/bmj.38859.531354.7C.

55. Dukes P, Gibson WC, Gashumba JK, et al. Absence of the LiTat 1.3 (CATT antigen) gene in Trypanosoma brucei gambiense stocks from Cameroon. Acta Trop 1992. DOI: 10.1016/0001-706x(92)90054-2.

56. Van Meirvenne N, Magnus E and Buscher P. Evaluation of variant specific trypanolysis tests for serodiagnosis of human infections with Trypanosoma brucei gambiense. Acta Trop 1995. DOI: 10.1016/0001-706x(95)00127-z.

57. Chappuis François, Louis L, Pere S, et al. Options for Field Diagnosis of Human African Trypanosomiasis. Clin.Microbiol.Rev. 2005. DOI: 10.1128/cmr.18.1.133-146.2005.